

Vulnerability of Maize Farming Systems to Climate Change: Farmers' Opinions Differ about the Relevance of Adaptation Strategies

Marine Albert, Jacques-Eric Bergez, Magali Willaume, Stéphane Couture

▶ To cite this version:

Marine Albert, Jacques-Eric Bergez, Magali Willaume, Stéphane Couture. Vulnerability of Maize Farming Systems to Climate Change: Farmers' Opinions Differ about the Relevance of Adaptation Strategies. Sustainability, 2022, 14 (14), pp.8275. 10.3390/su14148275. hal-03752727

HAL Id: hal-03752727 https://hal.inrae.fr/hal-03752727v1

Submitted on 17 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







Article

Vulnerability of Maize Farming Systems to Climate Change: Farmers' Opinions Differ about the Relevance of Adaptation Strategies

Marine Albert 1,*, Jacques-Eric Bergez 10, Magali Willaume 20 and Stéphane Couture 3

- ¹ INRAE, UMR1248 AGIR, Université de Toulouse, F-31320 Castanet-Tolosan, France; iacques-eric.bergez@inrae.fr
- ² INRAE, UR MIAT, Université de Toulouse, F-31320 Castanet-Tolosan, France; magali.willaume@toulouse-inp.fr
- 3 INPT ENSAT, UMR1248 AGIR, Université de Toulouse, F-31320 Castanet-Tolosan, France; stephane.couture@inrae.fr
- * Correspondence: marine.albert@inrae.fr

Abstract: Climate change has negative impacts on maize cultivation in southwestern France, such as soil erosion and water stress. The vulnerability of maize farming systems to climate change must be assessed before considering potential adaptation strategies. This study focused on eliciting and understanding criteria that maize growers use to assess the vulnerability of their farming systems to climate change. To this end, we surveyed maize growers in two consecutive stages: a qualitative stage, to elicit vulnerability criteria, and a quantitative stage, to test the genericity of criteria related to the adaptation strategies. The qualitative stage identified 144 criteria that farmers used to assess vulnerability to climate change, while the quantitative stage showed that farmers' opinions about the adaptation strategies differed. Many factors explained these differences, including structural (e.g., soil type) and psychological factors (e.g., interest in agroecology). Our typology of farmers revealed that their interest in agroecology and technology, as well as their perceptions of the risks of climate change and their attachment to their production systems, influence the type of adaptations they identify as relevant (i.e., intensification strategies, slight adjustments or agroecological innovations). Farmers' perceptions should be considered when providing individual advice and assessing vulnerability, by including criteria related to their psychological characteristics.

Keywords: evaluation; vulnerability; maize grower; climate change; farmers' perceptions; adaptation



Citation: Albert, M.; Bergez, J.-E.; Willaume, M.; Couture, S. Vulnerability of Maize Farming Systems to Climate Change: Farmers' Opinions Differ about the Relevance of Adaptation Strategies. Sustainability 2022, 14, 8275. https:// doi.org/10.3390/su14148275

Academic Editors: Nasir Mahmood and Irfan Ahmad Baig

Received: 19 May 2022 Accepted: 29 June 2022 Published: 6 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

Farmers must increasingly address the global increase in temperature and annual variations in rainfall [1]. In France, a 6 °C increase in the mean summer temperature is expected by the end of the 21st century; although annual variations in rainfall are not expected, large monthly variations are predicted [2,3]. Higher temperatures will increase soil evaporation and crop transpiration, which will increase water demand and consumption, which will in turn influence the surface water balance and the potential for severe drought [4–6]. In addition to thermal and hydric stresses, the increased frequency and intensity of extreme climatic events can result in soil erosion due to floods, droughts or storms [6]. Droughts would have a particularly severe impact on summer-irrigated crops. In southwestern France, maize cropping systems already suffer from these climate phenomena [7,8]. Consequently, farmers must adapt their farming systems to reduce their farms' vulnerability to climate change [9]. Farmers have access to many adaptation strategies [10], such as changing varieties or crop species, changing sowing density and modifying the schedule of cropping operations [7,11–15]. However, adaptation decisions can fail, either by not achieving the objectives, or worse, by increasing vulnerability [16].

Sustainability **2022**, 14, 8275 2 of 23

Therefore, the vulnerability of maize farming systems must be assessed before considering adaptation strategies to address climate change. More specifically, a better understanding of maize farming systems and the determinants of adaptive capacity can allow public policies and agricultural advising to enhance the agroecological transition.

The Intergovernmental Panel on Climate Change (IPCC) defines climate change vulnerability as "the degree to which a system is susceptible to, and unable to cope with, adverse effects of climate change, including climate variability and extremes" [1]. Vulnerability is a function of (i) exposure (i.e., characterized by the intensity, frequency and duration of perturbations), (ii) sensitivity (i.e., degree to which the exposure influences the system) and (iii) adaptive capacity (i.e., ability to implement adaptations to address perturbations) [17]. This conceptual framework combines biophysical and socio-economic factors to determine vulnerability [18].

The need to assess vulnerability has already been discussed [19–21] but is rarely rendered operational due to its complexity. Two main approaches are used to assess the vulnerability, or the related concept of resilience, of agricultural systems [22]: (i) the quantitative analysis of agricultural system outputs, in which vulnerability is assessed by studying dynamics related to the perturbations of agricultural outputs, such as yield or economic net return [20,23,24], and (ii) the quantitative or qualitative evaluation of predefined properties associated with vulnerability or resilience [25–27]. In the latter approach, vulnerability is assessed by considering the properties of the system identified using expert knowledge or the literature [28,29].

Most vulnerability and resilience assessment studies are based on a dynamic performance approach. However, this approach is limited to easily recordable components, and often focuses on only one type of performance, mainly yield [22,30,31], whereas assessing multiple types of performance is essential [19]. Moreover, these studies rarely focus on assessing vulnerability at the farm scale, even though all the dimensions of a farming system must be considered. Identifying the determinants of vulnerability to climate change requires a systemic approach, since vulnerability is a complex problem and can be influenced by diverse factors such as crop management and the financial status of the farm. Moreover, decisions about adaptation strategies are made at the farm level [30,32]. The predefined property approach can focus on a broader scale by using indicators to consider different components of agricultural systems, including adaptive capacity [28]. Despite this advantage, few agricultural studies have used a predefined property approach [33].

Many studies considered only the dimensions of exposure and sensitivity when assessing climate change vulnerability, ignoring the adaptive capacity of farmers [34]. However, Marshall et al. [35] showed that farmers' perceptions of their skills, their satisfaction with the adaptations they implement and their willingness to change, strongly influence their adaptive capacity. Consequently, farmers' perceptions will influence the vulnerability of their farms [35]. Previous studies have stressed the importance of cognition (i.e., the administration and implementation of information) on vulnerability [36–38]. For example, Marshall et al. [35] showed that less vulnerable farmers were well integrated into social networks. In the literature, there are several empirical studies studying farmers' adaptive capacity specifically regarding climate change impacts [39-43]. Some of these studies use individual interviews with farmers to obtain their perceptions about climate change [39,40,43], while other studies base their data collection on participative methods such as focus groups [41,42]. The data analysis conducted to identify the determinants of adoption can be based on Pearson correlation [40], mathematical models [39] such as the logit model [43], and factorial analysis using PCA and ANOVA [41]. The results of these studies show that adaptive capacity is influenced by the individual characteristics of the farmers, such as access to extension services [39,42], perceptions of risks [40,43], access to information [39,43], or the level of knowledge [39,41].

Ultimately, studying adaptive capacity is essential to identify criteria for assessing the vulnerability of farming systems.

Sustainability **2022**, 14, 8275 3 of 23

Most assessment methods are based on an objective approach (i.e., external judgements [44]) to identify performance indicators that measure the dynamics of vulnerability or identify properties that influence vulnerability. Although scientists have taken responsibility for designing these indicators, farmers are able to identify the determinants of vulnerability on their own farms [44] and can understand their situations [45] in relation to climate change. Using the determinants of vulnerability identified by farmers will help legitimize this set of indicators [44,46]. Moreover, considering farmers' personal characteristics (i.e., cognitive and psychological factors) when assessing farm vulnerability is crucial for advisors to provide specific guidelines based on each farmer's situation [35]. Jones [44] highlighted that subjective approaches can consider farmers' knowledge and experience of resilience, along with the factors that contribute to them, and can complement objective approaches.

This study aimed to identify and understand the criteria that maize growers use to assess the vulnerability of their farming systems to climate change. The criteria for assessing vulnerability were elicited (i) at the farm level (ii) to include the adaptive-capacity dimension of vulnerability, using (iii) a predefined property approach and (iv) a subjective approach based on farmers' perceptions. We conducted a two-stage survey with maize growers: one group of farmers was surveyed to elicit vulnerability criteria, and another group was surveyed to test the genericity of adaptation strategy criteria and understand farmers' opinions about the relevance of adaptations. We assumed that farmers' psychological and cognitive factors would explain their opinions. This study will help understand farming with a more systemic approach, by knowing to what extend the farmer's subjectivity needs to be considered when investigating vulnerability to climate change.

2. Materials and Methods

2.1. Conceptual Framework

We developed a conceptual framework that combines the Drivers–Pressures–State-Impacts–Responses (DPSIR) model developed by Smeets et al. [47] and the three dimensions of vulnerability defined by the IPCC (i.e., exposure, sensitivity and adaptive capacity) [17]. The DPSIR model represents causal interactions between a system and its environment: a driving force (D) creates pressure (P) on a system in a given state (S), which creates impacts (I). In response to these impacts, the system responds (R) with adaptations [47,48].

We developed the following framework to explore farm vulnerability to climate change (Figure 1): climate change variables are driving forces (D) (e.g., rainfall) that produce pressures (P) (e.g., excess water) that influence the state of cropping systems (S). This degradation of cropping systems has impacts on the farm (I) (e.g., soil erosion) that require adaptation strategies (R) (e.g., reduction in soil tillage). We added a supplementary component to the DPSIR model in our framework, since implementing adaptation strategies requires the resources of the farm (e.g., availability of equipment, financial situation) and of the farmer (e.g., knowledge, perceptions) (F). Along with climate change pressures, other driving forces can pressure the system, such as market volatility, regulations and citizens' opinions. We considered only the negative aspects of climate change in our conceptual framework, although it could include positive aspects (e.g., increasing CO₂ concentrations increase plant growth).

2.2. Overview of the Survey Design

We used an empirical approach that had two consecutive stages to include qualitative and quantitative approaches (Table 1). The qualitative stage aimed to elicit vulnerability criteria from a group of expert farmers. We considered "criteria" instead of "indicators" since we focused on identifying the determinants of vulnerability and not on measuring them. The three dimensions of vulnerability were addressed in this stage. The four-step interviews were based directly on our conceptual framework, since we asked each farmer to describe the following information: (i) pressures related to climate change (P) and impacts (I) the farmer observes on the farm; (ii) ways to measure the state of the cropping system (S); (iii) adaptation strategies the farmer implements or wishes to implement (R) and

Sustainability **2022**, 14, 8275 4 of 23

(iv) farm and farmer resources required to implement the adaptations (F). The discussion was facilitated using visual aids (i.e., a board and climate graphs) and the encouragement of the interviewer.




Figure 1. Conceptual framework for assessing vulnerability that combines the Drivers (D)–Pressures (P)–State (S)–Impacts (I)–Responses (R) model and the concept of vulnerability.

We designed the questionnaire for the second stage based on the results of the first stage. The first survey revealed a diversity of adaptation strategy criteria elicited from expert farmers. Thus, the second stage of the survey focused on the "adaptation strategies" dimension of our conceptual framework. We tested the genericity of the adaptation strategy criteria obtained in the qualitative survey by asking a larger sample of farmers to select criteria and indicate their relevance. To this end, we created a card game, with each card representing one adaptation strategy criterion (e.g., reduce soil tillage). We then used statistical analysis to explain farmers' opinions via explanatory factors such as the characteristics of the farm and farmer.

Table 1. Characteristics of the qualitative and quantitative stages of the survey.

Qualitative Stage	Quantitative Stage
	2
Elicit criteria for assessing vulnerability	Test the genericity of the adaptation strategy criteria and explain farmers' opinions
Expert farmers (13)	Farmers representative of the Occitanie region (32)
Semi-structured interviews on the farm	Semi-structured interviews on the farm
Poster/board representing farm resources; climate graphs	Cards for adaptation strategy criteria
All dimensions (i.e., exposure, sensitivity and adaptive capacity)	Adaptive capacity
Monography; expert classification	Data cleaning; statistical analysis (regressions, clustering)
	assessing vulnerability Expert farmers (13) Semi-structured interviews on the farm Poster/board representing farm resources; climate graphs All dimensions (i.e., exposure, sensitivity and adaptive capacity)

Sustainability **2022**, 14, 8275 5 of 23

2.3. Case Study

The study was conducted in the Occitanie region in southwestern France, within a 100 km radius around the city of Toulouse. Southwestern France is known for its maize production, as it supplies 39% of the national production of the crop [49]. In 2020, Occitanie counted 131,706 hectares of grain maize, including 102,094 irrigated hectares, representing 8% and 15% of the national area of grain maize, respectively [50]. The temperature in Occitanie continues to increase, which has a negative impact on maize cultivation. Higher temperatures in May and June cause problems during reproductive stages, such as pollen degradation and difficulty with absorbing nutrients [5,6]. An increase in summer droughts creates the need for more irrigation water for maize, meaning that maize farms economically depend on irrigation in 6 out of 10 years, on average [49]. Farmers interviewed in the qualitative and quantitative stages came from several departments in the Occitanie region—Tarn, Tarn et Garonne, Gers and Haute-Garonne (Figure 2)—to ensure that we included a diversity of maize farming systems in southwestern France. Access to irrigation differs between the departments, as does soil type, historical types of farm production, climate conditions and the social environment. All farmers interviewed grew irrigated maize (i.e., popcorn, waxy, grain or seed) on at least one field. The Chamber of Agriculture in the region provided contacts for farmers for both stages of the survey.




Figure 2. Location of the farms of the maize growers interviewed during the first, qualitative stage (initial surveys) (blue) and the second, quantitative stage (deeper surveys) (orange). Map data: 2020, Google.

To elicit criteria in the qualitative stage, we recruited a small sample of expert farmers (13, all male) who were sensitive to climate change and already adopting innovations for adaptation. The interviews, conducted from February to April 2021, lasted 0.75–3 h. For the quantitative stage, we interviewed a sample of 32 maize farmers (all male) representative of the study area in terms of areas and types of production. The interviews, conducted in June and July 2021, lasted 1–3 h.

2.4. Data Collection

For the qualitative stage of the survey, two visual aids were used to support the interviews. First, the interviewer showed climate graphs and discussed them with the

Sustainability **2022**, 14, 8275 6 of 23

farmer, and then encouraged the farmer to express his opinions about the climate pressures and impacts he observed on his farm. After discussing the graphs, the farmer was asked to observe a board representing components of the farming system (Figure 3) and explain how climate change manifests and influences each component. The facilitator encouraged the farmer to identify sensitive elements in his farming system and adaptations he had implemented (or wanted to implement). For each adaptation the farmer identified, the facilitator asked which resources it required. At the end of the interview, the facilitator asked the farmer to summarize the strengths and weaknesses of his farming system in relation to climate change. All interviews were audio recorded to ensure that data were collected accurately.




Figure 3. The board used in the qualitative survey that represented the components of a farm.

To design the questionnaire for the quantitative survey, we identified a list of explanatory variables and made assumptions about their influence on farmers' opinions about adaptation strategy criteria. We then summarized the extensive list of adaptation strategy criteria obtained in the qualitative stage to (i) identify more general criteria, (ii) avoid redundant criteria and (iii) reduce the number of criteria. We reduced the initial list of adaptation strategy criteria from 50 to 41. The data collection process was divided into three steps (Figure 4a):

- Farmers responded to a questionnaire by telephone to identify the farm characteristics
 that were potential explanatory variables. The questionnaire was divided into eight
 categories defined according to expert knowledge, including general information,
 material resources, water resources, soil resources, financial resources, crops and
 rotation, human resources and individual resources.
- Farmers were asked to assess the adaptation strategy criteria in a semi-structured interview using the 41 playing cards that represented the adaptation strategy criteria. The interviewer asked the farmer to select the four most relevant cards and the four least relevant cards for reducing vulnerability (Figure 4b).

Sustainability **2022**, 14, 8275 7 of 23

• The farmer's cognitive and psychological characteristics were identified using a face-to-face questionnaire (Table 2). This questionnaire supplemented the telephone questionnaire by adding new potential explanatory factors.




Figure 4. (a) Main steps of the quantitative survey, showing explanatory variables (red) and the variables explained (blue), and (b) a farmer playing the card game used in the quantitative survey.

Table 2. Cognitive and psychological variables and the associated elicitation technique in the questionnaire.

Cognitive and Psychological Factors	Variables	Elicitation Technique in the Questionnaire	Responses Analyzed
	Threat of climate change	Dichotomous question	Yes or no
Perceptions of climate change	Level of climate change pressure	Multiple-choice question	Low, medium or high
Agroecological practices	Degree of interest in agroecological practices	Self-assessment: score from 1–10 (high interest)	Low (1–3), medium (4–6) or high (7–10)
Resistance to change	Degree of attachment to the production system	Self-assessment: score from 1–10 (high interest)	Low (1–3), medium (4–6) or high (7–10)
Innovations Degree of interest in technology		Self-assessment: score from 1–10 (high interest)	Low (1–3), medium (4–6) or high (7–10)
Reactivity	Degree of planning	Self-assessment: score from 1–10 (high planning)	Low (1–3), medium (4–6) or high (7–10)
Assistance	Favorite information source	Multiple-choice question	Advisors, farmers, technology or laboratory
Risk aversion	Degree of risk aversion	Lottery game	High (1–3), medium (4–6) or low (7–10)

2.5. Data Processing and Analysis

For the qualitative survey, each interview was transcribed into monographs. We then identified, via farmers' comments, elements that were related to the dimensions of

Sustainability **2022**, 14, 8275 8 of 23

the conceptual framework. We collected and classified these elements into a database, and for each element, we determined the associated criteria using agronomic expertise. Finally, we removed redundant criteria and counted the number of times that each criterion was mentioned.

The data collected in the second stage of the survey yielded a database composed of 32 rows (farmers) \times 156 columns (explanatory variables) \times 41 criteria (explained variables). The statistical challenge was to analyze a database that had many more variables than individuals. Before analyzing the data, we cleaned them in several steps (Appendix A). The cleaning procedure left 25 explanatory variables (Appendix B) for the 32 farmers and 41 explained variables.

To study the distribution of the adaptation strategy criteria among farmers, we developed a typology of these criteria based on farmers' opinions and the number of times they had been mentioned. Classification consisted of multiple correspondence analysis (MCA) followed by hierarchical clustering on principal components (HCPC). We used three additional statistical methods to understand differences in the criteria chosen (Figure 5). We developed regression trees for criteria that were mentioned by at least five farmers. The regression tree method allows for the consideration of local interactions among variables, and is relevant for samples with many variables compared to the number of individuals [6]. We then performed a logistic regression of each criterion and its associated first explanatory variable identified by the regression tree. Finally, we practiced a classification of farmers based on their assessment of the adaptation strategy criteria and studied the distribution of explanatory variables in each cluster. The objective was to create a typology of farmers according to their choice of adaptations, and thus identify explanatory variables for these choices.

The statistical analysis was performed using R software [51], with the MCA and HCPC functions of the FactoMineR package [52] for classifications. The rpart [53] and rpart. plot [54] packages were used for regression trees, and the mlogit package [55] was used for logistic regressions.




Figure 5. Statistical analysis of the quantitative survey.

3. Results

3.1. Participants

The utilized agricultural area (UAA) varied among the 13 expert farmers involved in the qualitative survey. The mean UAA of 198 ha (standard deviation = 89 ha) was much larger than that of field-crop farms in the region (i.e., 83 ha) [56]. The mean area under

Sustainability **2022**, 14, 8275 9 of 23

irrigated maize was 52 ha (\pm 32 ha), which was also much larger than that of field-crop farms in the region (i.e., 28 ha).

The mean age and number of labor units of the 32 farmers interviewed in the quantitative survey were representative of field-crop farms in the region. Their farms had a much larger mean UAA and irrigated area (199 and 60 ha, respectively) than those of field-crop farms in the region (99 and 21 ha, respectively). More than 50% of their revenue came from maize production, while the mean was 36% for field-crop farms in the region. A larger percentage of them practiced organic agriculture than the mean percentage of field-crop farms in the region.

3.2. Distribution of the Criteria Elicited from Farmer Interviews

The qualitative stage of the survey resulted in a list of 144 criteria distributed among all dimensions of the conceptual framework (Figure 6). The least represented sub-dimension of vulnerability was "farm resources" (nine criteria). Most of its criteria concerned traditional financial assessments of a farm, such as income, gross profit and debt, which all farmers mentioned. The most represented sub-dimension was "adaptation strategies" (50 criteria), which was the most diversified sub-dimension among the 13 expert farmers interviewed.




Figure 6. Distribution of the criteria in the conceptual framework.

Among the 50 adaptation strategy criteria, only three were mentioned by all farmers: "increase water storage", "diversify water sources" and "irrigate winter crops". All of these criteria were related to water use, which was not surprising in a context of water scarcity. Conversely, 22 adaptation strategy criteria were mentioned only once, such as "partner with a beekeeper", "plant mixed crops" and "introduce livestock". Appendix C shows the number of times that each of the 50 adaptation strategy criteria was mentioned.

3.3. Differing Opinions about Adaptation Strategy Criteria among Farmers

The MCA and HCPC of the 41 adaptation strategy criteria according to their relevance for farmers (dimension 1) and the number of times they were mentioned (dimension 2) identified four clusters (Figure 7). Cluster 1 grouped criteria that tended to be considered irrelevant and that were mentioned often, while cluster 4 grouped criteria that tended to be considered relevant and which were mentioned often. Farmers considered the following adaptation strategy criteria to be particularly irrelevant for reducing vulnerability to climate change: "introduce livestock production unit" (A14), "convert to organic farming" (A19), "buy new land" (A17) and "stop growing maize" (A7). Criteria considered particularly relevant for reducing vulnerability to climate change included "use sensors" (A38), "use a maize variety resistant to hydric stress" (A11), "return harvest residues to the soil" (A27), "increase irrigation efficiency" (A34) and "plant cover crops" (A3). A11, A27 and A34 concerned technology or strategies for managing water, while A27 and A3 concerned strategies and crop practices for managing the soil. Cluster 2 grouped criteria that were

Sustainability **2022**, 14, 8275 10 of 23

mostly irrelevant and rarely mentioned (A10, A18, A23, and A29) along with criteria that generated highly contrasting opinions and were mentioned often (A22, A31). For example, farmers disagreed greatly about the criteria "stop plowing" (A22) and "build or develop a reservoir" (A31). Some farmers considered "stop plowing" a solution to improve soil fertility and control erosion (mentioned as relevant seven times), while other farmers considered it a high risk for crop productivity due to potential problems with weed management and crop establishment (mentioned as irrelevant six times). Cluster 3 grouped criteria that were rarely mentioned and about which farmers' opinions differed, such as "use mechanical weeding" (A24), "practice strip tillage" (A26), "practice direct seeding" (A25) and "diversify water sources" (A33). For example, several farmers agreed that direct seeding was a way to maintain soil quality (mentioned as relevant three times), while other farmers mentioned that equipment costs and uncertainty in productivity made this practice too risky for the financial situation of their farm (mentioned as irrelevant three times).




Figure 7. Classification of adaptation strategy criteria according to farmers' opinions. Numbers refer to the 41 adaptation strategy criteria (Appendix D).

3.4. Explaining Differences in Opinions about the Adaptation Strategy Criteria

For six criteria, logistic regressions confirmed the significant (p < 0.05) influence of the first explanatory variable revealed in the regression tree (Table 3). Of the six explanatory factors identified, three were related to farmer characteristics (i.e., age, perceptions of climate change risks, and interest in agroecology) and the other three were related to farm characteristics (i.e., soil types, number of crops).

Sustainability **2022**, 14, 8275 11 of 23

Table 3. Results of regression trees and logistic regressions, with detailed results from logistic regressions. *** p < 0.001; ** p < 0.05; * p < 0.1.

	Assessed	First Explanatory Variable in the	Logistic Regression Results	
Criterion	Relevance Regression Tree		Direction	Significance
A11: variety resistant to hydric stress	Relevant	Perceptions of climate change risks: low	+	0.0368 **
A27: return harvest residues to the soil	Relevant	Silty clay soil	+	0.0292 **
A29: buy more irrigation quota	Irrelevant	1–3 crops 4–6 crops (in the rotation)	- -	0.03279 ** 0.00324 ***
A31: build or develop a reservoir	Irrelevant	"Boulbènes" soil (clayey sand)	+	0.0184 **
A34: increase irrigation efficiency	Relevant	Age > 50 Age 35–50	-	0.0207 ** 0.0601 *
A35: irrigate winter crops	Relevant	Interest in agroecology: high	-	0.0334 **

To supplement the regression methods, we performed the HCPC of farmers based on their opinions of the adaptation strategy criteria (i.e., relevant, not selected, irrelevant). The four clusters it identified contained significant differences in farmers' opinions about the adaptation strategy criteria (Table 4). For each cluster, we identified the adaptation strategy criteria that characterized the cluster the most (Appendix E) and studied the distribution of the 25 explanatory variables (Table 4). Farmers in cluster 1, who adopted slight adaptation strategies, had a similar medium-sized UAA for the sample. These farmers perceived climate change as a threat but had little interest in agroecological practices, which was consistent with the results of the regressions (farmers with high interest in agroecology did not perceive "irrigate winter crops" as relevant). All farmers in cluster 2, who were associated with intensification strategies, practiced conventional agriculture. All of these farmers used more than 2000 m³ ha⁻¹ to irrigate and were highly attached to their production system, which explained why they perceived conversion to organic agriculture as irrelevant, as well as their choice of intensification adaptations. Cluster 3 could not be explained by the structural and cognitive variables tested. Cluster 4 (attend training courses and diversify crops) grouped farms with few labor units and clay-limestone soil. They did not plow, had a high interest in agroecological practices and technologies, and had a moderate attachment to their production system. These farmers grew at least four crops in 2020, which was consistent with their opinion that diversifying crops reduced vulnerability to climate change.

Table 4. Distribution of the 20 significant explanatory variables within each cluster. Shaded cells identify variables for which farmers' opinions differed within a cluster. The variables of age, water source, the precocity of the maize variety, reactivity and sources of information were tested but are not shown because they were non-significant (i.e., different in all clusters).

	Cluster 1: Slight Adjustments	Cluster 2: Intensification Strategies	Cluster 3: Diversification of Activities	Cluster 4: Agroecological Innovations
Number of farmers	2	11	14	4
Relevant adaptations to reduce vulnerability to climate change	A5: Balance summer and winter crops; A36: Irrigate winter crops	A30: Renew irrigation equipment; A31: Build or develop a reservoir	A16: Diversify paid activities	A4: Diversify crops; A40: Attend training courses
Irrelevant adaptations to reduce vulnerability to climate change		A19: Convert to organic agriculture; A24: Perform mechanical weeding	A17: Buy new land; A32: Build or develop a reservoir	A23: Lime the soil

Sustainability **2022**, 14, 8275 12 of 23

Table 4. Cont.

	Cluster 1: Slight Adjustments	Cluster 2: Intensification Strategies	Cluster 3: Diversification of Activities	Cluster 4: Agroecological Innovations
Utilized agricultural area (ha)	100–200 (medium size)			
Production system		Conventional		
Labor units (full-time equivalent)	1			<2
Education	Bachelor's degree			
Volume of water $(m^3 ha^{-1})$		>2000		<3000
Irrigable area	Non-irrigable fields			Non-irrigable fields
Soil structure				Clay-limestone
Number of pivots	None			
Volume of grain storage (t)	<1000			
Number of paid activities	2			
Percentage of revenue from maize	30–50%			
Distance between plots (km)	>14			
Number of crops	4–6			≥4
Soil tillage	Reduced tillage			No plowing
Climate change threat	Yes			Yes
Perception of risks from climate change	Medium to high			Medium to high
Interest in agroecology	Low			High
Attachment to production system		Medium to high		Medium
Interest in technology	Low			High
Risk aversion	Very high			

4. Discussion

4.1. Psychological Factors That Influence Opinions about the Adaptation Strategy Criteria

The statistical analysis showed that psychological factors explained some of the differences in farmers' opinions about the relevance of the adaptation strategy criteria. Regressions revealed that farmers who perceived the risks of climate change as low perceived the criterion "variety resistant to hydric stress" (A11) as relevant to reducing vulnerability to climate change. Since these farmers were not threatened by climate change, they perceived that slight adjustments, such as changing the variety, were sufficient. Conversely, farmers who perceived high risks prioritized larger adaptations, such as changing the crop pattern. He et al. [57] demonstrated that risk cognition (i.e., individual perception of risks) has a positive influence on adaptive behavior toward climate change. Willaume et al. [58] also related farmers' perceptions of climate change to the type of adaptations they implemented; regarding the "efficiency—substitution—redesign" transition model, farmers with low risk perception tended to choose "substitution" strategies.

Gbetibouo et al. [39] highlighted that farmers' perceptions of climate change are partly influenced by their access to information. Several studies highlighted the importance of cognition, such as the implementation of information [36,38]. Farmers' involvement in a social network would improve adaptive capacity [59]. However, our results indicated that the variable "favorite information source" was not a factor that influenced farmers' opinions about adaptation strategies.

Sustainability **2022**, 14, 8275 13 of 23

The regressions also highlighted that farmers with high interest in agroecological practices did not perceive "irrigate winter crops" as a relevant adaptation. These farmers may prioritize the conservation of resources, which is a fundamental principle of agroecology. Overall, "interest in agroecology" and commitment to change are determinants of adaptive capacity [35]. Farmers with low interest in agroecology and technology, and with very high risk aversion, tended to perceive slight adjustments as relevant for reducing vulnerability to climate change. Similarly, farmers who practiced conventional agriculture and were highly attached to their production system prioritized intensification adaptations. Conversely, farmers with a high interest in agroecology and technology, and a moderate attachment to their production system, were more likely to perceive innovative agroecological adaptations as relevant to reduce vulnerability to climate change. This confirms that perceptions of innovation (either agroecological practices or technology) [60], attachment to place [35] and resistance to change [60] are key factors that influence the adoption of strategies. Based on our results, high risk aversion seemed to be associated with slight adjustments. Indeed, high risk aversion is a barrier to adaptive behavior [61,62]. Other studies have demonstrated the influence of risk aversion on adaptive behavior [60,63].

Our results confirm that farmers' perceptions influence their adaptive capacity and thus the vulnerability of their farming systems [35]. Adaptation strategies that farmers implement to address climate therefore vary and depend on their cognitive and psychological profiles. Our study could be replicated with a bigger sample in order to confirm our results. Other factors, such as moral concerns (including environmental awareness), intuition and personality [60,64], can also influence the adaptive capacity of farmers and should be considered in future studies to more fully understand farmers' opinions about the relevance of adaptation strategies.

4.2. An Original Method Based on Combined Approaches

Our approach for assessing farm vulnerability is original, since we based the identification of vulnerability properties on farmers' expertise, while predefined property approaches are usually based on the literature and/or scientific expertise [28,29]. Thus, our approach was more likely to yield a comprehensive set of indicators that can be managed and adopted easily by farmers and which are appropriate for the farming context. As Wienroth [65] explained in the "let's RULE" model, an innovation should be reliable, useful and legitimate. Another original aspect of our study is the participative and interdisciplinary approach involving behavioral economics.

Similar to Dessart et al. (2019), who proposed a key theoretical framework for our study, our conceptual framework and the choice of the psychological and cognitive factors investigated do not rely on one specific theoretical framework, but are guided by various theories or models of behavior (such as the theory of planned behavior or the theory of expected utility). Indeed, since there is no unified theory of behavior to date and most theories cover only a certain aspect of decision-making [66], our approach allowed us to gather the different behavioral factors that are fundamental to explaining decision-making and, more specifically, the adoption of coping strategies.

One strength of our survey was its combination of qualitative and quantitative approaches, for which we developed original methods to render the concept of vulnerability operational. In the qualitative stage of the survey, criteria were elicited using climate scenarios and a board that represented major components of a farm (e.g., water resources, the farmer's network, equipment). In the quantitative stage of the survey, using cards to represent criteria made the interview playful and interesting for the farmers interviewed.

For the statistical analysis, using a variety of statistical models (i.e., regression trees, logistic regression and classification) enabled us to obtain robust and complementary results. The regression trees were able to consider non-additive effects, combined effects and interactions [6,67] by sequentially dividing responses according to the most relevant explanatory variabale (i.e., by minimizing the locally explained variance). In comparison, the logistic regressions isolated the effects of each variable [68]. The classification pro-

Sustainability **2022**, 14, 8275 14 of 23

vided results that were complementary to those of the regressions and thus improved the understanding of farmers' opinions about adaptation strategy criteria.

The novelty of this study regarding our results is that we highlighted the important role of the psychological and cognitive profiles of maize growers in their choice of adaptation strategies to address climate change.

The main disadvantage of our study was the small sample size of the quantitative survey (32 farmers). Although the sample was representative of the region, our results cannot be considered general. However, our goal was to test the genericity of adaptation strategy criteria within a group of farmers and not to describe farmers' opinions about the adaptation strategy criteria in the region. We successfully met this goal, since we demonstrated farmers' differing opinions about the adaptation strategy criteria and identified explanatory variables.

4.3. The Need to Reconsider Advising and Support Strategies for Farmers

Our results showed that farmers use different adaptation strategy criteria to assess farm vulnerability. Similar to Jones et al. [44], who demonstrated that resilience is not the same for everyone, we determined that vulnerability is also not the same for everyone. Our quantitative focus on adaptations demonstrates that relevant options depend strongly on individual choice, which indicates the need to challenge the genericity of commonly accepted adaptation strategies in the literature. They are presented as universal, but few studies have focused on the conditions of the success or failure of these adaptations in farming systems [10,69]. Misusing adaptations can negatively impact farming systems [16]. Our results indicate that farmers perceive that certain adaptations might increase the vulnerability of their farms. In this case, several studies mention "maladaptations" [10,69–73], which occur when adaptation decisions are made using inaccurate assumptions or failing to consider the potential negative external effects of adaptations (e.g., the degradation of biodiversity, increase in emissions, new or higher costs that farmers did not consider) [74]. This indicates the need to reconsider advising and support strategies for farmers, since public policies and advisor support are often allocated and applied the same way within a region. More individual advising could ensure that farmers adopt adaptation strategies better, as long as farmers' perceptions are considered when developing the recommendations. To this end, discussions with farmers about the vulnerability of their farms are essential before recommending any adaptation strategy. Based on our results, the adoption of agroecological practices is influenced by farmers' interest in agroecology and perception of the risks of climate change. Therefore, agricultural advising and public policies could improve access to information through specific awareness and training campaigns to increase the adoption of agroecological practices.

4.4. From Theoretical Results to More Operational Aspects

Various actions could be performed in order to enhance agroecological transition among the maize farming systems in southwestern France. Regarding agricultural advising, there is a need for individual support and groups of discussions and trainings in order to help farmers improve their knowledge on adaptation strategies. Small groups of farmers could be initiated by institutes and private companies advising farmers. A material support for discussion could be a serious game on adaptation strategies and their effects on farms' vulnerability to climate change. Simulated tests of adaptation strategies in virtual conditions could help farmers make decision to adopt specific adaptations and be aware of issues they could face. Farmers' networks that share values and stakes, and which are specifically dedicated to climate adaptation and managed by the Agriculture Chamber, could be created, as it was in the context of pesticide use, with the Defi Zero Phyto network. These networks could offer workshops and visits on farms in order to share experiences and show agroecological innovations. On-farm experiments could be carried out via collaborations between research institutes and farmers, in order to obtain both viewpoints regarding the relevance of the tested adaptations. Finally, public policies

Sustainability **2022**, 14, 8275 15 of 23

could enhance the agroecological transition by allocating subsidies for farmers who wish to implement agroecological innovations (such as subsidies to buy cover crop seeds or a direct seeder).

4.5. Toward an Assessment Method That Includes the Adaptive Capacity of Farmers

In the context of a vulnerability assessment, the dimension of adaptive capacity regarding farmers' internal resources (i.e., psychological and cognitive) requires a specific focus, whereas existing assessment methods often ignore this aspect of vulnerability. One issue is to consider the diversity of farmers' perceptions when developing a tool based on indicators, since farmers who have the same farming system characteristics will not implement the same strategies if their perceptions differ. Our study showed that assessments of adaptive capacity should include criteria and indicators related to psychological factors such as the perception of the risks of climate change, attachment to the production system and interest in agroecological practices. Farmers' opinions need to be compared to objective viewpoints when considering the relevance of adaptation strategies to reduce vulnerability to climate change. Farmers' opinions about adaptation strategy criteria might not agree with scientists' viewpoints due to differences in knowledge, experience and perceptions. This subjective approach for assessing adaptation strategies should be supplemented by consulting the literature and scientific experts to create a typology of relevant adaptations depending on the farming system. Combined with a method for assessing vulnerability, this typology would help advisors provide farmers with effective adaptations, which would result in the development of resilient farming systems. The criteria identified in our qualitative survey related to sensitivity and exposure should also be compared to scientific knowledge to design a multicriteria method for assessing vulnerability to climate change.

5. Conclusions

Our study rendered the concept of vulnerability operational by comprehensively identifying its determinants. We developed a conceptual framework that combines the DPSIR model and the concept of vulnerability defined by the IPCC. We identified criteria that farmers use to assess the vulnerability of their farming systems. The criteria of adaptation strategies are diverse and differ among farmers. The statistical analysis using complementary methods showed that farmers' opinions about adaptation strategies are influenced by structural factors (e.g., soil structure), the characteristics of the farmer (e.g., age), the cropping system (e.g., number of crops), as well as cognitive and psychological factors such as risk aversion, attachment to the production system and the perceptions of the risks of climate change. Our study highlighted that farmers' opinions of adaptation strategies are not general, even within the same region. This implies that agricultural advising should be more individualized. The results confirmed our hypothesis that farmers' cognitive and psychological resources influence their adaptive capacity. The relevance of adaptations does not only depend on agronomic or economic performance, but also on farmers' perceptions. Therefore, futures studies dealing with the performance assessment of adaptations should include indicators regarding the level of congruence with farmers' psychological and cognitive profiles. Finally, our study helped to understand farmers' perceptions of the vulnerability assessment, which is the initial step in designing a method to assess the vulnerability of maize farming systems to climate change. Future studies should compare farmers' perceptions of adaptation strategies, sensitivity and exposure criteria to scientific expertise in order to develop a set of indicators.

Author Contributions: Conceptualization and methodology, M.A., S.C., J.-E.B. and M.W.; writing—original draft preparation, M.A., S.C. and J.-E.B.; investigation, M.A.; formal analysis, M.A.; writing—review and editing, M.A., S.C., J.-E.B. and M.W.; funding acquisition, M.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by INRAE as part of the VACCARM project of the ACCAF metaprogram, project under S2I 00000285, MP-P10182.

Sustainability **2022**, 14, 8275 16 of 23

Informed Consent Statement: Informed consent to participate and publish was obtained from each participant in the study.

Data Availability Statement: Survey instruments and codes used in this study are available from the authors upon request.

Acknowledgments: The authors thank Hélène Raynal, who helped design the statistical analysis, and the trainees who helped collect data. The authors also thank the English proofreading reviewers.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analysis or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Appendix A

The data were cleaned in four major steps (Figure A1). We removed (i) 56 explanatory variables with low heterogeneity (e.g., sex), (ii) 46 explanatory variables that were poorly understood or confusing during the interviews and (iii) 10 strongly correlated variables. In the last step, we sorted the 46 remaining explanatory variables into the eight categories that were previously identified (i.e., general information, water resources, soil resources, human resources, financial resources, crops and rotation, material resources, individual resources) and selected no more than four variables per category to keep those that were the most general. This final step left 25 explanatory variables for the 32 farmers and 41 explained variables.




Figure A1. Diagram of the data cleaning procedure.

Sustainability **2022**, 14, 8275

Appendix B

Description of the 25 explanatory variables.

Table A1. Structural and material variables in the questionnaire.

Category	Variable	Values	
General information	Utilized agricultural area	Small_farm (<100 ha); medium_farm (100-200 ha); large_farm (200-300 ha); very_large_farm (>300 ha)	
	Production system	All conventional; Organic	
Human resources	Labor units (LU)	1_LU; 2_LU; 3_and_more	
Individual resources	Education	CAP_BEP; Bachelor; Engineer_Masters; Education_Other	
	Age	<35; 35–50; >50	
	Type of water source	Watercourse; Lake; Well; Watercourse_lake; Watercourse_lake_well	
Water resources	Volume for maize irrigation in 2020	<2000 m ³ ; 2000–3000 m ³ ; >3000 m ³	
	Irrigable area	All_irrigable; Fields_non_Irrigable	
E-visas at assessed	Number of pivots on the farm	0_pivot; 1_3 Pivots; >3Pivots	
Equipment resources	Volume of grain storage	None; <1000 T; 1000–2000 T; >2000 T	
Soil resources	Soil type	AC (Clay-limestone); AL (Clay-loam); SL (Sandy loam); B (Boulbènes); Others	
	Number of paid activities	1; 2; 3	
Financial resources	Percentage of revenue from maize in 2020	<15%; 15–30%; 30–50%; >50%	
	Distance between fields	<1 km; 1–5 km; 6–14 km; >14 km	
	Number of crops in 2020	1–3; 4–6; >6	
Crops and rotation	Soil tillage for maize	Deep tillage with inversion; Deep tillage without inversion; Reduced tillage (including no tillage)	
	Precocity index for maize	IP_Early (<400); IP_Late (>400); IP_Unknown	

Appendix C

Table A2. Number of times each of the 50 adaptation criteria elicited in the qualitative survey was mentioned. Total number of mentions: 170.

Criterion	Number of Mentions
Increase water storage	13
Diversify water sources	13
Irrigate winter crops	13
Plant cover crops	12
Stop plowing	11
Use sensors	10
Use a weather station	7
Extend the rotation	6
Advance the sowing date for maize	6

Sustainability **2022**, 14, 8275

Table A2. Cont.

Criterion	Number of Mentions
Diversify paid activities	6
Self-sufficiency in water	5
Reduce the precocity for maize	5
Diversify crops	4
Partner with a livestock farmer to obtain manure	4
Buy new land	3
Buy more irrigation quota	3
Diversify production	3
Balance winter and summer crops	3
Reduce soil tillage	3
Buy new seeding equipment	2
Convert to organic agriculture	2
Practice mechanical weeding	2
Increase irrigation efficiency	2
Hire an employee	2
Use modulation	2
Shorten water turns	2
Return harvest residues to the soil	2
Modify the irrigation strategy	2
Improve grain storage	1
Reduce the frequency of field operations	1
Stop growing maize	1
Plant mixed crops	1
Increase grain storage	1
Advance the date of the first irrigation	1
Lime the soil	1
Bury reels for irrigation	1
Scaring practices for wells	1
Practice green tillage	1
Join a group of employers	1
Plant legume crops	1
Use decision-support tools	1
Use a decision-support tool for irrigation	1
Partner with a beekeeper	1
Introduce a livestock production unit	1
Practice sylviculture	1
Use a specific modulation for fertilizer	1

Sustainability **2022**, 14, 8275

Table A2. Cont.

Criterion	Number of Mentions
Use a tall maize variety	1
Use a maize variety resistant to hydric stress	1
Practice direct selling	1
Sell stored maize before summer	1

Appendix D

 $\textbf{Table A3.} \ \ \textbf{The 41} \ \ \textbf{selected adaptation strategy criteria for the quantitative stage of the survey}.$

	Extend the rotation	A1
	Plant mixed crops	A2
Crop pattern and rotation	Plant cover crops	A3
	Diversify crops	A4
	Balance winter and summer crops	A5
	Plant legumes	A6
	Stop maize cultivation	A7
	Advance the sowing date for maize	A8
Maize cultivation	Reduce the precocity for maize	A9
	Use a tall maize variety	A10
	Use a maize variety resistant to hydric stress	A11
	Improve grain storage and commercialization	A12
	Diversify commercialization modes	A13
	Introduce a livestock production unit	A14
	Make a partnership with a neighboring farmer	A15
Farm scale strategy	Diversify production units and/or paid activities	A16
	Buy new lands	A17
	Hire an employee	A18
	Convert to organic farming	A19
	Buy new equipment	A20
	Reduce the frequency of field operations	A21
	Stop plowing	A22
	Lime the soil	A23
Cultural practices	Perform mechanical weeding	A24
-	Practice direct seeding	A25
	Practice strip-tillage	A26
	Return harvest residues to the soil	A27
	Use modulation for inputs	A28

Sustainability **2022**, 14, 8275 20 of 23

Table A3. Cont.

	Buy more quota for irrigation	A29
	Improve/renew equipment for irrigation	A30
	Build or enlarge a reservoir	A31
Water resource	Advance the date of the first irrigation for maize	A32
	Diversify water sources	A33
	Increase irrigation efficiency	A34
	Irrigate winter crops	A35
	Modify the frequency and/or number of water turns	A36
	Use decision-support tools	A37
	Use sensors	A38
Sources of information	Use a weather station	A39
	Attend training courses	A40
	Confront sources of information	A41

Appendix E

Table A4. Distribution of criteria within each of the four clusters identified by hierarchical clustering on principal components. See Table 3 in the article for definitions of the criteria codes.

Cluster	Criteria	Cla/Mod	Mod/Cla	Global	p Value	V Test
1	A36 relevant	50	100	12.90323	0.01290323	2.486429
1	A5 relevant	50	100	12.90323	0.01290323	2.486429
	A32 relevant	83.33333	45.45455	19.354839	0.0138045121	2.462310
2	A31 relevant	100.00000	27.27273	9.677419	0.0367074527	2.089003
2	A24 irrelevant	100.00000	36.36364	12.903226	0.0104878436	2.559316
	A19 irrelevant	58.33333	63.63636	38.709677	0.0485031501	1.972933
	A32 irrelevant	100.00000	57.14286	25.80645	0.0003806699	3.553134
3	A16 relevant	100.00000	28.57143	12.90323	0.0318131257	2.146751
	A17 irrelevant	76.92308	71.42857	41.93548	0.0037953858	2.894685
4	A41 relevant	75.000000	75	12.903226	0.003495948	2.920389
	A4 relevant	60.000000	75	16.129032	0.008580963	2.628313
	A23 irrelevant	100.000000	50	6.451613	0.012903226	2.486429

Cla/mod: percentage of individuals with the modality inside the class (or cluster). Mod/cla: percentage of individuals of the class (or cluster) with the modality.

References

- 1. *IPCC Climate Change* 2021: *The Physical Science Basis*; Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; Cambridge University Press: Cambridge, MA, USA, 2021.
- 2. Chaouche, K.; Neppel, L.; Dieulin, C.; Pujol, N.; Ladouche, B.; Martin, E.; Salas, D.; Caballero, Y. Analyses of precipitation, temperature and evapotranspiration in a French Mediterranean region in the context of climate change. *Comptes Rendus Geosci.* **2010**, 342, 234–243. [CrossRef]
- 3. Olesen, J.E.; Trnka, M.; Kersebaum, K.C.; Skjelvåg, A.O.; Seguin, B.; Peltonen-Sainio, P.; Rossi, F.; Kozyra, J.; Micale, F. Impacts and adaptation of European crop production systems to climate change. *Eur. J. Agron.* **2011**, *34*, 96–112. [CrossRef]
- 4. Peters, K.; Breitsameter, L.; Gerowitt, B. Impact of climate change on weeds in agriculture: A review. *Agron. Sustain. Dev.* **2014**, *34*, 707–721. [CrossRef]
- 5. Zhang, J.T.; Yang, J.; An, P.L.; Ren, W.; Pan, Z.H.; Dong, Z.Q.; Han, G.L.; Pan, Y.Y.; Pan, S.F.; Tian, H.Q. Enhancing soil drought induced by climate change and agricultural practices: Observational and experimental evidence from the semiarid area of northern China. *Agric. For. Meteorol.* **2017**, 243, 74–83. [CrossRef]
- 6. Praveen, B.; Sharma, P. A review of literature on climate change and its impacts on agriculture productivity. *J. Public Aff.* **2019**, 21, e2483. [CrossRef]

Sustainability **2022**, 14, 8275 21 of 23

7. Senthilkumar, K.; Bergez, J.E.; Leenhardt, D. Can farmers use maize earliness choice and sowing dates to cope with future water scarcity? A modelling approach applied to south-western France. *Agric. Water Manag.* **2015**, *152*, 125–134. [CrossRef]

- 8. Caubel, J.; Garcia de Cortazar-Atauri, I.; Vivant, A.C.; Launay, M.; de Noblet-Ducoudré, N. Assessing future meteorological stresses for grain maize in France. *Agric. Syst.* **2018**, *159*, 237–247. [CrossRef]
- 9. Howden, S.M.; Soussana, J.F.; Tubiello, F.N.; Chhetri, N.; Dunlop, M.; Meinke, H. Adapting agriculture to climate change. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 19691–19696. [CrossRef]
- 10. Juhola, S.; Glaas, E.; Linnér, B.-O.; Neset, T.-S.S. Redefining maladaptation. Environ. Sci. Policy 2016, 55, 135–140. [CrossRef]
- 11. Bindi, M.; Olesen, J.E. The responses of agriculture in Europe to climate change. Reg. Environ. Chang. 2011, 11, 151–158. [CrossRef]
- 12. Cammarano, D.; Payero, J.; Basso, B.; Stefanova, L.; Grace, P. Adapting wheat sowing dates to projected climate change in the Australian subtropics: Analysis of crop water use and yield. *Crop Pasture Sci.* **2012**, *63*, 974–986. [CrossRef]
- 13. Moradi, R.; Koocheki, A.; Nassiri Mahallati, M.; Mansoori, H. Adaptation strategies for maize cultivation under climate change in Iran: Irrigation and planting date management. *Mitig. Adapt. Strateg. Glob. Chang.* **2013**, *18*, 265–284. [CrossRef]
- 14. Brunel-Saldias, N.; Martinez, I.; Seguel, O.; Ovalle, C.; Acevedo, E. Structural characterization of a compacted alfisol under different tillage systems. *J. Soil Sci. Plant Nutr.* **2016**, *16*, 689–701. [CrossRef]
- 15. Holzkämper, A. Varietal adaptations matter for agricultural water use—A simulation study on grain maize in Western Switzerland. *Agric. Water Manag.* **2020**, 237, 106202. [CrossRef]
- Barnett, J.; O'Neill, S. Maladaptation. Glob. Environ. Chang. 2010, 20, 211–213. [CrossRef]
- 17. IPCC Climate Change 2007: Impacts, Adaptations and Vulnerability; Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change; Cambridge University Press: Cambridge, UK, 2007; ISBN 9780521880107.
- 18. Füssel, H.M.; Klein, R.J.T. Climate change vulnerability assessments: An evolution of conceptual thinking. *Clim. Chang.* **2006**, 75, 301–329. [CrossRef]
- 19. Darnhofer, I.; Bellon, S.; Dedieu, B.; Milestad, R. Adaptiveness to enhance the sustainability of farming systems. A review. *Agron. Sustain. Dev.* **2010**, *30*, 545–555. [CrossRef]
- 20. Martin, G.; Magne, M.-A.; Cristobal, M.S. An Integrated Method to Analyze Farm Vulnerability to Climatic and Economic Variability According to Farm Configurations and Farmers' Adaptations. *Front. Plant Sci.* **2017**, *8*, 1483. [CrossRef]
- 21. Sneessens, I.; Randrianasolo, H.; Sauvee, L.; Ingrand, S. A quantitative approach to assess farming systems vulnerability: An application to mixed crop-livestock systems. In Proceedings of the 11th Research Seminar on Social Science, Lyon, France, 14–15 December 2017; pp. 1–20.
- 22. Dardonville, M.; Bockstaller, C.; Therond, O. Review of quantitative evaluations of the resilience, vulnerability, robustness and adaptive capacity of temperate agricultural systems. *J. Clean. Prod.* **2021**, *286*, 125456. [CrossRef]
- 23. Andrés Ferreyra, R.; Podestá, G.P.; Messina, C.D.; Letson, D.; Dardanelli, J.; Guevara, E.; Meira, S. A linked-modeling framework to estimate maize production risk associated with ENSO-related climate variability in Argentina. *Agric. For. Meteorol.* **2001**, 107, 177–192. [CrossRef]
- 24. Barkaoui, K.; Roumet, C.; Volaire, F. Mean root trait more than root trait diversity determines drought resilience in native and cultivated Mediterranean grass mixtures. *Agric. Ecosyst. Environ.* **2016**, 231, 122–132. [CrossRef]
- 25. Biggs, R.; Schlüter, M.; Biggs, D.; Bohensky, E.L.; Burnsilver, S.; Cundill, G.; Dakos, V.; Daw, T.M.; Evans, L.S.; Kotschy, K.; et al. Toward principles for enhancing the resilience of ecosystem services. *Annu. Rev. Environ. Resour.* **2012**, *37*, 421–448. [CrossRef]
- 26. Cabell, J.F.; Oelofse, M. An Indicator Framework for Assessing Agroecosystem Resilience. Ecol. Soc. 2012, 17, 18. [CrossRef]
- 27. Wiréhn, L.; Danielsson, Å.; Neset, T.S.S. Assessment of composite index methods for agricultural vulnerability to climate change. *J. Environ. Manag.* **2015**, *156*, 70–80. [CrossRef]
- 28. Urruty, N.; Tailliez-Lefebvre, D.; Huyghe, C. Stability, robustness, vulnerability and resilience of agricultural systems. A review. *Agron. Sustain. Dev.* **2016**, *36*, 1–15. [CrossRef]
- 29. Dardonville, M.; Urruty, N.; Bockstaller, C.; Therond, O. Influence of diversity and intensification level on vulnerability, resilience and robustness of agricultural systems. *Agric. Syst.* **2020**, *184*, 102913. [CrossRef]
- 30. Reidsma, P.; Ewert, F.; Lansink, A.O.; Leemans, R. Adaptation to climate change and climate variability in European agriculture: The importance of farm level responses. *Eur. J. Agron.* **2010**, *32*, 91–102. [CrossRef]
- 31. Dong, Z.; Pan, Z.; An, P.; Wang, L.; Zhang, J.; He, D.; Han, H.; Pan, X. A novel method for quantitatively evaluating agricultural vulnerability to climate change. *Ecol. Indic.* **2015**, *48*, 49–54. [CrossRef]
- 32. Nicholas, K.A.; Durham, W.H. Farm-scale adaptation and vulnerability to environmental stresses: Insights from winegrowing in Northern California. *Global Environ. Chang.* **2012**, 22, 483–494. [CrossRef]
- 33. Gil, J.; Cohn, A.S.; Duncan, J.; Newton, P.; Vermeulen, S. The resilience of integrated agricultural systems to climate change. *WIREs Clim. Chang.* **2017**, *8*, e461. [CrossRef]
- 34. Bouttes, M.; San Cristobal, M.; Martin, G. Vulnerability to climatic and economic variability is mainly driven by farmers' practices on French organic dairy farms. *Eur. J. Agron.* **2018**, *94*, 89–97. [CrossRef]
- 35. Marshall, N.A.; Stokes, C.J.; Webb, N.P.; Marshall, P.A.; Lankester, A.J. Social vulnerability to climate change in primary producers: A typology approach. *Agric. Ecosyst. Environ.* **2014**, *186*, 86–93. [CrossRef]
- 36. Acosta-Michlik, L.; Rounsevell, M.D.A. From Generic Indices to Adaptive Agents: Shifting Foci in Assessing Vulnerability to the Combined Impacts of Climate Change and Globalization. *IHDP Update* **2005**, 14–16.

Sustainability **2022**, 14, 8275 22 of 23

37. Acosta-Michlik, L.; Espaldon, V. Assessing vulnerability of selected farming communities in the Philippines based on a behavioural model of agent's adaptation to global environmental change. *Glob. Environ. Chang.* **2008**, *18*, 554–563. [CrossRef]

- 38. Callo-Concha, D.; Ewert, F. Using the Concepts of Resilience, Vulnerability and Adaptability for the Assessment and Analysis of Agricultural Systems. *Chang. Adapt. Socio-Ecol. Syst.* **2014**, *1*, 1–11. [CrossRef]
- 39. Gbetibouo, G. *Understanding Farmers' Perceptions and Adaptations to Climate Change and Variability;* International Food Policy Research Institute Internship: Washington, DC, USA, 2009.
- 40. Marshall, N.A.; Park, S.E.; Adger, W.N.; Brown, K.; Howden, S.M. Transformational capacity and the influence of place and identity. *Environ. Res. Lett.* **2012**, *7*, 034022. [CrossRef]
- 41. Baca, M.; Läderach, P.; Haggar, J.; Schroth, G.; Ovalle, O. An integrated framework for assessing vulnerability to climate change and developing adaptation strategies for coffee growing families in mesoamerica. *PLoS ONE* **2014**, *9*, e88463. [CrossRef]
- 42. Comoé, H.; Finger, R.; Barjolle, D. Farm management decision and response to climate variability and change in Côte d'Ivoire. *Mitig. Adapt. Strateg. Glob. Chang.* **2014**, *19*, 123–142. [CrossRef]
- 43. Saha, M.K.; Abdul, A.; Biswas, A.; Meandad, J.; Ahmed, R.; Prokash, J.; Sakib, F.M. Factors Affecting to Adoption of Climate-smart Agriculture Practices by Coastal Farmers' in Bangladesh. *Am. J. Environ. Sustain. Dev.* **2019**, *4*, 113–121.
- 44. Jones, L. Resilience isn't the same for all: Comparing subjective and objective approaches to resilience measurement. *Wiley Interdiscip. Rev. Clim. Chang.* **2019**, *10*, e552. [CrossRef]
- 45. Perrin, A.; Cristobal, M.S.; Milestad, R.; Martin, G. Identification of resilience factors of organic dairy cattle farms. *Agric. Syst.* **2020**, *183*, 102875. [CrossRef]
- 46. Binder, C.R.; Feola, G.; Steinberger, J.K. Considering the normative, systemic and procedural dimensions in indicator-based sustainability assessments in agriculture. *Environ. Impact Assess. Rev.* **2010**, *30*, 71–81. [CrossRef]
- 47. Smeets, E.; Weterings, R.; Centre, T.N.O.; Bosch, P.; Büchele, M.; Gee, D. *Environmental Indicators: Typology and Overview. Technical Report N*°25; European Environment Agency: Copenhagen, Denmark, 1999.
- 48. Lairez, J.; Feschet, P.; Aubin, J.; Bockstaller, C.; Bouvarel, I. Agriculture et Développement Durable: Guide Pour L'évaluation Multicritère. Available online: https://books.google.fr/books?hl=fr&lr=&id=RpxmCwAAQBAJ&oi=fnd&pg=PA7&dq=guide+d%27%C3%A9valuation+multicrit%C3%A8re+d%C3%A9finition+crit%C3%A8re&ots=-0hN6CaMtX&sig=-askjH7E1gpT3Ra3 xwOkhm0say4 (accessed on 10 May 2021).
- 49. DRAAF. Analyse Économique des Exploitations Agricoles Irriguant du Maïs Grain; DRAAF: Marseille, France, 2017; Volume 2014.
- 50. DRAAF. Occitanie Bilan 2020 Grandes Cultures; DRAAF: Marseille, France, 2020.
- 51. Lopez, O.; Milhaud, X.; Thérond, P. Arbres de régression et de classification (CART). L'actuariel 2015, 15, 42-44.
- 52. R Core Team. R: The R Project for Statistical Computing. Available online: https://www.r-project.org/ (accessed on 10 August 2020).
- 53. Husson, A.F.; Josse, J.; Le, S.; Mazet, J.; Husson, M.F. Package 'FactoMineR'. Available online: http://factominer.free.fr (accessed on 10 August 2020).
- 54. Therneau, T.M.; Atkinson, E.J. *An Introduction to Recursive Partitioning Using the RPART Routines*; Mayo Foundation: Rochester, MI, USA, 2022.
- 55. Milborrow, A.S. Package 'rpart.plot'. Available online: http://www.milbo.org/rpart-plot/index.html (accessed on 10 August 2021).
- 56. Croissant, Y. Mlogit: Random utility models in r. J. Stat. Softw. 2020, 95, 1–41. [CrossRef]
- 57. DRAAF. Occitanie L'irrigation Contribue à 18% de la Valeur de la Production Agricole; DRAAF: Marseille, France, 2018.
- 58. He, R.; Jin, J.; Kuang, F.; Zhang, C.; Guan, T. Farmers' risk cognition, risk preferences and climate change adaptive behavior: A structural equation modeling approach. *Int. J. Environ. Res. Public Health* **2020**, *17*, 85. [CrossRef]
- 59. Willaume, M.; Rollin, A.; Casagrande, M. Farmers in southwestern France think that their arable cropping systems are already adapted to face climate change. *Reg. Environ. Chang.* **2014**, *14*, 333–345. [CrossRef]
- 60. Martin-Clouaire, R. Modelling Operational Decision-Making in Agriculture. Agric. Sci. 2017, 8, 527–544. [CrossRef]
- 61. Dessart, F.J.; Barreiro-Hurlé, J.; Van Bavel, R. Behavioural factors affecting the adoption of sustainable farming practices: A policy-oriented review. *Eur. Rev. Agric. Econ.* **2019**, *46*, 417–471. [CrossRef]
- 62. Roussy, C.; Ridier, A.; Chaib, K. Adoption d'innovations par les agriculteurs: Rôle des perceptions et des préférences. *Work. Pap. SMART—LERECO* **2015**, 15–03, 1–22.
- 63. Albert, M.; Couture, S.; Willaume, M.; Bergez, J.É.; Faivre, R. Decision-Making Process Factors Explain Some of the Heterogeneity of Irrigation Practices among Maize Farmers in Southwestern France. *Water* **2021**, *13*, 3504. [CrossRef]
- 64. Reynaud, A.; Couture, S. Stability of risk preference measures: Results from a field experiment on French farmers. *Theory Decis.* **2012**, 73, 203–221. [CrossRef]
- 65. Nuthall, P.L.; Old, K.M. Intuition, the farmers' primary decision process. A review and analysis. *J. Rural. Stud.* **2018**, *58*, 28–38. [CrossRef]
- 66. Wienroth, M. Value beyond scientific validity: Let's RULE (Reliability, Utility, LEgitimacy). *J. Responsible Innov.* **2020**, *7*, 92–103. [CrossRef]
- 67. Schlüter, M.; Baeza, A.; Dressler, G.; Frank, K.; Groeneveld, J.; Jager, W.; Janssen, M.A.; McAllister, R.R.J.; Müller, B.; Orach, K.; et al. A framework for mapping and comparing behavioural theories in models of social-ecological systems. *Ecol. Econ.* **2017**, *131*, 21–35. [CrossRef]

Sustainability **2022**, 14, 8275 23 of 23

68. Breiman, L.; Friedman, J.; Olshen, R.; Group, C.S.-I. Classification and Regression Trees; Wadsworth: Belmont, CA, USA, 1984.

- 69. Larmarange, J. Régression Logistique Binaire, Multinomiale et Ordinale. Available online: https://larmarange.github.io/analyse-R/analyse-R.pdf (accessed on 10 May 2021).
- 70. Neset, T.-S.; Juhola, S.; Wiréhn, L.; Käyhkö, J.; Navarra, C.; Asplund, T.; Glaas, E.; Wibeck, V.; Linnér, B.-O. Supporting Dialogue and Analysis on Trade-Offs in Climate Adaptation Research With the Maladaptation Game. *Simul. Gaming* **2020**, *51*, 378–399. [CrossRef]
- 71. Rickards, L.; Howden, S.M. Transformational adaptation: Agriculture and climate change. *Crop Pasture Sci.* **2012**, *63*, 240–250. [CrossRef]
- 72. Magnan, A.K.; Schipper, E.L.F.; Burkett, M.; Bharwani, S.; Burton, I.; Eriksen, S.; Gemenne, F.; Schaar, J.; Ziervogel, G. Addressing the risk of maladaptation to climate change. *Wiley Interdiscip. Rev. Clim. Chang.* **2016**, *7*, 646–665. [CrossRef]
- 73. Holzkämper, A. Adapting agricultural production systems to climate change—What's the use of models? *Agriculture* **2017**, 7, 86. [CrossRef]
- 74. Wiréhn, L. Nordic agriculture under climate change: A systematic review of challenges, opportunities and adaptation strategies for crop production. *Land Use Policy* **2018**, 77, 63–74. [CrossRef]