



HAL
open science

Effect of sainfoin (*Onobrychis viciifolia*) on cyathostomin eggs excretion, larval development, larval community structure and efficacy of ivermectin treatment in horses

Joshua Malsa, Elise Courtot, Michel Boisseau, Bertrand Dumont, Pascale Gombault, Tetiana A. Kuzmina, Marta Basiaga, Jérôme Lluch, Gwenolah Annonay, Sophie Dhorne-Pollet, et al.

► To cite this version:

Joshua Malsa, Elise Courtot, Michel Boisseau, Bertrand Dumont, Pascale Gombault, et al.. Effect of sainfoin (*Onobrychis viciifolia*) on cyathostomin eggs excretion, larval development, larval community structure and efficacy of ivermectin treatment in horses. *Parasitology*, 2022, 149 (11), pp.1439-1449. 10.1017/S0031182022000853 . hal-03753744

HAL Id: hal-03753744

<https://hal.inrae.fr/hal-03753744>

Submitted on 18 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Effect of sainfoin (*Onobrychis viciifolia*) on cyathostomin eggs excretion, larval development, larval community structure and efficacy of ivermectin treatment in horses

Research Article

*These authors contributed equally to this work

Cite this article: Malsa J *et al* (2022). Effect of sainfoin (*Onobrychis viciifolia*) on cyathostomin eggs excretion, larval development, larval community structure and efficacy of ivermectin treatment in horses. *Parasitology* 1–11. <https://doi.org/10.1017/S0031182022000853>

Received: 5 April 2022

Revised: 25 May 2022

Accepted: 9 June 2022



Key words:

Fecal egg count; ITS-2; nemabiome; nematode; nutraceutical; strongylid; tannin

Author for correspondence:

Joshua Malsa,

E-mail: joshua.alexandre.malsa@gmail.com

Joshua Malsa¹ , Élise Courtot¹, Michel Boisseau¹, Bertrand Dumont², Pascale Gombault³, Tetiana A. Kuzmina⁴, Marta Basiaga⁵, Jérôme Lluch⁶, Gwennolah Annonay⁶, Sophie Dhorne-Pollet⁷, Nuria Mach⁸, Jean-François Sutra⁹, Laurence Wimel¹⁰, Cédric Dubois¹⁰, Fabrice Guégnard¹, Delphine Serreau¹, Anne Lespine⁹, Guillaume Sallé^{1,*}  and Géraldine Fleurance^{2,11,*}

¹INRAE, Université de Tours, UMR 1282 Infectiologie et Santé Publique, Nouzilly, France; ²INRAE, Université Clermont Auvergne, VetAgro Sup, UMR 1213 Herbivores, Saint-Genès Champanelle, France; ³Multifolia, Viâpres-le-Petit, France; ⁴Department of Parasitology, I. I. Schmalhausen Institute of Zoology NAS of Ukraine, Kyiv, Ukraine; ⁵Department of Zoology and Animal Welfare, Faculty of Animal Science, University of Agriculture in Kraków, 24/28 Mickiewiczza Av., 30-059 Kraków, Poland; ⁶GeT-PlaGe, INRAE, Genotoul, Castanet-Tolosan, France; ⁷Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350 Jouy-en-Josas, France; ⁸IHAP, Université de Toulouse, INRAE, ENVT, Toulouse Cedex 3, 31076, France; ⁹INTHERES, Université de Toulouse, INRAE, ENVT, Toulouse Cedex 3, 31076, France; ¹⁰IFCE, Plateau technique de la Station Expérimentale, Chamberet, France and ¹¹IFCE, Pôle Développement Innovation et Recherche, Exmes, France

Abstract

Alternative strategies to chemical anthelmintics are needed for the sustainable control of equine strongylids. Bioactive forages like sainfoin (*Onobrychis viciifolia*) could contribute to reducing drug use, with the first hints of *in vitro* activity against cyathostomin free-living stages observed in the past. We analysed the effect of a sainfoin-rich diet on cyathostomin population and the efficacy of oral ivermectin treatment. Two groups of 10 naturally infected horses were enrolled in a 78-day experimental trial. Following a 1-week adaptation period, they were either fed with dehydrated sainfoin pellets (70% of their diet dry matter) or with alfalfa pellets (control group) for 21-days. No difference was found between the average fecal egg counts (FECs) of the two groups, but a significantly lower increase in larval development rate was observed for the sainfoin group, at the end of the trial. Quantification of cyathostomin species abundances with an ITS-2-based metabarcoding approach revealed that the sainfoin diet did not affect the nemabiome structure compared to the control diet. Following oral ivermectin treatment of all horses on day 21, the drug concentration was lower in horses fed with sainfoin, and cyathostomin eggs reappeared earlier in that group. Our results demonstrated that short-term consumption of a sainfoin-rich diet does not decrease cyathostomin FEC but seems to slightly reduce larval development. Consumption of dehydrated sainfoin pellets also negatively affected ivermectin pharmacokinetics, underscoring the need to monitor horse feeding regimes when assessing ivermectin efficacy in the field.

Introduction

Small strongyles or cyathostomins are a large group of gastrointestinal parasitic nematodes infecting grazing equids. Cyathostomins encompass more than 50 species able to infect domestic and wild equids (Lichtenfels *et al.*, 2008), among which a limited group of 10–12 species emerges as a prevalent and abundant parasitic core consistently found worldwide (Reinemeyer *et al.*, 1984; Bucknell *et al.*, 1995; Lyons *et al.*, 2000; Collobert-Laugier *et al.*, 2002; Osterman Lind *et al.*, 2003, 2007; Kuzmina *et al.*, 2005, 2012). The main clinical effect of cyathostominosis is weight loss, and the massive simultaneous emergence of larval stages encysted in the colonic mucosa, can be fatal for horses especially for young animals (Love and Duncan, 1992; Love *et al.*, 1999; Peregrine *et al.*, 2014). This larval cyathostominosis syndrome, characterized by protein-losing enteropathy, diarrhoea and colic (Giles *et al.*, 1985), remains the leading cause of parasite-mediated death of young horses in some areas like France (Sallé *et al.*, 2020). Cyathostominosis control has long been achieved mainly through the use of chemical anthelmintics (Love, 2003; Peregrine *et al.*, 2014), but this has promoted the selection of isolates resistant to all anthelmintics used in equine medicine (Kaplan, 2002, 2004; Traversa *et al.*, 2012; Matthews, 2014). Macrocyclic lactones, including ivermectin (IVM) and moxidectin, remain the most effective drug class against small strongylids (Peregrine *et al.*, 2014; Sallé *et al.*, 2017; Nielsen *et al.*, 2020). They also remain the most widely used anthelmintic drugs in equine operations in the USA (Becher *et al.*, 2018; Nielsen *et al.*, 2020) or

© The Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial licence (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use.




Fig. 1. Experimental design. The figure depicts the time points and sampling done in this experiment.

in the UK, where IVM and moxidectin are included in drenching programmes in more than a third of cases (Stratford *et al.*, 2014; Tzelos *et al.*, 2019). However, reduced IVM efficacy has been evidenced (Nielsen *et al.*, 2020), including a reduced egg reappearance period (ERP) (Lyons *et al.*, 2011a, 2011b; Nielsen *et al.*, 2020). In addition, high concentrations of these molecules are found in the feces of treated animals, with major detrimental effects on coprophagous organisms (Lumaret *et al.*, 2012; Verdú *et al.*, 2018).

This context warrants alternative strategies to the use of chemical anthelmintics, and the prospect of using bioactive plant compounds as nutraceuticals for the control of gastrointestinal parasites in livestock has been an active research field (Paolini *et al.*, 2003; Heckendorn *et al.*, 2007; Rochfort *et al.*, 2008; Manolaraki *et al.*, 2010; Sandoval-Castro *et al.*, 2012; Hoste *et al.*, 2015). Tannin-rich plants are studied for their anthelmintic activity and their use as bioactive forage (Hoste *et al.*, 2006, 2012; Gaudin *et al.*, 2016; Peña-Espinoza *et al.*, 2018).

Among tannin-rich plants, sainfoin (*Onobrychis viciifolia*) is often used as a model fodder (Hoste *et al.*, 2012). Some *in vitro* tests on ruminant gastrointestinal nematodes have shown that sainfoin and the tannins reduce larval migration (Paolini *et al.*, 2004; Manolaraki *et al.*, 2010). *In vivo* studies have reported a decrease in fecal egg counts (FECs), associated with a reduction in worm fertility or counts in goats or sheep fed with sainfoin (Paolini *et al.*, 2003, 2005; Heckendorn *et al.*, 2006; Manolaraki *et al.*, 2010; Gaudin *et al.*, 2016). In horses, a previous study (Collas *et al.*, 2018) evaluated the anthelmintic activity of dehydrated sainfoin pellets on equine strongyles. While strongyle FEC did not differ between horses fed with sainfoin compared to the control group, an *in vitro* approach using sainfoin pellet water solutions revealed a decrease in egg hatching and larval development (Collas *et al.*, 2018). However, it remains unclear whether the latter effects can be reproduced *in vivo* and if the putative effects of sainfoin extracts vary across various cyathostomin species. The recent development of an ITS-2-based metabarcoding approach has allowed the simultaneous identification and quantification of strongyle abundances in domestic horses (Mitchell *et al.*, 2019; Poissant *et al.*, 2021) or wild equids (Tombak *et al.*, 2021). This approach has been used to characterize the response to anthelmintic treatment of trichostrongylids in ruminants (Queiroz *et al.*, 2020; Halvarsson and Höglund, 2021) or the monitoring of drug resistance in these species (Queiroz

et al., 2020). It has never been applied yet to study a plant extract effect on any gut parasite community.

In addition to their anthelmintic activity, tannin-rich plants contain flavonoids. These molecules modulate ATP binding cassette (ABC) transporters (Morris and Zhang, 2006), thereby affecting the *in vitro* and *in vivo* pharmacodynamics and activity of macrocyclic lactones (Lespine *et al.*, 2008). For instance, quercetin significantly increases exposure to moxidectin in lambs (Dupuy *et al.*, 2003). The tannin-rich plant redberry juniper (*Juniperus pinchotii*) used in combination with oral IVM treatment has also increased the treatment efficacy against *Haemonchus contortus* in lambs (Whitney *et al.*, 2013). However, an *in vivo* trial in sheep reported a significant decrease in IVM efficacy in sainfoin-fed animals due to a concomitant reduction in plasma IVM concentrations (Gaudin *et al.*, 2016). Such interaction remains uncharacterized in horses to date.

To bridge this knowledge gap, the current study aimed (i) to evaluate the *in vivo* effect of dehydrated sainfoin pellets on cyathostomins (fecal egg excretion and larval development), (ii) to quantify any alterations of the cyathostomin larval community structure using a nemabiome approach, and (iii) to establish how the sainfoin diet may affect the efficacy of an oral IVM treatment in horses.

Materials and methods

The *in vivo* experiment was conducted from September 8th to December 15th 2020, at the French Horse and Riding Institute experimental farm (IFCE) in Chamberet, France. The experimental procedure received approval from the French Ministry of Research under protocol number APAFIS#26140-2020062216271790v2. The experimental design is summarized in Fig. 1.

Animal condition

Naturally infected saddle horses (Anglo-Arab breed, 2 years old) left undrenched for 137 days (last anthelmintic administered on 24 April 2020; 200 μg IVM kg^{-1} body weight (BW) and 1 mg praziquantel kg^{-1} BW; EQVALAN® DUO, France) were allocated into 2 groups of 10 horses. Horses were mainly infected by small strongylids (>95%), as revealed by larval culture (Collas *et al.*, 2018) and the herein reported nemabiome approach which did not disclose the presence of *Strongylus* spp. Both groups were balanced for

sex (8 females and 2 geldings in each group), FEC data before housing (September 8th 2020; sainfoin group FEC = 1875 ± 1253 eggs g^{-1} (EPG); control group FEC = 1940 ± 1064 EPG, mean \pm s.d.), and for their BW measured on September 15th (average BW of 454.8 ± 28.1 kg and 457.6 ± 25.6 kg in the sainfoin and control diet groups, respectively). From September 8th to 21th (day-21 to day-8; see Fig. 1), all horses were housed in a single stall and collectively fed with a transition diet composed of 77% grassland hay and 23% of concentrate (made of 61.5% of barley, 35% of soya bean meal, 3.5% of minerals and vitamins), designed to meet their energy (6.0 UFC day^{-1}) (UFC; horse feed unit) and protein requirements (318 g MADC day^{-1}) (MADC; horse digestible crude protein), as previously defined for this breed (INRA, 2015).

Experimental diet

Horses were adapted (adaptation period) to their experimental diet for a week – from day-7 to day 0 (Fig. 1), i.e. the horses were fed individually with a decreasing proportion of the transition diet and an increasing proportion of the experimental diet. They subsequently received their experimental diet for the 3 following weeks (day 0–21, Fig. 1). The horses were housed in individual stalls during these 2 periods (from day-7 to day 21; see Fig. 1). This duration was in line with successful experiments in ruminants (Paolini *et al.*, 2005; Heckendorn *et al.*, 2006) and with practical implementation in the field. The sainfoin diet contained 2.3% dry matter (DM) of condensed tannins as determined by an acetone–butanol–HCl assay (Grabber *et al.*, 2013); it included 70% DM Equifolia dehydrated sainfoin pellets (*Onobrychis viciifolia*; provided by Multifolia, Viapres-le-Petit, France) and 30% grassland hay (on DM bases). We worked with a production chain for dehydrated sainfoin pellets, which ensured appropriate agronomic conditions for the plant cultivation, a good conservation of sainfoin characteristics over time and the standardization and characterization of the batches before use. The control diet consisted of 60% DM of alfalfa (*Medicago sativa*) pellets and 40% DM of grassland hay (Table 1; Supplementary Table S1). Proportions of diet components were chosen to maximize the amount of condensed tannins in the sainfoin diet (while taking care not to cause digestive problems) and to ensure each diet covered horse energy and protein requirements. The energy requirements (UFC) were covered at $108\% \pm 1.3$ and $103\% \pm 1.5$ on average for the sainfoin and control groups, respectively. This difference was statistically significant between the 2 groups ($P < 10^{-4}$), but our previous observations have shown that energy requirements were not affecting the outcome of the comparison (Collas *et al.*, 2014). The protein requirements (MADC) were covered at $250\% \pm 3.7$ and $252\% \pm 5.5$ on average for the sainfoin and control groups, respectively ($P = 0.13$). These values were safe for the duration of the treatment and showed no significant correlation with FEC or larval development rate ($P = 0.45$ and $P = 0.89$, respectively), in good agreement with past experimental findings (Collas *et al.*, 2018). Individual requirements were estimated based on INRA (2015) tables for UFC and MADC as

$$\begin{aligned} \text{UFC requirements (per kg BW}^{0.75}) \\ = 0.0594 + 0.0252 \times \text{BW gain}^{1.4} \end{aligned}$$

$$\begin{aligned} \text{MADC requirement (in g day}^{-1}) &= 2.8 \times \text{BW}^{0.75} + 270 \\ &\times \text{BW gain, with BW gain} = 0.15 \text{ kg day}^{-1} \text{ for 2-year-old horses.} \end{aligned}$$

Horses received half of their diet at 8:00 am and the other half at 4:00 pm. Quantities offered were adjusted every Tuesday based

Table 1. Chemical composition and nutritive value of foodstuffs offered to the 2 groups (sainfoin diet, control diet) of horses during the experimental period^a

	Sainfoin pellets ^b	Alfalfa pellets ^c	Grassland hay ^d
DM (g kg^{-1})	921	941	976
CP (g kg^{-1} DM) ^e	169	184	52
CF (g kg^{-1} DM) ^f	231	287	354
UFC (kg ⁻¹ DM) ^g	0.67	0.62	0.47
MADC (g kg^{-1} DM) ^g	99	110	14

DM, dry matter; CP, crude protein; CF, crude fibre; UFC, horse feed unit; MADC, digestible protein.

^aAnalyses were performed by UpScience, Saint-Nolff, France, on 4 samples of each foodstuff collected all along the experimental period.

^{b,d}In sainfoin diet.

^{c,d}In control diet.

^eDumas method.

^fWeende method.

^gFrom INRA (INRA, 2015) equations.

on BW changes and the DM content of diet components. If an individual horse lost weight, its BW measured the week before was used to determine its requirements. Individual refusals were weighed every morning, and they never exceeded 5% DM of the offered diet for more than 3 consecutive days, thereby warranting any particular adaptation.

Fecal sample analysis

Fecal samples were collected individually from the rectum of each horse every Monday of the experimental period (day 0, day 7, day 14 and day 21). Samples were stored at $+4$ °C and shipped to the INRAE Centre Val de Loire facilities (Nouzilly, France) for further processing. Individual FEC data were determined using a modified McMaster technique (Raynaud *et al.*, 1970) based on 5 g of fecal matter diluted with a dilution factor of 5. Eggs were then counted using optical microscopy ($\times 150$ magnification), with a minimum detection limit of 50 EPG. To evaluate the effect of the diet on larval development, the remaining fecal matter (40–90 g) was incubated individually for each horse for 12 days at $+25$ °C and 60% relative humidity during 2 weeks as suggested by Roberts and O'Sullivan (1950). The correlation between the quantity of fecal matter cultured and the larval development was not significant (Spearman's $\rho = 0.08$, $P = 0.4$). Infective third-stage larvae (L3) of cyathostomins were then collected using a Baermann apparatus after 24 and 48 h of sedimentation, and pooled together for each horse and time point. To count larval concentration, 30 drops of $5 \mu\text{L}$ were taken from a homogenized larval suspension (using a bar magnet in a glass beaker) and inspected using optical microscopy. The average number of larvae across the 30 drops was related to the total volume of larval suspension collected after Baermann. Following Collas *et al.* (2018), the larval development rate was then derived as:

$$\left(\frac{\text{Counted L3}}{\text{FEC} \times \text{quantity of fecal matter}} \right) \times 100$$

Gastrointestinal nemabiome

To identify the putative effects of sainfoin on the gastrointestinal nemabiome, a metabarcoding approach was applied to cyathostomin larval populations (using pools of 20 000 L3) harvested from fecal samples collected on day 0 and day 21 of the experiment. Larvae were incubated with $10 \mu\text{L}$ of a 20 mg mL^{-1} proteinase K

(Qiagen) solution at 56 °C for 3 h; DNA was subsequently extracted using a phenol–chloroform protocol (Sambrook and Russell, 2006) and eluted in a 30 µL TE (Tris-EDTA) buffer solution. For metabarcoding, the ITS-2 gene region was PCR amplified using the NC1 (5'-ACGTCTGGTTCAGGGTTGTT-3') and NC2 (5'-TTAGTTTCTTTTCCTCCGCT-3') primers (Gasser et al., 1993). Between 1 and 3 random bases were added to the 5' primer end to increase sequence complexity. Moreover, a 2-bp linker and an Illumina 28-bp overhang were added for the forward and reverse sequences, respectively, for subsequent ligation with Illumina adapters. PCR was run for 3 min at 95 °C for the first denaturation, then 30 cycles starting with 15 s at 98 °C, then 60 °C for 15 s and 72 °C for 15 s, followed by a final extension of 72 °C for 2 min. PCR products were then loaded on a 1% agarose gel to validate the presence of an amplicon product. After this step, 31 out of the 40 samples from 16 horses remained for library preparation. Because MiSeq enables paired 250-bp reads, the ends of each read are overlapped and can be stitched together to generate extremely high-quality, full-length reads of the entire region in a single run. Single multiplexing was performed using a homemade 6 bp index, which was added to reverse primer during a second PCR with 12 cycles using forward primer (AATGATACGGCGACCACCGAGATCTACACTC TTCCCTACACGAC) and reverse primer (CAAGCAGAAGA CGGCATACGAGAT-index-GTGACTGGAGTTCAGACGTGT). The resulting PCR products were purified and loaded onto the Illumina MiSeq cartridge according to the manufacturer's instructions. Libraries were further processed for a single run of MiSeq sequencing using 500 cycle reagent kit v3 (2 × 250 output; Illumina, USA), the raw data of which are registered under SRA bioproject number PRJNA840924. The quality of the run was checked internally using PhiX, and then each paired-end sequence was assigned to its sample with the help of the previously integrated index.

To assess the predictive ability of the nemabiome approach and determine the best set of parameters for data processing, five mock communities of known species composition were built from worms collected and morphologically identified in Ukraine (T. A. Kuzmina, personal communication) or Poland (M. Basiaga, personal communication). Two mock communities of 5 species (a single male worm each), including *Cylicocyclus nassatus*, *Cylicocyclus insigne*, *Cyathostomum catinatum*, *Cyathostomum pateratum* and *Coronocyclus labiatus* were considered using either raw or equimolar DNA concentrations. In addition, two mock communities comprising *C. pateratum* and *C. catinatum* (equivalent proportion or in a 1:4 ratio) were also considered to establish the ability of the ITS-2 sequence to delineate between these 2 phylogenetically close species (Hung et al., 2000). Raw reads were processed with Cutadapt v.1.14 (Martin, 2011) to remove bad quality bases at the 3' end of reads (-q 15), trim primer sequences, remove sequences with evidence of indels (--no-indels) or that showing no trace of primer sequence (--discard-untrimmed). Quality filtered ITS-2 amplicon sequences were subsequently handled with the dada2 algorithm (Callahan et al., 2016). A set of different parameters was tested to measure the total number of amplicon sequence variants (ASVs), the rate of taxonomic assignment, and the proportion of false-positive and false-negative detection in mock community samples. First, the maximum error rate (--max-ee) was set to 1 for both reads or 2 and 5 for forward and reverse reads, respectively. The read truncation parameter was either 200 bp or 217 bp, corresponding to the minimal average read length measured across all samples as determined with FastQC v.0.11.7 (Andrews, 2010). The BAND_SIZE parameter effect (that penalizes the number of insertions permitted when aligning 2 sequences) was tested using either the default or the ITS-2 recommended values

(16 and 32, respectively), or disabling banding (BAND_SIZE = -1) to perform a full Needleman–Wunsch alignment as described elsewhere (Poissant et al., 2021). Denoising was performed using the *pseudo-pool* option, and chimaera removal was performed under the default *consensus* mode. Taxonomic classification was subsequently done using the curated ITS-2 database (<https://www.nemabiome.ca/its2-database.html>), last accessed on 2 February 2022 (Workentine et al., 2020). Two taxonomic assignments were considered and compared, considering mock communities as a ground truth. First, the *IdTaxa* (Murali et al., 2018) function implemented in the DECIPHER package v2.18.1 (Wright, 2016) was applied with 100 bootstraps and a threshold of 50%. Second, the *dada2 assignTaxonomy* (Callahan et al., 2016) function was implemented as already described (Poissant et al., 2021), i.e. setting *tryRC* = TRUE, *minBoot* = 0, *outputBootstraps* = TRUE and retaining ASVs with 50% bootstrap support. In both cases, the number of false-positive and false-negative counts were determined.

Count tables data were further analysed in R v. 4.1 (R Team, 2020) using the phyloseq (McMurdie and Holmes, 2013) and vegan packages (Oksanen et al., 2015). Rare ASVs (<50 counts in total) were considered contaminants and removed. Samples with less than 50 read counts (negative water control; *n* = 1 at day 21) were also discarded.

Egg reappearance period

At the end of the sainfoin-fed diet (day 21), all horses were treated with oral IVM (Eqvalan, 200 µg kg⁻¹ BW, Boehringer Ingelheim, Lyon, France) and were still fed with their respective experimental diets for 4 days. In addition, the ERP was determined from measurements of individual FECR data on 5 occasions over a 42-day window (on day 36, day 50, day 63, day 71 and day 78). The ERP was defined by WAAVP (Nielsen et al., 2022) as the time when the upper confidence interval for the mean fecal egg count reduction (FECR) fell below the mean of FECR determined 2 weeks post-treatment minus 10%. FECR data were calculated according to the WAAVP guidelines (Coles et al., 1992, 2006) as

$$\left(\frac{FECd21 - FEC_i}{FECd21} \right) \times 100,$$

where *FECd21* – FEC data on day 21, *FEC_i* stands for the FEC measured at each timepoint *i* between day 36 and day 78.

Ivermectin dosage in plasma and pharmacokinetics parameters

To quantify horse exposure to IVM, blood samples (9 mL) were taken from the jugular vein of each horse in heparinized tubes at 0, 1, 2, 24, 48, 72 and 96 h after IVM treatment. Blood samples were kept at +4 °C before centrifugation (1500 g for 30 min), and the plasma was frozen at -20 °C until further shipment and processing. Plasmatic IVM concentration was determined by high-performance liquid chromatography (HPLC) with fluorescence detection according to previously described and validated methods (Alvinerie et al., 1999). Data were analysed using a non-compartmental approach with version 4.2 of the Kinetic Tm computer program (innaPhase, Philadelphia, USA). The partial area under the plasma concentration–time curve (AUC) was calculated from 1 to 96 h by the linear trapezoidal rule. *C_{max}*, maximal concentration was then determined. Data were expressed as geometric mean ± standard error of the mean.




Fig. 2. Arithmetic average of FEC (A) and larval development rate (B) measured over the experimental period. Weekly arithmetic average FEC (A) or larval development rate (B) measured throughout the experimental period in horses receiving the control (light grey) or sainfoin (dark green) diet. The average cyathostomin FEC on day 21 (represented by the bar) was significantly different from the average FEC on day 0 ($P = 0.017$) (A). *Statistically significant difference in larval development rate from day 14 to day 21 between the sainfoin and control groups ($P = 0.02$). The bars indicate a significant difference in the mean larval development at day 7 ($P = 0.04$) and day 14 ($P = 9.8 \times 10^{-5}$), both compared to day 0 (B).

Statistical analysis

To test for the effect of the diet on FEC and larval development rate, a generalized estimating equation model was implemented with the *geeglm* function of the *geepack* v.1.3-1 package (Højsgaard *et al.*, 2006).

To test for differences in alpha diversity, a *t*-test was performed between the Shannon index values estimated for the sainfoin and control group on day 0 and day 21. To test for a differential temporal trend in species abundance between both groups during the experimental diet, species counts were regressed upon the experimental group and the day effects and setting horse as a random effect with the *nlme* package v.3.1-155. This analysis was restricted to 12 horses with data on both days ($n = 8$ in the sainfoin group, $n = 4$ in the control group) and to the most abundant cyathostomin species (overall abundance of 10 000 counts, including *C. nassatus*, *Cylicocycylus ashworthi*, *Cylicostephanus minutus* and *C. longibursatus*). Species counts were 4th-root transformed to better fit normality (Shapiro–Wilk test = 0.56 and 0.96 before and after transformation, respectively).

FEC measured after IVM treatment was modelled with a linear mixed-effects model using the *lme* function of the *nlme* package (Pinheiro *et al.*, 2013), fitting diet, time and diet \times time interaction terms as fixed effects and horse as a random effect. The FECR was measured according to the WAAVP guidelines (Coles *et al.*, 1992, 2006) with Bayesian hierarchical models using the *fecrCI* function on the *eggCounts* package v.3.2-3 (Wang *et al.*, 2022).

The average IVM concentration and C_{\max} for the sainfoin group were compared to the control group at every time point (+1, +2, +24, +48, +72 and +96 h). Data were analysed with a Mann–Whitney *U* test, and the *P* values were corrected for multiple tests using a Bonferroni procedure as implemented in the *wilcox_test* function of *rstatix* package v.0.6.0 (Kassambara, 2021). The pharmacokinetics (PK) parameters (AUC) for each group were compared with an unpaired *t*-test, and the *P* values were corrected by Bonferroni, as explained above.

Results

Sainfoin effect on FEC excretion

When horses received their experimental diet (day 0 to day 21), individual cyathostomin FECs significantly decreased by 35% from 1165 ± 545 EPG to 765 ± 396 EPG on average ($P = 0.017$, Fig. 2A). However, no significant difference between the sainfoin and the control group was found at any considered time point ($P = 0.90$).

Sainfoin effect on larval development rate

The average larval development rate (Fig. 2B) decreased between day 0 and day 14 of the experiment from an average development of $32.3 \pm 12.8\%$ at day 0 to $25.2 \pm 13.3\%$ at day 7 ($P = 0.04$) and $21.5 \pm 10.4\%$ at day 14 ($P = 9.8 \times 10^{-5}$). This decrease was similar between the 2 diets ($P = 0.27$ and 0.06 on day 7 and day 14, respectively). However, the subsequent increase in larval development rate from day 14 to day 21 was significantly lower in the sainfoin-fed group ($P = 0.02$, Fig. 2B).

Sainfoin effect on cyathostomin larval community structure

Among the combinations of parameters tested, a BAND_SIZE of -1 combined with a truncation length of 217 bp and maximal error rates of 2 and 5 for the forward and reverse reads were optimal. This combination yielded the highest fraction of assigned amplicon sequence variants without false-positive calls in the mock parasite community (83.6% assigned ASVs out of the 110 ASVs detected in total; Supplementary Table S2). Using this set of parameters and after filtering, 31 samples were left for analysis with an average of 6768 reads per sample (ranging between 2269 and 25 434 reads) that defined 110 ASVs. Out of these, 15 were considered contaminants (<50 occurrences), 75 ASVs were assigned to 13 identified cyathostomin species and 6 remained undetermined (amounting to 0.6% of total counts). Members of the *Cyathostomum* and *Cylicocycylus* genera accounted for the highest misassignment rate at the species level




Fig. 3. Cyathostomin larval community structure estimated using the metabarcoding approach across days and groups. Relative abundance of cyathostomin species in control (upper panels) and sainfoin-fed horses (lower panels) on days 0 and 21 of the experiment. Data are from 12 horses with samples successfully amplified.

(5.78 and 2.75% of total counts, respectively), while unassigned *Cylicostephanus* spp. represented 0.09% of total counts. To this respect, taxonomic assignment with the *Idtaxa* function performed better than the *assignTaxonomy* function that consistently introduced false-positive identification of *C. longibursatus* instead of *Cyathostomum* species. In line with this higher misassignment rate, the inferred proportions of *Cyathostomum* species in respective mock communities departed from the expected ratio, with a bias towards *C. catinatum* relative to *C. pateratum* (62.3 and 92.7% in lieu of 50 and 75% expected, Supplementary Figure S1).

On day 0, *Cylicostephanus minutus* was the most abundant species in the gastrointestinal nematobiome (13.83% of overall counts, Fig. 3), followed by *C. ashworthi* and *C. nassatus* (5.47 and 5.06% of total counts on that day, respectively, Fig. 3). At the community level, we could not evidence any differences in alpha diversity between larval communities of the sainfoin and control groups, neither at the beginning [average Shannon index difference of 0.18, 95% c. i. = (−0.71; 0.35), $P = 0.47$] nor at the end of the experimental period [average Shannon index difference of 0.16, 95% c. i. = (−0.72; 0.41), $P = 0.53$]. Similarly, PERMANOVA analyses did not evidence any difference between the larval community structures of the 2 groups on day 0 ($P = 0.47$

and $P = 0.2$ with the Bray-Curtis and Jaccard dissimilarity index, respectively) or day 21 ($P = 0.24$ and $P = 0.55$ with the Bray-Curtis and Jaccard dissimilarity index, respectively) of the experiment.

Analysis of species abundance trajectory from day 0 to day 21 was restricted to the top 4 most abundant species (Fig. 4). *Cylicostephanus longibursatus* contributed significantly fewer individuals than the other 3 species (differential 4th-root transformed count of 2.74 ± 1.27 relative to *C. ashworthi*, $P = 0.03$). No significant variation relative to the experimental diet was found for the top 4 most abundant species ($P = 0.59$ – 0.96 , Fig. 4).

Sainfoin effect on the ERP and ivermectin concentration in plasma following oral ivermectin treatment

Eggs reappeared 42 days after IVM treatment in the feces of both experimental groups (1 horse in each group). On day 78 (57 days after IVM treatment), sainfoin-fed horses excreted significantly more eggs than their counterparts fed with the control diet ($P = 0.04$, Fig. 5A). These excretion levels corresponded to FECR of 89.6% (75.4; 95.6) for the sainfoin group. This value was significantly lower than the FECR of the control group on the same day [95.8% (84.1; 98.9), Table 2]. However, the ERP was




Fig. 4. Most abundant species estimated using the metabarcoding approach in horses from the control and sainfoin groups. Evolution of the 4 most abundant species in horses from the control and sainfoin groups between day 0 and day 21. Data are from 12 horses with samples successfully amplified.

above 57 days in both groups. The plasmatic IVM concentration was also measured. The drug profiles followed a typical kinetic curve expected after an oral drench with a maximum IVM concentration (C_{max}) recorded at 24 h in both groups. Importantly, the sainfoin diet negatively impacted the IVM concentration in plasma. These IVM concentrations were twice lower in horses of the sainfoin group when compared with those fed the control diet (Fig. 5B). This was observed at 24 and 48 h after IVM treatment ($P=0.001$ and 0.006 , respectively, Fig. 5B, Table 3). Consequently, the average AUC, reflecting the animal exposure to the IVM drug, was significantly lower in horses fed with the sainfoin diet ($P<0.001$, Table 3) than in the control group.

Discussion

The present study evaluated the anthelmintic activity of a sainfoin diet as a possible alternative mode of control of cyathostomin populations in domestic horses and its interaction with IVM treatment. It provides evidence of a mild effect of dehydrated sainfoin pellets on larval development rate *in vivo* and their negative impact on the IVM pharmacokinetics. It also reports the first application of a nemabiome approach to study the effects of bioactive plants, which supports the lack of a systematic alteration of the cyathostomin larval community structure by sainfoin.

Our results agree with previous observations by Collas *et al.* (2018), who did not report any effect of dehydrated sainfoin pellets (3.6% DM condensed tannins in the diet) on strongyle FEC in horses. However, we observed a slight reduction of larval development rate in horses fed with sainfoin between day 14 and day 21 (9.7% vs 23.4% in the control group), i.e. the end of the distribution of experimental diets). On the contrary, Collas *et al.* (2018) demonstrated that sainfoin inhibited strongyle larvae development *in vitro*. This discrepancy between *in vitro* and *in vivo* studies may indicate that bioactive compounds, presumably condensed tannins (Hoste *et al.*, 2006, 2012), may not have reached sufficient levels in the gastrointestinal lumen of horses to affect the cyathostomin population. In small ruminants, it is suggested that a minimum threshold of 3–5% DM condensed tannins in the diet should be applied to observe an anthelmintic effect (Hoste *et al.*, 2012). Here, the concentration of condensed tannins in the diet was lower (i.e. 2.3% DM). Still, the proportion of dehydrated sainfoin pellets in the diet (i.e. 70% DM, as in Collas *et al.* (2018)) could not have been increased without compromising the proper functioning of the horse's digestive system. We also relied on the same treatment duration as implemented in successful experiments in ruminants (Paolini *et al.*, 2005; Heckendorn *et al.*, 2006). A more extended period of the sainfoin diet may substantially affect larval development.




Fig. 5. Average fecal egg count (A) and plasma ivermectin concentration (B) measured after treatment. Average cyathostomin FEC (A) or IVM concentration in plasma (B) measured after IVM treatment is represented for horses receiving the control (light grey) or sainfoin (dark green) diet. *Statistically significant difference of FEC between the 2 groups at day 78 ($P=0.04$) (A). **Statistically significant difference of average plasma IVM concentration between groups at 24 and 48 h post-treatment ($P<0.01$) (B).

Table 2. Average FECR data (%; arithmetic mean with the 95% confidence interval) in horses receiving the control or sainfoin-enriched diet after IVM treatment

	Day 36 (57 days a. t. ^a)	Day 50 (57 days a. t.)	Day 63 (57 days a. t.)	Day 71 (57 days a. t.)	Day 78 (57 days a. t.)
Control	100 (nd ^a ; nd ^b)	100 (nd ^a ; nd.)	98.33 (85.3; 99.8)	97.50 (88.2; 99.5)	95.83 (84.1; 98.9)
Sainfoin	100 (nd ^a ; nd.)	100 (nd ^a ; nd.)	99.2 (93.3; 99.9)	91.8 (77.8; 96.9)	89.55 (75.4; 95.6)

^aAfter treatment.

^bNot determined.

However, the excess protein intake (i.e. 250% MADC requirements) for more than 3 weeks could have caused health problems in horses.

Our study also evaluated the potential interaction between the tannin-rich diet and an oral IVM treatment, and showed a substantial decrease of IVM concentration in plasma when the drug was given with tannin-rich diet. In a previous experiment conducted on lambs, lower IVM plasma concentration after an oral treatment was reported in animal fed with sainfoin compared to their counterparts fed with a control diet (Gaudin *et al.*, 2016). This lower IVM plasma concentration was associated with the chelation of the molecule by the tannins *in vitro* in a dose-dependent manner (Gaudin *et al.*, 2016). Similarly, the plasma IVM concentration was reduced in horses fed with sainfoin from 24 h onwards in our study. It is generally reported that the T_{max} (time to peak plasma concentration) is 8 h for oral IVM treatment in horses (Gokbulut *et al.*, 2001; Saumell *et al.*, 2017; Vyniarska *et al.*, 2021). The measure at 8 h post-treatment was not practically feasible under our setting. In line with this lower IVM plasma concentration, measured fecal egg reduction was lower in the sainfoin group and egg excretion reappeared earlier in this group. Although the ERP was above 57 days in both groups, this observation is in favour of a decreased treatment efficacy associated with sainfoin.

This effect on IVM plasma concentration may be related to the presence of tannins in the intestinal fluid that could chelate the drug, and reduce the amount of drug available for intestinal absorption (Gaudin *et al.*, 2016). In addition, tannin-rich diets contain flavonoids, like the procyanidins (Quijada *et al.*, 2015), which are known to modulate the expression of ABC transporters (Morris and Zhang, 2006), like the P-glycoproteins (P-gps), pump the drug out of tissues and the organisms, lowering plasma IVM concentration (Bartley *et al.*, 2009; Dupuy *et al.*, 2010). Their inhibition increases IVM efficacy against susceptible *H. contortus*, a trichostrongylid of small ruminants (Kerboeuf *et al.*, 2003; Bartley *et al.*, 2009; Lespine *et al.*, 2012; Peachey *et al.*, 2017). There is evidence that polyphenols can decrease the parasite ATP-binding cassette (ABC) transporter activity or directly affect the parasite (Dupuy *et al.*, 2003; Whitney *et al.*, 2013). In addition, differential expression of these genes was found between IVM-resistant and IVM-susceptible isolates of *H. contortus* (Mate *et al.*, 2022) and cyathostomins (Peachey *et al.*, 2017) or *Parascaris* sp. (Janssen *et al.*, 2015). But our results failed to show a positive interaction between the tannin-rich diet and oral IVM treatment and in our experimental conditions, it is impossible to determine the origin of the low plasma IVM concentration in horse fed tannin-rich diet. The set-up of an equine fermenter able to mimic the hindgut conditions similar to that

Table 3. Pharmacokinetic parameters of ivermectin in plasma of horses receiving the sainfoin-rich or control diet

	Sainfoin	Control
C_{max}^a (ng mL ⁻¹)	3.8 ± 0.8***	6.8 ± 1.7
T_{max}^b (h)	24 ± 0.0 ^{ns}	24 ± 0.0
AUC_{0-t}^c (ng.h mL ⁻¹)	240.4 ± 77.5***	419.8 ± 111.9

The parameters were calculated from 7 time-point concentrations measured in each animal. Values are geometric mean ± s.e.m of 10 animals.

^{ns}Not significant.

***Statistically significant difference of kinetic parameters compared to control ($P \leq 0.001$).

^aMaximum plasma concentration.

^bTime of maximum concentration.

^cAUC from 0 to the last time point.

available for the foregut (Strauch *et al.*, 2017) could help validate this hypothesis in horses. In any case, these observations suggest that horse feed should be monitored in the field while measuring ERP, as variation in plant secondary metabolites like flavonoids may obscure the results or comparison across operations.

Because a slight difference in larval development rate was observed between the two groups, the metabarcoding approach was implemented to quantify the relative abundance of each species, with a specific interest for the *Cycticocycylus* genus. Previous records using a reverse line blot assay found that members of that genus reappeared first following IVM or pyrantel treatment (van Doorn *et al.*, 2014; Kooyman *et al.*, 2016). Here, the sainfoin diet administered had no detectable effects on the cyathostomin larval community structure. This would be in line with the limited effects observed on other traits or with non-specific effects that apply to every species equally. To our knowledge, this is the first attempt to apply the metabarcoding approach to analyse the activity of a plant product in a parasite system. In addition, the use of mock communities of known composition highlighted two strands of improvement for this approach. Suboptimal taxonomic assignment was detected for the members of the *Cyathostomum* genus, with a high rate of uncertainty when attempting to distinguish between *C. catinatum* and *C. pateratum* in mock communities. This limitation adds up to the similarity between *Cylicostephanus calicatus* ITS-2 sequences and that of *Coronocycylus coronatus* worms, and to the presence of cryptic cyathostomin species in the community (Bredtmann *et al.*, 2019; Louro *et al.*, 2021). Additional markers like the mitochondrial cytochrome oxidase I (CO1) barcode may help increase the specificity of this approach. However, its higher rate of evolution provides a within-species resolution that may be more difficult to handle (Ramünke *et al.*, 2018).

This study confirmed the absence of effects of dehydrated sainfoin pellets *in vivo* on cyathostomin FEC and reported a weak impact on cyathostomin larval development. This is despite sainfoin pellets amounting to 70% of the horse diet. Applying a metabarcoding approach with mock communities revealed substantial margins for improvement in the taxonomic assignment. Still, it did not evidence any effect of sainfoin on the cyathostomin larval population structure. In addition, sainfoin significantly reduced IVM plasma concentration in horses leading to an accelerated reappearance of eggs in the feces. Altogether, sainfoin, administered as in this study, does not appear to be able to control cyathostomin in the field.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182022000853>.

Data availability. The raw data files and associated R scripts are available under the following repository: <https://github.com/Joshua-Malsa/Sainfoin-paper.git>.

Acknowledgements. We are grateful to the GenoToul bioinformatics platform (<https://doi.org/10.15454/1.5572369328961167E12>) for providing computing and storage resources.

Author contributions. G. Fleurance, G. Sallé, B. Dumont and J. Malsa conceived and designed the study; G. Fleurance, L. Wimel, C. Dubois, E. Courtot, M. Boisseau, F. Guégnard, D. Serreau and J. Malsa gathered the data; A. Lespine and J-F. Sutra produced and analysed ivermectin plasma concentration data; J. Lluch, G. Annonay conducted the sequencing of cyathostomin ITS-2 amplicon; T. A. Kuzmina, M. Basiaga collected the samples used for the mock community preparation; G. Sallé performed the analyses of the cyathostomin ITS-2 amplicon sequencing data; J. Malsa performed statistical analyses; J. Malsa, G. Fleurance, G. Sallé, E. Courtot, T. A. Kuzmina, M. Basiaga, S. Dhorne-Pollet, N. Mach, A. Lespine drafted the paper; P. Gombault prepared the sainfoin pellets.

Financial support. This work benefited from the financial support of the Institut Français du Cheval et de l'Équitation (IFCE) and Institut Carnot France Future Élevage (F2E). Joshua Malsa is a grateful recipient of a joint Fond Eperon and IFCE fellowship.

Conflict of interest. The authors declare no conflict of interest.

Ethical standards. The experimental procedure received approval from the French Ministry of Research under protocol number APAFIS#26140-2020062216271790v2.

References

- Alvinerie M, Sutra JF, Galtier P, Lifschitz A, Virkel G, Sallovitz J and Lanusse C (1999) Persistence of ivermectin in plasma and faeces following administration of a sustained-release bolus to cattle. *Research in Veterinary Science* **66**, 57–61.
- Andrews S (2010) *FastQC A Quality Control Tool for High Throughput Sequence Data*.
- Bartley DJ, McAllister H, Bartley Y, Dupuy J, Ménez C, Alvinerie M, Jackson F and Lespine A (2009) P-glycoprotein interfering agents potentiate ivermectin susceptibility in ivermectin sensitive and resistant isolates of *Teladorsagia circumcincta* and *Haemonchus contortus*. *Parasitology* **136**, 1081–1088.
- Becher AM, van Doorn DC, Pfister K, Kaplan RM, Reist M and Nielsen MK (2018) Equine parasite control and the role of national legislation – a multinational questionnaire survey. *Veterinary Parasitology* **259**, 6–12.
- Bredtmann CM, Krücken J, Kuzmina T, Louro M, Madeira de Carvalho LM and von Samson-Himmelstjerna G (2019) Nuclear and mitochondrial marker sequences reveal close relationship between *Coronocycylus coronatus* and a potential *Cylicostephanus calicatus* cryptic species complex. *Infection Genetics and Evolution* **75**, 103956.
- Bucknell DG, Gasser RB and Beveridge I (1995) The prevalence and epidemiology of gastrointestinal parasites of horses in Victoria, Australia. *International Journal for Parasitology* **25**, 711–724.
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA and Holmes SP (2016) DADA2: high-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583.
- Coles GC, Bauer C, Borgsteede FH, Geerts S, Klei TR, Taylor MA and Waller PJ (1992) World Association for the Advancement of Veterinary Parasitology (W.A.V.P.) methods for the detection of anthelmintic resistance in nematodes of veterinary importance. *Veterinary Parasitology* **44**, 35–44.
- Coles GC, Jackson F, Pomroy WE, Prichard RK, von Samson-Himmelstjerna G, Silvestre A, Taylor MA and Vercruyse J (2006) The detection of anthelmintic resistance in nematodes of veterinary importance. *Veterinary Parasitology* **136**, 167–185.
- Collas C, Fleurance G, Cabaret J, Martin-Rosset W, Wimel L, Cortet J and Dumont B (2014) How does the suppression of energy supplementation affect herbage intake, performance and parasitism in lactating saddle mares? *Animal: An International Journal of Animal Bioscience* **8**, 1290–1297.
- Collas C, Sallé G, Dumont B, Cabaret J, Cortet J, Martin-Rosset W, Wimel L and Fleurance G (2018) Are sainfoin or protein supplements alternatives to control small strongyle infection in horses? *Animal* **12**, 359–365.
- Collobert-Laugier C, Hoste H, Sevin C and Dorchies P (2002) Prevalence, abundance and site distribution of equine small strongyles in Normandy. *Veterinary Parasitology* **110**, 77–83.

- Dupuy J, Larrieu G, Sutra JF, Lespine A and Alvinerie M (2003) Enhancement of moxidectin bioavailability in lamb by a natural flavonoid: quercetin. *Veterinary Parasitology* **112**, 337–347.
- Dupuy J, Alvinerie M, Ménez C and Lespine A (2010) Interaction of anthelmintic drugs with P-glycoprotein in recombinant LLC-PK1-mdr1a cells. *Chemo-Biological Interactions* **186**, 280–286.
- Gasser RB, Chilton NB, Hoste H and Beveridge I (1993) Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. *Nucleic Acids Research* **21**, 2525–2526.
- Gaudin E, Simon M, Quijada J, Schelcher F, Sutra J-F, Lespine A and Hoste H (2016) Efficacy of sainfoin (*Onobrychis viciifolia*) pellets against multi resistant *Haemonchus contortus* and interaction with oral ivermectin: implications for on-farm control. *Veterinary Parasitology* **227**, 122–129.
- Giles CJ, Urquhart KA and Longstaffe JA (1985) Larval cyathostomiasis (immature trichonema-induced enteropathy): a report of 15 clinical cases. *Equine Veterinary Journal* **17**, 196–201.
- Gokbulut C, Nolan AM and Mckellar QA (2001) Plasma pharmacokinetics and faecal excretion of ivermectin, doramectin and moxidectin following oral administration in horses. *Equine Veterinary Journal* **33**, 494–498.
- Grabber JH, Zeller WE and Mueller-Harvey I (2013) Acetone enhances the direct analysis of procyanidin- and prodelphinidin-based condensed tannins in lotus species by the butanol-HCl-iron assay. *Journal of Agricultural and Food Chemistry* **61**, 2669–2678.
- Halvarsson P and Höglund J (2021) Sheep nemabiome diversity and its response to anthelmintic treatment in Swedish sheep herds. *Parasites & Vectors* **14**, 114.
- Heckendorn F, Häring DA, Maurer V, Zinsstag J, Langhans W and Hertzberg H (2006) Effect of sainfoin (*Onobrychis viciifolia*) silage and hay on established populations of *Haemonchus contortus* and *Cooperia curticei* in lambs. *Veterinary Parasitology* **142**, 293–300.
- Heckendorn F, Häring DA, Maurer V, Senn M and Hertzberg H (2007) Individual administration of three tanniferous forage plants to lambs artificially infected with *Haemonchus contortus* and *Cooperia curticei*. *Veterinary Parasitology* **146**, 123–134.
- Hojsgaard S, Halekoh U and Yan J (2006) The R Package geePack for generalized estimating equations. *Journal of Statistical Software* **15**, 1–11.
- Hoste H, Jackson F, Athanasiadou S, Thamsborg SM and Hoskin SO (2006) The effects of tannin-rich plants on parasitic nematodes in ruminants. *Trends in Parasitology* **22**, 253–261.
- Hoste H, Martinez-Ortiz-De-Montellano C, Manolaraki F, Brunet S, Ojeda-Robertos N, Fourquaux I, Torres-Acosta JFJ and Sandoval-Castro CA (2012) Direct and indirect effects of bioactive tannin-rich tropical and temperate legumes against nematode infections. *Veterinary Parasitology* **186**, 18–27.
- Hoste H, Torres-Acosta JFJ, Sandoval-Castro CA, Mueller-Harvey I, Sotiraki S, Louvandini H, Thamsborg SM and Terrill TH (2015) Tannin containing legumes as a model for nutraceuticals against digestive parasites in livestock. *Veterinary Parasitology* **212**, 5–17.
- Hung G-C, Chilton NB, Beveridge I and Gasser RB (2000) A molecular systematic framework for equine strongyles based on ribosomal DNA sequence. *International Journal for Parasitology* **30**, 95–103.
- INRA (2015) *Equine Nutrition: INRA Nutrient Requirements, Recommended Allowances and Feed Tables*. Wageningen, the Netherlands: Wageningen Academic Publisher. Available at doi:10.3920/978-90-8686-855-1.
- Janssen IJ, Krücken J, Demeler J and von Samson-Himmelstjerna G (2015) Transgenically expressed Parascaris P-glycoprotein-11 can modulate ivermectin susceptibility in *Caenorhabditis elegans*. *International Journal for Parasitology: Drugs and Drug Resistance* **5**, 44–47.
- Kaplan RM (2002) Anthelmintic resistance in nematodes of horses. *Veterinary Research* **33**, 491–507.
- Kaplan RM (2004) Drug resistance in nematodes of veterinary importance: a status report. *Trends in Parasitology* **20**, 477–481.
- Kassambara A (2021) *rstatix: Pipe-Friendly Framework for Basic Statistical Tests*. Kerboeuf D, Blackhall W, Kaminsky R and von Samson-Himmelstjerna G (2003) P-glycoprotein in helminths: function and perspectives for anthelmintic treatment and reversal of resistance. *International Journal of Antimicrobial Agents* **22**, 332–346.
- Kooyman FNJ, van Doorn DCK, Geurden T, Mughini-Gras L, Ploeger HW and Wagenaar JA (2016) Species composition of larvae cultured after anthelmintic treatment indicates reduced moxidectin susceptibility of immature *Cylicocyclus* species in horses. *Veterinary Parasitology* **227**, 77–84.
- Kuzmina TA, Kharchenko VA, Starovir AI and Dvojnjos GM (2005) Analysis of the strongylid nematodes (Nematoda: Strongylidae) community after deworming of brood horses in Ukraine. *Veterinary Parasitology* **131**, 283–290.
- Kuzmina TA, Lyons ET, Tolliver SC, Dzeverin II and Kharchenko VA (2012) Fecundity of various species of strongylids (Nematoda: Strongylidae) – parasites of domestic horses. *Parasitology Research* **111**, 2265–2271.
- Lespine A, Alvinerie M, Vercruyse J, Prichard RK and Geldhof P (2008) ABC transporter modulation: a strategy to enhance the activity of macrocyclic lactone anthelmintics. *Trends in Parasitology* **24**, 293–298.
- Lespine A, Ménez C, Bourguinat C and Prichard RK (2012) P-glycoproteins and other multidrug resistance transporters in the pharmacology of anthelmintics: prospects for reversing transport-dependent anthelmintic resistance. *International Journal for Parasitology: Drugs and Drug Resistance* **2**, 58–75.
- Lichtenfels JR, Kharchenko VA and Dvojnjos GM (2008) Illustrated identification keys to strongylid parasites (Strongylidae: Nematoda) of horses, zebras and asses (Equidae). *Veterinary Parasitology* **156**, 4–161.
- Louro M, Kuzmina TA, Bredtmann CM, Diekmann I, de Carvalho LMM, von Samson-Himmelstjerna G and Krücken J (2021) Genetic variability, cryptic species and phylogenetic relationship of six *Cyathostomin* species based on mitochondrial and nuclear sequences. *Scientific Reports* **11**, 8245.
- Love S (2003) Treatment and prevention of intestinal parasite-associated disease. *Veterinary Clinics of North America: Equine Practice* **19**, 791–806.
- Love S and Duncan JL (1992) The development of naturally acquired cyathostome infection in ponies. *Veterinary Parasitology* **44**, 127–142.
- Love S, Murphy D and Mellor D (1999) Pathogenicity of cyathostome infection. *Veterinary Parasitology* **85**, 113–121; discussion 121–122, 215–225.
- Lumaret J-P, Errouissi F, Floate K, Römbke J and Wardhaugh K (2012) A review on the toxicity and non-target effects of macrocyclic lactones in terrestrial and aquatic environments. *Current Pharmaceutical Biotechnology* **13**, 1004–1060.
- Lyons ET, Drudge JH and Tolliver SC (2000) Larval cyathostomiasis. *Veterinary Clinics of North America: Equine Practice* **16**, 501–513.
- Lyons E, Tolliver S, Collins S, Ionita M, Kuzmina T and Rossano M (2011a) Field tests demonstrating reduced activity of ivermectin and moxidectin against small strongyles in horses on 14 farms in Central Kentucky in 2007–2009. *Parasitology Research* **108**, 355–360.
- Lyons ET, Tolliver SC and Collins SS (2011b) Reduced activity of moxidectin and ivermectin on small strongyles in young horses on a farm (BC) in Central Kentucky in two field tests with notes on variable counts of eggs per gram of feces (EPGs). *Parasitology Research* **108**, 1315–1319.
- Manolaraki F, Sotiraki S, Stefanakis A, Skampardonis V, Volanis M and Hoste H (2010) Anthelmintic activity of some Mediterranean browse plants against parasitic nematodes. *Parasitology* **137**, 685–696.
- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *European Molecular Biology Network* **17**, 10–12.
- Mate L, Ballent M, Cantón C, Lanusse C, Ceballos L, Alvarez L LI and Liron JP (2022) ABC-transporter gene expression in ivermectin-susceptible and resistant *Haemonchus contortus* isolates. *Veterinary Parasitology* **302**, 109647.
- Matthews JB (2014) Anthelmintic resistance in equine nematodes. *International Journal for Parasitology: Drugs and Drug Resistance* **4**, 310–315.
- McMurdie PJ and Holmes S (2013) Phyloseq: an R Package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* **8**, e61217.
- Mitchell CJ, O'Sullivan CM, Pinloche E, Wilkinson T, Morphey RM and McEwan NR (2019) Using next-generation sequencing to determine diversity of horse intestinal worms: identifying the equine 'nemabiome'. *Journal of Equine Science* **30**, 1–5.
- Morris ME and Zhang S (2006) Flavonoid–drug interactions: effects of flavonoids on ABC transporters. *Life Sciences* **78**, 2116–2130.
- Murali A, Bhargava A and Wright ES (2018) IDTAXA: a novel approach for accurate taxonomic classification of microbiome sequences. *Microbiome* **6**, 140.
- Nielsen MK, Banahan M and Kaplan RM (2020) Importation of macrocyclic lactone resistant cyathostomins on a US thoroughbred farm. *International Journal for Parasitology: Drugs and Drug Resistance* **14**, 99–104.
- Nielsen MK, von Samson-Himmelstjerna G, Kuzmina TA, van Doorn DCK, Meana A, Rehbein S, Elliott T and Reinemeyer CR (2022) World association for the advancement of veterinary parasitology (WAAVP): third edition of guideline for evaluating the efficacy of equine anthelmintics. *Veterinary Parasitology* **303**, 109676.
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin P, O'Hara B, Simpson G, Solymos P, Stevens H and Wagner H (2015) Vegan: community ecology package. *R Package Version 2.2-1* **2**, 1–2.
- Osterman Lind E, Eysker M, Nilsson O, Uggla A and Höglund J (2003) Expulsion of small strongyle nematodes (*Cyathostomin* spp) following

- deworming of horses on a stud farm in Sweden. *Veterinary Parasitology* 115, 289–299.
- Osterman Lind E, Kuzmina T, Uggla A, Waller PJ and Höglund J (2007) A field study on the effect of some anthelmintics on cyathostomins of horses in Sweden. *Veterinary Research Communications* 31, 53–65.
- Paolini V, Dorchie P and Hoste H (2003) Effects of sainfoin hay on gastrointestinal nematode infections in goats. *The Veterinary Record* 152, 600–601.
- Paolini V, Fouraste I and Hoste H (2004) *In vitro* effects of three woody plant and sainfoin extracts on 3rd-stage larvae and adult worms of three gastrointestinal nematodes. *Parasitology* 129, 69–77.
- Paolini V, De La Farge F, Prevot F, Dorchie P and Hoste H (2005) Effects of the repeated distribution of sainfoin hay on the resistance and the resilience of goats naturally infected with gastrointestinal nematodes. *Veterinary Parasitology* 127, 277–283.
- Peachey LE, Pinchbeck GL, Matthews JB, Burden FA, Lespine A, von Samson-Himmelstjerna G, Krücken J and Hodgkinson JE (2017) P-glycoproteins play a role in ivermectin resistance in cyathostomins. *International Journal for Parasitology: Drugs and Drug Resistance* 7, 388–398.
- Peña-Espinoza M, Valente AH, Thamsborg SM, Simonsen HT, Boas U, Enemark HL, López-Muñoz R and Williams AR (2018) Antiparasitic activity of chicory (*Cichorium intybus*) and its natural bioactive compounds in livestock: a review. *Parasites & Vectors* 11, 475.
- Peregrine AS, Molento MB, Kaplan RM and Nielsen MK (2014) Anthelmintic resistance in important parasites of horses: does it really matter? *Veterinary Parasitology* 201, 1–8.
- Pinheiro J, Bates D, DebRoy S and Sarkar D (2013) Nlme: linear and nonlinear mixed effects models. *R Package Version 3.1-110* 3, 1–113.
- Poissant J, Gavriluc S, Bellow J, Redman EM, Avramenko RW, Robinson D, Workentine ML, Shury TK, Jenkins EJ, McLoughlin PD, Nielsen MK and Gilleard JS (2021) A repeatable and quantitative DNA metabarcoding assay to characterize mixed strongyle infections in horses. *International Journal for Parasitology* 51, 183–192.
- Queiroz C, Levy M, Avramenko R, Redman E, Kearns K, Swain L, Silas H, Uehlinger F and Gilleard JS (2020) The use of ITS-2 rDNA nemabiome metabarcoding to enhance anthelmintic resistance diagnosis and surveillance of ovine gastrointestinal nematodes. *International Journal for Parasitology: Drugs and Drug Resistance* 14, 105–117.
- Quijada J, Fryganas C, Ropiak HM, Ramsay A, Mueller-Harvey I and Hoste H (2015) Anthelmintic activities against *Haemonchus contortus* or *Trichostrongylus colubriformis* from small ruminants are influenced by structural features of condensed tannins. *Journal of Agricultural and Food Chemistry* 63, 6346–6354.
- Ramünke S, de Almeida Borges F, von Son-de Fernex E, von Samson-Himmelstjerna G and Krücken J (2018) Molecular marker sequences of cattle *Cooperia* species identify *Cooperia spatulata* as a morphotype of *Cooperia punctata*. *PLoS One* 13, e0200390.
- Raynaud J-P, William G and Brunault G (1970) Etude de l'efficacité d'une technique de coproscopie quantitative pour le diagnostic de routine et le contrôle des infestations parasitaires des bovins, ovins, équins et porcins. *Annales de Parasitologie Humaine et Comparée* 45, 321–342.
- Reinemeyer CR, Smith SA, Gabel AA and Herd RP (1984) The prevalence and intensity of internal parasites of horses in the U.S.A. *Veterinary Parasitology* 15, 75–83.
- Roberts FHS and O'Sullivan PJ (1950) Methods for egg counts and larval cultures for strongyles infesting the gastro-intestinal tract of cattle. *Australian Journal of Agricultural Research* 1, 99–102.
- Rochfort S, Parker AJ and Dunshea FR (2008) Plant bioactives for ruminant health and productivity. *Phytochemistry* 69, 299–322.
- R Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Sallé G, Cortet J, Bois I, Dubès C, Guyot-Sionest Q, Larrieu C, Landrin V, Majorel G, Wittreck S, Woringe E, Couroucé A, Guillot J, Jacquet P, Guégnard F, Blanchard A and Leblond A (2017) Risk factor analysis of equine strongyle resistance to anthelmintics. *International Journal for Parasitology: Drugs and Drug Resistance* 7, 407–415.
- Sallé G, Guillot J, Tapprest J, Foucher N, Sevin C and Laugier C (2020) Compilation of 29 years of postmortem examinations identifies major shifts in equine parasite prevalence from 2000 onwards. *International Journal for Parasitology* 50, 125–132.
- Sambrook J and Russell DW (2006) Purification of nucleic acids by extraction with phenol:chloroform. *CSH Protocols* 2006, pdb.prot4455.
- Sandoval-Castro CA, Torres-Acosta JFJ, Hoste H, Salem AZM and Chan-Pérez JI (2012) Using plant bioactive materials to control gastrointestinal tract helminths in livestock. *Animal Feed Science and Technology* 176, 192–201.
- Saumell C, Lifschitz A, Baroni R, Fusé L, Bistoletti M, Sagües F, Bruno S, Alvarez G, Lanusse C and Alvarez L (2017) The route of administration drastically affects ivermectin activity against small strongyles in horses. *Veterinary Parasitology* 236, 62–67.
- Stratford CH, Lester HE, Morgan ER, Pickles KJ, Relf V, McGorum BC and Matthews JB (2014) A questionnaire study of equine gastrointestinal parasite control in Scotland. *Equine Veterinary Journal* 46, 25–31.
- Strauch S, Wichert B, Greef JM, Hillegeist D, Zeyner A and Liesegang A (2017) Evaluation of an *in vitro* system to simulate equine foregut digestion and the influence of acidity on protein and fructan degradation in the horse's stomach. *Journal of Animal Physiology and Animal Nutrition* 101, 51–58.
- Tombak KJ, Hansen CB, Kinsella JM, Pansu J, Pringle RM and Rubenstein DI (2021) The gastrointestinal nematodes of plains and Grevy's zebras: phylogenetic relationships and host specificity. *International Journal for Parasitology: Parasites and Wildlife* 16, 228–235.
- Traversa D, Castagna G, von Samson-Himmelstjerna G, Meloni S, Bartolini R, Geurden T, Pearce MC, Woringe E, Besognet B, Milillo P and D'Espois M (2012) Efficacy of major anthelmintics against horse cyathostomins in France. *Veterinary Parasitology* 188, 294–300.
- Tzelos T, Morgan ER, Easton S, Hodgkinson JE and Matthews JB (2019) A survey of the level of horse owner uptake of evidence-based anthelmintic treatment protocols for equine helminth control in the UK. *Veterinary Parasitology* 274, 108926.
- van Doorn DCK, Ploeger HW, Eysker M, Geurden T, Wagenaar JA and Kooyman FNJ (2014) *Cylicocyclus* species predominate during shortened egg reappearance period in horses after treatment with ivermectin and moxidectin. *Veterinary Parasitology* 206, 246–252.
- Verdú JR, Lobo JM, Sánchez-Piñero F, Gallego B, Numa C, Lumaret J-P, Cortez V, Ortiz AJ, Tonelli M, García-Teba JP, Rey A, Rodríguez A and Durán J (2018) Ivermectin residues disrupt dung beetle diversity, soil properties and ecosystem functioning: an interdisciplinary field study. *Science of The Total Environment* 618, 219–228.
- Vyniarska A, Ziolkowski H, Madej-Śmiechowska H and Jaroszewski JJ (2021) The pharmacokinetics and antiparasitic activity of ivermectin in Hutsul and Toric horses. *Journal of Veterinary Pharmacology and Therapeutics* 44, 11–17.
- Wang C, Paul M, Isler T and Furrer R (2022) *eggCounts: Hierarchical Modelling of Faecal Egg Counts*.
- Whitney TR, Wildeus S and Zajac AM (2013) The use of redberry juniper (*Juniperus pinchotii*) to reduce *Haemonchus contortus* fecal egg counts and increase ivermectin efficacy. *Veterinary Parasitology* 197, 182–188.
- Workentine ML, Chen R, Zhu S, Gavriluc S, Shaw N, de Rijke J, Redman EM, Avramenko RW, Wit J, Poissant J and Gilleard JS (2020) A database for ITS2 sequences from nematodes. *BMC Genetics* 21, 74.
- Wright ES (2016) Using DECIPHER v2.0 to analyze big biological sequence data in R. *The R Journal* 8, 352–359.