



**HAL**  
open science

## Characterization of sodium relaxation in food: a mandatory step to reach quantitative sodium images

Sylvie Clerjon, Guilhem Pagès, El Sabbagh, Amidou Traoré, J.-M. Bonny

### ► To cite this version:

Sylvie Clerjon, Guilhem Pagès, El Sabbagh, Amidou Traoré, J.-M. Bonny. Characterization of sodium relaxation in food: a mandatory step to reach quantitative sodium images. 15. International Conference on the Applications of Magnetic Resonance in Food Science, Jun 2022, Aarhus, Denmark. hal-03757144

**HAL Id: hal-03757144**

**<https://hal.inrae.fr/hal-03757144>**

Submitted on 22 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Characterization of sodium relaxation in food: a mandatory step to reach quantitative sodium images


Clerjon, S., Pagès, G., El Sabbagh, N., Traoré, A., Bonny, JM.

INRAE, UR QuaPA, F-63122 St Genes Champanelle, France  
INRAE, PROBE research infrastructure, AgroResonance facility, F-63122 St Genes Champanelle, France

Local quantification of  $^{23}\text{Na}$  is critical to (i) understand the relations between salt distribution (and relaxation) and sensory properties and (ii) construct mathematical models of the salting processes. The challenge of quantitative sodium MRI deals with (i) the nuclei low gyromagnetic ratio, (ii) the quadrupolar interactions, (iii) the short  $T_2$  relaxation times and (iv) similarly to other nuclei,  $B_0$  and  $B_1^+$  inhomogeneities. Sodium quantification usually assumes that a single population is present in a voxel and that 3/5 of its population is invisible due to short  $T_2$  relaxation times. This hypothesis is true if all the sodium nuclei experienced, at the NMR timescale, quadrupolar interactions [1]. However, even considering this invisibility factor, significant errors may persist [2]. That is why, we describe here an original approach to check that all sodium presents weak quadrupolar interactions, i.e. type c [1]. When valid, this information was used to map the sodium content.

Filets were salted on the upper side with 6% in weight of dry salt during 6 hours at  $8^\circ\text{C}$ . Then, they underwent 6 hours at  $24^\circ\text{C}$  (smoking) and were stored at  $5^\circ\text{C}$ . A 16 g sample and an entire filet were used for the relaxometry and imaging experiments, respectively. Relaxometry experiments were performed on a Bruker 9.4 T magnet with a 30 mm volumetric insert. A double quantum filter experiment (DQF) was first acquired. Then, a CPMG ( $TE=175\ \mu\text{s}$ , 256 echoes,  $TR=500\ \text{ms}$ ) was recorded and the signal decay was adjusted using a biexponential model. Imaging experiments were performed on a horizontal Biospec 4.7 T system equipped with a DOTY 20 cm  $^1\text{H}/^{23}\text{Na}$  quadrature-polarized coil. Sodium acquisitions were achieved with a CSI ( $TR=100\ \text{ms}$ , voxel volume =  $3 \times 2 \times 8\ \text{mm}^3$ , acquisition time = 1h). The FIDs were fitted voxelwise with a single Lorentzian lineshape for estimating the relaxation-corrected amplitudes and correcting  $B_0$  inhomogeneities. Emission RF-field variations were not corrected as the  $B_1^+$  maps showed weak variations.

A DQF signal was detected. Moreover, the CPMG decay showed a biexponential behavior with relative amplitudes of 2/5 and 3/5 for the slow and fast components, respectively. Both demonstrated that all the sodium was of type c, the same type than gelatin (ref. tubes) [3].  $^{23}\text{Na}$  CSI maps showed that at 24h, salt is not yet homogenized (Figure). The sodium profile exhibited the sodium gradient concentration in the thick part of the filet. The sodium content matched with sodium content measured by ion chromatography (black lines on the profile).



Surprisingly, analysis on other foods, e.g. carrot and pasta cooked in salted water, exhibited a different relaxation behavior: slow and fast fractions ratio differing from 2/5 and 3/5. These results suggest the presence of different sodium pools and may explain the errors previously observed.

Funded by the European FEAMP project InnoSalt driven by the CITPPMP (FEA470018FA1000012 2019-2022)

## References

1. Rooney, W. D. and C. S. Springer (1991). *Nmr in Biomedicine* 4(5).
2. Veliyulin, E. and I. G. Aursand (2007). *Journal of the Science of Food and Agriculture* 87(14).
3. Andrasko, J. (1974). *Journal of Magnetic Resonance* (1969) 16(3).