



**HAL**  
open science

# Nitrate Transport and Retention in Western European Catchments Are Shaped by Hydroclimate and Subsurface Properties

Sophie Ehrhardt, Pia Ebeling, Rémi Dupas, Rohini Kumar, Jan Fleckenstein, Andreas Musolff

## ► To cite this version:

Sophie Ehrhardt, Pia Ebeling, Rémi Dupas, Rohini Kumar, Jan Fleckenstein, et al.. Nitrate Transport and Retention in Western European Catchments Are Shaped by Hydroclimate and Subsurface Properties. *Water Resources Research*, 2021, 57 (10), <10.1029/2020WR029469>. <hal-03758864>

**HAL Id: hal-03758864**

**<https://hal.inrae.fr/hal-03758864v1>**

Submitted on 30 Mar 2026

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



# Nitrate Transport and Retention in Western European Catchments Are Shaped by Hydroclimate and Subsurface Properties

### Key Points:

- Time lags of nitrogen transport in Western European catchments were 5 years on average and varied in space with hydroclimatic conditions
- Large parts of the diffuse N input was retained in the catchments with differences explained by subsurface properties and specific discharge
- Biogeochemical legacy likely exceeded hydrologic legacy in most of the 238 analyzed catchments

Sophie Ehrhardt<sup>1</sup> , Pia Ebeling<sup>1</sup> , Rémi Dupas<sup>2</sup> , Rohini Kumar<sup>3</sup> , Jan H. Fleckenstein<sup>1,4</sup> , and Andreas Musolff<sup>1</sup> 

<sup>1</sup>Department of Hydrogeology, Helmholtz-Centre for Environmental Research, Leipzig, Germany, <sup>2</sup>UMR SAS, INRAE, Institute Agro, Rennes, France, <sup>3</sup>Department of Computational Hydro Systems, Helmholtz-Centre for Environmental Research, Leipzig, Germany, <sup>4</sup>Bayreuth Center of Ecology and Environmental Research, University of Bayreuth, Bayreuth, Germany

### Supporting Information:

Supporting Information may be found in the online version of this article.

### Correspondence to:

S. Ehrhardt,  
sophie.ehrhardt@ufz.de

### Citation:

Ehrhardt, S., Ebeling, P., Dupas, R., Kumar, R., Fleckenstein, J. H., & Musolff, A. (2021). Nitrate transport and retention in Western European catchments are shaped by hydroclimate and subsurface properties. *Water Resources Research*, 57, e2020WR029469. <https://doi.org/10.1029/2020WR029469>

Received 15 DEC 2020

Accepted 23 SEP 2021

**Abstract** Excess nitrogen (N) from anthropogenic sources deteriorates freshwater resources. Actions taken to reduce N inputs to the biosphere often show limited or only delayed effects in receiving surface waters hinting at large legacy N stores built up in the catchments' soils and groundwater. Here, we quantify transport and retention of N in 238 Western European catchments by analyzing a unique data set of long-term N input and output time series. We find that half of the catchments exhibited transport times with an average riverine concentration peak arriving 5 years after application. Longer transport times were evident in catchments with high potential evapotranspiration and low precipitation seasonality. On average the catchments retained 72% (Interquartile Range: 18%) of the N from diffuse sources with retention efficiency being specifically high in catchments with low discharge and thick, unconsolidated aquifers. The estimated transport time scales do not explain the observed N retention, suggesting a dominant role of biogeochemical legacy in the catchments' soils rather than a legacy store in the groundwater. Future water quality management should account for the accumulated biogeochemical N legacy by fostering denitrifying conditions or soil N recycling to avoid long-term leaching and water quality deteriorations for decades to come.

**Plain Language Summary** Despite different regulations that limit anthropogenic nitrate input to the biosphere, there is in many cases no or only delayed improvement in groundwater or surface water contamination. We assessed long-term data covering nitrogen in- and output for Western-European catchments to quantify (a) the needed transport time until reappearance in the river and (b) the quantity of reappeared nitrate. The excess N transport through the Western European catchments had an average time lag of 5 years; and the variation in this transport time was mainly controlled by hydrological parameters with high seasonality in precipitation favoring faster transports. Furthermore large parts of the nitrate was retained in the catchments, with differences explained by subsurface characteristics such that thick and unconsolidated material favored retention either by holding nitrate in the soil or by supporting a bacterial process that released nitrate to the atmosphere. We hypothesized that most of the retained nitrate is accumulated in the soil. This huge pool has on the one hand the potential of being recycled and on the other hand the danger of being leached down slowly, which would constitute a future long-lasting contamination source for groundwater and surface waters.

## 1. Introduction

Nitrogen (N) can be a limiting nutrient in terrestrial, freshwater, and marine ecosystems (Webster et al., 2003). However, the N cycling in these ecosystems is modified and disturbed by humans through inputs from atmospheric deposition, agricultural fertilizers and wastewater. High N inputs especially in economically developed countries have led to increased riverine nitrogen fluxes, causing ecological degradation in aquatic systems and posing a threat to drinking water safety (Dupas et al., 2016; Sebilo et al., 2013; Wassenaar, 1995). Diffuse agricultural sources (mineral fertilizer and manure) constitute most of the N emissions into waters in European countries (Bouraoui & Grizzetti, 2011; Dupas et al., 2013).

© 2021. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Several regulations at federal, national or international levels have been implemented, for example, the EU Nitrate Directive (CEC, 1991) or the Clean Water Act (EPA, 1972) in the US—aiming particularly at reducing N inputs to the terrestrial system. While these measures have proven successful in selected areas (e.g., Kroonvang et al., 2008), many catchments have seen no or only little improvement in water quality (Bouraoui & Grizzetti, 2011; Meals et al., 2010; Vero et al., 2017). The mismatch between implemented measures such as reductions in the nutrient inputs and the observed water quality in streams can be related to transport processes and retention within the catchments. The latter is closely connected to a legacy accumulation of N (e.g., Thomas & Abbott, 2018; Van Meter & Basu, 2015; Wang & Burke, 2017)—a buildup of large N stores in the catchment that are not or only slowly exported. This legacy acts as long-term memory of catchments and has been hypothesized to buffer stream concentration variability (Basu et al., 2010).


N legacies can be attributed to two major components: the biogeochemical and the hydrologic N storage. The first one is related to biogeochemical transformation processes of N in the unsaturated (vadose) zone, often leading to a large buildup of a persistent organic N pool in the soil matrix and only slowly converting to mobile nitrate ( $\text{NO}_3^-$ ; Van Meter & Basu, 2017). Hydrologic legacy describes the pool of dissolved N in the groundwater and unsaturated zone, subjected to very slow transport processes (Van Meter & Basu, 2015). This transport is controlled by the travel time, that is, the time rainfall needs to travel through a catchment (Kirchner et al., 2000). The diversity of subsurface flow paths in a catchment creates a distribution of travel times (Kirchner et al., 2000) varying from days to decades (e.g., Howden et al., 2011; Jasechko et al., 2016; McMahon et al., 2006; Sebiló et al., 2013) also integrating information on timing, amount, storage and mixing of water and thus solutes (Heidbüchel et al., 2020). Therefore, long travel times and a resulting temporary storage of reactive N in the soil, unsaturated zone and in groundwater (Ascott et al., 2017; Ehrhardt et al., 2019; Kumar et al., 2020), can create similar time lags as the biogeochemical legacy of N stored in the soil N pool (Bingham & Cotrufo, 2016; Bouwman et al., 2013; Sebiló et al., 2013). Due to the high complexity of hydrological and biogeochemical processes in catchments, a good understanding of the share of the two different legacy storages and the fate of N remains challenging.

Data-based approaches that jointly characterize N transport timescales and retention under different land-use and management practices can help better understanding the N trajectories for reactive N dynamics at a catchment scale (e.g., Ehrhardt et al., 2019; Van Meter & Basu, 2015). More specifically, comparing quantity and temporal patterns of diffuse N input and riverine N concentrations from catchments allow to estimate N transport time (TT) scales as well as retention (Dupas et al., 2020; Ehrhardt et al., 2019). Retention is defined here as the “missing N” that is either stored in a catchment due to the buildup of multi-annual legacies or permanently removed by denitrification. The estimated TT of N integrates time delays by biogeochemical immobilization and mobilization in the soils and the TT through the vadose zone and groundwater. Although the identification and quantification of legacy effects is of critical importance for predicting future N dynamics and for implementing effective restoration efforts (Bain et al., 2012), only a few studies have investigated retention and TTs simultaneously as availability of long-term data often limits the number of studied catchments (e.g., Dupas et al., 2020; Ehrhardt et al., 2019; Howden et al., 2010; Van Meter et al., 2017, 2018). Here we analyze a large-sample database of 238 Western European catchments with different geophysical and hydro-climatological characteristics and at least 20 years of observations with regards to observed nitrogen (a) TT scales and (b) retention. Furthermore, we connect both to catchment characteristics to discuss their (c) main controlling factors. These results are used to improve the understanding of catchment responses to changes in input and the fate of retained N being associated with different legacy stores and/or denitrification.

## 2. Materials and Methods

### 2.1. Study Area

For data on water quantity and quality, we relied on three national data sets. Water quality data for French catchments are publicly available at <http://naiades.eaufrance.fr/>, while water quantity data are available at <http://hydro.eaufrance.fr/>. For Germany, Musolff (2020) provided a database for water quality and water quantity.



**Figure 1.** Study catchments ( $n = 238$ ) based on the quality criteria with selected catchment characteristics: (a) Elevation (EEA, 2013), (b) Land cover (CLC, 2000), (c) Precipitation seasonality, (d) Lithology (BCR & UNESCO, 2014).

From this joint database we selected catchments where the following conditions were given: riverine  $\text{NO}_3\text{-N}$  concentration observations available for at least 20 years of data with data gaps less than 2 years and the total number of observations being more than 150. Given these criteria, 238 catchments were selected (Figure 1a). The time series covered data between 1971 and 2015 with a median length of 30 years and in total 96,443 measurements for  $\text{NO}_3\text{-N}$ .

By focusing on  $\text{NO}_3\text{-N}$  loads, here we do not account for riverine N export in other mineral ( $\text{NO}_2\text{-N}$ ,  $\text{NH}_4\text{-N}$ ) or organic N forms. In recent studies by Ebeling et al. (2021) and Dupas et al. (2015)  $\text{NO}_3\text{-N}$  was found to be the dominant form of exported N across the study catchments. Moreover, for the German catchments presented here, 71% of samples did not have information on all N species available. Calculating the loads of

exported N for samples where full information was available revealed that across the German catchments, a median of 92.4% of mineral N species was obtained as  $\text{NO}_3\text{-N}$ . And compared to total N (mineral and organic species) a median of 81.4% was in form of  $\text{NO}_3\text{-N}$ . Assuming similar conditions for other catchments, we acknowledge that our analysis based on only  $\text{NO}_3\text{-N}$  may miss 18.6% of loads that is exported by other N-species.

Overall the data selection covered 40% of the total land area of both countries (i.e., around 361,000 km<sup>2</sup>, taking nested catchments into account). The selected catchments encompass contrasting settings in terms of morphology, climate, geological properties, and land use attributes (Tables S1.1 and S1.2 in Supporting Information S1). More than half of the study catchments have a size of less than 1,000 km<sup>2</sup> (max. 62,500 km<sup>2</sup>). The median altitude ranges from 15 to 1,848 m with a median slope of 3°. Climatic settings of the sites reach from Atlantic to Continental climate with aridity indices ranging between 0.4 and 1.5. The median annual precipitation across the sites is around 816 mm, and the estimated base flow index (BFI) ranges from 29% to 97% with a median of 65%.

Most catchments (>90%) are dominated by sandy soils with a median value of the catchment average sand content for the upper 1 m of soil to be around 44.6% (median estimated across all catchments). The bedrock mainly covers fissured and hard rock geology with the latter being predominant in most of the catchments. The geology is characterized by crystalline rocks in the Armorican Massif, the Pyrenees and the Massif Central and in some of the German mountainous catchments; and younger sedimentary rocks in most parts of France and Germany (Allain, 1951; BGR & CGMW, 2005). Quaternary sediments are found in the Northern German Lowlands, the Alpine foothills and north of the Pyrenees (Allain, 1951; BGR & CGMW, 2005).

Regarding land use, 87% of the catchments had at least one-third of their area covered by agriculture that mainly incorporates non-irrigated arable land and pastures (EEA, 2016; Figure 1b). Riverine  $\text{NO}_3\text{-N}$  concentrations in these areas are therefore predominantly impacted by diffuse agricultural N sources (EEA, 2018). The median share of forest cover across the study catchments is 37%. Although the fraction of artificial surfaces was small, the median population density with 92 inhabitants km<sup>-2</sup> in the study catchments is almost three-times the average European value (Worldometers.info, 2020).

## 2.2. Nitrogen Input

The N input was selected as diffuse N stemming from agricultural N surplus, atmospheric deposition, and biological fixation in non-agricultural areas. The N surplus consists of agricultural N input that is in excess of crop and forage exports (also known as land nitrogen budget; de Vries et al., 2011). Here, we relied on two national scale data sets. Agricultural N contribution and atmospheric N deposition for the French catchments were provided by Poisvert et al. (2017). The annual agricultural N surplus for German catchments was provided by Bach and Frede (1998) as well as Häußermann et al. (2019). It consists of two data sets available at a (coarser) state level (NUTS2) for 1950–1999 and at finer county level (NUTS3) for 1995–2015. Both data sets were harmonized to produce a consistent long-term data set. The atmospheric N deposition for German catchments is based on Europe-wide gridded data from a chemical transport model of the Meteorological Synthesizing Center-West (MSC-W) of the European Monitoring and Evaluation Program (EMEP) (Bartnicky & Fagerli, 2006; Bartnicky & Benedictow, 2017).

In agricultural areas, biological fixation was already included as an input in the N budgets. The biologically fixed N fluxes to non-agricultural land use types, for example, to green lands (urban and natural) or forests, for France and Germany were calculated using the European Corine Land Cover data set from the year 2000 (EEA, 2020), which is most representative regarding the water quality time series. Terrestrial biological N mean uptake rates were set for forest (to 16.04 kg N ha<sup>-1</sup> year<sup>-1</sup>; Cleveland et al., 1999), for natural and urban grassland (to 2.7 kg N ha<sup>-1</sup> year<sup>-1</sup>; Cleveland et al., 1999) and other land use (wetlands, water bodies, open space with little or no vegetation to 0.75 kg N ha<sup>-1</sup> year<sup>-1</sup>; Van Meter et al., 2017). A comparison of the two national long-term data sets for diffuse N with a Europe-wide benchmark estimation for 1997–2003 (West et al., 2014) indicated an acceptable offset (see Section S2 in Supporting Information S1 for further information).

Due to the lack of spatially and temporally reliable long-term data on N input by wastewater, we did not consider this point source. For France, Dupas et al. (2015) estimated the contribution from domestic and

industrial point sources to total N loads to be 3% during the period 2005–2009, and we hypothesized that the negligible contribution of point sources also held for Germany. However, we acknowledge a high uncertainty of wastewater inflow that may have been higher in the past (Westphal et al., 2020).

### 2.3. Nitrogen Output as Riverine NO<sub>3</sub>-N Concentrations and Loads

Continuous (and regular) time series of discharge and concentrations are needed for adequately calculating the retention parameters based on the N surplus and in-stream NO<sub>3</sub>-N loads. Gaps in the discharge time series at 30 runoff stations in Germany were therefore filled through the support of simulations from the grid-based distributed mesoscale hydrological model mHM (Kumar et al., 2013; Samaniego et al., 2010). Here, only model simulations resulting in an  $R^2$  greater than 0.6 when compared with the observed discharge were accepted. A piecewise linear regression was utilized to correct for potential biases in the modeled data. These bias-corrected modeled discharge data were finally used to gap-fill the original data to obtain a continuous daily time series. In France, no such national hydrological model existed and therefore, we only included catchments with nearly continuous daily discharge monitoring for which short gaps in the discharge (max. 7 days) were interpolated by a fixed-interval smoothing via a state-space model using the R software package “Baytrends.”

The irregularly sampled, riverine NO<sub>3</sub>-N concentrations were used to estimate daily concentrations by using the software package *Exploration and Graphics for RivErTrends* (EGRET) in the R environment by Hirsch and de Cicco (2019). The applied *Weighted Regressions on Time, Discharge, and Season* (WRTDS) uses a flexible statistical representation for every day of the discharge record and has been proven to provide robust estimates (Hirsch et al., 2010; Van Meter & Basu, 2017). As we focus on changes in concentrations and fluxes independent of inter-annual discharge variability (Hirsch et al., 2010), we used flow-normalized concentrations and fluxes for further analyses. For each catchment median annual flow-normalized NO<sub>3</sub>-N concentrations and annual summed NO<sub>3</sub>-N fluxes were calculated and scaled to the catchment area.

### 2.4. Nitrogen Transport Time

Travel time distributions are commonly derived as the transfer function between rainfall concentration time series and stream concentrations of a conservatively transported solute or water isotope (e.g., Kirchner et al., 2000). We adopted this concept to reactive N transport with the N input as an incoming time series with annual resolution that is assumed to yield the median annual riverine NO<sub>3</sub>-N concentration, when convolved with a fitted distribution. This transport time distribution (TTD) can be based on different theoretical probability distribution functions. To represent the long memory of past inputs, long-tailed distributions are most suitable at catchment scales (Kirchner et al., 2000). Therefore, the N input was convolved using a log-normal distribution (Equation 1; Ehrhardt et al., 2019; Musolff et al., 2017) to find the optimal fit to riverine NO<sub>3</sub>-N concentrations. We alternatively used a gamma distribution (Equation 2; Godsey et al., 2010; Fiori et al., 2009; Kirchner et al., 2000) as a transfer function, and we compared the quality of fit ( $R^2$ ) with both methods.

$$f(t) = \frac{1}{t\sigma\sqrt{2\pi}} \exp\left(-\frac{(\ln t - \mu)^2}{2\sigma^2}\right) \quad (1)$$

$$f(t) = t^{-\alpha} \frac{\varepsilon^{-t/\beta}}{\beta^\alpha \Gamma(\alpha)} \quad (2)$$

The two parameters mu ( $\mu$ ) and sigma ( $\sigma$ ) for the log-normal and shape ( $\alpha$ ) and scale ( $\beta$ ) for the gamma distribution, respectively, were calibrated through optimization based on minimizing the sum of squared errors between the normalized annual diffuse N input and normalized annual median riverine NO<sub>3</sub>-N concentrations. For this purpose we used the Particle Swarm Optimization (using the R package “hydroPSO” by Zambrano-Bigiarini & Rojas, 2013) algorithm in 30 independent runs. We estimated the mode of the selected best fitted TTD (with max.  $R^2$ ) to represent the peak TT and at the same time to resemble the peak N export of the mobile, inorganic N.

### 2.5. Nitrogen Retention and Its Temporal Change

The total cumulative diffuse N input load was compared to the respective riverine NO<sub>3</sub>-N load (assumed as N load) to analyze the N retention in the catchment (Equation 3). The difference between the two is the load being retained in the catchment as biogeochemical legacy, as hydrologic legacy or being removed by denitrification. This mass calculation assumes that catchment delineation based on topography approximates subsurface catchments and that no significant amounts of N are bypassing the surface water gauging station. The cumulative flux differences were calculated based on two approaches: (a) using the annual frames of the overlapping years in in- and outflux, while disregarding time shifts; and (b) applying the derived TTs, to compare the convolved inputs with the corresponding annual exported load.

$$\text{Retention} = 1 - \frac{N_{\text{out}}}{N_{\text{in}}} = 1 - \frac{\sum_{i=ts}^{te} \text{NO}_3 - N_{\text{flux } i}}{\sum_{i=ts}^{te} N_{\text{input } i}} \quad (3)$$

To further characterize the catchment's reaction to N input changes, we compared the median diffuse N input in the 1980s (median year of max. N input: 1986) with the one in the last years of the time series ( $\geq 2010$ ) for a subset of stations ( $n = 120$ ) that sufficiently covered the 1980s and 2010s. The same was done with the exported riverine NO<sub>3</sub>-N loads in the 1980s and the 2010s. To gain robust estimates for the size of difference, we calculated the bootstrapped ( $n = 10,000$ ) median differences between the 1980s and 2010s (for N input and N output) with their corresponding 95% confidence intervals.

### 2.6. Statistical Analysis for Controls in Catchment Response and Retention

We applied a Partial Least Squares Regression (PLSR) to identify the main factors controlling N TTs and N retention in a catchment. PLSR is an established multivariate regression approach to analyze data sets that are strongly correlated among predictors and noisy (Wold et al., 2001). The PLSR model finds the variables (catchment characteristics) that best predict the response variables (TT and retention; Ai et al., 2015). The importance of each predictor for the dependent variable is indicated by the measure Variable Importance in the Projection (VIP). Factors with VIPs larger than 1 are considered to be significantly important for explaining the dependent variable (Ai et al., 2015; Shi et al., 2013). The corresponding regression coefficient is used to explain the direction of influence of each independent variable (Shi et al., 2013). The predictor variables used in this study characterize the topography, land cover, climate, hydrology, lithology, soils, and population density of the studied catchments (Supporting Information S1).

## 3. Results


### 3.1. Nitrogen Transport Time Scales

Using the gamma distribution yielded comparable results to the results for log-normal distribution (both with median  $R^2 = 0.8$ ), but less catchments with an acceptable fit ( $R^2 \geq 0.6$ ) between the convolved annual N inputs and riverine concentrations. Therefore, we only report the results using a log-normal distribution as a transfer function.

In some catchments ( $n = 72$ ) no acceptable fit of TTDs could be obtained. According to a Wilcoxon rank sum test, the variability in annual NO<sub>3</sub>-N concentrations in these catchments (CV: 0.08) is significantly different ( $p \leq 0.01$ ) to the ones in the other catchments (CV: 0.12 with  $n = 166$ ). A low temporal variability in the input or output makes it challenging to derive a reliable transfer function connecting them.

The median mode (peak) of the TTs for the 166 selected catchments with an acceptable fit was 5.4 years (Tables S3 and S4 in Supporting Information S1). Although the mode ranged from 0.2 to 34.1 years, the majority (70%) had a mode TT less than 10 years (Figure 2c). Only a few catchments (10%) showed a mode of at least 20 years, most of them (11/17) located in the Massif Central (Figure 2a).

Although the TT derivation was not mass conform, on average across the study catchments, 75% (75%-percentile) of the N input should have been exported after 18 years (range: 1.4–38.2).



**Figure 2.** (a) Spatial variation of the TT modes in the 166 catchments with an  $R^2 \geq 0.6$ . (b) Spatial variation of the overlapping retention in all of the analyzed catchments ( $n = 238$ ). (c) Histogram of the mode TTs. (d) Histogram of the retention for the overlapping time (beige curve) and the convolved retention (gray curve) with their corresponding medians (dashed lines). (e) Scatter plot of the overlapping retention versus the mode TTs, with the corresponding medians for both measures (dashed lines). Excluding one outlier with negative retention.

### 3.2. Nitrogen Retention

The median N retention of the selected catchments ( $n = 238$ ) was 72% (sd: 16%; Table S3 in Supporting Information S1; Figure 2b), meaning that a large part of N was retained as legacy or denitrified. Despite the wide range (−24%–96%, with one negative outlier Figure 2b and Text S5 in Supporting Information S1), 48% of the catchments had retention between 50% and 75%. A convolution of the N inputs according to the corresponding TT resulted in a slightly lower retention with a median of 70% ( $n = 238$ ; 71% with  $n = 166$ ; Figure 2d).

N retention and TT did not correlate in the study catchments. Almost the same amount of catchments with retention above the median had TTs below and above the median (Figure 2e).

The median diffuse N input in the 1980s was  $62.6 \text{ kg N ha}^{-1} \text{ year}^{-1}$  (Interquartile Range, IQR:  $42.0 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ), decreasing by around 36%, when assuming the bootstrapped difference in medians of  $22.6 \text{ kg N ha}^{-1} \text{ year}^{-1}$  (95% CI:  $20.5\text{--}25.6 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ) in comparison to the 2010s. Diffuse N input in the 2010s was around  $38.4 \text{ kg N ha}^{-1} \text{ year}^{-1}$  (IQR:  $23.1 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ). The median N load in the 1980s was  $12.4 \text{ kg N ha}^{-1} \text{ year}^{-1}$  (IQR:  $6.1 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ) with a bootstrapped difference of medians of  $1.2 \text{ kg N ha}^{-1} \text{ year}^{-1}$  (95% CI:  $0.8\text{--}1.6 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ) to the 2010s (median N load:  $11.2 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ; IQR:  $5.7 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ).

**Table 1**

*Ranked Drivers for Explaining the Variability in Transport Times and N Retention Across the Study Catchments Based on the Partial Least Squares Regression (PLSR) Analysis, Along With the Corresponding Explained Variability ( $R^2$ ), VIPs and Signs (See Also Section S6 in Supporting Information S1 for Details)*

Transport time $R^2 = 0.49$			Retention $R^2 = 0.69$		
Variable	VIP	Sign	Variable	VIP	Sign
Potential evapotranspiration	2.1	+	Depth to bedrock	1.9	+
Precipitation seasonality	2.1	–	Consolidated aquifer material	1.5	–
Coefficient of variation of discharge	1.9	–	Porous aquifer material	1.5	–
Topographic wetness index	1.5	–	Specific discharge	1.5	–
Median winter discharge	1.3	–	Precipitation frequency	1.4	–

*Note.* The sign indicates the direction of influence.

$\text{ha}^{-1} \text{ year}^{-1}$ ). The mismatch between N input and riverine N export decreased from an annual excess of  $50.2 \text{ kg N ha}^{-1}$  in the 1980s to  $27.2 \text{ kg N ha}^{-1}$  in the 2010s, also reflecting a decrease in apparent retention from 80% to 71%.

### 3.3. Controls of Catchment's Response and Retention

The PLSR for predicting the mode TTs in the selected catchments with a good fit ( $R^2 \geq 0.6$ ) explained 49% of the total variance (Table 1). Variables that are connected to the catchment's hydroclimatological characteristics were found to be most important (see also Figure S6.1 in Supporting Information S1). Potential evapotranspiration (PET) was analyzed as the most important variable indicating longer mode TTs with higher PET. The seasonality index of precipitation (P\_SI, see Supporting Information S1 for detailed description) was with an almost same VIP value the second most influential predictor. The higher the mean difference between monthly P averages and the annual average, the shorter the mode TT. The other three most important parameters indicate shorter TT related with (a) higher coefficients of variation of discharge, (b) higher topographic wetness indices (TWI) and (c) higher median winter discharges.

The N retention across the catchments was well predicted by the PLSR ( $R^2 = 0.69$ , excluding one negative outlier, Table 1). Three of the five most important parameters (see also Figure S6.2 in Supporting Information S1) referred to subsurface characteristics, while two predictors were hydrological descriptors. Depth to bedrock (i.e., distance from ground surface to continuous bedrock) was connected to high retention and was the most important predictor. At the same time, a high share of consolidated and porous aquifer materials were associated with low retention, while vice versa catchments with a high share of unconsolidated aquifers (VIP = 1.4, rank 6) favored higher retention. In context of hydrological parameters: high specific discharge and precipitation frequency, both favoring low retention were ones appear as most informative ones associated with the N retention across the study catchments.

## 4. Discussion

The results of a comparison of diffuse N input and riverine N output time series from 238 West European catchments revealed the median peak TTs of 5 years and a median retention of 72%. While differences in estimated TTs among the catchments were mainly explained by hydroclimatic properties, differences in the retention were mainly explained by hydrologic and subsurface properties.

### 4.1. Nitrogen Transport Times and Its Controlling Parameters

The high number of catchments showing a good fit between N input and riverine N export using a log-normal TTD indicate that the applied methodology is appropriate for the analyzed Western European catchments. This also shows that the temporal pattern of annual flow-weighted  $\text{NO}_3\text{-N}$  concentrations observed in the streams is mainly controlled by the pattern of the diffuse N input.

The PLSR that explained 49% of the variability of mode TTs between the catchments, reveals the importance of hydroclimatic variables (via PET, precipitation and discharge variability, winter discharge) and morphology (via TWI), which is partly in line with previous knowledge that stated recharge rate (besides aquifer porosity and thickness) as a major control for mean groundwater travel times (Haitjema, 1995). We note the close connection between hydroclimatic descriptors (e.g., between long-term mean precipitation, PET, discharge; Figure S7 in Supporting Information S1; as established through the Budyko (1974) framework), but only discuss here the ones ranked as most important for TTs according to the PLSR.

Especially regions with highest intra-annual precipitation seasonality (Figure 1c) like in the Armorican Massif and the Alpine foothills showed short TTs with modes shorter than 5 years. Precipitation seasonality, entailing changing wetness conditions, can cause changing aquifer connectivity (Blume & Van Meerveld, 2015; Roa-García & Weiler, 2010), which is known as a major control of  $\text{NO}_3$  export from catchments (Molénat et al., 2008; Ocampo et al., 2006; Wriedt et al., 2007). In terms of hydrological connectivity, Birkel et al. (2015) and Yang et al. (2018) stated that the activation of shallow flow paths during runoff events favors young water ages. Hence, we hypothesize that these high-flow events efficiently export young  $\text{NO}_3$  from the shallow subsurface to the stream and thus lowers N TT scales. High median winter discharge as another VIP, common in the Alpine foothills favoring short TTs, is in line with our hypothesis and the previous findings by Wriedt et al. (2007). The correlation between high TWI values and short TTs for N may be also attributed to a prevalence of N exports by shallow subsurface flow paths in areas with high groundwater levels (e.g., Grabs et al., 2012): lowland catchments, characterized by higher TWI's, show strong seasonal changes of discharging streams and the artificial drainage network (Van der Velde et al., 2009). As these drains favor rapid, shallow subsurface flows, their temporal connection during high-flow events favor short travel times (Van der Velde et al., 2009). As previous studies that indicated no signs of dilution patterns in a concentration-discharge relationship during high-flow conditions across the majority of the analyzed catchments in either France (Moatar et al., 2017) or Germany (Ebeling et al., 2021), we assume rather a mobilization of N from shallow nitrate sources than a bypass of N, for example, by overland flow.

Long N TTs were found in the western Massif Central and south of it where PET was highest among the study catchments and recharge likely low, corroborating Haitjema's (1995) finding for groundwater travel times.

A clear link between TTs for N and hydroclimatic settings make catchment N transport vulnerable to the changing future climate. Based on past observations since the 1960s, the intensity of extreme weather has been predicted to increase in most parts of Europe (EC, 2009). Hydroclimatic projection studies in general suggest generally drier conditions in Atlantic climatic zones in Europe in terms of longer drought durations and lower low-flows under warming climates (Marx et al., 2018; Samaniego et al., 2018). Both extremes, heavy precipitation events and longer droughts, are more likely. According to the discussed influence of precipitation and discharge variability on N dynamics, TTs are supposed to decrease in the future. The pronounced changes in evapotranspiration with increasing temperature (Donnelly et al., 2017) will likely counteract this trend by favoring for longer TTs with more drying conditions. Since the climate is expected to manifest differently within Europe, reliable predictions of future N TTs on regional scales need further research.

Despite a high number of catchments with a good fit using our TT estimations, we acknowledge the inherent uncertainties and limitations of the database as well as of the method itself. With better knowledge on the temporal evolution of waste water inputs and anthropogenic modifications in the catchment hydrology, like damming, more reliable TT estimations and a potentially better explainability among the catchments may have been possible (e.g., Goyette et al., 2019; Powers et al., 2015). Furthermore the method, assuming a constant (static) log-normal TTD, is only supposed to mirror the dominant long-term TT behavior, disregarding known temporal variability of water travel times in catchments (Benettin et al., 2013; Botter et al., 2011; Harman, 2015; Kumar et al., 2020; Van der Velde et al., 2010). Moreover, we estimated TTs from the small fraction of total N inputs that left the catchment as  $\text{NO}_3$ -N (median 28%). We thus do not capture N that may leave the catchment by other mineral or organic N species (see Section 2); and as well we cannot exclude the possible role of long flow pathways that do not return to the rivers within the observation period. Long-term tracer studies using labeled  $^{15}\text{N}$  compounds (e.g., Sebilo et al., 2013) hold promising avenues for a more detailed and hedged evaluation of the fate of N.

#### 4.2. Nitrogen Retention and Controlling Parameters

According to the PLSR, the variability in retention among the catchments was mainly explained by subsurface properties that can be connected to biogeochemical conditions and the specific discharge. This finding was in line with Merz et al. (2009) and Nolan et al. (2002), who stated that spatial differences in  $\text{NO}_3$  retention or contamination, respectively, result from a combination of the geochemical environment and the hydraulic conditions. We argue that the highly-ranked subsurface predictors describe favorable biogeochemical conditions for either permanent removal by denitrification or storage in the soils as biogeochemical legacy.

Areas with a high depth to bedrock and an unconsolidated aquifer (Figure 1d), which showed retention above 75%, were particularly common in the Northern German Lowlands and in the Alpine foothills. This is in line with Ebeling et al. (2021), who attributed areas with large depth to bedrock and unconsolidated (sedimentary) aquifers to natural attenuation or retention processes based on riverine  $\text{NO}_3$ -N concentration-discharge relationships. Unconsolidated deposits in the terrestrial subsurface, like in the Northern German Lowlands, are often associated with iron sulfide minerals (pyrite; Bouwman et al., 2013). The pyrite oxidation acts as electron donor for denitrification under anaerobic conditions (Zhang et al., 2009). For the unconsolidated aquifers in northern Germany, a recent study (Knoll et al., 2020) connected the high denitrification potential to strongly anaerobic redox conditions in the groundwater. Although denitrification permanently removes N from the catchment, it can be a source for  $\text{N}_2\text{O}$ , an important greenhouse gas, being 300-fold more effective in trapping heat than carbon dioxide (Griffis et al., 2017). Lastly, long-term consumption of reactants via denitrification can alter the reduction capacity of the aquifer (Merz et al., 2009), decreasing the catchment's N retention over time.

In contrast to northern Germany, for the unconsolidated sediments in the Alpine foothills different studies (BMU, 2003; Knoll et al., 2020) proposed aerobic subsurface conditions, hindering denitrification. Also in a recent study, Ebeling et al. (2021) found a very limited evidence for denitrification in these (German) areas. Excluding denitrification and long TTs (see Section 4.1), we hypothesize biogeochemical legacy as a likely process of the high retention in the Alpine foothills. In comparison to northern Germany, soils here contain higher degrees of silt and clay. These grain sizes are prone to microaggregate formation and anion sorption, both sequestering organic N in the mineral subsoil for long periods of time (Bingham & Cotrufo, 2016; von Lützwitz et al., 2006). Also mineral N fixed on clays can make a significant contribution to the soil N stock (Allred et al., 2007; Stevenson, 1986). We note that so far there is no clear soil data evidence for a buildup of a soil-legacy in the Alpine foothills comparable to Dupas et al. (2020) findings for the Armorican Massif of to the large scale-evidence presented by Van Meter et al. (2016).

In contrast, areas with a high share of consolidated subsurface materials and a small depth to bedrock, like the Armorican Massif, parts of the Massif Central or the Harz Mountains showed N retention below 75%. In general, denitrification and biogeochemical legacies can only evolve if favorable biogeochemical conditions in soils and groundwater are abundant in the catchment. An important part for denitrification is the contact area and contact time with organic-rich soils (providing electron donors, Bouwman et al., 2013). Due to abundant crystalline rocks, water moves along fissures in the weathered zone (Wyns et al., 2004), while it is dependent on joints and fractures in deeper depth (Wendland et al., 2007). Hence, there is only a limited reactive surface for  $\text{NO}_3$  within the areas dominated by consolidated materials (Wendland et al., 2007). Furthermore, Knoll et al. (2020) showed oxic conditions in consolidated units for Germany that do not allow for denitrification in groundwater.

One hydrological predictor for N retention was the specific discharge. High specific discharges were found in the Armorican Massif, the western part of the Massif Central, in the Harz Mountains and the southern Alpine foothills, were often spatially connected to areas with consolidated subsurface materials and had N retention below 75%. High discharge areas connect to short residence times in the catchment compartments like root zone, aquifer or riparian zone and therefore decreases denitrification efficiency through a reduced contact time (e.g., Howarth et al., 2006; Kumar et al., 2020; Kunkel & Wendland, 2006; Wendland et al., 2007). This assumption is in line with a recent study by Dupas et al. (2020), arguing that higher runoff lowers denitrification. Tesoriero et al. (2017) and Knoll et al. (2020) stated high recharge rates as important predictors for aerobic conditions. Furthermore, high discharge may be driven by a high degree of shallow

flow paths (Birkel et al., 2015; Yang et al., 2018), favoring a fast wash-out of N or an export before immobilization, thus decreasing retention as well. The other hydrologic predictor for retention was the precipitation frequency  $p$ -lambda. According to Botter et al. (2013), a high frequency of flow-producing rainfall events will increase the likelihood of a more persistent discharge regime. In connection to N transport we argue that catchments with a high frequency of runoff events and a more persistent discharge regime are supposed to more permanently export N from soil sources, counteracting the N retention.

With regard to climate change, the increase in European rainfall erosivity is estimated in the range from 10% to 15% until 2050 (Panagos et al., 2015). Especially in southern France and Germany, this may cause soil loss in arable lands up to  $10 \text{ t ha}^{-1} \text{ year}^{-1}$  (Panagos et al., 2015). We argue that such mobilization of soils with high biogeochemical legacy (e.g., Alpine foothills) can contribute to further deterioration of downstream river water quality. Furthermore, increased temperature stress and precipitation variability under future climate is likely to affect plant N uptake and subsequent unintentional  $\text{NO}_3$ -N leaching (e.g., Hatfield et al., 2011; Jabloun et al., 2015; Wang et al., 2015).

### 4.3. Joint Analysis of Nitrogen Transport Times and Retention

The joint analyses of N TT estimations and N retention (Figure 2e) revealed a discrepancy between the two across the studied catchments. The observed rather short TTs indicate that the largest part (75th-percentile) of N input should have been exported after at least 20 years. In contrast, the observed retention indicates that 72% of total N input was not exported. The retention was similarly high (70%) when convolving N input taking into consideration estimated TTs. It is worth mentioning here that these numbers may be lower as we refer to the N exported as  $\text{NO}_3$ -N that though accounts for the majority (81.4% in the German catchments, see Section 2) but not all riverine N loads. The missing relation between TTs and retention as well as the different predictors for both through the PLSR, indicate that hydrologic legacies of N alone could not explain the failure of measures to improve water quality in Western European catchments (e.g., Bouraoui & Grizzetti, 2011), despite decreasing N inputs. We rather assume a dominance of non-hydrologic retention, namely biogeochemical legacy and denitrification.

In line with the above discussed subsurface and hydrological catchment characteristics favoring biogeochemical legacy, and due to the specific conditions required for effective denitrification that are only fulfilled in a few areas, we argue that biogeochemical legacy is the dominant retention process in most of the study catchments. We explain the missing catchment response for decreasing N inputs with a buffer effect, which stems from the accumulated biogeochemical legacy – acting as a secondary source and constituting a system inert to declining N inputs. A similar kind of biogeochemical dominance was also reported in a recent study for northwestern French catchments (Dupas et al., 2020). They concluded two-third of the retention being stored in the subsoil with the potential to recycle this N in the agroecosystem. Also Ascott et al. (2017) concluded that the vadose zone is globally a significant  $\text{NO}_3$  store. If not being recycled and in light of limited denitrification potential, the stored N would further leach to the deeper subsurface (or groundwater), when being mineralized again (Van Meter & Basu, 2015). The missing export of three-quarters of the past N inputs in the study catchments therefore constitutes a huge challenge for efforts to reach effective water quality improvements now and in the future.

After the implementation of regulations such as the EU Nitrate Directive (CEC, 1991), the diffuse N input decreased between the 1980s and 2010s by more than  $20 \text{ kg N ha}^{-1} \text{ year}^{-1}$  (36%) in the studied Western European catchments. The responses of riverine N loads to this decrease in input was limited ( $<1.5 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ). Hence, the retention decreased but catchments still received (in the 2010s) excess N of almost  $30 \text{ kg N ha}^{-1}$  every year, which is two-thirds of the diffuse input. These observations imply either a hindered substantial exploitation of the (already massive) biogeochemical legacy by mineralization and/or an ongoing exhaustion of the catchment's denitrification potential.

## 5. Conclusions and Implications

In this study we used long-term time series of N input and riverine  $\text{NO}_3$ -N output from 238 Western European catchments to estimate the N TTs, retention amount as well as the controlling catchment characteristics for both.

1. The analysis of catchment responses revealed peak TTs around 5 years with 70% of the catchments showing a peak export within the first 10 years after N enters the system. Hence, when assessing the effectiveness of measures, catchment managers have to be aware of this time lag in the response of N concentrations to changing N inputs.
2. Our analyses suggest a minor role of hydrologic legacy to explain the high retention of N in catchments (around three-fourths of incoming N). We rather see evidence for a widespread biogeochemical legacy of N, while biogeochemical conditions for a permanent removal by denitrification are only rarely achieved. Therefore, decreasing concentrations within the first 10 years mean neither that most of the N was already exported nor that restoration efforts can be reduced. Management in such cases would need rather long-term strategies to reduce ongoing leaching from soil N pools, for example by recycling the retained N within the soil or by fostering denitrifying conditions (i.e., increase plant-availability of soil N, catch-crops, supporting riparian rehabilitation, decreasing land drainage or by increasing levels of bio-available organic carbon; Abbott et al., 2018; Hunter et al., 2006).
3. While TTs were mainly controlled by hydroclimatic parameters with low PET and high precipitation seasonality favoring more rapid transport of N to the streams, retention was mainly controlled by specific discharge and subsurface parameters as low specific discharge and a high share of thick, unconsolidated aquifers in the catchments favor high retention. Thus, catchment managers can estimate from subsurface and hydroclimatic data, the natural conditions for retention and the dimension of TTs, which can be a helpful tool to explain the failure of measures or to advise a realistic management plan.
4. From a management perspective, a better spatial and temporal knowledge of denitrification efficiency at larger scales should be aimed at. Being associated with this, research on long-term changes of N storage capacities in agricultural soils is required. These data-driven analyses can be used to support or complement modeling approaches assisting different large scale water quality management activities.

### Data Availability Statement

Please note that the used data base adheres to Enabling FAIR Data Project requirements and is referenced in the manuscript linking to the data bases and repositories. Water quality and quantity data for French catchments are available at <http://www.hydroshare.org/resource/d8c43e1e8a5a4872bc0b75a45f350f7a> (Ehrhardt et al., 2021). Diffuse N input data for France were derived from Poisvert et al. (2017). Water quality and quantity data for Germany are available at <https://www.hydroshare.org/resource/a42addcbd59a466a9aa56472d-fef8721/> (Musolff et al., 2020). Catchment characteristics for Germany and France are available at <https://www.hydroshare.org/resource/c7d4df3ba74647f0aa83ae92be2e294b/> (Ebeling & Dupas, 2021).

### Acknowledgments

The authors thank the Federal authorities for providing water sample data and all contributors to setting up the used data base. Open access funding enabled and organized by Projekt DEAL.

### References

- Abbott, B. W., Moatar, F., Gauthier, O., Fovet, O., Antoine, V., & Ragueneau, O. (2018). Trends and seasonality of river nutrients in agricultural catchments: 18 years of weekly citizen science in France. *The Science of the Total Environment*, 624, 845–858. <https://doi.org/10.1016/j.scitotenv.2017.12.176>
- Ai, L., Shi, Z. H., Yin, W., & Huang, X. (2015). Spatial and seasonal patterns in stream water contamination across mountainous watersheds: Linkage with landscape characteristics. *Journal of Hydrology*, 523, 398–408. <https://doi.org/10.1016/j.jhydrol.2015.01.082>
- Allain, M. (1951). *France GEOLOGIQUE. Atlas universel Quillet physique, économique, politique. Tome 1: France et union française (1951)*. Retrieved from [https://www.mapmania.org/map/76435/geology\\_of\\_france\\_carte\\_geologique\\_de\\_la\\_france\\_5038\\_x\\_4687](https://www.mapmania.org/map/76435/geology_of_france_carte_geologique_de_la_france_5038_x_4687)
- Allred, B. J., Bigham, J. M., & Brown, G. O. (2007). The impact of clay mineralogy on nitrate mobility under unsaturated flow conditions. *Vadose Zone Journal*, 6(2), 221–232. <https://doi.org/10.2136/vzj2006.0064>
- Ascott, M. J., Gooddy, D. C., Wang, L., Stuart, M. E., Lewis, M. A., Ward, R. S., & Binley, A. M. (2017). Global patterns of nitrate storage in the vadose zone. *Nature Communications*, 8(1), 1–7. <https://doi.org/10.1038/s41467-017-01321-w>
- Bach, M., & Frede, H. G. (1998). Agricultural nitrogen, phosphorus and potassium balances in Germany-Methodology and trends 1970 to 1995. *Zeitschrift für Pflanzenernährung und Bodenkunde*, 161, 8. <https://doi.org/10.1002/jpln.1998.3581610406>
- Bain, D. J., Green, M. B., Campbell, J. L., Chamblee, J. F., Chaoka, S., Fraterrigo, J. M., et al. (2012). Legacy effects in material flux: Structural catchment changes predate long-term studies. *BioScience*, 62(6), 575–584. <https://doi.org/10.1525/bio.2012.62.6.8>
- Bartnicky, J., & Benedictow, A. (2017). *Atmospheric deposition of nitrogen to OSPAR convention waters in the period 1995–2014*. Bartnicky, J., & Fagerli, H. (2006). *Atmospheric nitrogen in the OSPAR convention area in the period 1990–2004*.
- Basu, N. B., Destouni, G., Jawitz, J. W., Thompson, S. E., Loukinova, N. V., Darracq, A., et al. (2010). Nutrient loads exported from managed catchments reveal emergent biogeochemical stationarity. *Geophysical Research Letters*, 37(23). <https://doi.org/10.1029/2010gl045168>
- Benettin, P., Rinaldo, A., & Botter, G. (2013). Kinematics of age mixing in advection-dispersion models. *Water Resources Research*, 49(12), 8539–8551. <https://doi.org/10.1002/2013wr014708>
- BGR and CGMW. (2005). *The 1:5 million international geological map of Europe and adjacent areas (IGME 5000)*. Retrieved from [https://geoviewer.bgr.de/mapapps4/resources/apps/geoviewer/index.html?cover=geologie\\_igme5000\\_ags&tab=geologie&lang=de](https://geoviewer.bgr.de/mapapps4/resources/apps/geoviewer/index.html?cover=geologie_igme5000_ags&tab=geologie&lang=de)

- BGR and UNESCO. (2014). *International hydrogeological map of Europe 1: 1,500,000 (IHME1500)*. Digital map data v1.1. Retrieved from <http://www.bgr.bund.de/ihme1500/http://www.bgr.bund.de/ihme1500/>
- Bingham, A. H., & Cotrufo, M. F. (2016). Organic nitrogen storage in mineral soil: Implications for policy and management. *The Science of the Total Environment*, 551–552, 116–126. <https://doi.org/10.1016/j.scitotenv.2016.02.020>
- Birkel, C., Soulsby, C., & Tetzlaff, D. (2015). Conceptual modelling to assess how the interplay of hydrological connectivity, catchment storage and tracer dynamics controls nonstationary water age estimates. *Hydrological Processes*, 29(13), 2956–2969. <https://doi.org/10.1002/hyp.10414>
- Blume, T., & van Meerveld, H. J. (2015). From hillslope to stream: Methods to investigate subsurface connectivity. *Wiley Interdisciplinary Reviews-Water*, 2(3), 177–198. <https://doi.org/10.1002/wat2.1071>
- BMU, B. f. U., Naturschutz und Reaktorsicherheit. (2003). Hydrologischer Atlas von Deutschland. Grundwasser. Retrieved from [http://www.hydrology.uni-freiburg.de/forsch/had/had\\_bezug.htm](http://www.hydrology.uni-freiburg.de/forsch/had/had_bezug.htm)
- Botter, G., Basso, S., Rodriguez-Iturbe, I., & Rinaldo, A. (2013). Resilience of river flow regimes. *Proceedings of the National Academy of Sciences*, 110(32), 12925–12930. <https://doi.org/10.1073/pnas.1311920110>
- Botter, G., Bertuzzo, E., & Rinaldo, A. (2011). Catchment residence and travel time distributions: The master equation. *Geophysical Research Letters*, 38(11), a–n. <https://doi.org/10.1029/2011gl047666>
- Bourauoi, F., & Grizzetti, B. (2011). Long term change of nutrient concentrations of rivers discharging in European seas. *The Science of the Total Environment*, 409(23), 4899–4916. <https://doi.org/10.1016/j.scitotenv.2011.08.015>
- Bouwman, A. F., Beusen, A. H., Griffioen, J., Van Groenigen, J. W., Hefting, M. M., Oenema, O., et al. (2013). Global trends and uncertainties in terrestrial denitrification and N<sub>2</sub>O emissions. *Philosophical Transactions of the Royal Society of London B Biological Sciences*, 368(1621), 20130112. <https://doi.org/10.1098/rstb.2013.0112>
- Buyko, M. I. (1974). *Climate and life*. Academic Press.
- CEC. (1991). Council Directive 91/676/EEC of 12 December 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources. (No L 375/1). Official Journal of the European Communities. Retrieved from <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A31991L0676>
- CLC. (2000). CORINE land cover 2000. Retrieved from <https://land.copernicus.eu/pan-european/corine-land-cover>
- Cleveland, C. C., Townsend, A. R., Schimel, D. S., Fisher, H., Howarth, R. W., Hedin, L. O., et al. (1999). Global patterns of terrestrial biological nitrogen (N<sub>2</sub>) fixation in natural ecosystems. *Global Biogeochemical Cycles*, 13(2), 623–645. <https://doi.org/10.1029/1999gb900014>
- de Vries, W., Leip, A., Reinds, G. J., Kros, J., Lesschen, J. P., Bouwman, A. F., et al. (2011). Geographical variation in terrestrial nitrogen budgets across Europe. In M. A. Sutton, C. M. Howard, J. W. Erisman, G. Billen, A. Bleeker, P. Grennfelt, H. van Grinsven, & B. Grizzetti (Eds.), *The European nitrogen assessment. Sources, effects and policy perspectives* (pp. 317–344): Cambridge University Press.
- Donnelly, C., Greuell, W., Andersson, J., Gerten, D., Pisacane, G., Roudier, P., & Ludwig, F. (2017). Impacts of climate change on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level. *Climatic Change*, 143(1–2), 13–26. <https://doi.org/10.1007/s10584-017-1971-7>
- Dupas, R., Curie, F., Gascuel-Oudou, C., Moatar, F., Delmas, M., Parnaudeau, V., & Durand, P. (2013). Assessing N emissions in surface water at the national level: Comparison of country-wide vs. regionalized models. *The Science of the Total Environment*, 443, 152–162. <https://doi.org/10.1016/j.scitotenv.2012.10.011>
- Dupas, R., Delmas, M., Dorioz, J. M., Garnier, J., Moatar, F., & Gascuel-Oudou, C. (2015). Assessing the impact of agricultural pressures on N and P loads and eutrophication risk. *Ecological Indicators*, 48, 396–407. <https://doi.org/10.1016/j.ecolind.2014.08.007>
- Dupas, R., Ehrhardt, S., Musolff, A., Fovet, O., & Durand, P. (2020). Long-term nitrogen retention and transit time distribution in agricultural catchments in western France. *Environmental Research Letters*, 15(11), 115011. <https://doi.org/10.1088/1748-9326/abbe47>
- Dupas, R., Jomaa, S., Musolff, A., Borchardt, D., & Rode, M. (2016). Disentangling the influence of hydroclimatic patterns and agricultural management on river nitrate dynamics from sub-hourly to decadal time scales. *The Science of the Total Environment*, 571, 791–800. <https://doi.org/10.1016/j.scitotenv.2016.07.053>
- Ebeling, P., & Dupas, R. (2021). *CCDB—catchment characteristics data base France and Germany*. Retrieved from <https://www.hydroshare.org/resource/c7d4df3ba74647f0aa83ae92be2e294b/>
- Ebeling, P., Kumar, R., Weber, M., Knoll, L., Fleckenstein, J. H., & Musolff, A. (2021). Archetypes and controls of riverine nutrient export across German catchments. *Water Resources Research*, 57(4). <https://doi.org/10.1029/2020wr028134>
- EC. (2009). *Regions 2020. The climate change challenge for European regions*. Retrieved from [http://ec.europa.eu/regional\\_policy/sources/docoffic/working/regions2020/pdf/regions2020\\_climat.pdf](http://ec.europa.eu/regional_policy/sources/docoffic/working/regions2020/pdf/regions2020_climat.pdf)
- Ehrhardt, S., Ebeling, P., & Dupas, R. (2021). *Exported French water quality and quantity data*. Retrieved from <https://www.hydroshare.org/resource/d8c43e1e8a5a4872bc0b75a45f350f7a/>
- Ehrhardt, S., Kumar, R., Fleckenstein, J. H., Attinger, S., & Musolff, A. (2019). Trajectories of nitrate input and output in three nested catchments along a land use gradient. *Hydrology and Earth System Sciences*, 23(9), 3503–3524. <https://doi.org/10.5194/hess-23-3503-2019>
- EPA, U. S. E. P. A. (1972). *Federal water pollution control amendments of 1972. Clean water act. 33 U.S.C. § 1251 et seq.* Retrieved from <https://www3.epa.gov/npdes/pubs/cwatxt.txt>
- European Environment Agency. (2016). *CORINE land cover 2012 v18.5*. Retrieved from <https://land.copernicus.eu/pan-european/corine-land-cover>
- European Environment Agency. (2017). *Waterbase—UWWTD: Urban waste water treatment directive—reported data*. Retrieved from <https://www.eea.europa.eu/data-and-maps/data/waterbase-uwtd-urban-waste-water-treatment-directive-5#tab-european-data>
- European Environment Agency. (2018). *European waters. Assessment of status and pressure*. Retrieved from <https://www.eea.europa.eu/publications/state-of-water>
- European Environment Agency. (2020). *CORINE land cover 2000 v2020u1. From EEA*. Retrieved from <https://land.copernicus.eu/pan-european/corine-land-cover>
- Fiori, A., Russo, D., & Di Lazzaro, M. (2009). Stochastic analysis of transport in hillslopes: Travel time distribution and source zone dispersion. *Water Resources Research*, 45(8). <https://doi.org/10.1029/2008wr007668>
- Godsey, S. E., Aas, W., Clair, T. A., Wit, H. A. d., Fernandez, I. J., Kahl, J. S., et al. (2010). Generality of fractal 1/f scaling in catchment tracer time series, and its implications for catchment travel time distributions. *Hydrological Processes*, 24(12), 1660–1671. <https://doi.org/10.1002/hyp.7677>
- Goyette, J.-O., Bennett, E. M., & Maranger, R. (2018). Differential influence of landscape features and climate on nitrogen and phosphorus transport throughout the watershed. *Biogeochemistry*, 142(1), 155–174. <https://doi.org/10.1007/s10533-018-0526-y>

- Grabs, T., Bishop, K., Laudon, H., Lyon, S. W., & Seibert, J. (2012). Riparian zone hydrology and soil water total organic carbon (TOC): Implications for spatial variability and upscaling of lateral riparian TOC exports. *Biogeosciences*, 9(10), 3901–3916. <https://doi.org/10.5194/bg-9-3901-2012>
- Griffis, T. J., Chen, Z., Baker, J. M., Wood, J. D., Millet, D. B., Lee, X., et al. (2017). Nitrous oxide emissions are enhanced in a warmer and wetter world. *Proceedings of the National Academy of Sciences of the United States of America*, 114(45), 12081–12085. <https://doi.org/10.1073/pnas.1704552114>
- Haitjema, H. M. (1995). On the residence time distribution in idealized groundwatersheds. *Journal of Hydrology*, 172, 20. [https://doi.org/10.1016/0022-1694\(95\)02732-5](https://doi.org/10.1016/0022-1694(95)02732-5)
- Harman, C. J. (2015). Time-variable transit time distributions and transport: Theory and application to storage-dependent transport of chloride in a watershed. *Water Resources Research*, 51(1), 1–30. <https://doi.org/10.1002/2014wr015707>
- Hatfield, J. L., Boote, K. J., Kimball, B. A., Ziska, L. H., Izaurralde, R. C., Ort, D., et al. (2011). Climate impacts on agriculture: Implications for crop production. *Agronomy Journal*, 103(2), 351–370. <https://doi.org/10.2134/agnonj2010.0303>
- Häußermann, U., Bach, M., Klement, L., & Breuer, L. (2019). *Stickstoff-Flächenbilanzen für Deutschland mit Regionalgliederung Bundesländer und Kreise—Jahre 1995 bis 2017—Methodik, Ergebnisse und Minderungsmaßnahmen*. Retrieved from [https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2019-10-28\\_texte\\_131-2019\\_stickstofflaeichenbilanz.pdf](https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2019-10-28_texte_131-2019_stickstofflaeichenbilanz.pdf)
- Heidbüchel, I., Yang, J., Musolff, A., Troch, P., Ferré, T., & Fleckenstein, J. H. (2020). On the shape of forward transit time distributions in low-order catchments. *Hydrology and Earth System Sciences*, 24(6), 2895–2920. <https://doi.org/10.5194/hess-24-2895-2020>
- Hirsch, R. M., & de Cicco, L. (2019). *EGRET: Exploration and graphics for river trends, R package version 3.0.2*.
- Hirsch, R. M., Moyer, D. L., & Archfield, S. A. (2010). Weighted regressions on time, discharge, and season (WRTDS), with an application to Chesapeake Bay river inputs. *Journal of the American Water Resources Association*, 46(5), 857–880. <https://doi.org/10.1111/j.1752-1688.2010.00482.x>
- Howarth, R. W., Swaney, D. P., Boyer, E. W., Marino, R., Jaworski, N., & Goodale, C. (2006). The influence of climate on average nitrogen export from large watersheds in the Northeastern United States. *Biogeochemistry*, 79(1–2), 163–186. <https://doi.org/10.1007/s10533-006-9010-1>
- Howden, N. J. K., Burt, T. P., Worrall, F., Mathias, S., & Whelan, M. J. (2011). Nitrate pollution in intensively farmed regions: What are the prospects for sustaining high-quality groundwater? *Water Resources Research*, 47(6). <https://doi.org/10.1029/2011wr010843>
- Howden, N. J. K., Burt, T. P., Worrall, F., Whelan, M. J., & Bierzoza, M. (2010). Nitrate concentrations and fluxes in the River Thames over 140 years (1868–2008): Are increases irreversible? *Hydrological Processes*, 24(18), 2657–2662. <https://doi.org/10.1002/hyp.7835>
- Hunter, H., Fellows, C., Rassam, D., DeHayr, R., Pagendam, D., Conway, C., et al. (2006). *Managing riparian lands to improve water quality: Optimising nitrate removal via denitrification*. Retrieved from [http://www.ozcoasts.gov.au/pdf/CRC/57-riparian\\_guidelines.pdf](http://www.ozcoasts.gov.au/pdf/CRC/57-riparian_guidelines.pdf)
- Jabloun, M., Schelde, K., Tao, F., & Olesen, J. E. (2015). Effect of temperature and precipitation on nitrate leaching from organic cereal cropping systems in Denmark. *European Journal of Agronomy*, 62, 55–64. <https://doi.org/10.1016/j.eja.2014.09.007>
- Jasechko, S., Kirchner, J. W., Welker, J. M., & McDonnell, J. J. (2016). Substantial proportion of global streamflow less than three months old. *Nature Geoscience*, 9(2), 126–129. <https://doi.org/10.1038/ngeo2636>
- Kirchner, F., Neal, X., & Neal, C. (2000). Fractal stream chemistry and its implications for contaminant transport in catchments. *Nature*, 403(6769), 524–527. <https://doi.org/10.1038/35000537>
- Knoll, L., Breuer, L., & Bach, M. (2020). Nation-wide estimation of groundwater redox conditions and nitrate concentrations through machine learning. *Environmental Research Letters*, 15(6), 064004. <https://doi.org/10.1088/1748-9326/ab7d5c>
- Kronvang, B., Andersen, H. E., Børgesen, C., Dalgaard, T., Larsen, S. E., Bøgestrand, J., & Blicher-Mathiasen, G. (2008). Effects of policy measures implemented in Denmark on nitrogen pollution of the aquatic environment. *Environmental Science & Policy*, 11(2), 144–152. <https://doi.org/10.1016/j.envsci.2007.10.007>
- Kumar, R., Hesse, F., Rao, P. S. C., Musolff, A., Jawitz, J. W., Sarrazin, F., et al. (2020). Strong hydroclimatic controls on vulnerability to subsurface nitrate contamination across Europe. *Nature Communications*, 11(1), 6302. <https://doi.org/10.1038/s41467-020-19955-8>
- Kumar, R., Samaniego, L., & Attinger, S. (2013). Implications of distributed hydrologic model parameterization on water fluxes at multiple scales and locations. *Water Resources Research*, 49(1), 360–379. <https://doi.org/10.1029/2012wr012195>
- Kunkel, R., & Wendland, F. (2006). Diffuse Nitrateinträge in die Grund- und Oberflächengewässer von Rhein und Ems- Ist-Zustands- und Maßnahmenanalysen. In F. J. GmbH (Ed.), *Schriften des Forschungszentrums Jülich* (Vol. 62, p. 143). Reihe Umwelt/Environment Jülich.
- Marx, A., Kumar, R., Thober, S., Rakovec, O., Wanders, N., Zink, M., et al. (2018). Climate change alters low flows in Europe under global warming of 1.5, 2, and 3 °C. *Hydrology and Earth System Sciences*, 22(2), 1017–1032. <https://doi.org/10.5194/hess-22-1017-2018>
- McMahon, P. B., Dennehy, K. F., Bruce, B. W., Böhlke, J. K., Michel, R. L., Gurdak, J. J., & Hurlbut, D. B. (2006). Storage and transit time of chemicals in thick unsaturated zones under rangeland and irrigated cropland, High Plains, United States. *Water Resources Research*, 42(3). <https://doi.org/10.1029/2005wr004417>
- Meals, D. W., Dressing, S. A., & Davenport, T. E. (2010). Lag time in water quality response to best management practices: A review. *Journal of Environmental Quality*, 39(1), 85–96. <https://doi.org/10.2134/jeq2009.0108>
- Merz, C., Steidl, J., & Dannowski, R. (2009). Parameterization and regionalization of redox based denitrification for GIS-embedded nitrate transport modeling in Pleistocene aquifer systems. *Environmental Geology*, 58(7). <https://doi.org/10.1007/s00254-008-1665-6>
- Moatar, F., Abbott, B. W., Minaudo, C., Curie, F., & Pinay, G. (2017). Elemental properties, hydrology, and biology interact to shape concentration-discharge curves for carbon, nutrients, sediment, and major ions. *Water Resources Research*, 53(2), 1270–1287. <https://doi.org/10.1002/2016wr019635>
- Molenat, J., Gascuel-Oudou, C., Ruiz, L., & Gruau, G. (2008). Role of water table dynamics on stream nitrate export and concentration in agricultural headwater catchment (France). *Journal of Hydrology*, 348(3–4), 363–378. <https://doi.org/10.1016/j.jhydrol.2007.10.005>
- Musolff, A. (2020). *WQQDB—Water quality and quantity data base Germany: Metadata*. HydroShare.
- Musolff, A., Fleckenstein, J. H., Rao, P. S. C., & Jawitz, J. W. (2017). Emergent archetype patterns of coupled hydrologic and biogeochemical responses in catchments. *Geophysical Research Letters*, 44(9), 4143–4151. <https://doi.org/10.1002/2017gl072630>
- Musolff, A., Grau, T., Weber, M., Ebeling, P., Samaniego-Eguiguren, L., & Kumar, R. (2020). *WQQDB: Water quality and quantity data base Germany*. Retrieved from <http://www.ufz.de/record/dmp/archive/7754>
- Nolan, B. T., Hitt, K. J., & Ruddy, B. C. (2002). Probability of nitrate contamination of recently recharged groundwaters in the conterminous United States. *Environmental Science and Technology*, 36(10), 2138–2145. <https://doi.org/10.1021/es0113854>
- Ocampo, C. J., Sivapalan, M., & Oldham, C. (2006). Hydrological connectivity of upland-riparian zones in agricultural catchments: Implications for runoff generation and nitrate transport. *Journal of Hydrology*, 331(3–4), 643–658. <https://doi.org/10.1016/j.jhydrol.2006.06.010>

- Panagos, P., Borrelli, P., Poesen, J., Ballabio, C., Lugato, E., Meusburger, K., et al. (2015). The new assessment of soil loss by water erosion in Europe. *Environmental Science & Policy*, *54*, 438–447. <https://doi.org/10.1016/j.envsci.2015.08.012>
- Poisvert, C., Curie, F., & Moatar, F. (2017). Annual agricultural N surplus in France over a 70-year period. *Nutrient Cycling in Agroecosystems*, *107*(1), 63–78. <https://doi.org/10.1007/s10705-016-9814-x>
- Powers, S. M., Tank, J. L., & Robertson, D. M. (2015). Control of nitrogen and phosphorus transport by reservoirs in agricultural landscapes. *Biogeochemistry*, *124*(1–3), 417–439. <https://doi.org/10.1007/s10533-015-0106-3>
- Roa-García, M. C., & Weiler, M. (2010). Integrated response and transit time distributions of watersheds by combining hydrograph separation and long-term transit time modeling. *Hydrology and Earth System Sciences*, *14*(8), 1537–1549. <https://doi.org/10.5194/hess-14-1537-2010>
- Samaniego, L., Kumar, R., & Attinger, S. (2010). Multiscale parameter regionalization of a grid-based hydrologic model at the mesoscale. *Water Resources Research*, *46*(5). <https://doi.org/10.1029/2008wr007327>
- Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., et al. (2018). Anthropogenic warming exacerbates European soil moisture droughts. *Nature Climate Change*, *8*(5), 421–426. <https://doi.org/10.1038/s41558-018-0138-5>
- Sebilo, M., Mayer, B., Nicolardot, B., Pinay, G., & Mariotti, A. (2013). Long-term fate of nitrate fertilizer in agricultural soils. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(45), 18185–18189. <https://doi.org/10.1073/pnas.1305372110>
- Shi, Z. H., Ai, L., Li, X., Huang, X. D., Wu, G. L., & Liao, W. (2013). Partial least-squares regression for linking land-cover patterns to soil erosion and sediment yield in watersheds. *Journal of Hydrology*, *498*, 165–176. <https://doi.org/10.1016/j.jhydrol.2013.06.031>
- Stevenson, F. J. (1986). *Cycles of soil: Carbon, nitrogen, phosphorus, sulfur, micronutrients*: Wiley.
- Tesoriero, A. J., Gronberg, J. A., Juckem, P. F., Miller, M. P., & Austin, B. P. (2017). Predicting redox-sensitive contaminant concentrations in groundwater using random forest classification. *Water Resources Research*, *53*(8), 7316–7331. <https://doi.org/10.1002/2016wr020197>
- Thomas, Z., & Abbott, B. W. (2018). Hedgerows reduce nitrate flux at hillslope and catchment scales via root uptake and secondary effects. *Journal of Contaminant Hydrology*, *215*, 51–61. <https://doi.org/10.1016/j.jconhyd.2018.07.002>
- Van der Velde, Y., de Rooij, G. H., & Torfs, P. J. J. F. (2009). Catchment-scale non-linear groundwater-surface water interactions in densely drained lowland catchments. *Hydrology and Earth System Sciences*, *13*(10), 1867–1885. <https://doi.org/10.5194/hess-13-1867-2009>
- Van der Velde, Y., Rooij, G. H. d., Rozemeijer, J. C., van Geer, F. C., & Broers, H. P. (2010). Nitrate response of a lowland catchment: On the relation between stream concentration and travel time distribution dynamics. *Water Resources Research*, *46*(11), 1–17. <https://doi.org/10.1029/2010wr009105>
- Van Meter, K. J., & Basu, N. B. (2015). Catchment legacies and time lags: A parsimonious watershed model to predict the effects of legacy storage on nitrogen export. *PLoS One*, *10*(5), e0125971. <https://doi.org/10.1371/journal.pone.0125971>
- Van Meter, K. J., & Basu, N. B. (2017). Time lags in watershed-scale nutrient transport: An exploration of dominant controls. *Environmental Research Letters*, *12*(8), 084017. <https://doi.org/10.1088/1748-9326/aa7bf4>
- Van Meter, K. J., Basu, N. B., & van Cappellen, P. (2017). Two centuries of nitrogen dynamics: Legacy sources and sinks in the Mississippi and Susquehanna River Basins. *Global Biogeochemical Cycles*, *31*(1), 2–23. <https://doi.org/10.1002/2016gb005498>
- Van Meter, K. J., Basu, N. B., Veenstra, J. J., & Burras, C. L. (2016). The nitrogen legacy: Emerging evidence of nitrogen accumulation in anthropogenic landscapes. *Environmental Research Letters*, *11*(3). <https://doi.org/10.1088/1748-9326/11/3/035014>
- Van Meter, K. J., Van Cappellen, P., & Basu, N. B. (2018). Legacy nitrogen may prevent achievement of water quality goals in the Gulf of Mexico. *Science*, *360*(6387), 427–430. <https://doi.org/10.1126/science.aar4462>
- Vero, S. E., Basu, N. B., Van Meter, K., Richards, K. G., Mellander, P.-E., Healy, M. G., & Fenton, O. (2017). Review: The environmental status and implications of the nitrate time lag in Europe and North America. *Hydrogeology Journal*, *26*(1), 7–22. <https://doi.org/10.1007/s10040-017-1650-9>
- von Lutzow, M., Kogel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., & Flessa, H. (2006). Stabilization of organic matter in temperate soils: Mechanisms and their relevance under different soil conditions—A review. *European Journal of Soil Science*, *57*(4), 426–445. <https://doi.org/10.1111/j.1365-2389.2006.00809.x>
- Wang, L., & Burke, S. P. (2017). A catchment-scale method to simulating the impact of historical nitrate loading from agricultural land on the nitrate-concentration trends in the sandstone aquifers in the Eden Valley, UK. *The Science of the Total Environment*, *579*, 133–148. <https://doi.org/10.1016/j.scitotenv.2016.10.235>
- Wang, Z., Qi, Z., Xue, L., Bukovsky, M., & Halmers, M. J. (2015). Modeling the impacts of climate change on nitrogen losses and crop yield in a subsurface drained field. *Climatic Change*, *129*(1–2), 323–335. <https://doi.org/10.1007/s10584-015-1342-1>
- Wassenaar, L. I. (1995). Evaluation of the origin and fate of nitrate in the Abbotsford Aquifer using the isotopes. *Applied Geochemistry*, *10*, 15–405. [https://doi.org/10.1016/0883-2927\(95\)00013-A](https://doi.org/10.1016/0883-2927(95)00013-A)
- Webster, J. R., Mulholland, P. J., Tank, J. L., Valett, H. M., Dodds, W. K., Peterson, B. J., et al. (2003). Factors affecting ammonium uptake in streams – An inter-biome perspective. *Freshwater Biology*, *48*(8), 1329–1352. <https://doi.org/10.1046/j.1365-2427.2003.01094.x>
- Wendland, F., Blum, A., Coetsiers, M., Gorova, R., Griffioen, J., Grima, J., et al. (2007). European aquifer typology: A practical framework for an overview of major groundwater composition at European scale. *Environmental Geology*, *55*(1), 77–85. <https://doi.org/10.1007/s00254-007-0966-5>
- West, P. C., Gerber, J. S., Engstrom, P. M., Mueller, N. D., Brauman, K. A., Carlson, K. M., et al. (2014). Leverage points for improving global food security and the environment. *Science*, *345*(6194), 325–328. <https://doi.org/10.1126/science.1246067>
- Westphal, K., Musolff, A., Graeber, D., & Borchardt, D. (2020). Controls of point and diffuse sources lowered riverine nutrient concentrations asynchronously, thereby warping molar N:P ratios. *Environmental Research Letters*, *15*(10). <https://doi.org/10.1088/1748-9326/ab98b6>
- Wold, S., Sjostrom, M., & Eriksson, L. (2001). PLS-regression: A basic tool of chemometrics. *Chemometrics and Intelligent Laboratory Systems*, *58*(2), 109–130. Doi: [https://doi.org/10.1016/S0169-7439\(01\)00155-1](https://doi.org/10.1016/S0169-7439(01)00155-1)
- Worldometers.info. (2020). *Europe-population*. Retrieved from <https://www.worldometers.info/world-population/europe-population/>
- Wriedt, G., Spindler, J., Neef, T., Meißner, R., & Rode, M. (2007). Groundwater dynamics and channel activity as major controls of in-stream nitrate concentrations in a lowland catchment system? *Journal of Hydrology*, *343*(3–4), 154–168. <https://doi.org/10.1016/j.jhydrol.2007.06.010>
- Wyns, R., Mathieu, F., Vairon, J., Legchenko, A., Lachassagne, P., & Baltassat, J.-M. (2004). Application of proton magnetic resonance soundings to groundwater reserve mapping in weathered basement rocks (Brittany, France). *Bulletin de la Societe Geologique de France*, *175*(1), 21–34. <https://doi.org/10.2113/175.1.21>
- Yang, J., Heidbüchel, I., Musolff, A., Reinstorf, F., & Fleckenstein, J. H. (2018). Exploring the dynamics of transit times and subsurface mixing in a small agricultural catchment. *Water Resources Research*, *54*(3), 2317–2335. <https://doi.org/10.1002/2017wr021896>

- Zambrano-Bigiarini, M., & Rojas, R. (2013). A model-independent particle swarm optimisation software for model calibration. *Environmental Modelling & Software*, 43, 5–25. <https://doi.org/10.1016/j.envsoft.2013.01.004>
- Zhang, Y.-C., Slomp, C. P., Broers, H. P., Passier, H. F., & Cappellen, P. V. (2009). Denitrification coupled to pyrite oxidation and changes in groundwater quality in a shallow sandy aquifer. *Geochimica et Cosmochimica Acta*, 73(22), 6716–6726. <https://doi.org/10.1016/j.gca.2009.08.026>

## References From the Supporting Information

- Ballabio, C., Lugato, E., Fernandez-Ugalde, O., Orgiazzi, A., Jones, A., Borrelli, P., et al. (2019). Mapping LUCAS topsoil chemical properties at European scale using Gaussian process regression. *Geoderma*, 355, 113912. <https://doi.org/10.1016/j.geoderma.2019.113912>
- CIESIN, C. f. I. E. S. I. N.-C. U. (2017). *Gridded population of the world, version 4 (GPWv4): Population density, Revision 10*. <https://doi.org/10.7927/H4DZ068D>
- CLC. (2016). CORINE land cover 2012 v18.5. Retrieved from <https://land.copernicus.eu/pan-european/corine-land-cover>
- Cornes, R. C., van der Schrier, G., van den Besselaar, E. J. M., & Jones, P. D. (2018). An ensemble version of the E-OBS temperature and precipitation data sets. *Journal of Geophysical Research: Atmospheres*, 123(17), 9391–9409. <https://doi.org/10.1029/2017jd028200>
- European Environment Agency. (2013). *DEM over Europe from the GMES RDA project (EU-DEM, resolution 25 m)—Version 1*. Retrieved from <https://sdi.eea.europa.eu/catalogue/srv9008075/api/records/66fa7dca-8772-4a5d-9d56-2caba4ecd36a>
- FAO/IIASA/ISRIC/ISSCAS/JRC. (2012). *Harmonized world soil database*. Retrieved from <https://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>
- Shangguan, W., Hengl, T., Mendes de Jesus, J., Yuan, H., & Dai, Y. (2017). Mapping the global depth to bedrock for land surface modeling. *Journal of Advances in Modeling Earth Systems*, 9(1), 65–88. <https://doi.org/10.1002/2016ms000686>
- Vigiak, O., Grizzetti, B., Zanni, M., Aloe, A., Dorati, C., Bouraoui, F., & Pistocchi, A. (2019). *Domestic waste emissions to European freshwaters in the 2010s (v. 1.0)*. Retrieved from <https://data.jrc.ec.europa.eu/dataset/0ae64ac2-64da-4c5e-8bab-ce928897c1fb>
- Vigiak, O., Grizzetti, B., Zanni, M., Aloe, A., Dorati, C., Bouraoui, F., & Pistocchi, A. (2020). Domestic waste emissions to European waters in the 2010s. *Scientific Data*, 7(1), 33. <https://doi.org/10.1038/s41597-020-0367-0>