



HAL
open science

Deregulation of apoplastic or symplastic steps in sugar loading and transport in *Arabidopsis thaliana*

Françoise Vilaine, Laurence Bill, Rozenn Le Hir, Beate Hoffmann, Catherine Bellini, Sylvie Dinant

► **To cite this version:**

Françoise Vilaine, Laurence Bill, Rozenn Le Hir, Beate Hoffmann, Catherine Bellini, et al.. Deregulation of apoplastic or symplastic steps in sugar loading and transport in *Arabidopsis thaliana*. Workshop sugar allocation in plants, Oct 2021, Versailles, France. . hal-03760335

HAL Id: hal-03760335

<https://hal.inrae.fr/hal-03760335>

Submitted on 25 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title: Deregulation of apoplastic or symplastic steps in sugar loading and transport in *Arabidopsis thaliana*

Poster number :

Authors

Françoise Vilaine^{1,*}, Laurence Bill¹, Rozenn Le Hir¹, Beate Hoffmann¹, Catherine Bellini^{1,2}, and Sylvie Dinant¹

Affiliations

¹ Institut Jean-Pierre Bourgin, INRAE, AgroParisTech, Université Paris-Saclay, 78000 Versailles, France

² Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, 90187 Umeå, Sweden

Abstract

In addition to their role as nutrients, sugars perform a multitude of functions and influence many processes. They play a crucial role in establishing sap flow and long-distance transport, in adjusting plant development in response to the environment or during stress. The composition of the phloem sap is a determining parameter for yield and biomass.

This work aimed to dissociate different steps of sugar loading and transport in *Arabidopsis*, involving either active transport (apoplastic component) or passive flow through plasmodesmata (symplastic component). Depending on the plant species, the environmental conditions, the stage of development, it would seem that both pathways are operational, although it is not well known what makes one or the other really used.

We generated different *Arabidopsis* lines deregulated specifically in certain apoplastic and symplastic components of the loading and transport of sugars. It made it possible to highlight distinct impacts, depending on the component affected, on the growth and physiology of the plant, on the homeostasis of sugars, on the composition of phloem sap and apoplast content, on seed filling, and on the expression of genes involved in the transport and metabolism of sugars.

The main questions and perspectives of this work, concern the compensatory mechanisms set up to maintain the phloem sap composition and the regulation of the balance between apoplastic and symplastic pathways during development and stress conditions.

Ultimately, it can be assumed that the knowledge acquired will make it possible to develop innovative solutions to improve the distribution and use of resources, and thus the yield and tolerance to biotic and abiotic stresses of plants.