



**HAL**  
open science

## Impairment of sugar transport and changes in phloem exudate contents in phytoplasma-infected plants

Federica de Marco, Brigitte Batailler, Rozenn Le Hir, Françoise Vilaine, Alain Bouchereau, Sandrine Eveillard, Sylvie Dinant

### ► To cite this version:

Federica de Marco, Brigitte Batailler, Rozenn Le Hir, Françoise Vilaine, Alain Bouchereau, et al.. Impairment of sugar transport and changes in phloem exudate contents in phytoplasma-infected plants. Workshop sugar allocation in plants, Oct 2021, Versailles, France. hal-03760344

**HAL Id: hal-03760344**

**<https://hal.inrae.fr/hal-03760344>**

Submitted on 25 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Title: Impairment of sugar transport and changes in phloem exudate contents in phytoplasma-infected plants**

Poster number :

## **Authors**

Federica De Marco <sup>1</sup>, Brigitte Batailler <sup>2</sup>, Rozenn Le Hir <sup>1</sup>, Françoise Vilaine <sup>1</sup>, Alain Bouchereau <sup>3</sup>, Sandrine Eveillard<sup>2</sup>, Sylvie Dinant<sup>1</sup>

## **Affiliations**

<sup>1</sup>UMR1318, Institut Jean-Pierre Bourgin, Versailles.

<sup>2</sup>UMR 1332 Biologie du Fruit et Pathologie, Villenave d'Ornon

<sup>3</sup> UMR IGEPP, INRAE, Agrocampus Ouest, Université de Rennes 1, Le Rheu.

## **Abstract**

Phytoplasmas inhabit phloem sieve elements, and cause abnormal growth and altered sugar partitioning. But how they interact with phloem functions is not clearly known. The phloem responses were investigated in tomato infected by 'Candidatus Phytoplasma solani', at the beginning of the symptomatic stage, the first symptoms appearing in the newly-emerged leaf at the stem apex. Antisense lines impaired in the phloem sucrose transporters SUT1 and SUT2 were included. In symptomatic sink leaves, leaf curling was associated with higher starch accumulation and expression of defense genes. The analysis of leaf midribs of symptomatic leaves indicated that transcript levels for genes acting in the glycolysis and peroxisome metabolism differed from these in non-infected plants. The phytoplasma also multiplied in the three lower source leaves even if it was not associated with symptoms. In these leaves, the rate of phloem sucrose exudation was lower for infected plants. Metabolite profiling of phloem sap-enriched exudates revealed that glycolate and aspartate levels were affected by the infection. Their levels were also affected in the non-infected SUT1- and SUT2- antisense lines. The findings suggest a role of sugar transporters in the responses to the infection and describe the consequences of impaired sugar transport on primary metabolism.