



HAL
open science

Response of soil and vegetation in a warm-temperate Pine forest to intensive biomass harvests, phosphorus fertilisation, and wood ash application

Laurent Augusto, Florent Beaumont, Christophe Nguyen, Jean-Yves Fraysse, Pierre Trichet, Céline Meredieu, David Vidal, Valérie Sappin-Didier

► To cite this version:

Laurent Augusto, Florent Beaumont, Christophe Nguyen, Jean-Yves Fraysse, Pierre Trichet, et al.. Response of soil and vegetation in a warm-temperate Pine forest to intensive biomass harvests, phosphorus fertilisation, and wood ash application. *Science of the Total Environment*, 2022, 850, pp.157907. 10.1016/j.scitotenv.2022.157907 . hal-03763508

HAL Id: hal-03763508

<https://hal.inrae.fr/hal-03763508v1>

Submitted on 27 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title**

2 Response of soil and vegetation in a warm-temperate Pine forest to intensive biomass harvests,
3 phosphorus fertilisation, and wood ash application.

4

5

6

7 **Authors**

8 Laurent Augusto ^{a,*}

9 Florent Beaumont ^a

10 Christophe Nguyen ^a

11 Jean-Yves Fraysse ^b

12 Pierre Trichet ^a

13 Céline Meredieu ^{c,d}

14 David Vidal ^a

15 Valérie Sappin-Didier ^a

16

17 a INRAE, Bordeaux Sciences Agro, ISPA, F-33140 Villenave d'Ornon, France

18 b FCBA, F-33610 Cestas, France

19 c INRAE, UEFP, F-33610 Cestas, France

20 d INRAE, Univ. Bordeaux, BIOGECO, F-33610 Cestas, France

21

22

23 **Abstract**

24 Background and Aims: Concerns about climate change and carbon economy have prompted the
25 promotion of alternative energy sources, including forest-based bioenergy. An evaluation of the
26 environmental consequences of intensive harvests (stumps and roots, and also branches and foliage) for
27 energy wood supply, and use of wood-ash recycling as a compensatory practice, helps in the evaluation
28 of the use of forest biomass for energy production.

29 Methods: We made use of records from a split-plot experimental site crossing four different intensities
30 of biomass harvesting (Stem-Only Harvest [SOH], Aboveground Additional Harvest [AAH],
31 Belowground Additional Harvest [BAH], and Whole-Tree Harvest [WTH]) and three compensation
32 methods (control [C], wood ash application [A] and phosphorus fertilisation [P]) to evaluate, in the 11-
33 years-old stand (maritime pine: *Pinus pinaster*) that followed the biomass exports of the former stand,
34 their effects on nutrient budgets, tree growth, soil fertility, chemical properties and soil carbon. This site
35 is located in a forest on a poor soil, under a warm temperate climate (SW France).

36 Key results: Despite their low additional biomass exports (+10% for AAH to +34% for WTH), the non-
37 conventional harvest practices exported much higher quantities of nutrients than the conventional SOH
38 technique (for example +145% for N and K in WTH). Consequently, these treatments had negative
39 effects on the soil nutritive status. Additional biomass harvests impacted the soil organic matter content,
40 with negative effects on P_{-organic}, soil cation exchange capacity, exchangeable Ca, and most extractible
41 nutrients. However, data suggested that tree growth and foliage nutrient content had not yet been
42 significantly impacted by harvest treatments, whereas tree nutritional status was improved by P-fertiliser
43 or wood ash. As expected, we observed a positive effect of wood ash application on soil pH and nutrient
44 content but, like additional harvests, wood ash application decreased the pool of soil organic carbon
45 (~10% of the initial stock with ~7% of N_{-total} losses).

46 Conclusions: Overall, this factorial experiment showed that exporting more forest biomass due to the
47 additional harvesting of tree canopies, stumps and roots had negative consequences on the ecosystem
48 biogeochemistry. Additional harvests have aggravated the poverty of the already oligotrophic soil, and
49 decreased the soil organic carbon content. Importantly, applying nutrients as fertiliser or wood ash did
50 not compensate for all the negative impacts of biomass exports and the method of wood ash recycling
51 in forests could even decrease the soil organic carbon.

52 **Introduction**

53 Although there is a consensus that forests constitute an important leverage for climate change mitigation
54 (IPCC 2014), there is also a heated debate on how forests should be managed for mitigation purposes
55 (Lindner and Karjalainen 2007). In particular, it is debated to what extent additional harvesting of forest
56 biomass, such as foliage, branches, stumps, and roots (Nicholls et al. 2009) should be promoted for
57 energy production. Indeed, this practice can impoverish forest ecosystems because these tree
58 compartments dedicated to energy wood supply –the so-called *harvest residues* or *logging residues*– are
59 rich in nutrients (Andre et al. 2010; Achat et al. 2015a; Augusto et al. 2015a). Review studies indicate
60 that intensive biomass harvesting –as compared with conventional harvesting– can have negative
61 consequences on ecosystem functioning, such as soil nutrients and organic matter pools, and
62 consequently on future forest growth (Thiffault et al. 2011; Wall 2012; Achat et al. 2015a). Extracting
63 more biomass from forests without any compensatory fertilisation seems to be an unsustainable leverage
64 for climate change mitigation (Garcia et al. 2018). Therefore, it has been proposed that intensive harvests
65 should be associated with compensatory practices (Nohrstedt 2001; Ranius et al. 2018; Ventura et al.
66 2019).

67 Applying wood ash is often promoted as a good method to compensate for the negative effects of
68 intensive harvests on the ecosystem nutrient budget (Hannam et al. 2018; Ranius et al. 2018; Ventura et
69 al. 2019). Indeed, wood ash has a high nutrient content (Aronsson and Ekelund 2004), can reduce soil
70 acidity (Reid and Watmough 2014), and contributes to the concept of *circular bioeconomy* as wood ash
71 returns some of the nutrients that were exported by the removal of logging residues (Pitman 2006). On
72 the other hand, applying wood ash might contaminate ecosystems because of its relatively high content
73 in several micronutrients or non-essential metals (Vance 1996; Nnadi et al. 2019). Unfortunately, while
74 several meta-analyses have been carried out on intensive biomass harvests (Thiffault et al. 2011; Achat
75 et al. 2015a) or wood ash application (Augusto et al. 2008a; Reid and Watmough 2014), a combination
76 of these two practices have rarely been studied together to assess their interactive consequences
77 (Hagerberg and Wallander 2002; Wang et al. 2010). Therefore, we conducted a field experiment whose
78 main objective was to evaluate these possible interactions.

79 Located in south-western France, the *Landes de Gascogne* forest (where our study took place) has
80 several advantages for evaluating the environmental consequences of intensive harvests and their
81 compensatory practices. Firstly, it is a large man-made pine forest almost entirely dedicated to intensive
82 forestry. As such, the area is already subjected to collection of harvest residues (mainly stumps and
83 roots, and to a lesser extent also branches and needles (Mora et al. 2014; Banos and Dehez 2017)) for
84 energy wood consumption (Augusto et al. 2010, 2015a), which in turn produces large amounts of wood
85 ash (Alvarez-Alvarez et al. 2018). Secondly because this forest is strongly oligotrophic (Augusto et al.
86 2010), and particularly poor in phosphorus (Achat et al. 2009), it has been diagnosed to be particularly
87 sensitive to the potential negative consequences of intensive harvest (Durante et al. 2019). Finally, it is
88 a warm-temperate forest. Indeed, knowledge about wood ash application and removal of logging
89 residues is mainly based on studies carried out in boreal forests (Augusto et al. 2008a; Walmsley and
90 Godbold 2010), whereas the consequences of these management practices are probably climate-
91 dependent (Achat et al. 2015b). For instance, several Nordic studies indicated that stump harvests, or
92 wood ash application, had no effect on soil organic carbon (SOC) while studies conducted in warmer
93 latitudes concluded that the impact was highly negative (Zabowski et al. 2008; Stromgren et al. 2013;
94 Jurevics et al. 2016).

95 Our initial expectations were based on the current knowledge about intensive biomass harvest or wood
96 ash application, in comparison with the local conditions of climate, soils, and forestry. More explicitly,
97 we expected the following effects (Table 1).

98 Hypothesis H1: the soil phosphorus availability, the most limiting nutrient in the sandy acidic soils of
99 the study region, would be decreased by intensive harvests (Achat et al. 2015a) and would be only partly
100 compensated by wood ash that has a low phosphorus content and low bioavailability (Steenari and
101 Lindqvist 1997; Fransson et al. 1999).

102 H2: the sequestration rate of C into forest biomass (estimated as the growth rate of the subsequent forest
103 stand) would be decreased by intensive harvests (Achat et al. 2009) because the local soils are nutrient
104 poor (Achat et al. 2015b).

105 H3: the size of the soil organic carbon content would be decreased by intensive harvests as for most
106 temperate forests (Achat et al. 2015a, b), but might remain unaffected by wood ash application as
107 observed in boreal forests (Augusto et al. 2008a).

108 H4: the soil would be acidified by intensive harvest, but partly neutralised by wood ash (Reid and
109 Watmough 2014; Achat et al. 2015a).

110 H5: because the local pine trees contain small amounts of trace metals in their biomass compared to
111 stocks in the soil (Saur et al. 1992; Trichet et al. 2018), exports through biomass harvest and returns
112 through wood ash application would not significantly change the trace metal content in the ecosystem.

113

114

115 **Materials and Methods**

116 *Study region and experimental site*

117 The *Landes de Gascogne* area is a flat region ~1.3 Mha in south-western France (between 43.5 and
118 45.5°N, and 1.5°W and 0.3°E). The climate is warm oceanic (mean annual temperature = 13°C, with
119 mean monthly values ranging from 6.0 to 20.3°C), usually with a dry season in summer and a wet season
120 in winter and/or in spring (annual rainfall is around 900 mm per year). Soils are sandy podzols (9–14%
121 fine sands; 80–89% coarse sands), which are classified as more or less hydromorphic and humic (*entic*
122 *podzol* in the WRB soil classification), or with a hardpan (*ortsteinic podzol*; Augusto et al. (2010)). They
123 developed in Aeolian deposits of quaternary sands (the *Sables des Landes*) and are characterised by their
124 coarse particle size and their high quartz content (Bertran et al. 2011). Soils are acidic and poor in
125 nutrients, particularly in phosphorus (P) and they have been identified as being among the poorest soils
126 in P in the world (Achat et al. 2009; Bredoire et al. 2016). The presence of wetlands is the consequence
127 of the low drainage capacity of the landscape (low elevation plain with streams but only one river) that
128 makes the watertable close to the soil surface (Deirmendjian et al. 2018). Initially grown for its resin,
129 Maritime pine (*Pinus pinaster* Ait.) is now used for wood production and is often P-fertilised at
130 plantation to alleviate the main nutritional limitation of the region (Trichet et al. 2009), and stands are
131 surrounded by ditches to improve water drainage in rainy periods.

132

133 *Experimental design and field procedures*

134 The area used for this study is called the *Forêt du Nezer* (44.57°N, 1.05°W; elevation = 21 m asl;
135 slope < 1%; entic podzol). The experimental site is a split-plot factorial design, crossing four different
136 intensities of *Biomass Harvesting* (BH factor in main plots) and four *Compensation Methods* by
137 applying wood ash or phosphorus fertilisation (CM factor in subplots nested in BH plots), resulting in
138 sixteen treatments replicated in three blocks (Figure S1). Contrary to the CM treatments, the BH
139 treatments had to be applied to large plots (main plots) because they required the use of large machines.
140 In April 2007, the standing forest (45-year-old, 465 trees ha⁻¹; basal area = 29.7 m² ha⁻¹) was clear-cut.
141 In practice, twelve large square plots, 100 × 100 m (1 ha large), were established in order to apply the
142 three blocks × four *Biomass Harvesting* treatments, described below:

- 143 1- Stem-Only Harvest ('SOH'; stem harvest down to 7 cm of diameter above bark) without any residue
144 harvest (exports = stems: this has been the local conventional practice for decades),
- 145 2- Aboveground Additional Harvest ('AAH'), which included stem harvest with additional harvesting
146 of branches with part of their foliage (see below; exports = stems + canopies; canopy = branches +
147 twigs + foliage),
- 148 3- Belowground Additional Harvest ('BAH'), which included stem (without branches, twigs, and
149 foliage) harvest with additional harvesting of stumps and roots (exports = stems + stumps + roots),
- 150 4- Whole-Tree Harvest ('WTH') with additional harvest of all other tree compartments (exports = stems
151 + branches + twigs + foliage + stumps + roots).

152 The conventional harvesting (clear-cut) was carried out in April 2007 (see Figure S2 for more details
153 about the study sequence), with particular attention to the slash (branches and needles) which was
154 gathered in each of the four 1 ha plots (Figure S3a). In June 2007, the branches were bundled, forwarded
155 from the six main plots (AAH-WTH plots × 3 blocks; Figure S3b-d), and finally transported to the
156 power plant. The time-lapse between this harvest and the clear-cut has enabled a partial fall of the foliage
157 (needles, in unquantified proportions). Then, in July-August 2007, the stumps were excavated with their
158 main roots, fragmented into a few pieces, and left on the ground in the six main plots (BAH-WTH
159 plots × 3 blocks; Figure S3e-g), enabling the natural removal of the soil particles by the rain. These
160 fragments were harvested in February 2008. Two years after clear-cutting, in January 2009, the area was

161 strip ploughed to a depth of 0.3 m. The ploughed strips were repeated every 4 meters and dedicated to
162 seedling plantation. The soil surface of the ploughed strips (representing ~half of the total surface area)
163 was flatten with discs before the plantation of 1,250 seedlings per hectare (4×2 m) of one-year-old
164 maritime pine seedlings of local origin.

165 Forty-eight small sub-plots, 50×50 m (0.25 ha), were established within the two-year-old plantation.
166 Four sub-plots of compensation methods (CM factor) were nested in every main plot (dedicated to a BH
167 treatment; Figure S1). The *Compensation Methods* were applied in July 2011 for the wood ash and in
168 December 2011 for the other fertilisers, as follows:

169 1- Control: no nutrient application ('C'),

170 2- Phosphorus fertilisation ('P'), considered as standard in this forest area (application of 60 kg-P₂O₅
171 per hectare in the form of superphosphate (30% P₂O₅)),

172 3- Phosphate-potassium fertilisation (the same P treatment as above with the addition of 60 kg-K₂O per
173 hectare in the form of potassium sulphate (50% K₂O)). It should be noted that this CM treatment was
174 not included in this study.

175 4- Wood ash application ('A') consisted of the application of 5 Mg of loose ash per hectare (provided
176 by a local power biomass plant). The ash composition is shown in the Table S1.

177

178 *Plant measurements and field sampling*

179 Estimates of biomass and nutrient exports (*Biomass Harvesting* factor):

180 The quantity of biomass, and associated nutrients, exported from the plots due the Biomass Harvest
181 treatments were estimated, based on a combination of measurements and models. In practice, before
182 clear-cutting the standing pine stand, all trees were identified and each diameter was measured at breast
183 height (≈ 1.3 m).

184 Biomass estimates

185 Then, we selected 14 trees that represented the range of stem size of the forest. All these trees were cut
186 down and studied intensively:

187 (i) the stem diameter was measured, and wood disks were sampled, at several heights (0 m, 0.5 m, 1.3 m,
188 and then every 2 m); the wet mass and the dry mass of the stem disks (wood and bark, separately) were
189 measured,
190 (ii) the diameter of all branches was measured at 10 cm from their insertion level,
191 (iii) the total dry mass of three branch classes (living branches, treetop [*i.e.* the upper part of the stem,
192 at diameter < 7 cm], dead branches) were weighed (all branches of each class were pooled),
193 (iv) the dry mass of 5 living branches (wood + bark + foliage) per tree was weighed, and
194 (v) the dry mass values of the stem wood, and the stem bark, were measured separately.
195 The total dry mass by compartment of each sampled tree was then calculated. The tree size
196 measurements (diameter at breast height) were used to estimate the aboveground biomass of the different
197 tree compartments using allometric relationships (unpublished FCBA models, calibrated with 45 trees
198 distributed in 5 other pine stands of a similar age). The estimated biomass values of the tree
199 compartments were consistent with the measured values, so the models were used to calculate the
200 biomass at the plot scale, based on the inventory of all trees.
201 The effective export rate of the different tree compartments was estimated in different ways, depending
202 on the compartment studied:
203 (i) stem wood was assumed to be harvested in totality (export rate = 100%),
204 (ii) stem bark harvests were quantified by collecting all the bark that remained on the floor of a given
205 area, after the stems exported from five other stands of the region (FCBA unpublished data; export rate
206 \approx 80%),
207 (iii) the effective exports of the canopies (*i.e.* branches + twigs + foliage) and of the tree belowground
208 (stump + roots) were estimated by direct measurements of the harvests realised. In practice, the
209 harvested canopies were gathered into \sim 2.5 m long bundles (n = 465 in total), labelled per plot, and
210 taken to the local power biomass plant. Bundles were weighed individually (fresh weight = 225-253 kg;
211 mean weight = 240 kg), and 20 bundles were taken at random to estimate their dry matter content (mean
212 dry content = 67% of the initial mass). All bundles were crushed to provide pellets for the power plant.
213 During the crushing stage, 36 samples of biomass were taken at regular time intervals to estimate the
214 dry matter content. This procedure enabled us to estimate the effective harvest rate of the canopy.

215 Conversely, it was not logistically feasible to measure the effective rates for foliage, twigs, and large
216 branches. Based on visual observations and on field studies (Stupak et al. 2008), we assumed that (i) the
217 twig harvest rate was 30% higher than that of the foliage, and (ii) the branch harvest rate was 50% higher
218 than that of the twigs. Finally, based on these ratios and on the measured harvest rate of the whole
219 canopy, the harvest rate values were adjusted. The values of the nutrient content in the different canopy
220 compartments (*i.e.* needles, twigs, branches, stembark, stemwood) were the mean values measured on
221 samples from the 45 trees used to build allometric relationships (see above). These mean values were
222 within the ranges commonly observed in the study region (Augusto et al. 2008b; Trichet et al. 2018).
223 A similar approach to that used for branches was used for the belowground biomass. In practice, the
224 potentially available belowground biomass (*i.e.* stump + roots) was estimated for each tree based on
225 allometric relationships that use the stem diameter at breast height as predictive variable (Augusto et al.
226 2015a). The realised belowground biomass harvest was estimated from the weight of all the fresh
227 biomass per plot (by weighing the trucks, with and without the transported biomass). The water content
228 of the biomass was measured on 36 samples in order to calculate the dry biomass (Figure S3g; sampling
229 at regular intervals during the crushing flow; mean value per plot of the dry content = 65-72%). These
230 samples were used also for nutrient analyses. The measured values of nutrient content in stumps and
231 roots were within the ranges commonly observed (Augusto et al. 2015a), except for the potassium
232 content that was higher in the present study (1.50 versus 0.99 mg g⁻¹).

233 All these measurements and analyses enabled us to quantify the standing biomass before the Biomass
234 Harvest treatments, and the effective exports due to harvests in terms of biomass and nutrients.

235

236 *Field measurements in the subsequent forest plantation*

237 Tree growth and needle sampling:

238 The growth rate of the new plantation –established after biomass harvest– was assessed simply by
239 measuring the standing biomass of the young trees. Indeed, because the survival rate of the planted
240 seedlings was nearly 100%, and not affected by experimental treatments, the stand density was the
241 almost the same in all subplots. The individual size of trees was estimated by measuring 50 pines, chosen
242 at random, per subplot. Tree height and stem circumference at breast height were measured in December

243 2015 and 2019 (*i.e.* on 7- and 11-year-old trees). The aboveground biomass was then estimated based
244 on allometric relationships (Vidal et al. 2019).

245 Needles were sampled to assess the nutritive status of trees, and their degree of contamination by trace
246 elements. For this, 8 pines were first selected at random in each subplot. Current-year needles were
247 collected in December 2019 using a telescopic pruner. Needles were collected in the upper third of the
248 canopy, where sunlight is always directly available, and in two opposite directions. The needles were
249 cleaned in a bath of demineralised water, pooled by subplot, and then dried in an oven for ten days at a
250 temperature of 40°C.

251

252 Understorey survey:

253 The understorey composition was assessed in May 2019, using the “*phytovolume*” approach (Gonzalez
254 et al. 2013; Vidal et al. 2019). In practice, the same pair of operators surveyed all the surface area of the
255 studied subplot, and then estimated the soil cover percentage of the main plant functional types (*i.e.* the
256 perennial herb *Molinia caerulea*, the bracken fern *Pteridium aquilinum*, ericaceous shrubs (*Calluna*
257 *vulgaris* and *Erica scoparia*), and the common gorse *Ulex europaeus*). The mean height of each plant
258 functional type was estimated based on three height measurements representative of the plant height
259 range in the surveyed area (Figure S4). The phytovolume value was computed as the product of the
260 ground cover (estimated in squares of 16 m²) and the plant height. Finally, the phytovolume value was
261 converted into estimates of plant aboveground biomass using dedicated allometric relationships that
262 were previously calibrated with destructive measurements of the standing biomass (Gonzalez et al. 2013;
263 Vidal et al. 2019).

264

265 Soil sampling:

266 In December 2019, a composite sample of the topsoil was built in each of the 36 subplots. To do this, a
267 systematic grid of 8 sampling points was used per subplot: 4 points on the tree ridges, and 4 points in
268 the furrows, to capture the spatial distribution of the topsoil microtopography induced by the regular
269 design of the strip ploughing (Forrester et al. 2013) and tree plantation and by subsequent operations of
270 understorey control (*i.e.* a bladed roller passing in the furrows). Because of the soil preparation before

271 tree plantation and because stands were still young, the forest floor layer was thin at soil sampling. We
272 consequently did not sample it, although we recognize its importance for plant nutrition (Jonard et al.
273 2009). After having gently removed the forest floor layer, we sampled the topsoil (0-15 cm layer) using
274 a corer (diameter = 8 cm). All samples were bulked in the field, and continuously homogenised by hand
275 until the composite sample showed no heterogeneity. The soil bulk density (kg L^{-1}) was estimated based
276 on a pedo-transfer function specifically calibrated for the local soils (Augusto et al. 2010).

277

278 *Laboratory analyses*

279 Plant analyses:

280 The needle samples were ground ($< 1\mu\text{m}$) using a mill in titanium to avoid contamination of the sample.
281 Aliquots of powdered samples (0.25g) were digested using a mixture of 1 ml HNO_3 and 4 ml H_2O_2 in
282 DigiPREP System in dry bath blocks. Then, the solution was filtered to $0.45\mu\text{m}$ and the concentrations
283 of K, Ca, Mg, Na, Fe, Mn, Al, Cr, Cu, Ni, Zn, and Co in the extracts were analysed by ICP-AES and
284 Cd, Pb, Tl, and Mo by ICP-MS. The validity and accuracy of the procedures were checked using a
285 standard reference material (SRM-1573a, NIST).

286

287 Soil characteristics:

288 Soils were sieved to $< 2\text{ mm}$ and air-dried. All analyses (except pH) were carried by INRAE soil testing
289 laboratory (INRAE-LAS, Arras, France), according to French standardised procedures or international
290 procedures. Soil pH was determined in a distilled water (1:5 ratio) and 0.01 M CaCl_2 extract (1:10 ratio;
291 standard NF ISO 10390:2005). The soil organic carbon content (SOC) and total nitrogen (N_{total}) were
292 determined by dry combustion after correction for carbonate (SOC: NF ISO 10694:1995, N_{total} : NF ISO
293 13878). The available phosphorus (P_{Olsen}) was determined using the Olsen method (NF ISO 11263).
294 The cation exchange capacity (CEC) and exchangeable cations (K^+ , Ca^{2+} , Mg^{2+} , Na^+ , Fe^{2+} , Al^{3+} , and
295 Mn^{2+}) were measured at soil pH using the cobaltihexamine chloride method (NF X 31–130:1999). The
296 soil texture was determined (NF X 31–107:1983), but only on three soil samples (composite samples
297 corresponding to the three blocks of the site).

298 Total soil mineral element contents (major and trace) were quantified after solubilisation by fluorhydric
299 and perchloric acids (NF X 31–147:1996). After complete dissolution, the concentrations of elements
300 were determined by ICP-AES and ICP-MS (see above).

301

302 Soil solution extraction and analyses:

303 Two chemical extractions were realised for the soils to estimate the availability of trace metals: ultra-
304 pure water and ammonium nitrate 1M (NH₄-NO₃). Water extractions make it possible to estimate the
305 content of elements that would be found in the soil solution. Ammonium nitrate 1 M extractions quantify
306 the elements weakly adsorbed onto the solid phase by ion exchange at pH=6.

307 Ultra-pure water extraction was performed at the 1:5 soil to solution ratio, stirring on a roller-shaking
308 table for one hour. At the end of the shaking, the solution was filtered (0.22 µm) and then part of the
309 solution was acidified at 2% with HNO₃ for trace metal analysis. The second part was reserved at 4°C
310 to measure: pH, the concentration of dissolved organic (DOC) and inorganic (IC) carbon (by oxidative
311 combustion; TOC-VCSH, Shimadzu), and anions (PO₄³⁻, NO₂⁻, NO₃⁻) and NH₄⁺ (colorimetric method;
312 Technicon auto analyser II).

313 The NH₄-NO₃ extractions were performed at 1:2.5 soil to solution ratio (Symeonides and McRae 1977).
314 The suspensions were shaken, filtered at 0.22 µm, and then acidified with 2 % HNO₃. The concentrations
315 of P, K, Ca, Mg, Na, Fe, Mn, and Zn in extracts from two extractions were assayed by ICP-AES and Cd
316 and that of Pb and Cd by GF-AAS.

317

318 *Input-output budgets and data analyses*

319 Estimating input-output budgets enables to evaluate if a forest management is sustainable (Ranger and
320 Turpault 1999). However, because our study was not designed to quantify biogeochemical fluxes, the
321 presented budgets should be considered with caution as they are only rough estimates, and further
322 research is needed to assess the consequences of intensive forestry on ecosystem biogeochemistry. In
323 practice, atmospheric deposition was estimated based on data from a nearby monitoring site (Croise et
324 al. 2005), and values were adjusted to take into account the tree height effect on deposition (de Schrijver
325 et al. 2012). Symbiotic nitrogen fixation at 11 years-old was estimated based on gorse biomass (see

326 above), and taking into account the nitrogen content and fixation rate of this species (Cavard et al. 2007;
327 Augusto et al. 2009). For periods after 11 years-old, which corresponds to the canopy closure of pine
328 stands, a flux a $2.8 \text{ kg}_N \text{ ha}^{-1} \text{ yr}^{-1}$ was retained (Augusto et al. 2005). The flux of nutrients released by
329 weathering of soil minerals was estimated based on soil mineralogy and the PROFILE model (Sverdrup
330 et al. 2006). Elements that were added to the ecosystem through application of wood ash and fertiliser
331 were quantified based on ash composition (Table S1) and fertiliser composition (unpublished data and
332 Kabata-Pendias 2000). Exports of nutrients through biomass harvest were estimated by direct
333 measurements (see above). Losses of nutrients through deep seepage were estimated based on an on-
334 going monitoring experiment (XyloSylve) with the same tree species and similar tree age, soil, climate,
335 and experimental treatments. Because input-output budgets should be calculated for a complete
336 silvicultural rotation (Kimmins 1974), we calculated them not only for a stand age of 11 years-old (when
337 we sampled plants and soil) but also at 20 and 40 years after plantation as it corresponds to the local
338 rotation duration for intensive and conventional management, respectively. We did not present values
339 of soil nutrient stocks, however it is possible to estimate soil pools (in $\text{kg}_{\text{element}} \text{ ha}^{-1}$) by multiplying the
340 soil content (in $\text{mg}_{\text{element}} \text{ g}^{-1}$, which is equivalent to $\text{kg}_{\text{element}} \text{ Mg}^{-1}$) by 1578 (which is the mean value of
341 soil mass of the 0-15 cm layer; in $\text{Mg}_{\text{soil}} \text{ ha}^{-1}$; (Augusto et al. 2010)).

342 Data processing was performed with R, version 3.6.1. (R Core Team 2019). The split-plot design was
343 analysed by a mixed effect model (*lme* function, package nlme 3.1-143). The biomass harvest factor and
344 the compensation method factor were fixed effects (main and interaction effects). The block and biomass
345 harvest factor nested in blocks were random effects associated with the intercept. When detected,
346 heteroscedasticity was corrected by modelling the variance with the *varIdent()* function (nlme package)
347 which allows different variances by level of the biomass harvest and compensation methods. Least-
348 squares means were calculated (emmeans package 1.4.2) and linear contrasts were used to test for
349 significant differences between treatments and their corresponding controls, namely SOH for the
350 biomass harvest intensity (BH factor) and no fertilisation (C) for the compensation methods (CM factor).
351 When necessary, the $P=0.05$ significance probability was adjusted for multiple pairwise
352 comparisons (Tukey's test).

353 Preliminary analyses of the results showed that the interactions between the BH factor and the CM factor
354 were almost never statistically significant, as already been reported (Hagerberg and Wallander 2002),
355 and that is why we analysed these factors separately.

356 Because of the lack of significant interactions between the two studied factors, the mean values of the
357 Biomass Harvest treatments (n = 3 blocks = 3 replicates) pooled the values of the Compensation
358 Methods subplots. Similarly, the mean values of the Compensation Methods treatments corresponded
359 to 12 replicates (3 blocks × 4 BH treatments). The difference of number of replicates (3 versus 12), due
360 to the split-plot design, implied that the statistical power is greater for the CM factor nested in BH
361 compared to the BH factor. The main consequence of this split-plot design was consequently a greater
362 difficulty in detecting significant effects of the BH factor.

363 Mean values are shown ± one standard error. The mean values of the main response variables are shown
364 in Appendix 1, for the all 12 experimental combinations of treatments (4 BH × 3 CM). In addition, all
365 data and scripts are available at <https://doi.org/10.15454/LCU6OZ>.

366

367

368 **Results**

369 *Initial stand biomass and nutrient content, and effective harvests*

370 The standing biomass of the stand prior to harvests was $179.1 \pm 3.5 \text{ Mg ha}^{-1}$ of which $40.1 \pm 0.7 \text{ Mg ha}^{-1}$
371 ¹ were belowground (Table S2). The standing biomass was distributed homogeneously among the
372 experimental plots, with a coefficient of variation of 1% among the WTH plots to 8% among the SOH
373 plots (CV=6% at the site scale). This biomass contained $24.7 \pm 0.5 \text{ kg-P ha}^{-1}$ (phosphorus) and $311 \pm$
374 6 kg-N ha^{-1} (nitrogen; Table S2). The harvesting treatments did not export all this biomass and its
375 associated nutrients: the effective rate of harvest was measured as 71% for belowground biomass (in
376 BAH and WTH treatments) and 91% for aboveground biomass (in AAH and WTH treatments). Within
377 the aboveground biomass pool, the measured harvest rate of canopies (foliage + twigs + branches) was
378 53%, with estimated rates of 34%, 44%, and 66% for foliage, twigs and branches, respectively. The
379 realised export of biomass was $115.32 \text{ Mg ha}^{-1}$ for the conventional harvest (SOH), with moderately
380 increased values for more intensive harvests (from +10% for AAH to +34% for WTH; Figure 1a).

381 Despite their low additional biomass exports, due to the differences in nutrient contents between tree
382 parts, the non-conventional harvest practices exported much more nutrients than the conventional SOH
383 technique. This was particularly the case for the canopy export of nitrogen (+74% in AAH; Figure 1b)
384 and the belowground export of potassium (+123% in BAH). In the most impacting treatment (WTH),
385 the increase in nutrient exports was between +64% for P (Figure 1c) and +145% for N and K (Figure 1b
386 and 1d).

387

388 *Soil properties and fertility (hypothesis H1)*

389 The main soil characteristics were representative of the sandy podzols of the region. For instance, the
390 mean soil particle size distribution showed the dominance of sand ($85.7\pm 0.2\%$, $7.5\pm 0.5\%$, $0.7\pm 0.3\%$,
391 $2.4\pm 0.2\%$ and $3.6\pm 0.3\%$, for the coarse sand, fine sand, coarse silt, fine silt and clay fractions,
392 respectively), and a low initial value of the total soil P content of $44\pm 17 \mu\text{g-P g}^{-1}$ (see Augusto et al.
393 (2010) for comparisons).

394 The biomass harvesting treatments had many effects on the soil nutritive status, but the effects depended
395 on the form of the nutrients and micronutrients. While the AAH treatment showed increases in values
396 of K, Na, Fe and Mn total contents (Table 2), the pattern was quite different for exchangeable
397 cations (Table 3) and extractable nutrients (Table 4). Indeed, the nutrients that constitute the so-called
398 *basic cations* of the CEC (*i.e.* K^+ , Ca^{2+} , Mg^{2+} , and Na^+) were depleted to varying extents (except for K^+)
399 by the non-conventional harvest practices (AAH, BAH, and WTH) as compared with the SOH
400 treatment (Table 3). The same pattern applied to Ca, Mg, and to a lesser extent K, that were $\text{NH}_4\text{-NO}_3$
401 extracted (Table 4). The losses in these nutrients (Ca, Mg, Mn, and Zn) were generally compensated by
402 the application of wood ash (Table S1), as shown by the significant increases in their soil content as
403 compared with the control treatment (Tables 3 and 4).

404 The effects of the treatments on P also depended on the chemical form considered. The soil P content
405 as measured using the sodium bicarbonate extractant ($\text{P}_{\text{-Olsen}}$) showed no significant differences, even if
406 the non-conventional harvest treatments resulted in a decreasing trend in this soil content (Table 2). This
407 not-significant trend became significant when considering the P content as measured using the $\text{NH}_4\text{-}$

408 NO₃ extractant (Table 4). As expected, the application of wood ash did not significantly modify the soil
409 P status. Surprisingly the P-fertiliser treatment did not improve this P status either.

410 The soil N_{-total} content followed the same, but insignificant, trend as for P with low values in the
411 treatments that involved canopy harvesting (Table 2). The C/N ratio was remarkably irresponsive to all
412 experimental treatments, but the soil N content was reduced by wood ash application.

413

414 *Vegetation response (H2)*

415 The total aboveground biomass of the spontaneous vegetation remained unaffected by the *Biomass*
416 *Harvesting* treatments, but changed due to the *Compensatory Methods* treatments with an increasing
417 standing biomass following the ranking: Control ≤ P fertilisation ≤ Ash (Figure 2). For both factors, the
418 proportion of the prominent plant species was influenced by treatments. For the *Biomass Harvesting*
419 factor (BH), the proportion of species adapted to oligotrophic-acidic moorlands (*i.e. Molinia caerulea*
420 and ericaceous species) was increased by the intensive harvests as compared with the conventional SOH
421 treatment (Figure 2). The opposite pattern was observed in the wood ash treatment (*Compensation*
422 *Methods* factor, CM), which strongly decreased the abundance of those moorland-adapted species. The
423 wood ash application and, to a lesser extent, the P fertilisation treatments enhanced the growth of the N-
424 fixers (almost entirely composed of the spiny shrub *Ulex europaeus*).

425 Measuring the trees at 7 years-old suggested that treatments did not affect the initial stand growth as no
426 significant differences were observed (data not presented). At 11 years-old, the most severe treatments
427 of the *Biomass Harvesting* factor (*i.e.* AAH and WTH) tended to depress the tree growth (field
428 observations and Figure 3), but these trends remained statistically not-significant. Conversely we
429 measured an unexpected significant difference between two treatments of the *Compensatory Methods*
430 factor, with a negative effect of P fertilisation, and a positive trend of wood ash application (Figure 3;
431 Table S3). The wood ash trend can be related to the improvement of the nutritional status of foliage (N,
432 P, K, Mg, B, Cu, and Fe; Table 5).

433

434 *Soil organic matter and CEC (H3)*

435 The soil Cation Exchange Capacity (CEC) was strongly and linearly correlated with the quantity of
436 elements that are commonly found as organic matter in acidic soils, which are total nitrogen (N_{total}) and
437 organic carbon (SOC; $CEC = 0.088 \times SOC$; $r^2 = 0.76$). Due to the tight proportional relationship
438 between CEC and SOC, the observed effects of the experimental treatments on these variables were
439 quite similar. The harvest of the belowground biomass and the application of wood ash both tended to
440 decrease the soil organic matter content and the soil CEC (Tables 2 and 3). The dissolved organic carbon
441 (DOC) followed a similar pattern with a significant decrease in concentration after whole-tree harvesting
442 (WTH) and wood ash application (Table 2).

443

444 *Soil acidity status (H4)*

445 Harvesting more biomass than conventionally practiced did not modify any of the soil variables used to
446 assess the acidic status of a soil (pH-H₂O, pH-CaCl₂, Base Saturation of the CEC; Table 3). The absence
447 of effect on the Base Saturation was the consequence of the concomitant decrease in *basic cations* (K^+ ,
448 Ca^{2+} , Mg^{2+} , and Na^+) and CEC. Conversely to biomass harvesting, applying wood ash modified the soil
449 acidity status. The soil pH-H₂O value increased by +0.24 unit, and the Base Saturation increased from
450 39% to 46% of the CEC (Table 3). The P fertilisation had no effect on soil acidity.

451

452 *Ecosystem content in trace metals (H5)*

453 Overall, the *Biomass Harvesting* treatments generally had no significant effect on the trace elements
454 (As, Cd, Mo, Ni, Pb, Tl, and Zn) in the ecosystem (Tables 2, 4, 5, 6, S4). The exceptions were a decrease
455 in soil Pb and Zn contents due to WTH.

456 The *Compensatory Methods* also had only a few significant effects on the trace element distribution in
457 the ecosystem (Tables 2, 4, 5, 6, S4). The most noticeable exceptions were increases of soil Zn content
458 due to wood ash application (Tables 4 and S4), an effect of P fertilisation on Cd distribution between
459 the soil (decrease of Cd soil content) and trees (increase of Cd foliage content), and an improvement of
460 Cu content of tree foliage after P or wood ash application (Table 6). This lack of effects can be explained
461 by comparing the applied amounts with the initial soil stocks: for Fe, Mn, and Pb, the doses were 1-2 %

462 of the initial stocks (0-15 cm soil layer). For Cd and Zn, the doses applied represented 8-10% of the
463 initial stocks (Cd: 8% and 10% by fertilisation and wood ash, respectively; Zn: 8% by ash).

464

465 *Long term perspective: estimates of input-output budgets*

466 The effects of treatments were highly nutrient-dependent. For nitrogen, the budget values were positive
467 or close to equilibrium for the SOH, but were negative at 11 years for the other levels of biomass
468 harvest (Figure S5). The budget values became positive over time, mainly due to atmospheric deposition
469 and nitrogen symbiotic fixation. Applying P-fertiliser or wood ash directly improved budgets by
470 enhancing the growth of the spontaneous N-fixers (Figure 2). In the case of phosphorus, because most
471 external sources of P were negligible (*i.e.* $< 1 \text{ kg ha}^{-1} \text{ yr}^{-1}$), only applying P-fertiliser enabled to balance
472 the biomass harvests.

473 The BAH and WTH treatments had a strong impact on K input-output budgets because the belowground
474 biomass of trees was particularly rich in this nutrient. In these cases, applying wood ash and keeping
475 long silvicultural rotations enabled to balance the exports (Figure S5). The same pattern was observed
476 for calcium and magnesium but to different extents, depending on the biomass nutrient content and the
477 compensatory fluxes that are atmospheric deposition and mineral weathering.

478

479 **Discussion**

480 *Importance of the study design and the study region for data interpretation*

481 Implementing a careful harvest of adult tree compartments over small areas was hardly feasible because
482 the machines that collect stumps or canopies need space to work (Figure S3). Consequently, the *Biomass*
483 *Harvesting* factor could be studied only by using large plots (*i.e.* one ha per plot). With 16 experimental
484 treatments and three blocks, building a complete factorial experimental design (*i.e.* with 48 large plots)
485 was not possible because the surface area of 48 ha required was incompatible with the premise of an
486 area that was initially homogeneous in its properties and past-management. Therefore, we built our
487 experiment based on a split-plot design. Such a design had one major consequence on data analysis.
488 Indeed, the number of the subplots –dedicated to the *Compensation Methods* treatments– were four-fold
489 more numerous than the experimental areas dedicated to *Biomass Harvesting* treatments (Figure S1).

490 The power of the statistical tests was consequently much lower for the *Biomass Harvesting* factor, so
491 isolating significant effects was sometimes difficult.

492 The forest region where our experiment took place is atypical from a biogeochemical perspective as the
493 soils are extremely poor in phosphorus. While in natural systems the soil P_{total} content ranges between
494 ~10 and 2,000 µg g⁻¹ (Achat et al. 2009; Yang and Post 2011; Augusto et al. 2017), the study region has
495 a mean value of ~50 µg g⁻¹ and is within a range of ~10-100 µg g⁻¹ (Achat et al. 2009; Augusto et al.
496 2010). Being also acidic and poor in other nutrients and micronutrients (Augusto et al. 2010; Trichet et
497 al. 2018), the soils of the study region have been identified as having the highest level of vulnerability
498 to biomass exports according to the national system of forest evaluation (Durante et al. 2019). Poorest
499 of the poor, the study region is also expected to respond strongly and negatively to additional harvests
500 of biomass, which implies that results should be extrapolated to other regions with caution.

501

502 *Consequences of harvesting more biomass on the biogeochemistry of an oligotrophic forest*

503 Exporting tree canopies generally enhances the amount of harvested biomass (~ +20% to +50%), but
504 this is at the expense of losing huge quantities of nutrients (~ +100% to +250%; (Achat et al. 2015a)).
505 Our results (10-34% extra biomass, and 37-145% extra nutrients) fit well with this general pattern. On
506 the other hand, the quantities of exported nutrients of this field experiment are about half of the values
507 used in previous modelling studies (Augusto et al. 2015b; Achat et al. 2018). This was the direct
508 consequences of the harvesting techniques that were purposely used to reduce the losses of nutrients:
509 letting branches shed their needles (Stupak et al. 2008), and harvesting only stumps and coarse roots
510 (Augusto et al. 2015a). Despite these precautions, harvesting more biomass, and above all harvesting
511 the whole-tree, reduced the soluble pools of nutrients (K, Ca, Mg, and P) in the soil (-15% to -58%;
512 Table 4). Additional biomass harvests likely also impacted the soil content in organic matter, with
513 negative ripple effects on organic P and CEC, the latter being entirely dependent on organic matter in
514 these sandy soils (Augusto et al. 2010). In the study region the organic forms of P are important for
515 seedling and tree nutrition (Jonard et al. 2009; Achat et al. 2013), which may explain why we observed
516 a depressive –but not significant– effect of harvesting canopies on tree growth in the subsequent stand

517 (see Table S3). A longer monitoring is needed to confirm the observed trend. The absence of change in
518 the tree foliage composition might be considered as surprising as foliage nutrient content is often used
519 to assess plant nutrition (CSIRO 1997; Mellert and Göttlein 2012). Nevertheless, maintaining the
520 nutrient content of foliage at the expense of growth is a response commonly observed after the harvest
521 of additional biomass (Achat et al. 2015a), and our results are consistent with this general pattern.

522

523 *Fertilisation and wood ash application as possible mitigation practices*

524 The major effect of applying wood ash was expected to be a change in the acidity status of soils (Reid
525 and Watmough 2014). Our experiment was not an exception and the soil pH, base saturation and
526 concomitant available calcium and magnesium, were improved by wood ash application (Gomez-Rey
527 et al. 2013). The available soil micronutrient contents of Mn and Zn also logically increased because
528 wood ash contains them in substantial quantities (Table S1).

529 Whereas the effects of wood ash were expected and straight forward to interpret, the P fertilisation had
530 surprising outcomes: the soil P status was not improved, and the soil contents of available Ca and Mg
531 actually decreased by P fertilisation. These results may be explained by the tree foliage composition that
532 was modified by both P fertilisation and wood ash application, with increases in concentration values
533 for most nutrients and micronutrients (and even some non-essential trace metals). Hence, it seemed that
534 the two compensatory methods improved tree nutrition but, for P fertilisation, it was at the expense of
535 the soil reserves in Ca and Mg.

536 In turn, these changes in the soil properties modified the dynamics of the spontaneous vegetation and of
537 the tree plantation. The vegetation experienced a shift of species prominence after the wood ash
538 application, with a decrease in the species typical of acidic moorlands and heathlands (*i.e. Molinia*
539 *caerulea* and ericaceous species) to the benefit of the local main N-fixer (the shrub *Ulex europaeus*).
540 The decrease of the abundance of the acidophilus species is logical since wood ash decreased the soil
541 acidity. The increase of the N-fixer shrub was expected as we are used to observing a positive response
542 of this leguminous species to P fertilisation (*e.g.* Delerue et al. (2015)), but although the N-fixers
543 abundance was increased by P fertilisation, it was relatively low compared with previous experiments
544 (Augusto et al. 2005; Vidal et al. 2019), and lower than results after the wood ash application. This ash

545 effect may be the positive consequence of reducing the ambient acidity on the symbionts of the
546 leguminous species (Slattery and Coventry 1995).

547 Trees that were planted responded differently to the *Compensatory Methods*. Firstly, trees that received
548 wood ash tended to grow the fastest. Since wood ash application improved the availability of many
549 nutrients and micronutrients, it suggests that plant growth in this site was limited by several elements at
550 the same time. On the other hand, the depressive effect of P fertilisation was surprising considering that
551 this practice has proven its efficiency in the study region for more than half a century (Trichet et al.
552 2009). As explained above, it seems that P fertilisation improved the tree foliage composition at the
553 expense of soil fertility, and we speculate that this soil impoverishment might be at the origin of
554 subsequent degraded tree growth. However, this atypical phenomenon requires further investigation.

555 Considering the positive effects of wood ash application on soil acidity and nutrient content, and its
556 positive repercussion on tree growth, one might conclude that this compensatory method should be
557 promoted in oligotrophic forests submitted to intensive biomass harvestings. Nevertheless, this
558 conclusion does not take into account one major drawback of wood ash application in our experiment,
559 which was a decrease of the soil organic matter content, with negative consequences on organic carbon
560 and total nitrogen contents and soil exchange capacity (losses \approx 7-11%, after correcting for the change
561 in soil bulk density). We interpret this decrease as the result of the acidity alleviation by wood ash, which
562 probably enhanced the soil microbial activity and respiration (Bååth and Arnebrant 1994; Jokinen et al.
563 2006; Omil et al. 2013). In a meta-analysis, we previously concluded that wood ash had no influence on
564 soil organic carbon (Augusto et al. 2008a), but this study was almost entirely based on Nordic, or cold
565 temperate experiments (mean annual temperature $<$ 8.5°C; e.g. Feldkirchner et al. (2003)). Under a
566 warm temperate climate, wood ash application in sites with a long history of forest occupation (*i.e.* high
567 SOC values and often low pH values) can decrease the soil content in organic carbon (Solla-Gullon et
568 al. 2006).

569

570 *Initial expectations and future anticipation*

571 Most, but not all, of our initial expectations were confirmed by the field experiment. Firstly, as
572 anticipated, harvesting more biomass aggravated the P scarcity of the local soils, and the application of

573 wood ash had only a minor influence on the soil P content (Hypothesis H1; Table S5). Still in line with
574 our expectations, the impoverishment of the soil caused by additional harvests of biomass could result
575 in reduced growth of the subsequent forest stand; wood ash application had positive effects on early tree
576 growth (H2). The fact that, unexpectedly, P application did not enable trees to overcome P limitation
577 suggested that other nutritional limitations –such as Ca and Mg– prevented the experimental fertilisation
578 from having an effect. If our two first hypotheses were fairly well supported by results, the third and
579 fourth hypotheses received mixed support. Indeed, both high rates of biomass harvesting and wood ash
580 application decreased the pool of soil organic carbon, but only the former treatment was supposed to do
581 so (H3). Similarly, while wood ash application logically improved the acidity status of the soil to a
582 moderate extent, intensive biomass exports did not deteriorate it, as in many other studies (H4; see above
583 for possible explanations). Finally, as expected, none of the experimental factors had major effects on
584 the trace metal distribution within the ecosystem (H5).

585 Overall, this factorial experiment showed that exporting more forest biomass through the additional
586 harvests of tree canopies, stumps, and roots had negative consequences on soil properties. Additional
587 harvests have aggravated the poverty of the already oligotrophic soil, which in turn may decrease tree
588 growth and the soil content of organic carbon in the future. Importantly, applying nutrients as fertiliser
589 or wood ash did not fully compensate for the negative impact of biomass exports. Indeed, our estimates
590 of input-output budgets suggests that, in addition to applying compensatory treatments, maintaining
591 quite long silvicultural rotations is more suitable to ensure the sustainability of the forest management.

592

593

594 **Acknowledgements**

595 We are grateful to the ADEME, GIS G-PMF and the Council of the Nouvelle-Aquitaine region for their
596 financial support (D.V. PhD grant; SYLVOGENE and PINASTER projects). Similarly, we thank the
597 ISPA lab for providing the F.B. Master grant. We are grateful to the Forest Experimental Facility
598 (UEFP-<https://doi.org/10.15454/1.5483264699193726E12>) for their assistance during the successive
599 field campaigns. We thank Sylvie Bussière, Cécile Coriou, and Catherine Lambrot for their help during

600 the sample analyses. Finally, we thank Jean-Luc Denou and Pierre Bordenave for providing data from
601 the XyloSylve monitoring experiment.

602

603

604 **References**

605 Achat DL, Bakker MR, Augusto L, et al (2009) Evaluation of the phosphorus status of P-deficient
606 podzols in temperate pine stands: combining isotopic dilution and extraction methods.

607 *Biogeochemistry* 92:183–200. doi: 10.1007/s10533-008-9283-7

608 Achat DL, Bakker MR, Augusto L, Morel C (2013) Contributions of microbial and physical-chemical
609 processes to phosphorus availability in Podzols and Arenosols under a temperate forest. *Geoderma*

610 211:18–27. doi: DOI 10.1016/j.geoderma.2013.07.003

611 Achat DL, Deleuze C, Landmann G, et al (2015a) Quantifying consequences of removing harvesting
612 residues on forest soils and tree growth—A meta-analysis. *For Ecol Manage* 348:124–141. doi: doi:

613 10.1016/j.foreco.2015.03.042

614 Achat DL, Fortin M, Landmann G, et al (2015b) Forest soil carbon is threatened by intensive biomass
615 harvesting. *Sci Rep* 5:15991

616 Achat DL, Martel S, Picart D, et al (2018) Modelling the nutrient cost of biomass harvesting under
617 different silvicultural and climate scenarios in production forests. *For Ecol Manage* 429:642–653.

618 doi: 10.1016/j.foreco.2018.06.047

619 Alvarez-Alvarez P, Pizarro C, Barrio-Anta M, et al (2018) Evaluation of tree species for biomass energy
620 production in Northwest Spain. *FORESTS* 9:. doi: 10.3390/f9040160

621 Andre F, Jonard M, Ponette Q (2010) Biomass and nutrient content of sessile oak (*Quercus petraea*
622 (Matt.) Liebl.) and beech (*Fagus sylvatica* L.) stem and branches in a mixed stand in southern

623 Belgium. *Sci Total Environ* 408:2285–2294. doi: DOI 10.1016/j.scitotenv.2010.02.040

624 Aronsson KA, Ekelund NGA (2004) Biological effects of wood ash application to forest and aquatic
625 ecosystems. *J Environ Qual* 33:1595–1605. doi: 10.2134/jeq2004.1595

626 Augusto L, Achat DL, Bakker MR, et al (2015a) Biomass and nutrients in tree root systems—sustainable
627 harvesting of an intensively managed *Pinus pinaster* (Ait.) planted forest. *Glob Chang Biol*

- 628 Bioenergy 7:231–243. doi: doi: 10.1111/gcbb.12127
- 629 Augusto L, Achat DL, Jonard M, et al (2017) Soil parent material - A major driver of plant nutrient
630 limitations in terrestrial ecosystems. *Glob Chang Biol* 23:3808–3824
- 631 Augusto L, Bakker MR, De Lavaissiere C, et al (2009) Estimation of nutrient content of woody plants
632 using allometric relationships: quantifying the difference between concentration values from the
633 literature and actuals. *Forestry* 82:463–477. doi: 10.1093/forestry/cpp019
- 634 Augusto L, Bakker MR, Meredieu C (2008a) Wood ash applications to temperate forest ecosystems -
635 potential benefits and drawbacks. *Plant Soil* 306:181–198. doi: 10.1007/s11104-008-9570-z
- 636 Augusto L, Bakker MR, Morel C, et al (2010) Is “grey literature” a reliable source of data to characterize
637 soils at the scale of a region? A case study in a maritime pine forest in southwestern France. *Eur J*
638 *Soil Sci* 61:807–822
- 639 Augusto L, Crampon N, Saur E, et al (2005) High rates of nitrogen fixation of *Ulex* species in the
640 understory of maritime pine stands and the potential effect of phosphorus fertilization. *Can J For*
641 *Res Can Rech For* 35:1183–1192. doi: 10.1139/x05-054
- 642 Augusto L, De Schrijver A, Vesterdal L, et al (2015b) Influences of evergreen gymnosperm and
643 deciduous angiosperm tree species on the functioning of temperate and boreal forests. *Biol Rev*
644 90:444–466
- 645 Augusto L, Meredieu C, Bert D, et al (2008b) Improving models of forest nutrient export with equations
646 that predict the nutrient concentration of tree compartments. *Ann For Sci* 65:. doi:
647 10.1051/forest:2008059
- 648 Bååth E, Arnebrant K (1994) Growth rate and response of bacterial communities to pH in limed and ash
649 treated forest soils. *Soil Biol Biochem* 26:995–1001
- 650 Banos V, Dehez J (2017) Le bois-énergie dans la tempête, entre innovation et captation ? Les nouvelles
651 ressources de la forêt landaise. *Natures Sci Sociétés* 25:122–133
- 652 Bertran P, Bateman MD, Hernandez M, et al (2011) Inland aeolian deposits of south-west France: facies,
653 stratigraphy and chronology. *J Quat Sci* 26:374–388
- 654 Bredoire F, Bakker MR, Augusto L, et al (2016) What is the P value of Siberian soils? Soil phosphorus
655 status in south-western Siberia and comparison with a global data set. *Biogeosciences* 13:2493–

- 656 2509. doi: 10.5194/bg-13-2493-2016
- 657 Bright RM, Cherubini F, Astrup R, et al (2012) A comment to “Large-scale bioenergy from additional
658 harvest of forest biomass is neither sustainable nor greenhouse gas neutral”: Important insights
659 beyond greenhouse gas accounting. *Glob Chang Biol Bioenergy* 4:617–619
- 660 Cavard X, Augusto L, Saur E, Trichet P (2007) Field effect of P fertilization on N₂ fixation rate of *Ulex*
661 *europaeus*. *Ann For Sci* 64:875–881. doi: 10.1051/forest:2007066
- 662 Croise L, Ulrich E, Duplat P, Jaquet O (2005) Two independent methods for mapping bulk deposition
663 in France. *Atmos Environ* 39:3923–3941
- 664 CSIRO (1997) Plant analysis: an interpretation manual. CSIRO Publishing, Reuter, D. J. and Robinson,
665 J. B. (Eds), Collingwood, Australia
- 666 de Schrijver A, de Frenne P, Staelens J, et al (2012) Tree species traits cause divergence in soil
667 acidification during four decades of postagricultural forest development. *Glob Chang Biol*
668 18:1127–1140
- 669 Deirmendjian L, Loustau D, Augusto L, et al (2018) Hydro-ecological controls on dissolved carbon
670 dynamics in groundwater and export to streams in a temperate pine forest. *Biogeosciences* 15:669–
671 691
- 672 Delerue F, Gonzalez M, Michalet R, et al (2015) Weak Evidence of Regeneration Habitat but Strong
673 Evidence of Regeneration Niche for a Leguminous Shrub. *PLoS One* 10:. doi:
674 10.1371/journal.pone.0130886
- 675 Durante S, Augusto L, Achat DL, et al (2019) Diagnosis of forest soil sensitivity to harvesting residues
676 removal - A transfer study of soil science knowledge to forestry practitioners. *Ecol Indic* 104:512–
677 523. doi: 10.1016/j.ecolind.2019.05.035
- 678 Feldkirchner DC, Wang C, Gower ST, et al (2003) Effects of nutrient and paper mill biosolids
679 amendments on the growth and nutrient status of hardwood forests. *For Ecol Manage* 177:95–116.
680 doi: 10.1016/S0378-1127(02)00318-3
- 681 Forrester DI, Pares A, O’hara C, et al (2013) Soil organic carbon is increased in mixed-species
682 plantations of *Eucalyptus* and nitrogen-fixing *Acacia*. *Ecosystems* 16:123–132
- 683 Fransson AM, Bergkvist B, Tyler G (1999) Phosphorus solubility in an acid forest soil as influenced by

- 684 form of applied phosphorus and liming. *Scand J For Res* 14:538–544. doi:
685 10.1080/02827589950154014
- 686 Garcia W de O, Amann T, Hartmann J (2018) Increasing biomass demand enlarges negative forest
687 nutrient budget areas in wood export regions. *Sci Rep* 8:. doi: 10.1038/s41598-018-22728-5
- 688 Gomez-Rey MX, Madeira M, Coutinho J (2013) Soil C and N dynamics, nutrient leaching and fertility
689 in a pine plantation amended with wood ash under Mediterranean climate. *Eur J For Res* 132:281–
690 295. doi: 10.1007/s10342-012-0674-x
- 691 Gonzalez M, Augusto L, Gallet-Budynek A, et al (2013) Contribution of understory species to total
692 ecosystem aboveground and belowground biomass in temperate *Pinus pinaster* Ait. forests. *For*
693 *Ecol Manage* 289:38–47. doi: DOI 10.1016/j.foreco.2012.10.026
- 694 Haberl H, Schulze ED, Korner C, et al (2013) Response: complexities of sustainable forest use. *Glob*
695 *Chang Biol Bioenergy* 5:1–2
- 696 Hagerberg D, Wallander H (2002) The impact of forest residue removal and wood ash amendment on
697 the growth of the ectomycorrhizal external mycelium. *FEMS Microbiol Ecol* 39:139–146. doi:
698 10.1016/S0168-6496(01)00207-0
- 699 Hannam KD, Venier L, Allen D, et al (2018) Wood ash as a soil amendment in Canadian forests: what
700 are the barriers to utilization? *Can J For Res* 48:442–450. doi: 10.1139/cjfr-2017-0351
- 701 IPCC (2014) *Climate Change 2014: Synthesis Report. Fifth Assessment Report*. Geneva, Switzerland
- 702 Jokinen H, Kiiikkilä O, Fritze H (2006) Exploring the mechanisms behind elevated microbial activity
703 after wood ash application. *Soil Biol Biochem* 38:2285–2291
- 704 Jonard M, Augusto L, Morel C, et al (2009) Forest floor contribution to phosphorus nutrition:
705 experimental data. *Ann For Sci* 66:510. doi: 10.1051/forest/2009039
- 706 Jurevics A, Peichl M, Olsson BA, et al (2016) Slash and stump harvest have no general impact on soil
707 and tree biomass C pools after 32-39 years. *For Ecol Manage* 371:33–41. doi:
708 10.1016/j.foreco.2016.01.008
- 709 Kabata-Pendias A (2000) *Trace elements in soils and plants*. CRC Taylor and Francis Group, London
710 NewYork
- 711 Kimmins JP (1974) *Sustained yield, timber mining, and the concept of ecological rotation, a British*


- 712 Columbian view. For Chron Feb 1974:27–31
- 713 Lindner M, Karjalainen T (2007) Carbon inventory methods and carbon mitigation potentials of forests
714 in Europe: a short review of recent progress. Eur J For Res 126:149–156
- 715 Luyssaert S, Marie G, Valade A, et al (2018) Trade-offs in using European forests to meet climate
716 objectives. Nature 562:259–262
- 717 Mellert KH, Göttlein A (2012) Comparison of new foliar nutrient thresholds derived from van den
718 Burg's literature compilation with established central European references. Eur J For Res
719 131:1461–1472
- 720 Mora O, Banos V, Regolini M, Carnus JM (2014) Using scenarios for forest adaptation to climate
721 change: a foresight study of the Landes de Gascogne Forest 2050. Ann For Sci 71:313–324
- 722 Nicholls D, Monserud RA, Dykstra DP (2009) International bioenergy synthesis-Lessons learned and
723 opportunities for the western United States. For Ecol Manage 257:1647–1655
- 724 Nnadi EO, Mbah CN, Nweke AI, Njoku C (2019) Physicochemical properties of an acid ultisol
725 subjected to different tillage practices and wood-ash amendment: Impact on heavy metal
726 concentrations in soil and Castor plant. Soil Tillage Res 194:104288
- 727 Nohrstedt HO (2001) Response of coniferous forest ecosystems on mineral soils to nutrient additions:
728 A review of Swedish experiences. Scand J For Res 16:555–573. doi: 10.1080/02827580152699385
- 729 Omil B, Pineiro V, Merino A (2013) Soil and tree responses to the application of wood ash containing
730 charcoal in two soils with contrasting properties. For Ecol Manage 295:199–212. doi:
731 10.1016/j.foreco.2013.01.024
- 732 Pitman RM (2006) Wood ash use in forestry - a review of the environmental impacts. FORESTRY
733 79:563–588. doi: 10.1093/forestry/cpl041
- 734 R Core Team (2019) R: A language and environment for statistical computing. R Foundation for
735 Statistical Computing; Vienna; Austria. URL <https://www.R-project.org>
- 736 Ranger J, Turpault MP (1999) Input-output nutrient budgets as a diagnostic tool for sustainable forest
737 management. For Ecol Manage 122:139–154
- 738 Ranius T, Hamalainen A, Egnell G, et al (2018) The effects of logging residue extraction for energy on
739 ecosystem services and biodiversity: A synthesis. J Environ Manage 209:409–425. doi:

- 740 10.1016/j.jenvman.2017.12.048
- 741 Reid C, Watmough SA (2014) Evaluating the effects of liming and wood-ash treatment on forest
742 ecosystems through systematic meta-analysis. *Can J For Res* 44:867–885. doi: 10.1139/cjfr-2013-
743 0488
- 744 Saur E, Ranger J, Lemoine B, Gelpe J (1992) Micronutrient Distribution in 16-Year-Old Maritime Pine.
745 *Tree Physiol* 10:307–316
- 746 Schulze ED, Korner CI, Law BE, et al (2012) Large-scale bioenergy from additional harvest of forest
747 biomass is neither sustainable nor greenhouse gas neutral. *Glob Chang Biol Bioenergy* 4:611–616
- 748 Slattery JF, Coventry DR (1995) Acid-tolerance and symbiotic effectiveness of *Rhizobium*-
749 *Leguminosarum* by *Trifolium* isolated from subterranean clover growing in permanent pastures.
750 *Soil Biol Biochem* 27:111–115. doi: 10.1016/0038-0717(94)00143-O
- 751 Solla-Gullon F, Santalla M, Rodriguez-Soalleiro RJ, Merino A (2006) Nutritional status and growth of
752 a young *Pseudotsuga menziesii* plantation in a temperate region after application of wood-bark
753 ash. *For Ecol Manage* 237:312–321. doi: 10.1016/j.foreco.2006.09.054
- 754 Steenari BM, Lindqvist O (1997) Stabilisation of biofuel ashes for recycling to forest soil. *Biomass*
755 *Bioenergy* 13:39–50. doi: 10.1016/S0961-9534(97)00024-X
- 756 Stromgren M, Egnell G, Olsson BA (2013) Carbon stocks in four forest stands in Sweden 25 years after
757 harvesting of slash and stumps. *For Ecol Manage* 290:59–66. doi: 10.1016/j.foreco.2012.06.052
- 758 Stupak I, Nordfjell T, Gundersen P (2008) Comparing biomass and nutrient removals of stems and fresh
759 and predried whole trees in thinnings in two Norway spruce experiments. *Can J For Res Can Rech*
760 *For* 38:2660–2673
- 761 Sverdrup H, Thelin G, Robles M, et al (2006) Assessing nutrient sustainability of forest production for
762 different tree species considering Ca, Mg, K, N and P at Bjornstorp Estate, Sweden.
763 *Biogeochemistry* 81:219–238
- 764 Symeonides C, McRae SG (1977) The assessment of plant-available cadmium in soils. *J Environ Qual*
765 6:120–123
- 766 Thiffault E, Hannam KD, Pare D, et al (2011) Effects of forest biomass harvesting on soil productivity
767 in boreal and temperate forests - A review. *Environ Rev* 19:278–309

- 768 Trichet P, Bakker MR, Augusto L, et al (2009) Fifty years of fertilization experiments on *Pinus pinaster*
769 in southwest France: the importance of phosphorus as a fertilizer. *For Sci* 55:390–402
- 770 Trichet P, Cheval N, Lambrot C, et al (2018) Using a dune forest as a filtering ecosystem for water
771 produced by a treatment plant - One decade of environmental assessment. *Sci Total Environ*
772 640:849–861. doi: 10.1016/j.scitotenv.2018.05.263
- 773 Vance ED (1996) Land application of wood-fired and combination boiler ashes: An overview. *J Environ*
774 *Qual* 25:937–944. doi: 10.2134/jeq1996.00472425002500050002x
- 775 Ventura M, Panzacchi P, Muzzi E, et al (2019) Carbon balance and soil carbon input in a poplar short
776 rotation coppice plantation as affected by nitrogen and wood ash application. *NEW For* 50:969–
777 990. doi: 10.1007/s11056-019-09709-w
- 778 Vidal DF, Trichet P, Puzos L, et al (2019) Intercropping N-fixing shrubs in pine plantation forestry as
779 an ecologically sustainable management option. *For Ecol Manage* 437:175–187
- 780 Wall A (2012) Risk analysis of effects of whole-tree harvesting on site productivity. *For Ecol Manage*
781 282:175–184
- 782 Walmsley JD, Godbold DL (2010) Stump harvesting for bioenergy - A review of the environmental
783 impacts. *Forestry* 83:17–38
- 784 Wang P, Olsson BA, Arvidsson H, Lundkvist H (2010) Short-term effects of nutrient compensation
785 following whole-tree harvesting on soil and soil water chemistry in a young Norway spruce stand.
786 *Plant Soil* 336:323–336. doi: 10.1007/s11104-010-0484-1
- 787 Yang X, Post WM (2011) Phosphorus transformations as a function of pedogenesis: A synthesis of soil
788 phosphorus data using Hedley fractionation method. *Biogeosciences* 8:2907–2916
- 789 Zabowski D, Chambreau D, Rotramel N, Thies WG (2008) Long-term effects of stump removal to
790 control root rot on forest soil bulk density, soil carbon and nitrogen content. *For Ecol Manage*
791 255:720–727
- 792
- 793

794
795
796

Figure 1 – Realised exports of tree biomass and nutrients from the cut forest




797
798
799
800
801
802
803
804
805

SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Error bars are standard errors.

All data (including data about calcium, and magnesium) are presented in Table S2.


The methods used to quantify the biomass and nutrient exports are detailed in the manuscript in the section “*Plant measurements and field sampling*”, in particular in the subsection “*Estimates of biomass and nutrient exports*”.

806 **Figure 2** – Understorey aboveground biomass as affected by biomass harvest and compensation methods
 807



808
 809 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control =
 810 no nutrient application.
 811 Mean values of the Biomass Harvesting treatments (n = 3 blocks) were calculated by pooling the Compensation Methods subplots. Mean values
 812 of the Compensation Methods treatments were calculated based on 12 replicates (3 blocks × 4 BH treatments).
 813 The understorey biomass was significantly higher in the Wood Ash treatment than in the Control treatment.
 814 The relative abundance of the species typical of oligotrophic-acidic conditions (herbaceous *Molinia caerulea* and ericaceous *Calluna vulgaris* &
 815 *Erica scoparia*) is highlighted by the red dotted line. Photos of the main species are given in the Figure S4.
 816
 817
 818

819 **Figure 3** – Tree stem volume as affected by biomass harvest and compensation methods
820



821 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground
822 additional harvest; WTH = whole-tree harvest. Control = no nutrient application. Error bars are
823 standard errors.

824 Mean values of the Biomass Harvesting treatments (n = 3 blocks) were calculated by pooling
825 the Compensation Methods subplots. Mean values of the Compensation Methods treatments
826 were calculated based on 12 replicates (3 blocks × 4 BH treatments).

827 P values for differences between treatments of reference (*i.e.* SOH and Control) on the one
828 hand, and other treatments on the other hand, are indicated in the graph.

829 Trees were measured at 11 years old.

830 Other tree metrics are shown in Table S3.

831

832

833

834 **Table 1** – Initial expected consequences of intensive biomass harvesting and wood ash
 835 application on the studied ecosystem (temperate oligotrophic forest)
 836 (→: means “has no effect on...”; ↓: means “decreases...”; ↗: means “increases...”; ?: means “unknown
 837 effect on...”)

Ecosystem trait		Intensive harvests	Wood ash
Soil fertility (phosphorus) {hypothesis H1}	<i>General knowledge</i>	↓ soil P	→ soil P
	<i>Local conditions</i>	P-limited soils	
	<i>Expected results</i>	↓ (aggravated P-limitation)	→ (or ↗) soil P
Tree growth (carbon sequestration) {H2}	<i>General knowledge</i>	↓ tree growth	→ (or ↗) tree growth
	<i>Local conditions</i>	oligotrophic P-limited ecosystem	
	<i>Expected results</i>	↓ tree growth	→ tree growth
Soil Organic Matter SOM (carbon and nitrogen) {H3}	<i>General knowledge</i>	→ boreal forests ↓ temperate forests	→ boreal forests ? temperate forests
	<i>Local conditions</i>	warm temperate	
	<i>Expected results</i>	↓ SOM	→ SOM
Soil acidity (pH, base saturation) {H4}	<i>General knowledge</i>	↓ acido-basic status	↗ acido-basic status
	<i>Local conditions</i>	acidic soils	
	<i>Expected results</i>	↓ acido-basic status	↗ acido-basic status
Trace metals in ecosystem (micronutrients: [B, Cu, Fe, Mn, Mo, Zn] and toxic trace metals [As, Pb, Cd]) {H5}	<i>General knowledge</i>	→ or ↓ trace metals (due to exports)	→ for podzol soils (boreal soils in Nordic areas)
	<i>Local conditions</i>	low content in tree biomass growing on podzol soils	
	<i>Expected results</i>	→ trace metals	→ or ↓ bioavailability at moderate ash dose + increase pH

838
839

840 **Table 2** – Soil content in carbon, nutrients, and micronutrients as affected by biomass harvest and compensation methods
 841

	SOC (mg g ⁻¹)	DOC (mg L ⁻¹)	N (mg g ⁻¹)	C/N unit less	P _{Olsen} (μg g ⁻¹)	K (mg g ⁻¹)	Ca (mg g ⁻¹)	Mg (mg g ⁻¹)	Na (mg g ⁻¹)	Fe (mg g ⁻¹)	Mn (μg g ⁻¹)	Mo (μg g ⁻¹)	Zn (μg g ⁻¹)
<i>Biomass Harvesting</i>													
SOH	38.6±3.9	44±2	1.41±0.15	27.7±0.8	29.3±2.2	4.07±0.2	0.82±0.06	0.17±0.09	1.17±0.09	1.8±0.4	41.5±3.5	0.16±0.01	6.60±0.29
AAH	35.4±3.9	42±4	1.46±0.15	24.5±0.8	25.3±2.5	5.40±0.2 *	0.98±0.06	0.25±0.09	1.65±0.08 *	3.1±0.3 *	62.0±3.5 *	0.21±0.01	7.37±0.48
BAH	30.2±3.9	39±2	1.14±0.15	26.3±0.8	25.3±2.9	4.31±0.2	0.71±0.04	0.15±0.08	1.22±0.08	2.2±0.3	43.6±3.5	0.15±0.01	5.77±0.36
WTH	31.1±3.9	31±2 *	1.10±0.15	28.2±0.8	21.0±3.5	4.09±0.2	0.72±0.03	0.11±0.04	1.11±0.05	1.5±0.3	35.4±3.5	0.13±0.01	3.85±0.54
<i>Compensation Methods</i>													
Control	37.1±2.9	39±1	1.36±0.11	27.2±0.7	27.1±2.4	4.47±0.13	0.85±0.03	0.18±0.10	1.31±0.05	2.1±0.3	46.6±3.2	0.16±0.01	5.69±0.31
Phosphorus	33.6±2.9	42±2	1.28±0.11	26.3±0.7	24.5±2.4	4.45±0.26	0.74±0.03 *	0.16±0.08	1.26±0.09	2.0±0.3	45.0±4.9	0.17±0.01	5.57±0.32
Ash	30.8±2.9 *	36±1 *	1.18±0.11 *	26.5±0.7	24.0±2.4	4.50±0.15	0.85±0.03	0.17±0.09	1.31±0.06	2.3±0.3	45.2±2.0	0.16±0.01	6.42±0.36

842 All values are for the total content of the elements in the soil (except for P). SOC = soil organic carbon; DOC = dissolved organic carbon. For phosphorus (P), values are
 843 extractable P using the Olsen method.
 844 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
 845 Treatments with an asterisk differ at $P < 0.05$ from their reference treatment (SOH or Control, respectively; see Methods).
 846 Mean values of the Biomass Harvesting treatments (n = 3 blocks) were calculated by pooling the Compensation Methods subplots. Mean values of the Compensation Methods
 847 treatments were calculated based on 12 replicates (3 blocks × 4 BH treatments).
 848
 849
 850

851 **Table 3** – Soil acidity and exchangeable cations as affected by biomass harvest and compensation methods

852

	pH-H ₂ O unit less	pH-CaCl ₂ unit less	CEC _m (cmol.c kg ⁻¹)	BS (%)	K (cmol.c kg ⁻¹)	Ca (cmol.c kg ⁻¹)	Mg (cmol.c kg ⁻¹)	Na (cmol.c kg ⁻¹)
<i>Biomass Harvesting</i>								
SOH	4.43±0.07	3.43±0.08	3.51±0.23	43±2	0.06±0.00	0.92±0.04	0.36±0.02	0.06±0.00
AAH	4.68±0.07	3.69±0.08	2.90±0.24	38±2	0.06±0.00	0.67±0.03 *	0.21±0.01 *	0.05±0.00
BAH	4.49±0.07	3.57±0.08	2.51±0.23 *	42±2	0.05±0.00	0.65±0.04 *	0.23±0.02 *	0.04±0.00 *
WTH	4.51±0.07	3.46±0.08	2.90±0.24	39±2	0.06±0.01	0.69±0.06	0.28±0.02 *	0.04±0.00 *
<i>Compensation Methods</i>								
Control	4.41±0.06	3.47±0.07	3.45±0.21	39±2	0.06±0.00	0.74±0.05	0.29±0.01	0.05±0.00
Phosphorus	4.53±0.05	3.51±0.07	2.58±0.19 *	38±2	0.06±0.00	0.57±0.04 *	0.22±0.02 *	0.04±0.00 *
Ash	4.65±0.05 *	3.64±0.07	2.84±0.19 *	46±2 *	0.06±0.00	0.89±0.04 *	0.30±0.02	0.05±0.00

853

854 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.

855 CEC_m = cationic exchange capacity (measured value); BS = base saturation of the CEC, which is the proportion of charges occupied by non-acidic exchangeable cations (K,
856 Ca, Mg, Na).

857 For consistency between the different terms of its equation, BS was calculated using the effective CEC (CEC_e: calculated as the sum all the charges of all the analysed cations),
858 whose values differ slightly from CEC_m values.

859 Treatments with an asterisk differ at $P < 0.05$ from their reference treatment (SOH or Control, respectively; see Methods).

860 Mean values of the Biomass Harvesting treatments (n = 3 blocks) were calculated by pooling the Compensation Methods subplots. Mean values of the Compensation Methods
861 treatments were calculated based on 12 replicates (3 blocks × 4 BH treatments).

862

863

864 **Table 4** – Soil composition in NH₄-NO₃ extractable nutrients and micronutrients as affected by biomass harvest and compensation methods
 865

	P (µg g ⁻¹)	K (µg g ⁻¹)	Ca (µg g ⁻¹)	Mg (µg g ⁻¹)	Fe (µg g ⁻¹)	Mn (µg g ⁻¹)	Zn (µg g ⁻¹)
<i>Biomass Harvesting</i>							
SOH	1.2±0.3	27±2	178±8	47±2	7.0±1.2	1.5±0.1	0.60±0.05
AAH	0.5±0.2 *	21±2 (*)	117±10 *	25±2 *	9.7±1.2	1.3±0.1	0.48±0.04
BAH	0.8±0.2	21±2 (*)	128±8 *	30±2 *	8.2±1.2	1.2±0.1	0.47±0.04
WTH	0.9±0.2	21±2 (*)	151±12	38±2	6.1±1.2	1.2±0.1	0.45±0.04
<i>Compensation Methods</i>							
Control	0.9±0.2	23±1	131±9	35±2	7.2±1.3	1.1±0.1	0.44±0.02
Phosphorus	0.7±0.2	22±1	117±9	30±2 *	8.5±0.9	1.1±0.1	0.48±0.03
Ash	0.9±0.1	22±1	183±9 *	40±1 *	7.5±0.5	1.8±0.1 *	0.58±0.06 *

866 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
 867 Treatments with an asterisk differ at $P < 0.05$ (except for K: P value < 0.10 ; symbol in brackets) from their reference treatment (SOH or Control, respectively; see Methods).
 868 Mean values of the Biomass Harvesting treatments (n = 3 blocks) were calculated by pooling the Compensation Methods subplots. Mean values of the Compensation Methods
 869 treatments were calculated based on 12 replicates (3 blocks × 4 BH treatments).
 870
 871
 872
 873

874 **Table 5** – Tree foliage composition (nutrients and micronutrients) as affected by biomass harvest and compensation methods
 875

	N (mg g ⁻¹)	P (mg g ⁻¹)	K (mg g ⁻¹)	Ca (mg g ⁻¹)	Mg (mg g ⁻¹)	B (μg g ⁻¹)	Cu (μg g ⁻¹)	Fe (μg g ⁻¹)	Mn (μg g ⁻¹)	Mo (μg g ⁻¹)	Zn (μg g ⁻¹)
<i>Biomass Harvesting</i>											
SOH	12.8±0.7	0.64±0.04	4.6±0.1	2.3±0.1	1.4±0.1	16.3±1.0	1.8±0.2	27.8±2.0	96.3±11.0	0.011±0.003	29.8±2.5
AAH	13.9±0.7	0.69±0.04	4.4±0.1	2.2±0.1	1.5±0.1	16.7±1.0	1.8±0.2	32.1±2.1 *	104.2±10.9	0.014±0.002	32.1±2.5
BAH	12.1±0.7	0.65±0.04	4.5±0.1	2.2±0.1	1.4±0.1	17.2±1.0	2.0±0.2	27.1±2.0	92.9±11.7	0.009±0.003	30.5±2.5
WTH	12.6±0.7	0.65±0.05	4.6±0.1	2.3±0.1	1.4±0.1	17.2±1.0	1.8±0.3	28.9±2.9	81.0±11.7	0.012±0.002	32.1±2.5
<i>Compensation Methods</i>											
Control	12.3±0.7	0.64±0.04	4.2±0.1	2.2±0.1	1.4±0.1	16.0±0.9	1.7±0.1	27.0±2.1	92.3±9.8	0.011±0.002	29.1±2.2
Phosphorus	13.6±0.7 *	0.66±0.04 *	4.5±0.1 *	2.3±0.1	1.5±0.1 *	18.0±0.9 *	1.9±0.1	30.4±2.1 *	100.1±9.4 *	0.012±0.002	32.7±2.2 *
Ash	12.7±0.7 *	0.67±0.04 *	4.8±0.1 *	2.3±0.1	1.4±0.1	16.6±0.9 *	2.0±0.1	29.6±2.0 *	88.4±9.8	0.012±0.002	31.7±2.2 *

876 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.
 877 Treatments with an asterisk differ at $P < 0.05$ from their reference treatment (SOH or Control, respectively; see Methods).
 878 Mean values of the Biomass Harvesting treatments (n = 3 blocks) were calculated by pooling the Compensation Methods subplots. Mean values of the Compensation Methods
 879 treatments were calculated based on 12 replicates (3 blocks × 4 BH treatments).
 880
 881
 882

883 **Table 6** – Ecosystem content in non-essential trace metals

884

	Cadmium			Lead			Arsenic
	<i>Soil</i> ($\mu\text{g-Cd kg}^{-1}$)	<i>Extractable</i> ($\mu\text{g-Cd kg}^{-1}$)	<i>Foliage</i> ($\mu\text{g-Cd g}^{-1}$)	<i>Soil</i> ($\mu\text{g-Pb g}^{-1}$)	<i>Extractable</i> ($\mu\text{g-Pb g}^{-1}$)	<i>Foliage</i> ($\mu\text{g-Pb g}^{-1}$)	<i>Foliage</i> ($\mu\text{g-As kg}^{-1}$)
<i>Biomass Harvesting</i>							
SOH	33±2	9.4±0.6	0.17±0.01	9.45±0.79	0.24±0.01	0.17±0.01	19±3
AAH	33±2	8.9±0.6	0.18±0.01	9.53±0.58	0.17±0.01	0.18±0.01	21±3
BAH	30±2	8.4±0.6	0.16±0.01	7.59±0.45	0.17±0.01	0.16±0.01	15±3
WTH	26±2	7.6±0.6	0.17±0.01	6.76±0.24 *	0.16±0.01	0.07±0.01	21±3
<i>Compensation Methods</i>							
Control	31±1	8.2±0.5	0.16±0.01	8.20±0.39	0.20±0.01	0.16±0.01	16±2
Phosphorus	28±1 *	8.2±0.4	0.18±0.01 (*)	8.24±0.39	0.18±0.01	0.18±0.01 (*)	20±4
Ash	32±1	9.4±0.7	0.17±0.01	8.57±0.39	0.17±0.01	0.17±0.01	20±4

885

886 “*Soil*” = soil substrate (total content of Cd and Pb); “*Extractable*” = soil extractable with 1M NH₄-NO₃ extractant; “*Foliage*” = tree foliage (*i.e.* pine needles) content. It should
887 be noted that the units used differ among Cd variables.

888 SOH = stem-only harvest; AAH = aboveground additional harvest; BAH = belowground additional harvest; WTH = whole-tree harvest. Control = no nutrient application.

889 Treatments with an asterisk differ at $P < 0.05$ (for P value < 0.10 , the symbol is in brackets) from their reference treatment (SOH or Control, respectively; see Methods).890 Mean values of the Biomass Harvesting treatments ($n = 3$ blocks) were calculated by pooling the Compensation Methods subplots. Mean values of the Compensation Methods
891 treatments were calculated based on 12 replicates (3 blocks \times 4 BH treatments).892 Total soil content of nickel (Ni) and thallium (Tl) are not presented. No significant differences were observed for these variables (Ni=1.49 $\mu\text{g g}^{-1}$; Tl=0.11 $\mu\text{g g}^{-1}$).

893

894

895

896