



HAL
open science

Biodiversity of Aflatoxigenic *Aspergillus* section *Flavi* Species According to Food Matrices and Geographic Areas

Amaranta Carvajal-Campos, Ama Lethicia Manizan, Didier Montet, Sophie
Lorber, Olivier Puel, Catherine Brabet

► **To cite this version:**

Amaranta Carvajal-Campos, Ama Lethicia Manizan, Didier Montet, Sophie Lorber, Olivier Puel, et al.. Biodiversity of Aflatoxigenic *Aspergillus* section *Flavi* Species According to Food Matrices and Geographic Areas. *Mycotoxins in Food and Beverages Innovations and Advances Part I Food Biology Series*, 2021, 978-0-367-42209-7. hal-03773672

HAL Id: hal-03773672

<https://hal.inrae.fr/hal-03773672v1>

Submitted on 9 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Biodiversity of Aflatoxigenic *Aspergillus* section *Flavi* Species According to Food Matrices and Geographic Areas

Carvajal-Campos Amaranta¹, Manizan Ama Lethicia², Didier Montet^{3,4},
Sophie Lorber⁵, Olivier Puel⁵ and Catherine Brabet^{3,4}

¹ Université de Montpellier, Faculté des Sciences, Campus Triolet, Place Eugène Bataillon, CC 023, 34095 Montpellier Cedex 5, France

² Laboratoire de Biotechnologie et Microbiologie des Aliments, UFR des Sciences et Technologie des Aliments, Université Nangui Abrogoua, 02 BP 801 Abidjan 02, Côte d'Ivoire

³ CIRAD, UMR QualiSud, 73 rue Jean-François Breton, 34398 Montpellier Cedex 5, France

⁴ Qualisud, Université Montpellier, CIRAD, Montpellier SupAgro, Université d'Avignon, Université de La Réunion, Montpellier, France

⁵ Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRAE, ENVT, INP-Purpan, UPS, Toulouse, France

Au: Pl. specify the corresponding author with his Email. address

1. Introduction

Aflatoxins (AFs) are polyketide-derived metabolites produced by fungi on a wide range of crops (cereals, oilseeds, tree nuts, spices, dried fruits, etc.), both in the field and the post-harvest. As chemical stable molecules, resistant to conventional thermal or technological processes, they will pass through the whole food or feed supply chain to the final processed products (Kumar et al. 2017, Pankaj et al. 2018). There are more than 20 known AFs and derivatives, yet the most hazardous AFs include the four naturally-occurring AFB1, AFB2, AFG1 and AFG2, and the hydroxylated metabolites of AFB1 and AFB2, and AFM1 and AFM2, which are produced through biotransformation in the liver and excreted in the milk of humans and mammals (Kumar et al. 2017). AFs are known for their high acute and chronic toxicity to both humans and animals, and are considered among the most dangerous mycotoxins with carcinogenic,

*Corresponding author:



Au: Email. address

hepatotoxic, immunotoxic, teratogenic and mutagenic effects (Kowalska et al. 2017). AFB1 is the most prevalent and toxic aflatoxin, and is classified as a Group 1 human carcinogens (IARC 2012). The intake of high amounts of AFs can cause acute intoxication (aflatoxicosis) associated with jaundice, vomiting, hemorrhages, abdominal pain, acute liver failure, problems with absorption of nutrients, and can be lethal (IARC 2015). Reported outbreaks in western India (1974) and in Kenya (2004) caused the death of 106 and 125 people, respectively (Lewis et al. 2005, Azziz-Baumgartner et al. 2005). Chronic exposure to low levels of AFs is associated with high risk of hepatocellular carcinoma (HCC), immunosuppression, teratogenic and mutagenic effects, reduction of nutrient absorption, child stunting, detrimental effects in the endocrinal system, and liver failure (Stack and Carlson 2003, Turner et al. 2005, Bbosa et al. 2013). More than five billion people worldwide are at risk of chronic exposure to AFs through contaminated foods (Wu and Glucu 2012). Humans' exposure to AFs mainly occurs by direct intake of contaminated foods of vegetal or animal origin (Bhat et al. 2010, IARC 2015). Hence, the presence of AFs in foodstuff and feedstuff is a public health issue associated with detrimental effects in economy. In addition, commodities are generally contaminated by several mycotoxins, and this co-occurrence may result in a greater toxicity to humans caused by the possible additive or synergistic effects of these compounds (Grenier and Oswald 2011). The main aflatoxin-producing fungi belong to *Aspergillus* section *Flavi* of the *Circumdati* subgenus, and only five AF producers do not belong to this section, *A. ochraceoroseus*, *A. rambellii* (*A. section Ochraceorosei*), *A. stellatus*, *A. olivicola*, and *A. venezuelensis* (*A. section Nidulantes*) (Varga et al. 2015). The aim of the present chapter is to describe the characteristics of the aflatoxigenic *Aspergillus* section *Flavi* species isolated from food or feed, and to address the biodiversity according to food matrices and geographic areas.

2. Characteristics of the Aflatoxigenic *Aspergillus* section *Flavi* Species

The major producers of AFs worldwide are *Aspergillus flavus* and *A. parasiticus* and for a while they were recognized as the only AF producers, and later, diversity surveys pointed *A. nomius* as the third main AF producer for its frequency in spoiled food (Kurtzman et al. 1987, Perrone et al. 2014). However, the implementation of polyphasic species identification showed a more complex story. Over the last two decades, this approach demonstrated a high biodiversity and plasticity within the section *Flavi* (Varga et al. 2011, Soares et al. 2012, Taniwaki et al. 2012, Frisvad et al. 2019). Currently this section encloses 34 species, from which 19 are aflatoxigenic (Frisvad et al. 2019).

2.1 Morphological Characteristics

Macro- and micro-morphological characters of the 19 aflatoxigenic *Aspergillus* section *Flavi* species on Malt Extract Agar (MEA) and Czapek Yeast Agar

(CYA) after seven days of incubation, in the dark at 25°C are summarized in Table 1. These two media are recommended as standard for *Aspergillus* and commonly used in taxonomic studies on this genus (Samson et al. 2014). Most aflatoxigenic *Aspergillus* species grow well on MEA and CYA at 25°C attaining colony diameter of more than 50 mm after seven days; *A. austwickii* is the slowest growing species (colony diameter MEA: 45–47 mm, CYA: 4648 mm). Colony surface is mostly deep, floccose, rarely plane, velvety or powdery. Most species sporulate and produce sclerotia with the exception of *A. pipericola* (no sporulation) and *A. arachidicola*, *A. luteovirescens*, *A. novoparasiticus*, *A. pseudocaelatus* and *A. pseudonomius* (no sclerotia). Conidia are mainly in shades of (dark) yellow-green, but also in shades of brown. Conidial heads are uniseriate with smooth or rough conidia, biseriate with rough conidia, or uniseriate to biseriate with smooth or slightly rough, smooth to rough or rough conidia. Sclerotia size varies between large (400–2000 µm), intermediate (*A. sergii*: 513–551 µm) and small (< 400 µm). Sclerotia morphology can be globose to ellipsoidal, and they become dark brown or black with time (Varga et al. 2011, Frisvad et al. 2019). Morphological characters are useful to differentiate some aflatoxigenic species within section *Flavi*, but for closely related species a proper description based only on these characters is a challenge. For this reason, a polyphasic approach that includes the morphological, chemical (mycotoxins and other extrolite production) and molecular characteristics is necessary to identify and characterize the *Aspergillus* section *Flavi* species (Varga et al. 2011).

2.2 Mycotoxins and Other Extrolites

Aflatoxigenic species from section *Flavi* produce an ample spectrum of secondary metabolites, besides of AFs some important and emergent mycotoxins include cyclopiazonic acid (CPA), tenuazonic acid and aflatrems (Table 2). From the nineteen aflatoxigenic species, three produce only AFB (AFB1 and AFB2: *A. flavus*, *A. pseudotamarii*; AFB1: *A. togoensis*), the other species produce both AFB (B1, B2) and AFG (G1, G2). *A. flavus* is the main AFB1 producer across the world and therefore the best known species from *Flavi* section, its AF production potential varies from non-aflatoxigenic to aflatoxigenic strains, with a high incidence of non-aflatoxigenic strains (60–70%) (Varga et al. 2011). AF production potential is not associated with virulence or competitive ability during crop infection. This plasticity and the lack of production of AFG are driven by several genetic differences, including single polymorphisms and large deletions in the AF biosynthesis gene cluster (Ehrlich 2004, Chang and Ehrlich 2010). *A. flavus* L-morphotype generally presents a 0.8 kb deletion in the *CypA*/*NorB* (*AflU*/*AflF*) region of the aflatoxin biosynthetic gene cluster (BGC), whereas *A. flavus* S-morphotype present of 1.5 kb (Ehrlich and Yu 2010). The CYP_A protein encoded by *AflU* is required for AFG production (Ehrlich et al. 2004). A third group, gathering the strains responsible of the lethal outbreak in Kenya, was highlighted with a 2.2 kb deletion (Probst et al. 2012). AF production by *A. flavus* L-morphotype isolates is variable, while *A. flavus* S-morphotype isolates produce high quantities

Table 1: Morphological characteristics of aflatoxigenic *Aspergillus* section *Flavi* species on MEA and CYA after seven days of incubation in the dark at 25°C

<i>Aflatoxigenic Aspergillus</i> species	Medium	Colony diameter (mm)	Colony surface and conidial color	Sclerotia (µm)	Conidial head morphology and conidial surface	References
<i>A. aflatoxiformans</i> *	MEA	47–50	Colonies moderately deep, mycelium floccose and white Conidia yellow-green, moderately dense on MEA and sparse on CYA	100–250	Uniseriate Smooth	Frisvad et al. 2019
	CYA	50–51				
<i>A. arachidicola</i>	MEA	60–65	Colonies velvety Conidia olive to olive brown, abundant	Not observed	Uniseriate or biseriate Rough	Pildain et al. 2008
	CYA					
<i>A. austroickii</i>	MEA	45–47	Colonies moderately deep, mycelium floccose and white Conidia yellow-green, moderately dense on MEA and sparse on CYA	100–300	Uniseriate Smooth	Frisvad et al. 2019
	CYA	46–48				
<i>A. flavus</i>	MEA	50–70	Colonies similar to those on CYA although usually less dense Colonies plane, sparse to moderately dense, velutinous in marginal areas at least, often floccose centrally, rare deep; mycelium white Conidia greyish green, yellow green or olive yellow, sometimes yellow	< 400 (S-type) 400–800 (L-type) or not observed	Typically biseriate, rare uniseriate Finely rough, rarely smooth	Pitt and Hocking 2009, Frisvad et al. 2019
	CYA	60–70				
<i>A. korlagoensis</i> *	MEA	37–60	Colonies deeply floccose with a dominant white aerial mycelium Conidia dull yellowish green	<400	Uniseriate or biseriate Smooth or slightly rough	Carvajal-Campos et al. 2017
	CYA	57–80				

(Contid.)

Table 1: (Contd.)

Aflatoxigenic <i>Aspergillus</i> species	Medium	Colony diameter (mm)	Colony surface and conidial color	Sclerotia (μm)	Conidial head morphology and conidial surface	References
<i>A. luteovirescens</i> *	MEA	65	Colonies loose and deep, yellow-green becoming darker at maturity	Generally not observed, when present large	Mostly biseriolate	Peterson et al. 2001, Frisvad et al. 2019
	MEA CYA	60-70	Colonies floccose, white mycelium Conidia light-greyish green, sparse Colonies velvety Conidia light-greyish green	150-300	Normally biseriolate, rare uniseriate Smooth to rough	Pildain et al. 2008
<i>A. mottiae</i>	MEA CYA	>70	Colonies plane, mycelium white Conidia yellow-green, scarce	249-371	Normally biseriolate, rare uniseriate Smooth to finely rough	Soares et al. 2012
	MEA CYA	40-70 52-60	Colonies mostly floccose. Conidia green Colonies greyish green	< 500	Mostly biseriolate Rough	Kurtzman et al. 1987, Doster et al. 2009
<i>A. novoparasiticus</i>	MEA CYA	56 - 60 58 - 63	Colonies powdery Greenish-yellow to olive	Not observed	Usually uniseriate, rarely biseriolate	Gonçaves et al. 2012
	MEA CYA	50-65 50-70	Colonies similar to those on CYA but usually less dense Colonies plane, low, dense and velutinous, mycelium white Conidia dark yellowish green	400-800	Mostly uniseriate Rough	Pitt and Hocking 2009

AQ : missing in references

<i>A. pipericola</i>	MEA	61-72	Colonies moderately deep, mycelium floccose and white Sporulation absent	75-250	Biseriate Rough	Frisvad et al. 2019
	CYA	58-72				
<i>A. pseudocaelatus</i>	MEA	60-65	Colonies velvety Conidia olive to olive brown, abundant	Not observed	Uniseriate or biseriate Rough	Varga et al. 2011
	CYA					
<i>A. pseudonomius</i>	MEA	60-65	Colonies floccose with dominant aerial mycelium Poor sporulation	Not observed	Uniseriate Rough	Varga et al. 2011
	CYA					
<i>A. pseudotamarii</i>	MEA	60-70	Colonies mostly floccose Conidia olive green	1000-2000	Biseriate Rough	Ito et al. 2001
<i>A. sergii</i>	MEA	55	Colonies plane, velvety and dense. Conidia in a uniform, dense layer but sparse in the areas of sclerotium production, light green	513-551	Uniseriate Rough	Soares et al. 2012
	CYA					
<i>A. texensis</i>	CYA	71	Colonies velvety. mycelium white Conidia yellow-green, sparse	130-300	-	Singh et al. 2018
<i>A. togoensis</i>	MEA	>50	Conidia yellow-brown to orange-brown	Large sclerotia	Biseriate	Samson and Seifert 1985, Frisvad et al. 2019
	CYA	>50				
<i>A. transmontanensis</i>	MEA	55-57	Colonies similar to growth on CYA with conidial heads more dense and floccose	458-609	Mostly uniseriate Rough	Soares et al. 2012
	CYA		Colonies dense and velutinous. Conidia in a uniform, dense layer but sparse in the areas of sclerotium production, dark yellow-green			

* *A. parvisclerotigenus*, *A. bombycis* and *A. korhogoensis* synonymized as *A. aflatoxiformans*, *A. liteovirescens* and *A. cerealis*, respectively, by Frisvad et al. 2019.

Table 2: Production of mycotoxins and other secondary metabolites by aflatoxigenic *Aspergillus* section *Flavi* species

<i>Aflatoxigenic Aspergillus species</i>	<i>Aflatoxins</i>	<i>Other mycotoxins</i>	<i>Other secondary metabolites</i>	<i>References</i>
<i>A. aflatoxiformans</i>	AFB1 AFB2 AFG1 AFG2	Aflatrem, cyclopiazonic acid, versicolorins	Aspergillic acid, aflavarins, aflavinines, aspirochlorin, kojic acid, paspaline, paspalimine, metabolite gfm	Frisvad et al. 2019
<i>A. arachidicola</i>		Versicolorins	Aspergillic acid, aspirochlorin, chrysoquine, ditryptophenaline kojic acid, miyakimides, parasiticolide, "NO2" metabolite, parasitocolides	Pildain et al. 2008, Varga et al. 2011, Frisvad et al. 2019, Iamanaka et al. 2019
<i>A. australickii</i>		Aflatrem, cyclopiazonic acid, versicolorins	Aflavarins, kojic acid, paspaline, paspalimine, metabolite gfm	Frisvad et al. 2019
<i>A. korhogoensis</i>		Aflatrem, cyclopiazonic acid, 3-O-methylsterigmatocystin, sterigmatocystin, versicolorins	Aspergillic acid, aflavarins, asparosones, asparosone A aflavinines, kojic acid, leporin B, norsolorinic acid, paspaline, paspalimine	Carvajal-Campos et al. 2017
<i>A. lutovirescens</i>		Tenuazonic acid (some strains)	Aspergillic acid, kojic acid For some strains: an altersolanol, chrysoquine	Pildain et al. 2008, Varga et al. 2011, Frisvad et al. 2019
<i>A. minisclerotigenes</i>		Aflatrem, cyclopiazonic acid	Aspergillic acid, aflavarins, aflavinines, kojic acid, paspaline, parasiticolides. For some strains: aflavazole	Pildain et al. 2008, Frisvad et al. 2019
<i>A. mottae</i>		Cyclopiazonic acid, 3-O-methylsterigmatocystin, versicolorins	Aspergillic acid, an aflavinin, kojic acid, parasiticol, paspalimine	Soares et al. 2012, Frisvad et al. 2019

<i>A. nomius</i>	3-O-methylsterigmatocystin tenuazonic acid, versicolorins	Aspergillilic acid, anominine, aspernomine, kojic acid, a miyakamide, pseurotin, parasiticol, paspaline, paspalinine, pseurotin A	Kurtzman et al. 1987, Frisvad et al. 2019
<i>A. notoparasiticus</i>	Tenuazonic acid	Aspergillilic acid, aspirochlorin, dirtryptophenaline, kojic acid, miyakamides, parasiticolide, crysogine, a tetracyclic compound, ustilagoidin	Goncalvez et al. 2012, Frisvad et al. 2019, Iamanaka et al. 2019
<i>A. parasiticus</i>		Aspergillilic acid, kojic acid, parasperone, parasiticol, parasiticolide A and B	Frisvad et al. 2019
<i>A. pipericola</i>	Aflatrem, cyclopiiazonic acid	Aflavinins, aflavarins, paspaline, paspalinine	Frisvad et al. 2019
<i>A. pseudoacaelatus</i>	Cyclopiiazonic acid, tenuazonic acid	Aspirochlorin, dirtryptophenaline, kojic acid	Varga et al. 2011, Frisvad et al. 2019
<i>A. pseudonomius</i>	Tenuazonic acid	Aspergillilic acid, chrysogine, kojic acid, a miyakamide	Varga et al. 2011, Frisvad et al. 2019
<i>A. sergii</i>	Aflatrem, cyclopiiazonic acid, 3-O-methylsterigmatocystin, sterigmatocystin, versicolorins,	Aspergillilic acid, aflavazole, an aflavarin, aflavinines, asperfuran, kojic acid, lepornin B, paspalinine	Soares et al. 2012, Frisvad et al. 2019
<i>A. texensis</i>	Cyclopiiazonic acid	Aspergillilic acid	Singh et al. 2018

(Contd.)

Table 2: (Contd.)

<i>Aflatoxigenic Aspergillus species</i>	<i>Aflatoxins</i>	<i>Other mycotoxins</i>	<i>Other secondary metabolites</i>	<i>References</i>
<i>A. transmontanensis</i>			Aspirochlorin, kojic acid, a miyakamide	Soares et al. 2012, Frisvad et al. 2019
<i>A. flavus</i>	AFB1 AFB2	Aflatremis (only in sclerotium producers), cyclopiazonic acid	Aspergillilic acid, asperfuran, aspirochlorin, citreoisocoumarin, ditryptophenaline, flavimin, kojic acid, miyakamides, ustilaginoidin C, ustiloxin B. Only in sclerotium producers: aflavarins, aflavinines, paspaline and paspalinine	Amaike and Keller 2011, Umemura et al. 2013, Arroyo-Manzanares et al. 2015, Cary et al. 2015a,b, Frisvad et al. 2019
<i>A. pseudotamarii</i>		Cyclopiazonic acid, tenuazonic acid	Aflavinines, kojic acid, For some strains: an altersolanol, aspirochlorin, paspaline and paspalinine	Ito et al. 1999, Varga et al. 2011, 2015, Frisvad et al. 2019
<i>A. togoensis</i>	AFB1	Sterigmatocystin	A bisiderin, paspaline, paspalinine. For some strains: paxilline	Frisvad et al. 2019

of AFs (Ehrlich 2004). Recently *A. pseudotamarii* genome was sequenced and BGC analysis showed it lacks the first 600 base pairs in the *AflU* gene encoding for CYP_A protein (Kjaerboelling et al. 2020). *A. parasiticus*, the second main AF producer, also shows AF production plasticity, 3 to 6% of strains are considered non-aflatoxigenic (Chang et al. 2007). For the other AFBG species, the production seems to occur in all isolates.

Besides of AFs, some of their biosynthetic intermediates, like versicolorins, 3-O-methylsterigmatocystin and sterigmatocystin, are considered to be potentially toxic and reported in several aflatoxigenic species (Table 2). Though, these species are weak producers, as the intermediates are mainly transformed into AFs. Sterigmatocystin is considered an emerging mycotoxin and classified as 2B carcinogen for its potentially carcinogenic, mutagenic and teratogenic effects (IARC 2012, Bertuzzi et al. 2017). It is reported to be produced by *A. korhogoensis*, *A. sergii* and *A. togoensis*. The latter species is the only species from *Flavi* section capable to storage sterigmatocystin (Varga et al. 2015). Versicolorin A was shown as cytotoxic and genotoxic for lungs (Jakšić et al. 2012), renal and hepatic cells (Theumer et al. 2018). Recently, versicolorin A was reported as more cytotoxic than AFB₁ for human intestinal cells. The toxic effects of 3-O-methylsterigmatocystin are less acute than AFs, sterigmatocystin and versicolorin A (Jakšić et al. 2012).

Cyclopiazonic acid is an important mycotoxin from section *Flavi* that has a synergetic effect with AFs. It might play a role in niche adaptation, providing an advantage in fungal fitness under specific environmental conditions (Georgianna et al. 2010). It is produced by *A. flavus* and closely related species, as well as *A. pseudocaelatus* and *A. pseudotamarii* (Table 2). Some species from section *Flavi* unable to produce CPA are believed to have deletions of the biosynthetic pathway silencing its production, such as *A. parasiticus*, *A. nomius* and their closely related species (Moore et al. 2016). Tenuazonic acid is a non-mutagenic mycotoxin that causes tremors, diarrhea, vomiting and hemorrhages. In section *Flavi*, it is reported in the clades *A. nomius* and *A. tamarii* (Table 3) and some strains of *A. novoparasiticus*. Aflatrem is another emergent mycotoxin, classified as a potent tremorgenic compound that causes neurological disorders. It seems that it might interfere with the release of neurotransmitters by receptors in the central and peripheral nervous systems (Zhang et al. 2004).

Two common extrolites produced by several aflatoxigenic species in section *Flavi* are kojic acid and aspergillic acid; the first organic acid is used in various industrial applications, especially in cosmetic and health care industries (Mohamad et al. 2010). Kojic acid is only not produced by *A. togoensis* and *A. pipericola*, whereas aspergillic acid is not produced by *A. austwickii*, *A. pipericola*, *A. pseudocaelatus*, *A. transmontanesis*, *A. pseudotamarii* and *A. togoensis* (Table 2). The lack of aspergillic acid production is characteristic of *A. tamarii* clade (Varga et al. 2011, Frisvad et al. 2019). Leporins are also found in several *Flavi* section species and are proven to have anti-insectan activity (Cary et al. 2015 a, b). Sclerotia extrolites include some mycelium extrolites and some unique extrolites; among them some reported in aflatoxigenic species are AFs

and aflatrems (already mentioned above), aflavazole, aflavinines, aflavarins, anominine, aspernomine, and paspalines (Table 2) (Cary et al. 2015b, Frisvad et al. 2019). Aflavinines and aflavarins are frequently reported, especially in species closely related with *A. flavus*. They have anti-insectan activity (Gloer et al. 1988), and are suggested to have key ecological roles in species survival (TePaske and Gloer 1992, Cary et al. 2015b).

2.3 Molecular Markers and Phylogenetic Analyses

The inclusion of molecular markers for species identification in *Aspergillus* section *Flavi* is proven to be advantageous, especially for cryptic species identification. It helped not only to unmask diversity but also to understand the relationships within *Aspergillus* section *Flavi*. Genes tested with fruitful results include *benA*, *cmdA* and *RPB2*. These markers are advantageous because they have conserved and variable regions and are widely described in literature (primers and sequences) (Varga et al. 2011, Frisvad et al. 2019). Phylogenetic studies showed that most aflatoxigenic species are derived species. A recent study performed by Frisvad et al. (2019) showed that aflatoxigenic species are grouped in three main clusters: *A. flavus*-clade, *A. tamaritii*-clade and *A. nomius*-clade (Table 3). *A. togoensis* is a basal species and is not considered as a food contaminant.

Species from *A. flavus*-clade can be sub-divided in two main clusters, one that includes species with *A. flavus* overlapping traits and the other that includes species with *A. parasiticus* overlapping traits, and *A. mottae* as its basal species (Table 3). *A. flavus* sub-clade species number has increased dramatically over the last decade, and as aforementioned, it is composed of cryptic species (overlapping morphological, genetic and chemotype traits) (Frisvad et al. 2005, 2019, Varga et al. 2011). Within this sub-group four clusters are formed: (1) *A. flavus*, (2) *A. minisclerotigenes* and *A. texensis*, (3) *A. pipericola* and (4) a cluster including *A. aflatoxiformans*, *A. korhogoensis* and *A. austwickii* (Varga et al. 2011, Carvajal-Campos et al. 2017, Singh et al. 2018, Frisvad et al. 2019). *A. oryzae*, considered as a domesticated variant of *A. flavus* for a long time, forms a subgroup based on the three above-mentioned genes. The origin of *A. oryzae* is controversial. A recent study based on 200 monocore genes supports an earlier hypothesis that it is closer to *A. aflatoxiformans* and *A. minisclerotigenes* (Kjaerbolling et al. 2020), thus, another study based on 82 genomes of *A. oryzae* strains used in the manufacture of Asian fermented foods confirmed that it formed a monophyletic cluster nested in *A. flavus* clade (Watarai et al. 2019). AFBG strains with small sclerotia are commonly misclassified as *A. flavus* S_{BG} morphotype; however phylogenetic surveys revealed that this classification includes several cryptic species related to *A. flavus*: *A. minisclerotigenes*, *A. texensis*, *A. aflatoxiformans*, *A. korhogoensis*, *A. pipericola* and *A. austwickii* (Pildain et al. 2008, Frisvad et al. 2005, 2019, Varga et al. 2011, Soares et al. 2012, Perrone et al. 2014, Carvajal-Campos et al. 2017, Singh et al. 2018, Singh and Cotty 2019). *A. parasiticus* sub-clade species are AFBG producers. Their phylogenetic relationships are more complicated to explain as they present slight differences depending on the molecular markers used. Anyhow, *A.*

Table 3: Main clades of aflatoxigenic *Aspergillus* section *Flavi*

Clade	Sub-clade	Species
<i>A. flavus</i> -clade	Group of species closely related to <i>A. flavus sensu stricto</i>	<i>A. flavus</i> , <i>A. aflatoxiformans</i> , <i>A. austwickii</i> , <i>A. korhogoensis</i> , <i>A. minisclerotigenes</i> , <i>A. pipericola</i> , <i>A. texensis</i>
	Group of species closely related to <i>A. parasiticus</i>	<i>A. parasiticus</i> , <i>A. arachidicola</i> , <i>A. novoparasiticus</i> , <i>A. sergii</i> , <i>A. transmontanensis</i>
	Basal species to the <i>A. flavus</i> and <i>A. parasiticus</i> groups	<i>A. mottae</i>
<i>A. tamarii</i> -clade	-	<i>A. tamarii</i> , <i>A. caelatus</i> , <i>A. pseudocaelatus</i> , <i>A. pseudotamarii</i>
<i>A. nomius</i> -clade	-	<i>A. nomius</i> , <i>A. luteovirenses</i> , <i>A. pseudonomius</i>

parasiticus and *A. sojae* (domesticated species) cluster together. Both species are closely related to *A. arachidicola* and *A. novoparasiticus*, the latter being morphologically similar to *A. parasiticus*. *A. sergii* and *A. transmontanensis* are the basal species of this sub-clade. Finally, *A. mottae* sets as the basal species of the *A. flavus* and *A. parasiticus* sub-clades, though it resembles the species from the *A. parasiticus* sub-clade, sharing morphological, genetic and chemical traits (Varga et al. 2011, Soares et al. 2012, Carvajal-Campos et al. 2017, Frisvad et al. 2019).

A. tamarii-clade species produce AFBG with the exception of *A. tamarii* and *A. caelatus*. Mature colonies of *A. pseudotamarii* resemble morphologically to the *A. tamarii* and *A. pseudocaelatus* overlap traits with its sister taxon, *A. caelatus* (Varga et al. 2011).

A. nomius-clade is composed only by AFBG species. *A. pseudonomius* (Varga et al. 2011) is the sibling species of *A. nomius*, and both overlap traits, risking misidentification (Peterson et al. 2001, Moore et al. 2016). *A. luteovirenses* also share traits with species from its cluster (Peterson et al. 2001). Although *A. luteovirenses* produces AFBG, it is not considered as a pathogen for humans or animals as it occurs less frequently in food and feedstuff (Varga et al. 2015, Moore et al. 2016). *A. nomius*, *A. pseudonomius*, and *A. luteovirenses* lack production of CPA, which has been suggested as a fixed trait in the cluster (Varga et al. 2011, Frisvad et al. 2015).

3. Aflatoxigenic *Aspergillus* section *Flavi* Biodiversity According to Food Matrices, Production Chain Stage and Geographical Areas

The presence of aflatoxigenic *Aspergillus* section *Flavi* species in various foodstuff (cereals, dried fruits, legumes, oilseeds, roots and tubers, spices, tree nuts, etc.) and on all continents (Africa, Asia, Oceania, Europe and America)

are highlighted by several studies (Table 4). Species distribution is not homogenous, a higher diversity is found in tropical and subtropical regions, as environmental conditions enhance their development (Schatzmayr and Streit 2013), *Aspergillus* section *Flavi* species physiological requirements of moisture are around 0.85 to 0.99 a_w , while optimal temperatures are between 28 and 42°C (Medina et al. 2015, Yogendrarajah et al. 2016). Aflatoxigenic species of section *Flavi* are mainly saprophytic and inhabit numerous ecological niches, like soil, decaying vegetation, hay and seeds, agricultural fields, and stored products. Fungal contamination of staples occurs during pre-harvest steps (crops and recollect) and/or in post-harvest steps (mainly storage and manufacturing production processes) (Rodrigues et al. 2012).

3.1 Biodiversity According to Food Matrices and Geographical Areas

The aflatoxigenic *Aspergillus* section *Flavi* species reported worldwide in the main food matrices are listed in Table 4. Species in this section were also reported in other food crops like *A. flavus* from acha in Nigeria, canola in Serbia, mustard seeds and chilgoza pine nuts in India (Tomar-Balhara et al. 2006, Sharma et al. 2013, Škrinjar et al. 2013, Ezekiel et al. 2014). In sugarcane, besides of *A. flavus*, *A. novoparasiticus* was reported in Egypt, Brazil, Japan and USA while *A. parasiticus* and *A. arachidicola* only in Asian countries and Brazil, respectively (Takahashi et al. 2004, Kumeda et al. 2004, Ahmed et al. 2010, Garber and Cotty 2014, Iamanaka et al. 2019, Abdallah et al. 2020). *A. flavus*, *A. parasiticus* and *A. nomius* were isolated from cocoa beans in Brazil (Copetti et al. 2011) and *A. aflatoxiformans* from edible mushrooms in Nigeria (Ezekiel et al. 2013a).

3.2 Cereals

AQ : 2003 in References

AQ ; missing in References

3.2.1 Maize

Worldwide, *A. flavus* is reported as the most common species from section *Flavi* through the maize production chain (field, storage, derived products) (Table 4), including aflatoxigenic and non-aflatoxigenic isolates (Probst et al. 2014, Kachapulula et al. 2017). Maize plants are sensitive to drought and temperature, by consequence prone to fungal infection under plant stress conditions. *A. flavus* is considered to have a commensal relationship with this crop (Kornher 2018, Taniwaki et al. 2018); nevertheless, it is not the only species from section *Flavi* contaminating maize, *A. parasiticus* is the second most important contaminant, and occasionally some other species are reported (Table 4). In Africa, the presence of aflatoxigenic species on maize are mainly reported at post-harvest steps, with *A. flavus*, *A. parasiticus* and S_{BC} isolates as the most important contaminates. *A. flavus* L-morphotype strains are more common, including aflatoxigenic and non-aflatoxigenic strains (Probst et al. 2014, Kachapulula et al. 2017). S_{BC} isolates in the region are more probably *A. aflatoxiformans* and *A. minisclerotigenes* (in this order). *A. minisclerotigenes*

Table 4: Aflatoxigenic *Aspergillus* section *Flavi* species according to food matrices, production chain stage and geographical areas

Crops	Production chain step	Geographic zone		Aflatoxigenic <i>Aspergillus</i> species	References
Cereals					
Barley	Post-harvest: Grains**	Asia	Pakistan	<i>A. flavus</i>	Fakhrunnisa et al. 2006
		Europe	Spain	<i>A. flavus</i> <i>A. parasiticus</i>	Mateo et al. 2011
Maize	Post-harvest: kernels from storage or market	Africa	Nigeria, South Africa	<i>A. flavus</i>	Atehnkeng et al. 2008, Chilaka et al. 2012, Ekwomadu et al. 2017
			Egypt, Kenya	<i>A. flavus</i> *	El-Shanshoury et al. 2014, Gachara et al. 2018, Abbas et al. 2020
			Cameroon	<i>A. flavus</i> <i>A. parasiticus</i>	Njobeh et al. 2009
			Kenya	<i>A. flavus</i> <i>A. minisclerotigenes</i> <i>A. parasiticus</i>	Okoth et al. 2018
			Ghana	<i>A. flavus</i> <i>A. parasiticus</i> Sub isolates	Agbetiamah et al. 2018
			Nigeria	<i>A. korhogoensis</i>	Frisvad et al. 2019
	Non-specified step: at harvest and kernels from market		Cameroon, Ghana, Mali, Rwanda, Sierra Leona, Tanzania, Uganda	<i>A. flavus</i>	Probst et al. 2014

(Contd.)

AQ : publishing date 2018 in References below

AQ : missing in references

Table 4: (Contd.)

Crops	Production chain step	Geographic zone	Malawi, Zambia	Aflatoxigenic Aspergillus species	References
				<i>A. flavus</i> <i>A. parasiticus</i>	Probst et al. 2014, Kachapulula et al. 2017
			Burkina Faso, Ethiopia, Senegal	<i>A. flavus</i> S _{BC} isolates	Diedhiou et al. 2011, Probst et al. 2014
			Congo, Kenya, Nigeria	<i>A. flavus</i> <i>A. minisclerotigenes</i> S _{BC} isolates	Probst et al. 2014
			Mozambique, Zambia, Zimbabwe	<i>A. flavus</i> <i>A. parasiticus</i> S _{BC} isolates	Probst et al. 2014
			Somalia	<i>A. minisclerotigenes</i>	Probst et al. 2014
	Pre-harvest: at harvest	Asia	China, Pakistan	<i>A. flavus</i>	Saleem et al. 2012, Mamo et al. 2018
	Post-harvest: fresh kernels, corn meal		South Korea	<i>A. flavus</i>	Frisvad et al. 2019
			China, Vietnam	<i>A. flavus</i> *	Gao et al. 2007, Trung et al. 2008, Tran-Dinh et al. 2009
			Pakistan, Philippines	<i>A. flavus</i> <i>A. parasiticus</i>	Ibrahim et al. 2016, Balendres et al. 2019
	Non-specified step		China	<i>A. flavus</i> <i>A. arachidicola</i> <i>A. novoparasiticus</i> <i>A. pseudonomius</i>	Rasheed et al. 2019

Pre-harvest: at harvest	Central and South America	Mexico, Venezuela	<i>A. flavus</i>	Mazzani et al. 2008, Ortega-Beltran et al. 2015
		Argentina	<i>A. flavus</i> <i>A. parasiticus</i>	Etcheverry et al. 1999, Nesci and Etcheverry 2002
Post-harvest: fresh kernels, corn meal		Colombia	<i>A. flavus</i>	Acuña et al. 2005
		Argentina, Ecuador	<i>A. flavus</i> <i>A. parasiticus</i>	Etcheverry et al. 1999, Pacin et al. 2003
Pre-harvest: at harvest	Europe	Argentina	<i>A. flavus</i> <i>A. parasiticus</i> S _{BC} isolates	Camiletti et al. 2017
		France	<i>A. flavus</i> <i>A. parasiticus</i>	Bailly et al. 2018
Post-harvest: kernels from storage and market		Hungary, Italy, Serbia	<i>A. flavus</i>	Giorni et al. 2007, Dobolyi et al. 2013, Mauro et al. 2013, Baranyi et al. 2015a
		France, Hungary	<i>A. flavus</i> <i>A. parasiticus</i>	Sebők et al. 2014, Bailly et al. 2018
Pre-harvest: at harvest	North America	Portugal	<i>A. flavus</i> <i>A. mottiae</i>	Soares et al. 2012
		Serbia	<i>A. pseudonomius</i>	Jakić-Dimić et al. 2009
		USA	<i>A. flavus</i>	Probst et al. 2014
			<i>A. flavus</i> , <i>A. parasiticus</i>	Jaime-Garcia and Cotty 2004, 2010

(Contd.)

Table 4: (Contd.)

Crops	Production chain step	Geographic zone		Aflatoxigenic Aspergillus species	References
	Post-harvest: kernels**			<i>A. flavus</i>	Frisvad et al. 2019
	Non-specified step: kernels			<i>A. texensis</i>	Singh et al. 2018
Millet	Post-harvest: kernels**	Oceania	Australia	<i>A. flavus</i>	Egmond and Jonker 2004
	Post-harvest: grains from storage or markets	Africa	Nigeria	<i>A. flavus</i>	Osamwonyi and Wakil 2012, Ezekiel et al. 2014
Rice	Post-harvest: grains from market	Africa	Egypt	<i>A. flavus</i> *	El-Shanshoury et al. 2014
		Asia	Nigeria	<i>A. aflatoxiformans</i> <i>A. australickii</i> <i>A. korlagoensis</i>	Frisvad et al. 2019
	Pre-harvest: field		China	<i>A. flavus</i>	Mamo et al. 2018
	Post-harvest: grains from market		Cambodia, India, Indonesia, Malaysia, Saudi Arabia, Thailand	<i>A. flavus</i>	Jayaraman and Kalyanasundaram 1990, Reddy et al. 2010, Al-Husnan et al. 2019
Non-specified step			South Korea, Pakistan	<i>A. flavus</i> <i>A. parasiticus</i>	Park et al. 2005, Ibrahim et al. 2016
			Thailand	<i>A. flavus</i>	Frisvad et al. 2019
			Philippines	<i>A. flavus</i> <i>A. parasiticus</i>	Sales and Takumi 2005
Pre-harvest: fields		South America	Brazil	<i>A. flavus</i> <i>A. arachidicola</i> <i>A. novoparasiticus</i> <i>A. pseudocaelatus</i>	Katsurayama et al. 2018

	Post-harvest: grains from market				<i>A. flavus</i>	Katsurayama et al. 2018
Sorghum	Post-harvest: stored grains	Africa	Ethiopia		<i>A. flavus</i> <i>A. parasiticus</i>	Weledesemayat et al. 2016
	Non-specified step: seeds	Asia	Turkey, India, Pakistan,		<i>A. flavus</i>	Fakhrunnisa et al. 2006, Turgay and Ünal 2009, Divakara et al. 2014
Wheat	Post-harvest: stored grains	Europe	Serbia		<i>A. flavus</i>	Jakić-Dimić et al. 2009
	Pre-harvest: kernels**	Africa	Algeria		<i>A. flavus</i>	Riba et al. 2008, 2010
	Post-harvest: stored grains, grains and flour from market		Algeria, Egypt, Morocco		<i>A. flavus</i>	Riba et al. 2008, 2010, Dahab et al. 2016, Ennouari et al. 2018
			Egypt		<i>A. flavus</i> *	El-Shanshoury et al. 2014
	Pre-harvest: grains**	Asia	Pakistan		<i>A. flavus</i>	Fakhrunnisa et al. 2006, Hussain et al. 2013
	Post-harvest: stored grains, grains and flour from market		India, Iran, Saudi Arabia, Turkey		<i>A. flavus</i>	Sinha and Sinha 1990, Saberi-Riseh et al. 2004, Doolotkeldieva 2010, Al-Wadai et al. 2013, Joshaghani et al. 2013
			Iran, Pakistan		<i>A. flavus</i> <i>A. parasiticus</i>	Kachuei et al. 2009, Ibrahim et al. 2016

(Contd.)

Table 4: (Contd.)

Crops	Production chain step	Geographic zone		Aflatoxigenic Aspergillus species	References
	Pre-harvest: grains**	South America	Argentina, Brazil	<i>A. flavus</i>	González et al. 2008, Savi et al. 2014
	Post-harvest: stored grains	Europe	Serbia	<i>A. flavus</i>	Jakić-Dimić et al. 2009
		North America	Slovakia	<i>A. parasiticus</i>	Dovičková et al. 2012
			Canada	<i>A. flavus</i>	Wallace and Sinha 1962
Dried Fruits					
Apricot	Post-harvest: dried fruit from market	Africa	Egypt	<i>A. flavus</i>	Zohri and Abdel-Gawad 1993
		Asia	Iraq	<i>A. flavus</i> <i>A. parasiticus</i>	Saadulah and Abdullah 2015
Figs	Post-harvest: dried fruit from market	Africa	Egypt	<i>A. flavus</i>	Zohri and Abdel-Gawad 1993
			Algeria	<i>A. flavus</i> , <i>A. parasiticus</i>	Ait Mimoune et al. 2018
		Asia	Iran, Turkey	<i>A. flavus</i>	Şenyuva et al. 2008, Javanmard 2010
Plum	Post-harvest: dried fruit from market		Iraq	<i>A. flavus</i> <i>A. parasiticus</i>	Saadulah and Abdullah 2015
		Africa	Egypt	<i>A. flavus</i>	Zohri and Abdel-Gawad 1993

	Asia	Iraq	<i>A. flavus</i> <i>A. parasiticus</i>	Saadulah and Abdullah 2015
Raisins	Africa	Egypt	<i>A. flavus</i>	Zohri and Abdel-Gawad 1993
	Asia	Iraq	<i>A. flavus</i> <i>A. parasiticus</i>	Saadulah and Abdullah 2015
Legumes				
Beans	Africa	Cameroon	<i>A. flavus</i> <i>A. parasiticus</i>	Njobeh et al. 2009
	South America	Brazil	<i>A. flavus</i> <i>A. parasiticus</i>	Costa and Scussel 2002
Soybeans	Africa	Cameroon	<i>A. flavus</i> <i>A. parasiticus</i>	Njobeh et al. 2009
Oilseeds				
Cotton	North America	USA	<i>A. flavus</i>	Cotty 1997
	Asia	India	<i>A. flavus</i>	Klich et al. 1986
	Africa	Egypt	<i>A. flavus</i>	Mazen et al. 1990, Tomar et al. 2012
Peanuts		Egypt	<i>A. flavus</i>	Embaby and Abdel-Galel 2014
		Ethiopia, Nigeria	<i>A. flavus</i> <i>A. parasiticus</i>	Mohamed and Chala 2014, Guchi 2015, Wartu et al. 2015

(Contd.)

Table 4: (Contd.)

Crops	Production chain step	Geographic zone		Aflatoxigenic Aspergillus species	References
	Post-harvest: dried peanuts, peanuts from storage; peanuts, butter or cake from markets		Côte d'Ivoire	<i>A. aflatoxiformans</i> <i>A. korhogoensis</i>	Carvajal-Campos et al. 2017, Manizan 2019
			Algeria, Benin, Cameroon, Congo, Côte d'Ivoire, Egypt, Ethiopia, Nigeria, South Africa	<i>A. flavus</i> <i>A. parasiticus</i>	Njobeh et al. 2009, Sultan and Magan 2010, Adjou et al. 2012, Boli et al. 2014, Kamika et al. 2014, Mohamed and Chala 2014, Warty et al. 2015, Ait Mimoune et al. 2018
			Botswana, Ghana, Nigeria	<i>A. flavus</i> <i>A. parasiticus</i> S _{BC} isolates	Mphande et al. 2004, Ezekiel et al. 2013b, Agbetiamah et al. 2018,
			Ethiopia	<i>A. flavus</i> <i>A. parasiticus</i> <i>A. nomius</i>	Guchi 2015
	Pre-harvest: fields	Asia	Nigeria	<i>A. aflatoxiformans</i>	Frisvad et al. 2019
			China, Indonesia, Vietnam	<i>A. flavus</i>	Tran-Dinh et al. 2009, Mamo et al. 2018, Frisvad et al. 2019
	Post-harvest: stored peanuts and peanut-based products		Malaysia	<i>A. flavus</i> <i>A. nomius</i>	Norlia et al. 2018

AQ : only 2012 in references

	Pre-harvest: plants and grains from field	South America	Argentina, Brazil	<i>A. flavus</i> <i>A. arachidicola</i> <i>A. minisclerotigenes</i> <i>A. parasiticus</i>	Vaamonde et al. 2003, Barros et al. 2003, Pildain et al. 2008, Martins et al. 2017
	Post-harvest: drying processes and storage		Argentina	<i>A. flavus</i>	Vaamonde et al. 2003, Barros et al. 2003, Pildain et al. 2008
	Pre-harvest: fields	North America	USA	<i>A. flavus</i> <i>A. parasiticus</i>	Jaime-Garcia and Cotty 2004, 2010 Horn 2007
	Post-harvest: drying processes and storage				
	Pre-harvest: fields	Oceania	Australia	<i>A. flavus</i> <i>A. minisclerotigenes</i> <i>A. parasiticus</i>	Pitt and Hocking 2006, Frisvad et al. 2019
	Post-harvest: drying and storage				Pitt and Hocking 2006, Frisvad et al. 2019
Sesame	Post-harvest: grains from storage and market	Africa	Nigeria	<i>A. flavus</i> <i>A. aflatoxiformans</i>	Ezekiel et al. 2014
			Nigeria	<i>A. aflatoxiformans</i> <i>A. austwickii</i>	Frisvad et al. 2019
			Senegal	<i>A. flavus</i> S _{BC} isolates	Diedhiou et al. 2011
			Mexico	<i>A. aflatoxiformans</i>	Frisvad et al. 2019
Roots and Tubers					
Cassava	Post-harvest: chips	Africa	Benin	<i>A. flavus</i>	Gnonlonfin et al. 2008

(Contd.)

Table 4: (Contd.)

Crops	Production chain step	Geographic zone	Geography	Aflatoxigenic Aspergillus species	References
Yam			Benin	<i>A. flavus</i> <i>A. aflatoxiformans</i> <i>A. novoparasiticus</i>	Adjovi et al. 2014
			Cameroon	<i>A. flavus</i> <i>A. parasiticus</i> <i>A. nomius</i>	Onana et al. 2013
		Africa	Benin, Nigeria	<i>A. flavus</i>	Bankole and Mabekeje 2004, Gnonlonfin et al. 2008
Spices					
Chili peppers	Post-harvest: market	Africa	Benin, Mali, Nigeria, Togo	<i>A. flavus</i>	Hell et al. 2009, Ezekiel et al. 2019
			Morocco	<i>A. flavus</i> <i>A. minisclerotigenes</i>	El Mahgubi et al. 2013
			Nigeria	<i>A. flavus</i> <i>A. aflatoxiformans</i> <i>A. minisclerotigenes</i> <i>A. parasiticus</i>	Singh and Cotty 2019
		Asia	Pakistan	<i>A. flavus</i> <i>A. parasiticus</i>	Ibrahim et al. 2016
			India, Lebanon	<i>A. flavus</i>	Sharfun-Nahar et al. 2004, Makhoul et al. 2019
Cumin	Post-harvest: market	Africa	Morocco	<i>A. flavus</i> <i>A. minisclerotigenes</i>	El Mahgubi et al. 2013

Curry	Post-harvest: market	Asia	Lebanon	<i>A. flavus</i>	Makhlouf et al. 2019
Pepper	Post-harvest: market	Africa	Morocco	<i>A. flavus</i> <i>A. minisclerotigenes</i>	El Mahgubi et al. 2013
	Post-harvest: market	Asia	Lebanon, Turkey	<i>A. flavus</i>	Probst et al. 2014, Makhlouf et al. 2019
	Post-harvest: market	South America	Brazil	<i>A. flavus</i>	Freire et al. 2000
Tree Nuts					
Almonds	Post-harvest: storage and market	Africa	Algeria	<i>A. flavus</i> <i>A. parasiticus</i>	Ait Mimoune et al. 2018
	Pre-harvest: orchards and at harvest	Europe	Portugal	<i>A. flavus</i> <i>A. minisclerotigenes</i> A. <i>parasiticus</i> <i>A. sergii</i>	Rodrigues et al. 2009, 2012, Soares et al. 2012
	Post-harvest: drying, storage			<i>A. flavus</i> <i>A. minisclerotigenes</i> A. <i>parasiticus</i> <i>A. sergii</i> <i>A. transmontanensis</i>	Rodrigues et al. 2009, 2012, Soares et al. 2012
	Pre-harvest: orchards and at harvest	North America	USA	<i>A. flavus</i> <i>A. parasiticus</i>	Purcell et al. 1980, Bayman et al. 2002, Doster et al. 2014
	Post-harvest: drying, storage				Bayman et al. 2002, Doster et al. 2014

(Contd.)

Table 4: (Contd.)

Crops	Production chain step	Geographic zone	Country	Aflatoxigenic Aspergillus species	References
Brazil nuts	Pre-harvest: at harvest	South America	Peru	<i>A. flavus</i>	Taniwaki et al. 2017
			Brazil	<i>A. flavus</i> <i>A. nomius</i>	Taniwaki et al. 2017
	Post-harvest: drying, storage		Brazil	<i>A. flavus</i> <i>A. arachidicola</i> <i>A. luteovirescens</i> <i>A. nomius</i> <i>A. parasiticus</i> <i>A. pseudocaelatus</i> <i>A. pseudonomius</i> <i>A. pseudotamarii</i>	Freire et al. 2000, Pacheco et al. 2010, Freitas-Silva et al. 2011, Reis et al. 2012, Andersson 2012, Baquião et al. 2013, Calderari et al. 2013, Massi et al. 2014, Taniwaki et al. 2017
Hazelnuts	Pre-harvest: orchards and at harvest	Europe	Italy	<i>A. flavus</i> <i>A. parasiticus</i>	Rodrigues et al. 2012, Prencipe et al. 2018
	Post-harvest: drying process, storage				
Nuts	Post-harvest	Europe	Croatia, Hungary	<i>A. flavus</i>	Baranyi et al. 2015a
	Pre-harvest: orchards	Asia	Turkey	<i>A. flavus</i>	Heperkan et al. 1994, Kabirian et al. 2011
			Iran	<i>A. flavus</i> * <i>A. parasiticus</i>	Rahimi et al. 2007

	Post-harvest: drying, storage		Turkey	<i>A. flavus</i>	Heperkan et al. 1994, Kabirian et al. 2011
	Pre-harvest: orchards	North America	USA	<i>A. flavus</i>	Bayman et al. 2002
	Post-harvest: market				Bayman et al. 2002

* The survey results may include other species closely related to *A. flavus* S_{BC} isolates correspond to *A. flavus* closely related species that produce AFBG and have small sclerotia (*A. minisclerotigenes*, *A. aflatoxiformans*, *A. korhogoensis* or an unknown non-characterized species) and were generally identified as *A. flavus*.

** Step not specified

was properly identified in Kenya, Congo, Nigeria and Somalia, whereas *A. korhogoensis* was reported in Nigeria (Table 4).

In Asia, *A. flavus* is the main contaminant but not all strains are AF producers; other species reported less frequently include *A. parasiticus*, *A. arachidicola*, *A. novoparasiticus* and *A. pseudonomius* (Table 4). Surveys in China and Vietnam may unmask diversity, as *A. flavus* isolates have been reported with atypical chemotypes and morphological traits (Gao et al. 2008, Trung et al. 2008, Tran-Dinh et al. 2009). Rasheed et al. (2019) showed that diversity in the region is sub-estimated by reporting *A. arachidicola*, *A. novoparasiticus* and *A. pseudonomius*; whereas in Philippines, Balendres et al. (2019) reported *A. flavus* and *A. parasiticus* from maize and soil field samples. Similarly, in Pakistan, surveys from maize samples reported *A. flavus* and *A. parasiticus* (Table 4), while in Iran samples from maize soil fields showed *A. flavus* as the most frequent species, followed by *A. parasiticus* and *A. nomius* (Razzaghi-Abyaneh et al. 2006). In Central and South America, *A. flavus* and *A. parasiticus* are pointed as the major aflatoxigenic species infecting maize (Table 4). Ortega-Beltran et al. (2015) reported *A. flavus* S- and L-morphotypes and a S_{BG} strain from maize soil in Sonora (Mexico). In Argentinean crops, *A. flavus* L-morphotype was frequent in pre-harvest steps, whereas *A. parasiticus* aflatoxigenic strains in pre-planted crops (soil with plant debris). Camiletti et al. (2017) suggested that infection by aflatoxigenic *Aspergillus* section *Flavi* species increases while the flowering period is hot in some Argentinean regions. It is highly possible that the S_{BG} isolate reported in Argentina corresponds to *A. minisclerotigenes* because it has been isolated from the same region in other staples. In Europe, over the last decade, maize contamination by aflatoxigenic species from section *Flavi* has become an increasing issue. Climate change in Europe has a seasonal impact, as seasons are becoming hotter and drier, especially in the Southern, Central and Eastern regions, increasing the chances of *A. Flavi* species colonization (EFSA 2013, Battilani et al. 2016, Moretti et al. 2019). Despite the fact that *Fusarium* spp. such as *F. verticillioides* and *F. graminearum* are still the main contaminants of maize in Europe, the reports of AFs are rising (Giorni et al. 2007, Covarelli et al. 2011, Gallo et al. 2012, Mauro et al. 2013, Dobolyi et al. 2013, Baranyi et al. 2015a, Bailly et al. 2018, Kos et al. 2018). In European maize, *A. flavus* and *A. parasiticus* are the most frequent species (Table 4), yet *A. parasiticus* infection may be favored as cooler temperature suits its physiological requirements better (Horn 2007, Bailly et al. 2018). Also, *Aspergillus* section *Flavi* diversity seems to rise in storage conditions, at present *A. flavus* (non-aflatoxigenic and aflatoxigenic strains) and *A. parasiticus* (mainly aflatoxigenic strains) have been reported in fields and at harvesting steps while *A. flavus* (non-aflatoxigenic and aflatoxigenic strains), *A. parasiticus*, *A. pseudonomius* and *A. mottae* in silos and storage samples (Table 4). In the USA, *A. flavus* is the main species recovered from maize fields (soil, debris and ears) and stored maize kernels, followed by *A. parasiticus* and less frequent *A. texensis*. *A. flavus* diversity in the region includes S- and L-morphotypes, and aflatoxigenic and non-aflatoxigenic strains (Bayman et al. 2002, Jaime-Garcia and Cotty 2004,

2010), being the S-morphotype more frequent and generally aflatoxigenic. In Australia, *Fusarium* spp. are prone to contaminate maize as environmental conditions favors its development, yet *A. flavus* has been reported to occur in soil and kernels (Egmond and Jonker 2004).

3.2.2 Rice

Globally, rice is the second most produced and consumed cereal after maize, and *A. flavus* the most frequent contaminant species from the section *Flavi*. In Asia, *A. flavus* is the most common species reported, followed by *A. parasiticus* (Table 4), yet its identification has been based principally in morphological characteristics. In Africa, Frisvad et al. (2019) reported the presence of *A. aflatoxiformans* and two rare species from Nigerian markets, *A. korhogoensis* and *A. austwickii*; while in South America, Katsurayama et al. (2018) reported *A. novoparasiticus*, *A. arachidicola*, *A. pseudocaelatus* and *A. flavus* from rice and rice plantation soil in Brazil (Table 4). The proportion of aflatoxigenic *A. flavus* strains reported was low (1.5% of the isolated *A. flavus*), yet most of the strains produced CPA (Katsurayama et al. 2018). In Europe, rice is produced in some southern regions, like Spain and France, and even though surveys targeting *Aspergillus* section *Flavi* diversity in this staple do not exist to our knowledge. A study by Suárez-Bonnet et al. (2013) isolated the four AFs B and G from Spanish rice samples and traces of AFGs from French rice samples, suggesting a contamination caused by other species than *A. flavus*.

3.2.3 Wheat and Barley

Wheat and barley are the third and fourth cereals more produced and consumed around the world. Both cereals are more prone to be infected by *Fusarium* spp. hence most studies target these species; despite some reports based on morphology include *A. flavus* and *A. parasiticus* (Table 4).

3.2.4 Millet and Sorghum

Millet and sorghum are better adapted to harsh environments than other cereals and are common crops in Asia and Africa (FAO 1995). The main aflatoxigenic species from section *Flavi* reported to occur in these staples is *A. flavus* and *A. parasiticus* was identified only in stored sorghum grains from Ethiopia (Table 4), but reports of AFBG occurrence in grains and byproducts suggest species misidentification.

3.3 Dried Fruits and Legumes

Contamination of these food matrices by aflatoxigenic species from *Flavi* section was reported from market samples. Based on morphological description, *A. flavus* is the most common identified species in dried apricot, figs, raisins and plums collected in North Africa and Asia, followed by *A. parasiticus*. Both *A. flavus* and *A. parasiticus* were reported in all dried beans and soybeans samples from Cameroon and Brazil (Table 4).

3.4 Oilseeds

3.4.1 Peanuts

A. parasiticus and *A. flavus* are commonly reported in peanuts through the production chain (Table 4), both reported as commensal species of peanut plant (Pitt et al. 2012). The presence of fungal propagules in soil increases infection risk, notably if peanut crop suffers drought stress or related factors, raising the risk of high levels of AF production before harvest. Depending on the region, other species were also isolated in high frequencies (Table 4). In addition, the methods used for identification might fall to underestimate cryptic species and interesting populations, like S_{BC} species (Table 4). In Africa, many studies indicated the presence of *A. flavus* and *A. parasiticus* in peanuts during pre- and post-harvest stages including manufactured products such as cake and butter. Other species reported to infect peanuts during post-harvest include *A. aflatoxiformans*, *A. korhogoensis* and *A. nomius* (Table 4). Slow drying and poor storage conditions are a serious issue in the humid tropics, as moisture absorption might favor aflatoxigenic fungal infection after harvest (Pitt et al. 2012, Taniwaki et al. 2018). In Asia, *A. flavus* was found to be the most common species isolated from peanuts along the production chain and the most common mycotoxigenic component from the mycobiota (Table 4). In Malaysia, *A. nomius* and an atypical *A. flavus* L-morphotype were isolated in post-harvest steps (Table 4). In South America, studies are scarce though peanuts are widely consumed. In Argentina, the principal infectious species during pre-harvest from *Aspergillus* section *Flavi* were *A. flavus* (aflatoxigenic and non-aflatoxigenic strains) and *A. parasiticus*. Yet, *A. arachidicola* and *A. minisclerotigenes* were also present, whereas only *A. flavus* was identified during post-harvest (Table 4). In Brazil, throughout the peanut production chain, *A. flavus* was the most frequent species from section *Flavi*, followed by *A. parasiticus* (Martins et al. 2017). Nevertheless, species prevalence varied among the production chain stages. *A. flavus* prevalence was higher at sorting stage (78.3%), followed by drying (63.2%), threshing (54.5%), blanching (47.6%), field (40.7%) and ready-to-eat products (31.3%), while *A. parasiticus* was reported at drying (12.5%) and in ready-to-eat (1.3%) samples. This study also tested the aflatoxigenic potential of *A. flavus* strains, around 50% of samples were aflatoxigenic, and were isolated more frequently from fields than from processing plants (83% and 31%, respectively) (Martins et al. 2017). In northern USA, *A. flavus* and *A. parasiticus* were reported as the main peanut contaminants, both occurring in fields and after harvest (Table 4). *A. flavus* follows the same trend observed for other crops in USA, being S-morphotype more common. In Australia, studies on peanuts have reported the presence of *A. flavus*, *A. parasiticus* and *A. minisclerotigenes* (the latter especially in soil samples), during pre- and post-harvest steps (Pitt and Hocking 2006).

3.4.2 Cotton and Sesame

Reports from these staples are poor, probably the best know from them is cotton from USA. Cotton is contaminated by *A. flavus* (Table 4), mainly by

A. flavus S-morphotype isolates (Cotty 1997). Sesame contamination during post-harvest steps in Africa are caused by *A. flavus* and S_{BC} isolates (*A. aflatoxiformans* and the newly described species *A. austwickii*) (Table 4).

3.5 Roots and Tubers

Cassava and yam are important staples in Africa. Cassava is ranked as the third most important food crop in tropical regions after rice and maize. The conditions of production and storage of cassava, yam, and their traditional derivatives favor contamination and development of fungi (Adjovi et al. 2014). *A. flavus* is the most frequent species isolated, though *A. aflatoxiformans*, *A. novoparasiticus*, *A. nomius* and *A. parasiticus* were also reported from cassava chips (Table 4).

3.6 Spices

Post-harvest practices and environmental conditions during growth make spices susceptible to *Aspergillus* section *Flavi* fungal infection, principally by *A. flavus* and S_{BC} species (Table 4). In Asia and South America (based on morphological identification), *A. flavus* is the only spice contaminant (pepper and curry); in Central America, *A. pipericola* was isolated from a Mexican pepper sample (Frisvad et al. 2019) whereas in Africa, besides this species, *A. minisclerotigenes* has been identified (pepper and cumin) (Table 4). Chili peppers (paprika and chili) are frequently reported to present AFs, among the aflatoxigenic species *A. flavus* is most reported, followed by *A. minisclerotigenes*, *A. parasiticus* and *A. aflatoxiformans* (Table 4). Singh and Cotty (2019) reported that the majority of aflatoxigenic fungi isolated from chili peppers in Nigeria were *A. flavus* L-morphotype (76.7%), followed by *A. aflatoxiformans* (8.3%), *A. minisclerotigenes* (8.0%), and a non-identified S_{BC} lineage (2.8%).

3.7 Tree Nuts

3.7.1 Brazil Nuts

Brazil nuts are produced in the Amazon rainforest in Brazil, Bolivia and Peru and are vulnerable to the contamination of several aflatoxigenic, *i.e.* *A. flavus*, *A. nomius*, *A. pseudonomius*, *A. parasiticus*, *A. luteovirescens*, *A. arachidicola*, *A. pseudocaelatus* and *A. pseudotamarii* (Table 4). Incidence of Brazil nut contamination by these species can reach 100% (Calderari et al. 2013, Taniwaki et al. 2017). *A. flavus* and *A. nomius* are suggested to contaminate Brazil nuts before they are collected in dry seasons, while other *Flavi* species contaminate them during rainy season, drying processes, storage and markets (Taniwaki et al. 2017).

3.7.2 Pistachios, Almonds, Hazelnuts and Nuts

The most common aflatoxigenic species identified in pistachios during pre- and post-harvest is *A. flavus*, both in Asia and USA. Only in the survey from Iran,

besides *A. flavus*, *A. parasiticus*, and an unknown AFBG strain were isolated (Table 4). *Aspergillus* section *Flavi* surveys on almonds showed higher diversity. In Europe (Portugal), the species reported are *A. flavus*, *A. minisclerotigenes*, *A. parasiticus*, *A. sergii* and *A. transmontanensis*; from these, *A. transmontanensis* was isolated only from plant processor environments (Table 4), while in USA and Algeria *A. flavus* and *A. parasiticus* have been identified in this crop (orchards and post-harvest steps). Hazelnuts in Europe (Italy) were contaminated by *A. flavus*, mostly non-aflatoxigenic strains, followed by *A. parasiticus* whereas *A. flavus* is the only section *Flavi* species reported from nuts in Europe (Table 4). Surveys showed differences between orchards and raw products in market outlets, the latter presented higher ratios of contamination, while considering the matrices pistachios are more sensitive, followed by walnuts and almonds (Bayman et al. 2002, Doster et al. 2014).

3.8 Biodiversity According to Food Production Chain Stages

In crops and pre-harvest stages, fungal infection is favored under humid and hot environmental conditions. In addition, cropping system may increase the incidence of fungal propagules in soil, if plant debris is left or incorporated into soil it becomes a source of infection and favors spore inoculation into the next planting cycle. The increase of fungal propagule number from year to year is reported for section *Flavi* species (Nesci and Etcheverry 2002, Jaime-Garcia and Cotty 2004, 2010). Insects also facilitate fungal infection by dispersing spores and wounding kernels and plants (Rychlik et al. 2014, Aiko and Mehta 2015). The main aflatoxigenic *Aspergillus* section *Flavi* species reported in the pre-harvest stages of the food production chain are *A. flavus* and *A. parasiticus*, which seem to have commensal relationships with certain crops (maize and peanuts) (Taniwaki et al. 2018). Some other species identified in these stages were *A. arachidicola*, *A. minisclerotigenes*, *A. nomius*, *A. novoparasiticus*, *A. pseudocaelatus* and *A. sergii* (Table 4).

In post-harvest a key step is the drying process, if the raw material is not properly dried, mold can easily develop. Moisture enhances *Aspergillus* section *Flavi* development, so its control is a delicate step in the food production chain to prevent AFs presence in final products. This control is easiest to be achieved in developed countries than in developing countries, especially for small producers (Marin et al. 2012, Gowda et al. 2013, Baranyi 2015b). The diversity of *Flavi* species is higher in post-harvest and first storage steps: *A. flavus*, *A. parasiticus*, *A. minisclerotigenes*, *A. aflatoxiformans*, *A. korhogoensis*, *A. austwickii*, *A. transmontanensis*, *A. sergii*, *A. luteovirescens*, *A. mottae*, *A. nomius*, *A. pipericola*, *A. pseudonomius*, *A. arachidicola*, *A. pseudocaelatus*, and *A. pseudotamarii* (Table 4).

A. flavus and *A. parasiticus* are the most frequent species found in commodities along the production chain, i.e. in pre- and post-harvest, and in a less extent *A. nomius* in Brazil nuts, *A. minisclerotigenes* in peanuts and almonds and *A. sergii* in almonds (Table 4).

4. Biodiversity of Aflatoxigenic *Aspergillus* section *Flavi* Species and Climate Change

Food security has become a very important issue across the globe and the potential effects of climate change on production and quality of food crops, including mycotoxins, is under scope, especially from a risk assessment perspective (Magan et al. 2011). Crop growth and its interaction with beneficiary and pathogenic and/or toxigenic microorganisms vary from year to year, mainly depending on local weather, making the agricultural sector particularly exposed to climate change (Moore and Lobell 2015). Climate change is a driver for distribution shifts in *Aspergillus* section *Flavi*, as it modifies the environmental conditions, resulting in new suitable niches favoring species development and also facilitating their colonization into temperate regions (Perrone et al. 2014); hence, climate change is an emerging issue worldwide for food and feed safety (Battilani et al. 2016). Projections of climate change suggest that in Africa and Oceania the suitable areas for agriculture will decrease, whereas some areas in Asia and Latin America tropical forest will become savanna (Paterson and Lima 2011). The possible change in patterns of AFs occurrence in crops due to climate change is a matter of concern that may require anticipatory actions, as these phenomena can lead to more health risk in affected areas (Battilani et al. 2016). However, it is difficult to create accurate projections as the knowledge of *Aspergillus* section *Flavi* diversity is poor (distribution, frequency and poor information of the life history and ecology). Although some species are found rarely, their total AF production is high and they produce other mycotoxins, like CPA that has additive effects. For instance, in Africa under climatic changing conditions there is a possibility that *A. minisclerotigenes*, *A. aflatoxiformans* and *A. korhogoensis* could expand their home ranges, as the scenarios of climate change suggest drier and warmer conditions, which may favor their frequency. The case of Africa is interesting, over the last three decades several studies performed in the region have increased the comprehension of the effects of AFs in human populations, unmasked species diversity of *Aspergillus* section *Flavi*, and helped to identify sensitive steps of production chain prone to mold contamination. These studies have shown a rapid diversification of species closely related to *A. flavus* (*A. minisclerotigenes*, *A. aflatoxiformans*, *A. korhogoensis* and *A. austwickii*). *A. flavus*, *A. minisclerotigenes* and *A. aflatoxiformans* contaminate staples in field, during harvesting and storage; whereas, *A. korhogoensis* and *A. austwickii* at storage steps (Cotty and Cardwell 1999, Carvajal-Campos et al. 2017, Frisvad et al. 2019). Rapid radiation in Africa may be a unique phenomenon linked with environmental conditions and evolutive pressures; or it can be just an artifact caused by the diversity underestimation in other world regions, originated by the reduced number of diversity surveys in Asia, Oceania (Pacific Islands) and Central and South America. Lack of studies is caused mainly by two factors: (1) diversity surveys are costly, time consuming and need special skills; (2) several studies are performed with the *a priori* that the only aflatoxigenic species are *A. flavus* and *A. parasiticus*. Likewise, several species identification surveys

are based only on morphology and AF production, so misidentification of species occurs; as *A. nomius* in Brazil, which was considered to be the main contaminant of Brazil nuts, and later corrected by *A. pseudonomius* (Baquião et al. 2014, Massi et al. 2014).

References

Author: To maintain consistency throughout the book, we have standardized the reference section of all the chapters by listing up to six names followed by et al.

- Abbas, A., Hussien, T. and Yli-Mattila, T. 2020. A polyphasic approach to compare the genomic profiles of aflatoxigenic and non-aflatoxigenic isolates of *Aspergillus* Section *Flavi*. *Toxins* 12(1): 56. DOI: 10.3390/toxins12010056.
- Abdallah, M.F., Audenaert, K., Lust, L., Landschoot, S., Bekaert, B., Haesaert, G. et al. 2020. Risk characterization and quantification of mycotoxins and their producing fungi in sugarcane juice: A neglected problem in a widely-consumed traditional beverage. *Food Control* 108: 106811. DOI: 10.1016/j.foodcont.2019.106811.
- Acuña, C.A., Díaz, G.J. and Espitia, M.E. 2005. Aflatoxinas en maíz: Reporte de caso en la costa atlántica colombiana. *Revista de la Facultad de Medicina Veterinaria y Zootecnia* 52(2): 156–162. DOI: 10.15446/rfmvz.
- Adjovi, Y.C.S., Bailly, S., Gnonlonfin, B.J.G., Tadríst, S., Querin, A., Sanni, A. et al. 2014. Analysis of the contrast between natural occurrence of toxigenic *Aspergilli* of the *Flavi* section and aflatoxin B1 in cassava. *Food Microbiology* 38: 151–159. DOI: 10.1016/j.fm.2013.08.005.
- Adjou, E.S., Yehouenou, B., Sossou, C.M., Soumanou, M.M. and de Souza, C.A. 2012. Occurrence of mycotoxins and associated mycoflora in peanut cake product (kulikuli) marketed in Benin. *African Journal of Biotechnology* 11(78): 14354–14360. DOI: 10.5897/AJB12.324.
- Agbetiaméh, D., Ortega-Beltrán, A., Awuah, R.T., Atehnkeng, J., Cotty, P.J. and Bandyopadhyay, R. 2018. Prevalence of aflatoxin contamination in maize and groundnut in Ghana: Population structure, distribution, and toxigenicity of the causal agents. *Plant Disease* 102(4): 764–772. DOI: 10.1094/PDIS-05-17-0749-RE.
- Aiko, V. and Mehta, A. 2015. Occurrence, detection and detoxification of mycotoxins. *Journal of Biosciences* 40(5): 943–954. DOI: 10.1007/s12038-015-9569-6.
- Ait Mimoune, N., Arroyo-Manzanares, N., Gámiz-Gracia, L., García-Campaña, A.M., Bouti, K., Sabaou, N. et al. 2018. *Aspergillus* section *Flavi* and aflatoxins in dried figs and nuts in Algeria. *Food Additives & Contaminants: Part B* 11(2): 119–125. DOI: 10.1080/19393210.2018.1438524.
- Ahmed, A., Dawar, S. and Tariq, M. 2010. Mycoflora associated with sugar cane juice in Karachi city. *Pakistan Journal of Botany* 42(4): 2955–2962.
- Al-Husnan, L., Al-Kahtani, M. and Farag, R.M. 2019. Bioinformatics analysis of aflatoxins produced by *Aspergillus* sp. in basic consumer grain (corn and rice) in Saudi Arabia. *Potravinárstvo Slovak Journal of Food Sciences* 13(1): 65–75. DOI: 10.5219/1020.
- Al-Wadai, A.S., Al-Othman, M.R., Mahmoud, M.A. and Abd El-Aziz, A.R.M. 2013. Molecular characterization of *Aspergillus flavus* and aflatoxin contamination

- of wheat grains from Saudi Arabia. *Genetics and Molecular Research* 12(3): 3335–3352. DOI: 10.4238/2013.September.3.10.
- Amaike, S. and Keller, N.P. 2011. *Aspergillus flavus*. *Annual Review of Phytopathology* 49: 107–133. DOI: 10.1146/annurev-phyto-072910-095221.
- Andersson, E. 2012. A small growth study of *Aspergillus* section *Flavi*, and potentially aflatoxigenic fungi and aflatoxin occurrence in Brazil nuts from local markets in Manaus, Brazil. M.S. Thesis, Swedish University of Agricultural Sciences, Uppsala.
- Arroyo-Manzanares, N., Di Mavungu, J.D., Uka, V., Malysheva, S.V., Cary, J.W., Ehrlich, K.C. et al. 2015. Use of UHPLC high-resolution Orbitrap mass spectrometry to investigate the genes involved in the production of secondary metabolites in *Aspergillus flavus*. *Food Additives & Contaminants: Part A* 32: 1656–1673. DOI: 10.1080/19440049.2015.1071499.
- Atehnkeng, J., Ojiambo, P.S., Donner, M., Ikotun, T., Sikora, R.A., Cotty, P.J. et al. 2008. Distribution and toxigenicity of *Aspergillus* species isolated from maize kernels from three agro-ecological zones in Nigeria. *International Journal of Food Microbiology* 122(1): 74–84. DOI: 10.1016/j.ijfoodmicro.2007.11.062.
- Azziz-Baumgartner, E., Lindblade, K., Gieseker, K., Rogers, H.S., Kieszak, S., Njapau, H. et al. 2005. Aflatoxin Investigative Group. Case-control study of an acute aflatoxicosis outbreak, Kenya, 2004. *Environmental Health Perspectives* 113: 1779–1783. DOI: 10.1289/ehp.8384.
- Bailly, S., Mahgubi, A.E., Carvajal-Campos, A., Lorber, S., Puel, O., Oswald, I.P. et al. 2018. Occurrence and identification of *Aspergillus* Section *Flavi* in the context of the emergence of aflatoxins in French maize. *Toxins* 10(12): 525. DOI: 10.3390/toxins10120525.
- Balendres, M.A.O., Karlovsky, P. and Cumagun, C.J.R. 2019. Mycotoxigenic fungi and mycotoxins in agricultural crop commodities in the Philippines: A Review. *Foods* 8(7): 249. DOI: 10.3390/foods8070249.
- Bankole, S.A. and Mabekoje, O.O. 2004. Mycoflora and occurrence of aflatoxin B1 in dried yam chips from markets in Ogun and Oyo States, Nigeria. *Mycopathologia* 157(1): 111–115. DOI: 10.1023/B:MYCO.0000012211.31618.18.
- Baquião, A.C., De Oliveira, M.M., Reis, T.A., Zorzete, P., Danielle D.A. and Correa, B. 2013. Monitoring and determination of fungi and mycotoxins in stored Brazil nuts. *Journal of Food Protection* 76(8): 1414–1420. DOI: 10.4315/0362-028X.JFP-13-005.
- Baranyi, N., Jaksić, D.D., Palágyi, A., Kiss, N., Kocsubé, S., Szekeres, A. et al. J. 2015a. Identification of *Aspergillus* species in Central Europe able to produce G-type aflatoxins. *Acta Biologica Hungarica* 66: 339–347. DOI: 10.1556/018.66.2015.3.9.
- Baranyi, N., Kocsubé, S. and Varga, J. 2015b. Aflatoxins: Climate change and biodegradation. *Current Opinion in Food Science* 5: 60–66. DOI: 10.1016/j.cofs.2015.09.002.
- Barros, G., Torres, A., Palacio, G. and Chulze, S. 2003. *Aspergillus* species from section *Flavi* isolated from soil at planting and harvest time in peanut-growing regions of Argentina. *Journal of the Science of Food and Agriculture* 83(13): 1303–1307. DOI: 10.1002/jsfa.1539.
- Battilani, P., Toscano, P., Van der Fels-Klerx, H.J., Moretti, A., Leggieri, M.C., Brera, C. et al. 2016. Aflatoxin B1 contamination in maize in Europe increases due to climate change. *Scientific Reports* 6: 24328. DOI: 10.1038/srep24328.

- Bayman, P., Baker, J.L. and Mahoney, N.E. 2002. *Aspergillus* on tree nuts: Incidence and associations. *Mycopathologia* 155(3): 161–169. DOI: 10.1023/A:1020419226146.
- Bbosa, G., Kitya, D., Lubega, A., Ogwal-Okeng, J., Anokbonggo, W.W. and Kyegombe, D.B. 2013. Chapter 12: Review of the biological and health effects of aflatoxins on body organs and systems. pp. 240–265. *In*: M. Razzaghi-Abyanedh (ed.). *Aflatoxins – Recent Advances and Future Prospects*. IntechOpen, Rijeka.
- Bertuzzi, T., Romani, M., Rastelli, S., Mulazzi, A. and Pietri, A. 2017. Sterigmatocystin occurrence in paddy and processed rice produced in Italy in the years 2014–2015 and distribution in milled rice fractions. *Toxins* 9(3): 86. DOI: 10.3390/toxins9030086.
- Bhat, R., Rai, R.V. and Karim, A.A. 2010. Mycotoxins in food and feed: Present status and future concerns. *Comprehensive Reviews in Food Science and Food Safety* 9(1): 57–81. DOI: 10.1111/j.1541-4337.2009.00094.x.
- Boli, Z.A., Zoue, L.T., Alloue-Boraud, W.M., Kakou, C.A. and Koffi-Nevry, R. 2014. Proximate composition and mycological characterization of peanut butter sold in retail markets of Abidjan (Côte d’Ivoire). *Journal of Applied Biosciences* 72(1): 5822–5829. DOI: 10.4314/jab.v72i1.99667.
- Calderari, T.O., Iamanaka, B.T., Frisvad, J.C., Pitt, J.I., Sartori, D., Pereira, J.L. et al. 2013. The biodiversity of *Aspergillus* section *Flavi* in Brazil nuts: From rainforest to consumer. *International Journal of Food Microbiology* 160(3): 267–272. DOI: 10.1016/j.ijfoodmicro.2012.10.018.
- Camiletti, B.X., Torrico, A.K., Maurino, M.F., Cristos, D., Magnoli, C., Lucini, E.I. et al. 2017. Fungal screening and aflatoxin production by *Aspergillus* section *Flavi* isolated from pre-harvest maize ears grown in two Argentine regions. *Crop Protection* 92: 41–48. DOI: 10.1016/j.cropro.2016.10.012.
- Carvajal-Campos, A., Manizan, A.L., Tadrst, S., Akaki, D.K., Koffi-Nevry, R., Moore, G.G. et al. 2017. *Aspergillus korhogoensis*: A novel aflatoxin producing species from the Côte d’Ivoire. *Toxins* 9(11): 353. DOI: 10.3390/toxins9110353.
- Cary, J.W., Uka, V., Han, Z., Buyst, D., Harris-Coward, P.Y., Ehrlich, K.C. et al. 2015a. A secondary metabolic gene cluster containing a hybrid PKS–NRPS is necessary for synthesis of the 2-pyridones, leporins. *Fungal Genetics and Biology* 81: 88–97. DOI: 10.1016/j.fgb.2015.05.010.
- Cary, J.W., Han, Z., Yin, Y., Lohmar, J.M., Shantappa, S., Harris-Coward, P.Y. et al. 2015b. Transcriptome analysis of *Aspergillus flavus* reveals veA-dependent regulation of secondary metabolite gene clusters, including the novel aflavarin cluster. *Eukaryotic Cell* 14: 983–997. DOI: 10.1128/EC.00092-15.
- Chang, P.K., Matsushima, K., Takahashi, T., Yu, J., Abe, K., Bhatnagar, D. et al. 2007. Understanding nonaflatoxigenicity of *Aspergillus sojae*: A windfall of aflatoxin biosynthesis research. *Applied Microbiology and Biotechnology* 76(5): 977–984. DOI: 10.1007/s00253-007-1116-4.
- Chang, P.K. and Ehrlich, K.C. 2010. What does genetic diversity of *Aspergillus flavus* tell us about *Aspergillus oryzae*? *International Journal of Food Microbiology* 138(3): 189–199. DOI: 10.1016/j.ijfoodmicro.2010.01.033.
- Chilaka, C.A., De Kock, S., Phoku, J.Z., Mulunda, M., Egbuta, M.A. and Dutton, M.F. 2012. Fungal and mycotoxin contamination of South African commercial maize. *Journal of Food, Agriculture & Environment* 10(2): 296–303.

- Copetti, M.V., Iamanaka, B.T., Pereira, J.L., Fungaro, M.H. and Taniwaki, M.H. 2011. Aflatoxigenic fungi and aflatoxin in cocoa. *International Journal of Food Microbiology* 148(2): 141–144. DOI: 10.1016/j.ijfoodmicro.2011.05.020
- Costa, L.L.F. and Scussel, V.M. 2002. Toxigenic fungi in beans (*Phaseolus vulgaris* L.) classes black and color cultivated in the state of Santa Catarina, Brazil. *Brazilian Journal of Microbiology* 33(2): 138–144. DOI: 10.1590/S1517-83822002000200008.
- Cotty, P.J. 1997. Aflatoxin-producing potential of communities of *Aspergillus* section *Flavi* from cotton producing areas in the United States. *Mycological Research* 101(6): 698–704. DOI: 10.1017/S0953756296003139.
- Cotty, P.J. and Cardwell, K.F. 1999. Divergence of West African and North American Communities of *Aspergillus* Section *Flavi*. *Applied and Environmental Microbiology* 65: 2264–2265.
- Covarelli, L., Beccari, G. and Salvi, S. 2011. Infection by mycotoxigenic fungal species and mycotoxin contamination of maize grain in Umbria, central Italy. *Food and Chemical Toxicology* 49(9): 2365–2369. DOI: 10.1016/j.fct.2011.06.047.
- Dahab, N.F.A., Abdel-Hadi, A.M., Abdul-Raouf, U.M., El-Shanawany, R.A. and Hassane, A.M.A. 2016. Qualitative detection of aflatoxins and aflatoxigenic fungi in wheat flour from different regions of Egypt. *IOSR Journal of Environmental Science, Toxicology and Food Technology* 10(7): 20–26.
- Diedhiou, P.M., Bandyopadhyay, R., Atehnkeng, J. and Ojiambo, P.S. 2011. *Aspergillus* colonization and aflatoxin contamination of maize and sesame kernels in two agro-ecological zones in Senegal. *Journal Phytopathology* 159: 268–275. DOI: 10.1111/j.1439-0434.2010.01761.x.
- Divakara, S.T., Aiyaz, M., Puttaswamy, H., Chandra, N.S. and Ramachandrapa, N.S. 2014. *Aspergillus flavus* infection and contamination in sorghum seeds and their biological management. *Archives of Phytopathology and Plant Protection* 47(17): 2141–2156. DOI: 10.1080/03235408.2013.869892
- Dobolyi, C.S., Sebők, F., Varga, J., Kocsubé, S., Szigeti, G., Baranyi, N. et al. 2013. Occurrence of aflatoxin producing *Aspergillus flavus* isolates in maize kernel in Hungary. *Acta Alimentaria* 42(3): 451–459. DOI: 10.1556/AAlim.42.2013.3.18.
- Doolotkeldieva, T.D. 2010. Microbiological control of flour-manufacture: Dissemination of mycotoxins producing fungi in cereal products. *Microbiology Insights* 3: 1–15. DOI: 10.4137/MBI.S3822.
- Doster, M.A., Cotty, P.J. and Michailides, T.J. 2009. Description of a distinctive aflatoxin-producing strain of *Aspergillus nomius* that produces submerged sclerotia. *Mycopathologia* 168: 193–201. DOI: 10.1007/s11046-009-9214-8.
- Doster, M.A., Cotty, P.J. and Michailides, T.J. 2014. Evaluation of the atoxigenic *Aspergillus flavus* strain AF36 in pistachio orchards. *Plant Disease* 98(7): 948–956. DOI: 10.1094/PDIS-10-13-1053-RE
- Dovičičová, M., Tančinová, D., Labuda, R. and Sulyok, M. 2012. *Aspergillus parasiticus* from wheat grain of Slovak origin and its toxigenic potency. *Czech Journal of Food Sciences* 30: 483–487.
- EFSA Panel on Contaminants in the Food Chain (CONTAM). 2013. Scientific opinion on the risk for public and animal health related to the presence of sterigmatocystin in food and feed. *EFSA Journal* 11: 3254. DOI: 10.2903/j.efsa.2013.3254.

- Egmond, H.P. and Jonker, M.A. 2004. Current situation on regulations for mycotoxins. *JSM Mycotoxins* 2003(Suppl 3): 1–15.
- Ehrlich, K.C., Chang, P.K., Yu, J. and Cotty, P.J. 2004. Aflatoxin biosynthesis cluster gene *cypA* is required for G aflatoxin formation. *Applied and Environmental Microbiology* 70: 6518–6524. DOI: 10.1128/AEM.70.11.6518-6524.2004.
- Ehrlich, K.C. and Yu, J. 2010. Aflatoxin-like gene clusters and how they evolved. pp. 65–75. *In*: M. Rai and A. Varma (eds.). *Mycotoxins in the Food, Feed and Bioweapons*. Berlin, Springer Verlag.
- Ekwomadu, T.I., Gopane, R.E. and Mwanza, M. 2018. Occurrence of filamentous fungi in maize destined for human consumption in South Africa. *Food Science & Nutrition* 6(4): 884–890. DOI: 10.1002/fsn3.561.
- El Mahgubi, A.E., Puel, O., Bailly, S., Tadriss, S., Querin, A., Ouadia, A. et al. 2013. Distribution and toxigenicity of *Aspergillus* section *Flavi* in spices marketed in Morocco. *Food Control* 32: 143–148. DOI: 10.1016/j.foodcont.2012.11.013.
- El-Shanshoury, A.R., El-Sabbagh, S.M., Emara, H.A. and Saba, H.E. 2014. Occurrence of moulds, toxicogenic capability of *Aspergillus flavus* and levels of aflatoxins in maize, wheat, rice and peanut from markets in central delta provinces, Egypt. *International Journal of Current Microbiology and Applied Sciences* 3(3): 852–865.
- Embaby, E.M. and Abdel-Galel, M.M. 2014. Detection of fungi and aflatoxins contaminated peanut samples (*Arachis hypogaea* L.). *Journal of Agricultural Technology* 10(2): 423–437.
- Ennouari, A., Sanchis, V., Rahouti, M. and Zinedine, A. 2018. Isolation and molecular identification of mycotoxin producing fungi in durum wheat from Morocco. *Journal of Materials and Environmental Sciences* 9: 1470–1479. DOI: 10.26872/jmes.2018.9.5.161.
- Etcheverry, M., Nesci, A., Barros, G., Torres, A. and Chulze, S. 1999. Occurrence of *Aspergillus* section *Flavi* and aflatoxin B1 in corn genotypes and corn meal in Argentina. *Mycopathologia* 147(1): 37–41. DOI: 10.1023/A: 1007040123181.
- Ezekiel, C.N., Sulyok, M., Frisvad, J.C., Somorin, Y.M., Warth, B., Houbraeken, J. et al. 2013a. Fungal and mycotoxin assessment of dried edible mushroom in Nigeria. *International Journal of Food Microbiology* 162(3): 231–236. DOI: 10.1016/j.ijfoodmicro.2013.01.025.
- Ezekiel, C.N., Sulyok, M., Babalola, D.A., Warth, B., Ezekiel, V.C. and Krska, R. 2013b. Incidence and consumer awareness of toxigenic *Aspergillus* section *Flavi* and aflatoxin B1 in peanut cake from Nigeria. *Food Control* 30(2): 596–601. DOI: 10.1016/j.foodcont.2012.07.048.
- Ezekiel, C.N., Udom, I.E., Frisvad, J.C., Adetunji, M.C., Houbraeken, J., Fapohunda, S.O. et al. 2014. Assessment of aflatoxigenic *Aspergillus* and other fungi in millet and sesame from Plateau State, Nigeria. *Mycology* 5(1): 16–22. DOI: 10.1080/21501203.2014.889769.
- Ezekiel, C.N., Ortega-Beltran, A., Oyediji, E.O., Atehnkeng, J., Kössler, P., Tairu, F. et al. 2019. Aflatoxin in chili peppers in Nigeria: Extent of contamination and control using atoxigenic *Aspergillus flavus* genotypes as biocontrol agents. *Toxins* 11(7): 429. DOI: 10.3390/toxins11070429.
- Fakhrunnisa, M.H. and Ghaffar, A. 2006. Seed-borne mycoflora of wheat, sorghum and barley. *Pakistan Journal of Botany* 38(1): 185–192.
- FAO. 2003. *Worldwide Regulations for Mycotoxins in Food and Feed in 2003*. FAO Nutr. Pap. Rome, pp. 1–165.

- Freire, F.C., Kozakiewicz, Z. and Paterson, R.R. 2000. Mycoflora and mycotoxins in Brazilian black pepper, white pepper and Brazil nuts. *Mycopathologia* 149(1): 13–19. DOI: 10.1023/a: 1007241827937.
- Freitas-Silva, O., Teixeira, A., da Cunha, F.Q., de Oliveira Godoy, R.L. and Venâncio, A. 2011. Predominant mycobiota and aflatoxin content in Brazil nuts. *Journal für Verbraucherschutz und Lebensmittelsicherheit* 6(4): 465–472. DOI: 10.1007/s00003-011-0703-6.
- Frisvad, J.C., Skouboe, P. and Samson, R.A. 2005. Taxonomic comparison of three different groups of aflatoxin producers and a new efficient producer of aflatoxin B1, sterigmatocystin and 3-O-methylsterigmatocystin, *Aspergillus rambellii* sp. nov. *Systematic and Applied Microbiology* 28(5): 442–453. DOI: 10.1016/j.syapm.2005.02.012.
- Frisvad, J.C., Hubka, V., Ezekiel, C.N., Hong, S.B., Nováková, A., Chen, A.J. et al. 2019. Taxonomy of *Aspergillus* section *Flavi* and their production of aflatoxins, ochratoxins and other mycotoxins. *Studies in Mycology* 93: 1–63. DOI: 10.1016/j.simyco.2018.06.001.
- Gachara, G.W., Nyamache, A.K., Harvey, J., Gnonlonfin, G.J.B. and Wainaina, J. 2018. Genetic diversity of *Aspergillus flavus* and occurrence of aflatoxin contamination in stored maize across three agro-ecological zones in Kenya. *Agriculture & Food Security* 7(1): 52. DOI: 10.1186/s40066-018-0202-4.
- Gallo, A., Stea, G., Battilani, P., Logrieco, A.F. and Perrone, G. 2012. Molecular characterization of an *Aspergillus flavus* population isolated from maize during the first outbreak of aflatoxin contamination in Italy. *Phytopathologia Mediterranea* 51(1): 198–206.
- Gao, J., Liu, Z. and Yu, J. 2007. Identification of *Aspergillus* section *Flavi* in maize in northeastern China. *Mycopathologia* 164(2): 91–95. DOI: 10.1007/s11046-007-9029-4.
- Garber, N.P. and Cotty, P.J. 2014. *Aspergillus parasiticus* communities associated with sugarcane in the Rio Grande Valley of Texas: Implications of global transport and host association within *Aspergillus* Section *Flavi*. *Phytopathology* 104: 462–471. DOI: 10.1094/PHYTO-04-13-0108-R
- Gauthier, T., Duarte-Hospital, C., Vignard, J., Boutet-Robinet, E., Sulyok, M., Snini, S.P. et al. 2020. Versicolorin A, a precursor in aflatoxins biosynthesis, is a food contaminant toxic for human intestinal cells. *Environment International* 137: 105568. DOI: 10.1016/j.envint.2020.105568.
- Georgianna, D.R., Fedorova, N.D., Burroughs, J.L., Dolezal, A.L., Bok, J.W., Horowitz-Brown, S. et al. 2010. Beyond aflatoxin: Four distinct expression patterns and functional roles associated with *Aspergillus flavus* secondary metabolism gene clusters. *Molecular Plant Pathology* 11(2): 213–226. DOI: 10.1111/j.1364-3703.2009.00594.x.
- Giorni, P., Magan, N., Pietri, A., Bertuzzi, T. and Battilani, P. 2007. Studies on *Aspergillus* section *Flavi* isolated from maize in northern Italy. *International Journal of Food Microbiology* 113(3): 330–338. DOI: 10.1016/j.ijfoodmicro.2006.09.007
- Gloer, J.B., TePaske, M.R., Sima, J.S., Wicklow, D.T. and Dowd, P.F. 1988. Antiinsectan aflavinine derivative from sclerotia of *Aspergillus flavus*. *Journal of Organic Chemistry* 53: 5457–5460. DOI: 10.1021/jo00258a011.
- Gnonlonfin, G.J.B., Adjovi, C.S.Y., Katerere, D.R., Shephard, G.S., Sanni, A. and Brimer, L. 2012. Mycoflora and absence of aflatoxin contamination of

AQ : 2008 missing
see p 23

AQ : Missing in author
citations

AQ : 2008 ?See p. 21

- commercialized cassava chips in Benin, West Africa. *Food Control* 23: 333–337. DOI: 10.1016/j.foodcont.2011.07.026.
- Gonçalves, J.S., Ferracin, L.M., Vieira, M.L.C., Iamanaka, B.T., Taniwaki, M.H. and Fungaro, M.H.P. 2012. Molecular analysis of *Aspergillus* section *Flavi* isolated from Brazil nuts. *Journal of Microbiology and Biotechnology* 28(4): 1817–1825. DOI: 10.1007/s11274-011-0956-3.
- Gowda, N.K.S., Swamy, H.V.L.N. and Mahajan, P. 2013. Recent advances for control, counteraction and amelioration of potential aflatoxins in animal feeds. pp. 129–140. *In: M. Razzaghi-Abyanedh (ed.). Aflatoxins – Recent Advances and Future Prospects. IntechOpen, Rijeka.*
- González, H.H.L., Moltó, G.A., Pacin, A., Resnik, S.L., Zelaya, M.J., Masana, M. et al. 2008. Trichotheceae and mycoflora in wheat harvested in nine locations in Buenos Aires province, Argentina. *Mycopathologia* 165(2): 105–114. DOI: 10.1007/s11046-007-9084-x.
- Grenier, B. and Oswald, I. 2011. Mycotoxin co-contamination of food and feed: Meta-analysis of publications describing toxicological interactions. *World Mycotoxin Journal* 4(3): 285–313. DOI: 10.3920/WMJ2011.1281.
- Guchi, E. 2015. Effect of storage time on occurrence of *Aspergillus* species in groundnut (*Arachis hypogaea* L.) in Eastern Ethiopia. *Journal of Applied & Environmental Microbiology* 3(1): 1–5. DOI: 10.12691/jaem-3-1-1
- Hell, K., Gnonlonfin, B.G.J., Kodjogbe, G., Lamboni, Y. and Abdourhamane, I.K. 2009. Mycoflora and occurrence of aflatoxin in dried vegetables in Benin, Mali and Togo, West Africa. *International Journal of Food Microbiology* 135(2): 99–104. DOI: 10.1016/j.ijfoodmicro.2009.07.039.
- Hepkerkan, D., Aran, N. and Ayfer, M. 1994. Mycoflora and aflatoxin contamination in shelled pistachio nuts. *Journal of the Science of Food and Agriculture* 66 (3): 273–278. DOI: 10.1002/jsfa.2740660302.
- Horn, B.W. 2007 Biodiversity of *Aspergillus* section *Flavi* in the United States: A review. *Food Additives and Contaminants* 24(10): 1088–1101. DOI: 10.1080/02652030701510012.
- Hussain, M., Ghazanfar, M.U., Hamid, M.I. and Raza, M. 2013. Seed borne mycoflora of some commercial wheat (*Triticum aestivum* L.) cultivars in Punjab, Pakistan. *International Journal of Phytopathology* 2(2): 97–101. DOI: 10.33687/phytopath.002.02.0198.
- Iamanaka, B.T., de Souza Lopes, A., Martins, L.M., Frisvad, J.C., Medina, A., Magan, N. et al. 2019. *Aspergillus* section *Flavi* diversity and the role of *A. novoparasiticus* in aflatoxin contamination in the sugarcane production chain. *International Journal of Food Microbiology* 293: 17–23. DOI: 10.1016/j.ijfoodmicro.2018.12.024
- IARC. 2012. Monographs on the evaluation of carcinogenic risks to humans: Chemical agents and related occupations. A review of human carcinogens. Lyon, France. 2012. International Agency for Research on Cancer 100F: 224–248.
- IARC. 2015. Mycotoxin control in low- and middle-income countries. *In: Wild, C.P., Miller, J.D. and Groopman, J.D. (Eds.). International Agency for Research on Cancer, Working Group Report Volume 9*, pp. 56.
- Ibrahim, F., Jalal, H., Khan, A.B., Asghar, M.A., Iqbal, J., Ahmed, A. et al. 2016. Prevalence of Aflatoxigenic *Aspergillus* in Food and Feed Samples from

- Karachi, Pakistan. Journal of Infection and Molecular Biology 4(1): 1–8. DOI: 10.14737/journal.jimb/2016/4.1.1.8.
- Ito, Y., Peterson, S.W., Wicklow, D.T. and Goto, T. 2001. *Aspergillus pseudotamarii*, a new aflatoxin producing species in *Aspergillus* section *Flavi*. Mycological Research 105: 233–239. DOI: 10.1017/S0953756200003385.
- Jaime-Garcia, R. and Cotty, P.J. 2004. *Aspergillus flavus* in soils and corncoobs in south Texas: Implications for management of aflatoxins in corn-cotton rotations. Plant Disease 88(12): 1366–1371. DOI: 10.1094/PDIS.2004.88.12.1366
- Jaime-Garcia, R. and Cotty, P.J. 2010. Crop rotation and soil temperature influence the community structure of *Aspergillus flavus* in soil. Soil Biology and Biochemistry 42(10): 1842–1847. DOI: 10.1016/j.soilbio.2010.06.025.
- Jakić-Dimić, D., Nešić, K. and Petrović, M. 2009. Contamination of cereals with aflatoxins, metabolites of fungi *Aspergillus flavus*. Biotechnol. Biotechnology in Animal Husbandry 25(5-6-2): 1203–1208.
- Jakšić, D., Puel, O., Canlet, C., Kopjar, N., Kosalec, I. and Klarić M.Š. 2012. Cytotoxicity and genotoxicity of versicolorins and 5-methoxysterigmatocystin in A549 cells. Archives of Toxicology 86: 1583–1591. DOI: 10.1007/s00204-012-0871-x.
- Javanmard, M. 2010. Occurrence of mould counts and *Aspergillus* species in Iranian dried figs at different stages of production and processing. Journal of Agricultural Science and Technology 12: 331–338.
- Jayaraman, P. and Kalyanasundaram, I. 1990. Natural occurrence of toxigenic fungi and mycotoxins in rice bran. Mycopathologia 110: 81–85. DOI: 10.1007/BF00446995.
- Joshaghani, H., Namjoo, M., Rostami, M., Kohsar, F. and Niknejad, F. 2013. Mycoflora of fungal contamination in wheat storage (Silos) in Golestan Province, north of Iran. Jundishapur Journal of Microbiology 6(4): e6334. DOI: 10.5812/jjm.6334.
- Kabirian, H.R., Afshari, H., Moghadam, M.M. and Hokmabadi, H. 2011. Evaluation of pistachio contamination to *Aspergillus flavus* in Semnan Province. Journal of Nuts 2(3): 01–06. DOI: 10.22034/jon.2011.515741.
- Kachuei, R., Hossein, Y.M., Sasan, R., Abdolamir, A., Naser, S., Farideh, Z. et al. 2009. Investigation of stored wheat mycoflora, reporting the *Fusarium* cf. *langsethiae* in three provinces of Iran during 2007. Annals of Microbiology 59(2): 383. DOI: 10.1007/BF03178344.
- Kachapulula, P.W., Akello, J., Bandyopadhyay, R. and Cotty, P.J. 2017. *Aspergillus* section *Flavi* community structure in Zambia influences aflatoxin contamination of maize and groundnut. International Journal of Food Microbiology 261: 49–56. DOI: 10.1016/j.ijfoodmicro.2017.08.014
- Kamika, I., Mngqawa, P., Rheeder, J.P., Teffo, S.L. and Katerere, D.R. 2014. Mycological and aflatoxin contamination of peanuts sold at markets in Kinshasa, Democratic Republic of Congo, and Pretoria, South Africa. Food Additives & Contaminants: Part B 7(2): 120–126. DOI: 10.1080/19393210.2013.858187.
- Katsurayama, A.M., Martins, L.M., Iamanaka, B.T., Fungaro, M.H.P., Silva, J.J., Frisvad, J.C. et al. 2018. Occurrence of *Aspergillus* section *Flavi* and aflatoxins in Brazilian rice: From field to market. International Journal of Food Microbiology 266: 213–221. DOI: 10.1016/j.ijfoodmicro.2017.12.008.
- Kjærboelling, I., Vesth, T., Frisvad, J.C., Nybo, J.L., Theobald, S., Kildgaard, S. et al.

2020. A comparative genomics study of 23 *Aspergillus* species from Section *Flavi*. *Nature Communications* 11: 1106. DOI: 10.1038/s41467-019-14051-y.
- Klich, M.A., Lee, L.S. and Huizar, H.E. 1986. The occurrence of *Aspergillus flavus* in vegetative tissue of cotton plants and its relation to seed infection. *Mycopathologia* 95(3): 171–174. DOI: 10.1007/BF00437123.
- Kornher, L. 2018. Maize markets in Eastern and Southern Africa (ESA) in the context of climate change. *The State of Agricultural Commodity Markets. (SOCO) 2018*. Rome. FAO: p. 58.
- Kos, J., Janić Hajnal, E., Šarić, B., Jovanov, P., Mandić, A., Đuragić, O. et al. 2018. Aflatoxins in maize harvested in the Republic of Serbia over the period 2012–2016. *Food Additives & Contaminants: Part B* 11(4): 246–255. DOI: 10.1080/19393210.2018.1499675.
- Kowalska, A., Walkiewicz, K., Koziel, P. and Muc-Wierzgoń, M. 2017. Aflatoxins: Characteristics and impact on human health. *Postepy higieny i medycyny doswiadczalnej* 71: 315–327. DOI: 10.5604/01.3001.0010.3816.
- Kumar, P., Mahato, D.K., Kamle, M., Mohanta, T.K. and Kang, S.G. 2017. Aflatoxins: A global concern for food safety, human health and their management. *Frontiers in Microbiology* 7: 2170. DOI: 10.3389/fmicb.2016.02170.
- Kumeda, Y., Asao, T., Takahashi, H. and Ichinoe, M. 2003. High prevalence of B and G aflatoxin-producing fungi in sugarcane field soil in Japan: Heteroduplex panel analysis identifies a new genotype within *Aspergillus* section *Flavi* and *Aspergillus nomius*. *FEMS Microbiology Ecology* 45: 229–238 DOI: 10.1016/S0168-6496(03)00154-5.
- Kurtzman, C.P., Horn, B.W. and Hesseltine, C.W. 1987. *Aspergillus nomius*, a new aflatoxin-producing species related to *Aspergillus flavus* and *Aspergillus tamaritii*. *Antonie van Leeuwenhoek* 53(3): 147–158. DOI: 10.1007/BF00393843.
- Lewis, L., Onsongo, M., Njapau, H., Schurz-Rogers, H., Lubber, G., Kieszak, S. et al. and the Kenya Aflatoxicosis Investigation Group. 2005. Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in eastern and central Kenya. *Environmental Health Perspectives* 113: 1763–1767. DOI: 10.1289/ehp.7998.
- Magan, N., Medina, A. and Aldred, D. 2011. Possible climate-change effects on mycotoxin contamination of food crops pre- and post-harvest: Mycotoxins and climate change. *Plant Pathology* 60(1): 150–163. DOI: 10.1111/j.1365-3059.2010.02412.x.
- Makhlouf, J., Carvajal-Campos, A., Querin, A., Tadriss, S., Puel, O., Lorber S. et al. 2019. Morphologic, molecular and metabolic characterization of *Aspergillus* section *Flavi* in spices marketed in Lebanon. *Scientific Reports* 9(1): 5263. DOI: 10.1038/s41598-019-41704-1.
- Mamo, F.T., Shang, B., Selvaraj, J.N., Wang, Y. and Liu, Y. 2018. Isolation and characterization of *Aspergillus flavus* strains in China. *Journal of Microbiology* 56(2): 119–127. DOI: 10.1007/s12275-018-7144-1.
- Manizan, A.L. 2019. Evaluation de la contamination par les mycotoxines des céréales et oléagineux les plus consommées en Côte d’Ivoire: cas du circuit post-récolte de l’arachide (*Arachis hypogea* L.). Ph.D. Thesis, Université Nangui Abrogoua, Abidjan, Côte d’Ivoire.
- Marin, S., Ramos, A.J. and Sanchis, V. 2012. Modelling *Aspergillus flavus* growth and aflatoxins production in pistachio nuts. *Food Microbiology* 32: 378–388. DOI: 10.1016/j.fm.2012.07.018

AQ : or 2004 as cited within text above?

- Martins, L.M., Sant'Ana, A.S., Fungaro, M.H.P., Silva, J.J., do Nascimento, M.D.S., Frisvad, J.C. et al. 2017. The biodiversity of *Aspergillus* section *Flavi* and aflatoxins in the Brazilian peanut production chain. *Food Research International* 94: 101–107. DOI: 10.1016/j.foodres.2017.02.006.
- Massi, F.P., Vieira, M.L.C., Sartori, D., Penha, R.E.S., de Freitas Munhoz, C., Ferreira, J.M. et al. 2014. Brazil nuts are subject to infection with B and G aflatoxin-producing fungus, *Aspergillus pseudonomius*. *International Journal of Food Microbiology* 186: 14–21. DOI: 10.1016/j.ijfoodmicro.2014.06.006.
- Mateo, E.M., Gil-Serna, J., Patiño, B. and Jiménez, M. 2011. Aflatoxins and ochratoxin A in stored barley grain in Spain and impact of PCR-based strategies to assess the occurrence of aflatoxigenic and ochratoxigenic *Aspergillus* spp. *International Journal of Food Microbiology* 149(2): 118–126. DOI: 10.1016/j.ijfoodmicro.2011.06.006.
- Mauro, A., Battilani, P., Callicott, K.A., Giorni, P., Pietri, A. and Cotty, P.J. 2013. Structure of an *Aspergillus flavus* population from maize kernels in northern Italy. *International Journal of Food Microbiology* 162(1): 1–7. DOI: 10.1016/j.ijfoodmicro.2012.12.021.
- Mazzani, C., Luzón, O., Chavarri, M., Fernández, M. and Hernández, N. 2008. *Fusarium verticillioides* y fumonisinas en maíz cosechado en pequeñas explotaciones y conucos de algunos estados de Venezuela. *Fitopatología Venezolana* 21(1): 18–22.
- Mazen, M.B., El-Kady, I.A. and Saber, S.M. 1990. Survey of the mycoflora and mycotoxins of cotton seeds and cotton seed products in Egypt. *Mycopathologia* 110(3): 133–138. DOI: 10.1007/BF00437536.
- Medina, A., Rodríguez, A., Sultan, Y. and Magan, N. 2015. Climate change factors and *Aspergillus flavus*: Effects on gene expression, growth and aflatoxin production. *World Mycotoxin Journal* 8 (2): 171–179. DOI: 10.3920/WMJ2014.1726.
- Mohamad, R., Mohamed, M.S., Suhaili, N., Salleh, M.M. and Ariff, A.B. 2010. Kojic acid: Applications and development of fermentation process for production. *Biotechnology and Molecular Biology Reviews* 5(2): 24–37.
- Mohamed, A. and Chala, A. 2014. Incidence of *Aspergillus* contamination of groundnut (*Arachis hypogaea* L.) in Eastern Ethiopia. *African Journal of Microbiology Research* 8(8): 759–765. DOI: 10.5897/AJMR12.2078.
- Moore, F.C. and Lobell, D.B. 2015. The fingerprint of climate trends on European crop yields. *Proceedings of the National Academy of Sciences* 112(9): 2670–2675. DOI: 10.1073/pnas.1409606112.
- Moore, G.G., Mack, B.M., Beltz, S.B. and Gilbert, M.K. 2016. Draft genome sequence of an aflatoxigenic *Aspergillus* species, *A. bombycis*. *Genome Biology and Evolution* 8: 3297–3300. DOI: 10.1093/gbe/evw238. DOI: 10.1093/gbe/evw238.
- Moretti, A., Pascale, M. and Logrieco, A.F. 2019. Mycotoxin risks under a climate change scenario in Europe. *Trends in Food Science & Technology* 84: 38–40. DOI: 10.1016/j.tifs.2018.03.008.
- Mphande, F.A., Siame, B.A. and Taylor, J.E. 2004. Fungi, aflatoxins, and cyclopiazonic acid associated with peanut retailing in Botswana. *Journal of Food Protection* 67(1): 96–102. DOI: 10.4315/0362-028X-67.1.96.
- Nesci, A. and Etcheverry, M. (2002). *Aspergillus* section *Flavi* populations from

- field maize in Argentina. *Letters in Applied Microbiology* 34(5): 343–348. DOI: 10.1046/j.1472-765X.2002.01094.x.
- Njobeh, P.B., Dutton, M.F., Koch, S.H., Chuturgoon, A., Stoev, S. and Seifert, K. 2009. Contamination with storage fungi of human food from Cameroon. *International Journal of Food Microbiology* 135(3): 193–198. DOI: 10.1016/j.ijfoodmicro.2009.08.001.
- Norlia, M., Jinap, S., Nor-Khaizura, M.A.R., Son, R. and Chin, C.K. 2018. Polyphasic approach to the identification and characterization of aflatoxigenic strains of *Aspergillus* section *Flavi* isolated from peanuts and peanut-based products marketed in Malaysia. *International Journal of Food Microbiology* 282 (3): 9–15. DOI: 10.1016/j.ijfoodmicro.2018.05.030
- Okoth, S., De Boevre, M., Vidal, A., Di Mavungu, J.D. Landschoot, S., Kyallo, M. et al. 2018. Genetic and toxigenic variability within *Aspergillus flavus* population isolated from maize in two diverse environments in Kenya. *Frontiers in Microbiology* 9: 57. DOI: 10.3389/fmicb.2018.00057.
- Onana, B., Essono, G., Bekolo, N., Ambang, Z. and Ayodele, M. 2013. Mycoflora associated with processed and stored cassava chips in rural areas of southern Cameroon. *African Journal of Microbiology Research* 7(43): 5036–5045. DOI: 10.5897/AJMR2013.6081.
- Ortega-Beltran, A., Jaime, R. and Cotty, P.J. 2015. Aflatoxin-producing fungi in maize field soils from sea level to over 2000 masl: A three-year study in Sonora, Mexico. *Fungal Biology* 119(4): 191–200. DOI: 10.1016/j.funbio.2014.12.006.
- Osamwonyi, U.O. and Wakil, S.M. 2012. Isolation of fungal species from fermentating pearl millet gruel and determination of their antagonistic activities against indicator bacterial species. *Nigerian Food Journal* 30(1): 35–42. DOI: 10.1016/S0189-7241(15)30011-4.
- Pacheco, A.M., Lucas, A., Parente, R. and Pacheco, N. 2010. Association between aflatoxin and aflatoxigenic fungi in Brazil nut (*Bertholletia excelsa* HBK). *Food Science and Technology* 30(2): 330–334. DOI: 10.1590/S0101-20612010000200007
- Pacin, A.M., Gonzalez, H.H.L., Etcheverry, M., Resnik, S.L., Vivas, L. and Espin, S. 2003. Fungi associated with food and feed commodities from Ecuador. *Mycopathologia* 156(2): 87–92. DOI: 10.1023/A: 1022941304447.
- Pankaj, S.K., Shi, H. and Keener, K.M. 2018. A review of novel physical and chemical decontamination technologies for aflatoxin in food. *Trends in Food Science & Technology* 71: 73–83. DOI: 10.1016/j.tifs.2017.11.007.
- Park, J.W., Choi, S.Y., Hwang, H.J. and Kim, Y.B. 2005. Fungal mycoflora and mycotoxins in Korean polished rice destined for humans. *International Journal of Food Microbiology* 103(3): 305–314. DOI: 10.1016/j.ijfoodmicro.2005.02.001.
- Paterson, R.R.M. and Lima, N. 2011. Further mycotoxin effects from climate change. *Food Res. Int.* 44: 2555–2566. DOI: 10.1016/j.foodres.2011.05.038.
- Perrone, G., Gallo, A. and Logrieco, A.F. 2014. Biodiversity of *Aspergillus* section *Flavi* in Europe in relation to the management of aflatoxin risk. *Frontiers in Microbiology* 5: 377. DOI: 10.3389/fmicb.2014.00377.
- Peterson, S.W., Ito, Y., Horn, B.W. and Goto, T. 2001. *Aspergillus bombycis*, a new aflatoxigenic species and genetic variation in its sibling species, *A. nomius*. *Mycologia* 93: 689–703. DOI: 10.2307/3761823.

- Pildain, M.B., Frisvad, J.C., Vaamonde, G., Cabral, D., Varga, J. and Samson, R.A. 2008. Two novel aflatoxin-producing *Aspergillus* species from Argentinean peanuts. *International Journal of Systematic and Evolutionary Microbiology* 58: 725–735. DOI: 10.1099/ij.s.0.65123-0.
- Pitt, J.I. and Hocking, A.D. 2006. Mycotoxins in Australia: Biocontrol of aflatoxin in peanuts. *Mycopathologia* 162(3): 233–243. DOI: 10.1007/s11046-006-0059-0.
- Pitt, J.I. and Hocking, A.D. 2009. *Fungi and Food Spoilage*, 3rd ed., Springer US: New York, USA. ISBN 978-0-387-92206-5.
- Pitt, J.I., Taniwaki, M.H. and Cole, M.B. 2012. Mycotoxin production in major crops as influenced by growing, harvesting, storage and processing, with emphasis on the achievement of Food Safety Objectives. *Food Control* 32(1): 205–215. DOI: 10.1016/j.foodcont.2012.11.023.
- Prencipe, S., Siciliano, I., Contessa, C., Botta, R., Garibaldi, A., Gullino, M.L. et al. 2018. Characterization of *Aspergillus* section *Flavi* isolated from fresh chestnuts and along the chestnut flour process. *Food Microbiology* 69: 159–169. DOI: 10.1016/j.fm.2017.08.004.
- Probst, C., Callicott, K.A. and Cotty, P.J. 2012. Deadly strains of Kenyan *Aspergillus* are distinct from other aflatoxin producers. *European Journal of Plant Pathology* 132: 419–429. DOI: 10.1007/s10658-011-9887-y
- Probst, C., Bandyopadhyay, R. and Cotty, P.J. 2014. Diversity of aflatoxin-producing fungi and their impact on food safety in sub-Saharan Africa. *International Journal of Food Microbiology* 174: 113–122. DOI: 10.1016/j.ijfoodmicro.2013.12.010.
- Purcell, S.L., Phillips, D.J. and Mackey, B.E. 1980. Distribution of *Aspergillus flavus* and other fungi in several almond-growing areas of California. *Phytopathology* 70(9): 926–929.
- Rahimi, P., Sharifnabi, B. and Bahar, M. 2007. Detection of aflatoxin in *Aspergillus* species isolated from pistachio in Iran. *Journal Phytopathology* 156(1): 15–20. DOI: 10.1111/j.1439-0434.2007.01312.x.
- Rasheed, U., Wu, H., Wei, J., Ou, X., Qin, P., Yao, X. et al. 2019. A polyphasic study of *Aspergillus* section *Flavi* isolated from corn in Guangxi, China – a hot spot of aflatoxin contamination. *International Journal of Food Microbiology* 310: 108307. DOI: 10.1016/j.ijfoodmicro.2019.108307.
- Razzaghi-Abyaneh, M., Shams-Ghahfarokhi, M., Allameh, A., Kazeroon-Shiri, A., Ranjbar-Bahadori, S., Mirzahoseini, H. et al. 2006. A survey on distribution of *Aspergillus* section *Flavi* in corn field soils in Iran: Population patterns based on aflatoxins, cyclopiazonic acid and sclerotia production. *Mycopathologia* 161(3): 183–192. DOI: 10.1007/s11046-005-0242-8.
- Reddy, K.R.N., Farhana, N.I., Wardah, A.R. and Salleh, B. 2010. Morphological identification of foodborne pathogens colonizing rice grains in South Asia. *Pakistan Journal of Biological Sciences* 13(16): 794–801.
- Reis, T.A.D., Oliveira, T.D., Baquião, A.C., Gonçalves, S.S., Zorzete, P. and Corrêa, B. 2012. Mycobiota and mycotoxins in Brazil nut samples from different states of the Brazilian Amazon region. *International Journal of Food Microbiology* 159(2): 61–68. DOI: 10.1016/j.ijfoodmicro.2012.08.005.
- Riba, A., Mokrane, S., Mathieu, F., Lebrihi, A. and Sabaou, N. 2008. Mycoflora and ochratoxin-A producing strains of *Aspergillus* in Algerian wheat. *International Journal of Food Microbiology* 122(1–2): 85–92. DOI: 10.1016/j.ijfoodmicro.2007.11.057.

- Riba, A., Bouras, N., Mokrane, S., Mathieu, F., Lebrihi, A. and Sabaou, N. 2010. *Aspergillus* section *Flavi* and aflatoxins in Algerian wheat and derived products. *Food and Chemical Toxicology* 48(10): 2772–2777. DOI: 10.1016/j.fct.2010.07.005.
- Rodrigues, P., Venâncio, A., Kozakiewicz, Z. and Lima, N. 2009. A polyphasic approach to the identification of aflatoxigenic and non-aflatoxigenic strains of *Aspergillus* section *Flavi* isolated from Portuguese almonds. *International Journal of Food Microbiology* 129(2): 187–193. DOI: 10.1016/j.ijfoodmicro.2008.11.023.
- Rodrigues, P., Venâncio, A. and Lima, N. 2012. Aflatoxigenic fungi and aflatoxins in Portuguese almonds. *The Scientific World Journal* vol n pp DOI: 10.1100/2012/471926.
- Rychlik, M., Humpf, H.U., Marko, D., Dänicke, S., Mally, A., Berthiller, F. et al. 2014. Proposal of a comprehensive definition of modified and other forms of mycotoxins including “masked” mycotoxins. *Mycotoxin Research* 30(4): 197–205. DOI: 10.1007/s12550-014-0203-5.
- Saadullah, A. and Abdullah, S. 2015. Mycobiota and incidence of toxigenic fungi in dried fruits from Duhok Markets, North Iraq. *Egyptian Academic Journal of Biological Sciences, G. Microbiology* 7(1): 61–68.
- Saberi-Riseh, R., Javan-Nikkhah, M., Heidarian, R., Hosseini, S. and Soleimani, P. 2004. Detection of fungal infectious agent of wheat grains in store-pits of Markazi province, Iran. *Communications in Agricultural and Applied Biological Sciences* 69(4): 541–544.
- Saleem, M.J., Hannan, A. and Qaisar, T.A. 2012. Occurrence of aflatoxins in maize seed under different conditions. *International Journal of Agriculture & Biology* 14(3): 473–476.
- Sales, A.C. and Takumi, Y. 2005. Updated profile of aflatoxin and *Aspergillus* section *Flavi* contamination in rice and its byproducts from the Philippines. *Food Additives & Contaminants* 22(5): 429–436. DOI: 10.1080/02652030500058387.
- Samson, R.A. and K.A. Seifert. 1985. The ascomycete genus *Penicillium* and its anamorphs. pp. 397–426. *In*: R.A. Samson and J.I. Pitt (eds.). *Advances in Penicillium and Aspergillus Systematics*. Plenum Press, New York and London.
- Samson, R.A., Visagie, C.M., Houbraken, J., Hong, S.B., Hubka, V., Klaassen, C.H. et al. 2014. Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology* 78: 141–173. DOI: 10.1016/j.simyco.2014.07.004.
- Savi, G.D., Piacentini, K.C., Tibola, C.S. and Scussel, V.M. 2014. Mycoflora and deoxynivalenol in whole wheat grains (*Triticum aestivum* L.) from Southern Brazil. *Food Additives & Contaminants: Part B* 7(3): 232–237. DOI: 10.1080/19393210.2014.898337.
- Schatzmayr, G. and Streit, E. 2013. Global occurrence of mycotoxins in the food and feed chain: Facts and figures. *World Mycotoxin Journal* 6(3): 213–222. DOI: 10.3920/WMJ2013.1572
- Sebök, F., Dobolyi, C., Hartman, M., Risa, A., Kritafon, C., Szoboszlai, S. et al. 2014. Occurrence of potentially aflatoxin producing *Aspergillus* species in maize fields of Abstract Book of the Meeting of the Hungarian Microbiological Society and the EU FP7 PROMISE Hungary. *In*: Regional Meeting, pp. 62–63.
- Şenyuva, H., Gilbert, J., Samson, R.A., Özcan, S., Öztürköğlü, Ş. and Önal, D. 2008. Occurrence of fungi and their mycotoxins in individual Turkish dried figs. *World Mycotoxin Journal* 1(1): 79–86. DOI: 10.3920/WMJ2008.x009

AQ : SP ? Saadulah in Table 4 p

- Sharfun-Nahar, S.N., Mushtaq, M. and Pathan, I.H. 2004. Seed-borne mycoflora of *Capsicum annuum* imported from India. Pakistan Journal of Botany 36(1): 191–198.
- Sharma, S., Gupta, D. and Sharma, Y.P. 2013. Aflatoxin contamination in chilgoza pine nuts (*Pinus gerardiana* wall.) commercially available in retail markets of Jammu, India. International Journal of Pharma and Bio. Sciences 4: 751–759.
- Singh, P. and Cotty, P.J. 2019. Characterization of *Aspergilli* from dried red chilies (*Capsicum* spp.): Insights into the etiology of aflatoxin contamination. International Journal of Food Microbiology 289: 145–153. DOI: 10.1016/j.ijfoodmicro.2018.08.025
- Sinha, A.K. and Sinha, K.K. 1990. Insect pests, *Aspergillus flavus* and aflatoxin contamination in stored wheat: A survey at North Bihar (India). Journal of Stored Products Research 26(4): 223–226. DOI: 10.1016/0022-474X(90)90026-O.
- Škrinjar, M.M., Miklič, V.J., Kocić-Tanackov, S.D., Jeromela-Marjanović, A.M., Maširević, S.N., Suturović, I.Z. et al. 2013. Xerophilic mycopopulations isolated from rapeseeds (*Brassica napus*). Acta Periodica Technologica 44: 115–124. DOI: 10.2298/APT1344115S.
- Soares, C., Rodrigues, P., Peterson, S.W., Lima, N. and Venâncio, A. 2012. Three new species of *Aspergillus* section *Flavi* isolated from almonds and maize in Portugal. Mycologia 104: 682–697. DOI: 10.3852/11-088.
- Stack, J. and Carlson, M. 2003. NF571 *Aspergillus flavus* and Aflatoxins in Corn. Historical Materials from University of Nebraska-Lincoln Extension.
- Suárez-Bonnet, E., Carvajal, M., Méndez-Ramírez, I., Castillo-Urueta, P., Cortés-Eslava, J., Gómez-Arroyo, S. et al. 2013. Aflatoxin (B1, B2, G1, and G2) contamination in rice of Mexico and Spain, from local sources or imported. J. Food Sci. 78(11): T1822-T1829.
- Sultan, Y. and Magan, N. 2010. Mycotoxigenic fungi in peanuts from different geographic regions of Egypt. Mycotoxin Research 26: 133–140. DOI: 10.1007/s12550-010-0048-5.
- Takahashi, H., Kamimura, H. and Ichinoe, M. 2004. Distribution of aflatoxin-producing *Aspergillus flavus* and *Aspergillus parasiticus* in sugarcane fields in the southernmost islands of Japan. Journal of Food Protection 67(1): 90–95. DOI: 10.4315/0362-028X-67.1.90.
- Taniwaki, M.H., Pitt, J.I., Iamanaka, B.T., Sartori, D., Copetti, M.V., Balajee, A. et al. 2012. *Aspergillus bertholletius* sp. nov. from Brazil Nuts. PLoS ONE 7: 42480. DOI: 10.1371/journal.pone.0042480.
- Taniwaki, M.H., Frisvad, J.C., Ferranti, L.S., de Souza Lopes, A., Larsen, T.O., Fungaro, M.H.P. et al. 2017. Biodiversity of mycobiota throughout the Brazil nut supply chain: From rainforest to consumer. Food Microbiology 61: 14–22. DOI: 10.1016/j.fm.2016.08.002
- Taniwaki, M.H., Pitt, J.I. and Magan, N. 2018. *Aspergillus* species and mycotoxins: Occurrence and importance in major food commodities. Current Opinion in Food Science 23: 38–43. DOI: 10.1016/j.cofs.2018.05.008.
- TePaske, M.R., Gloer, J.B., Wicklow, D.T. and Dowd, P.F. 1992. Aflavarin and beta-aflatrem: New anti-insectan metabolites from the sclerotia of *Aspergillus flavus*. Journal of Natural Products 55: 1080–1086. <http://dx.doi.org/10.1021/np50086a008>.
- Theumer, M.G., Henneb, Y., Khoury, L., Snini, S.P., Tadrist, S., Canlet, C. et al. 2018.

- Genotoxicity of aflatoxins and their precursors in human cells. *Toxicology Letters* 287: 100–107. DOI: 10.1016/j.toxlet.2018.02.007.
- Tomar, D.S., Shastry, P.P., Nayak, M.K. and Sikarwar, P. 2012. Effect of seed borne mycoflora on cotton seed (JK 4) and their control. *Journal Cotton Restitution of Development* 26(1): 105–108.
- Tran-Dinh, N., Kennedy, I., Bui, T. and Carter, D. 2009. Survey of Vietnamese peanuts, corn and soil for the presence of *Aspergillus flavus* and *Aspergillus parasiticus*. *Mycopathologia* 168(5): 257–268. DOI: 10.1007/s11046-009-9221-9.
- Trung, T., Tabuc, C., Bailly, S., Querin, A., Guerre, P. and Bailly, J. 2008. Fungal mycoflora and contamination of maize from Vietnam with aflatoxin B1 and fumonisin B1. *World Mycotoxin Journal* 1(1): 87–94. DOI: 10.3920/WMJ2008.x010.
- Turgay, E.B. and Ünal, F. 2009. Detection of seed borne mycoflora of sorghum in Turkey. *The Journal of Turkish Phytopathology* 38(1-3): 9–20. DOI:
- Turner, P.C., Sylla, A., Gong, Y.Y., Diallo, M.S., Sutcliffe, A.E., Hall, A.J. et al. 2005. Reduction in exposure to carcinogenic aflatoxins by postharvest intervention measures in West Africa: A community based intervention study. *Lancet* 365: 1950–1956. DOI: 10.1016/S0140-6736(05)66661-5.
- Umemura, M., Koike, H., Nagano, N., Ishii, T., Kawano, J., Yamane, N. et al. 2013. MIDDAS-M: Motif-independent de novo detection of secondary metabolite gene clusters through the integration of genome sequencing and transcriptome data. *PLoS One* 8: e84028. DOI: 10.1371/journal.pone.0084028.
- Vaamonde, G., Patriarca, A., Pinto, V.F., Comerio, R. and Degrossi, C. 2003. Variability of aflatoxin and cyclopiazonic acid production by *Aspergillus* section *Flavi* from different substrates in Argentina. *International Journal of Food Microbiology* 88(1): 79–84. DOI: 10.1016/S0168-1605(03)00101-6.
- Varga, J., Frisvad, J.C. and Samson, R.A. 2011. Two new aflatoxin producing species and an overview of *Aspergillus* section *Flavi*. *Studies in Mycology* 69: 57–80. DOI: 10.3114/sim.2011.69.05.
- Varga, J., Baranyi, N., Chandrasekaran, M., Vágvölgyi, C. and Kocsubé, S. 2015. Mycotoxin producers in the *Aspergillus* genus: An update. *Acta Biologica Szegediensis* 59: 151–167.
- Wallace, H.A.H. and Sinha, R.N. 1962. Fungi associated with hot spots in farm stored grain. *Canadian Journal of Plant Science* 42(1): 130–141. DOI: 10.4141/cjps62-016.
- Wartu, J.R., Whong, C.M.Z., Umoh, V.J. and Diya, A.W. 2015. Occurrence of aflatoxin levels in harvest and stored groundnut kernels in Kaduna State, Nigeria. *Journal of Environmental Science, Toxicology and Food Technology* 9(1): 62–66. DOI: 10.9790/2402-09126266.
- Watarai, N., Yamamoto, N., Sawada, K. and Yamada, T. 2019. Evolution of *Aspergillus Oryzae* before and after domestication inferred by large-scale comparative genomic analysis. *DNA Research* 26: 465–472 DOI: 10.1093/dnares/dsz024.
- Weledesemaya, G.T., Gezmu, T.B., Woldegiorgis, A.Z. and Gemele H.F. 2016. Study on *Aspergillus* species and aflatoxin levels in sorghum (*Sorghum bicolor* L.) stored for different period and storage system in Kewet Districts, Northern Shewa, Ethiopia. *JSOA Journal of Food Science and Nutrition* 2(1): 1–8. DOI: 10.24966/FSN-1076/100010.

- Wu, F. and Guclu, H. 2012. Aflatoxin regulations in a network of global maize trade. *PloS One* 7(9): 45151. DOI: 10.1371/journal.pone.0045151.
- Yogendrarajah, P., Vermeulen, A., Jacxsens, L., Mavromichali, E., De Saeger, S., De Meulenaer, B. et al. 2016. Mycotoxin production and predictive modelling kinetics on the growth of *Aspergillus flavus* and *Aspergillus parasiticus* isolates in whole black peppercorns (*Piper nigrum* L). *International Journal of Food Microbiology* 228: 44–57. DOI: 10.1016/j.ijfoodmicro.2016.03.015.
- Zhang, S., Monahan, J.B., Tkacz, J.S. and Berry, S. 2004. Indole diterpene gene cluster from *Aspergillus flavus*. *Applied and Environmental Microbiology* 70: 6875–6883. DOI: 10.1128/AEM.70.11.6875–6883.2004.
- Zohri, A.A. and Abdel-Gawad, K.M. 1993. Survey of mycoflora and mycotoxins of some dried fruits in Egypt. *Journal of Basic Microbiology* 33(4): 279–288. DOI: 10.1002/jobm.3620330413.