



Durable resistance or efficient disease control? Adult Plant Resistance (APR) at the heart of the dilemma

Loup Rimbaud, Julien Papaïx, Jean-François Rey, Benoît Moury, Luke Barrett, Peter Thrall

► To cite this version:

Loup Rimbaud, Julien Papaïx, Jean-François Rey, Benoît Moury, Luke Barrett, et al.. Durable resistance or efficient disease control? Adult Plant Resistance (APR) at the heart of the dilemma. 2022. hal-03775182

HAL Id: hal-03775182

<https://hal.inrae.fr/hal-03775182>

Preprint submitted on 12 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Durable resistance or efficient disease control?

Adult Plant Resistance (APR) at the heart of the dilemma

Loup Rimbaud^{1*}, Julien Papaïx², Jean-François Rey², Benoît Moury¹, Luke Barrett³, Peter Thrall⁴

¹ INRAE Pathologie Végétale, 84143 Montfavet, France

² INRAE BioSP, 84914 Avignon, France

³ CSIRO Agriculture and Food, Canberra 2601 ACT, Australia

⁴ CSIRO National Collections & Marine Infrastructure, Canberra 2601 ACT, Australia

* Corresponding author

Correspondence: loup.rimbaud@inrae.fr

<http://orcid.org/0000-0002-8098-9984>

ABSTRACT

Adult plant resistance (APR) is an incomplete and delayed protection of plants against pathogens. At first glance, such resistance should be less efficient than classical major-effect resistance genes, which confer complete resistance from seedling stage, to reduce epidemics. However, by allowing some 'leaky' levels of disease, APR genes are predicted to be more durable than major genes because they exert a softer selection pressure on pathogens towards adaptation to resistance. However, the impact of partial efficiency and delayed mode of action of APR on the evolutionary and epidemiological outcomes of resistance deployment has never been tested.

Using the demogenetic, spatially explicit, temporal, stochastic model *landsepi*, this study is a first attempt to investigate how resistance efficiency, time to resistance expression and target pathogenicity trait jointly impact resistance durability and disease control at the landscape scale. Our numerical experiments explore the deployment of APR in a simulated agricultural landscape, alone or together with a major resistance gene. As a case study, the mathematical model has been parameterised for rust fungi (genus *Puccinia*) of cereal crops, for which extensive data are available.

Our simulations confirm that weak efficiency and delayed expression of APR genes reduce the selection pressure applied on pathogens and their propensity to overcome resistance, but do not confer effective protection. On the other hand, stronger APR genes (which increase selection pressure on the pathogen) may be quickly overcome but have the potential to provide some disease protection in the short-term. This is attributed to strong competition between different pathogen genotypes and the presence of fitness costs of adaptation, especially when APR genes are deployed together with major resistance gene via crop mixtures or rotations.

Keywords: adaptation, adult plant resistance, disease control, immunity, mature plant resistance, ontogenic, puccinia, resistance, resistance durability, rust, simulation modelling.

34

Introduction

35 In plant pathology, durable resistance and efficient disease control are two important
36 considerations in the use of genetically controlled plant resistance to manage crop diseases (Burdon JJ
37 et al., 2016). Indeed, strategies to deploy plant resistance should first be as efficient as possible to
38 mitigate epidemics and preserve crop health. However, the high evolutionary potential of many plant
39 pathogens means that they can adapt and overcome such resistance, sometimes quickly after
40 deployment in the field (Johnson R, 1983; Parlevliet JE, 2002; García-Arenal F & BA McDonald, 2003).
41 Resistance breakdown results in potentially destructive epidemics and economic losses, leading to
42 increased reliance on pesticides and acceleration of associated environmental issues. In addition,
43 resistance breakdown also means the loss of precious and non-renewable genetic resources, and the
44 need to develop new resistant cultivars, a long and costly process (Zhan J et al., 2015). Therefore, in
45 addition to the provision of efficient crop protection in the short-term, resistance must also be durable,
46 even if these two goals are not necessarily compatible (van den Bosch F & CA Gilligan, 2003; Papaix J
47 et al., 2018; Rimbaud L et al., 2018a). In this context, simulation models provide powerful tools to
48 explore and evaluate different crop deployment strategies with respect to their epidemiological and
49 evolutionary outcomes, while circumventing the logistical challenges associated with field experiments
50 at large spatio-temporal scales (Rimbaud L et al., 2021).

51 Plant breeding has typically focused on resistance conferred by major-effect genes, which often
52 confer complete resistance, such that pathogens are unable to infect cultivars carrying those genes.
53 Most major genes encode for an immune receptor of the nucleotide-binding leucine-rich repeat (NLR)
54 protein family, which triggers the immune response (often involving a hypersensitive reaction) after
55 recognition of a pathogen effector (de Ronde D et al., 2014; Gallois J-L et al., 2018). Nevertheless,
56 pathogens may escape this recognition after mutation or suppression of this effector, leading to the
57 restoration of infectivity and resistance breakdown. In these cases, the plant-pathogen genetic
58 interaction is best described by the 'gene-for-gene' (GFG) model, according to which the occurrence
59 of disease depends on whether or not the plant carries a resistance gene, and whether or not the
60 pathogen possesses the matching effector (Flor HH, 1955). The scientific literature describes numerous
61 examples of major resistance genes being rapidly overcome by fungi (Johnson R, 1983, 1984;
62 McDonald BA & C Linde, 2002; Parlevliet JE, 2002; Stuthman DD et al., 2007; Park RF, 2008; Burdon JJ
63 & PH Thrall, 2014), bacteria (McDonald BA & C Linde, 2002; Parlevliet JE, 2002), viruses (García-Arenal
64 F & BA McDonald, 2003; Lecoq H et al., 2004; Moury B et al., 2010), and nematodes (McDonald BA &
65 C Linde, 2002), although some of them have maintained effectiveness for many years. Such resistance
66 breakdown results from the high selection pressure experienced by pathogen populations in the
67 presence of such resistance, since only adapted individuals can infect resistant hosts. 'Resistance-
68 breaking' mutants may be initially present in the population at low frequency, derive from other
69 pathogen genotypes by mutation or recombination, or be introduced from distant areas through
70 migration. In such cases, the frequency of the mutant genotype increases as it will be strongly favoured
71 by selection and the whole host population may end up infected (Johnson R, 1983, 1984; Lecoq H et
72 al., 2004; Moury B et al., 2010).

73 Resistance is, however, not always complete or continuous in time. Whether they may be
74 insufficiently expressed, dependent on environmental conditions or simply weak, resistance genes
75 sometimes confer only partial protection to pathogens. In this context, 'resistance efficiency' is a key
76 component of partial resistance, and describes how well the infectious cycle of the pathogen is

mitigated, i.e., the extent of reduction of one or several pathogenicity traits, such as infection rate, latent or infectious period durations, and reproduction rate (Parlevliet JE, 1979; Lannou C, 2012). Resistance may also be specific to certain host developmental phases (Barrett LG & M Heil, 2012), such as is the case for adult plant resistance (APR, also called ‘mature plant resistance’; Develey-Rivière M-P & E Galiana, 2007). APR genes are often described as being only expressed in adult plants (Burdon JJ et al., 2014; Niks RE et al., 2015), with an efficiency varying from 0% to 100% and depending on plant age and environment (Krattinger SG & B Keller, 2016). However, moderate levels of expression of APR genes can sometimes be detected in young plants (Park RF & RG Rees, 1989; Cromeley MG, 1992; Broers LHM, 1997; Sandoval-Islas JS et al., 2007; Qamar M et al., 2012). This expression tends to increase progressively and the date after which APR genes are fully active (referred to as ‘time to resistance expression’ hereafter) depends on the resistance gene and may occur as late as the anthesis stage (Ma H & RP Singh, 1996). Many APR genes against rust fungi have been documented in cereal crops (Burdon JJ, 1987 p56; McIntosh RA et al., 1995; Boyd LA, 2005). They can impact all pathogenicity traits associated with the pathogen infectious cycle: infection rate (e.g. Lr34-Yr18; Qamar M et al., 2012), latent period (Lr16-Lr18, Lr34-Yr18; Tomerlin JR et al., 1983; Elahinia SA & JP Tewari, 2005; Qamar M et al., 2012), sporulation rate (Lr16-Lr18; Tomerlin JR et al., 1983), sporulation duration (Lr16-Lr18; Tomerlin JR et al., 1983). Nonetheless, a wide panoply of molecular mechanisms may underpin APR resistance and these are poorly known (Develey-Rivière M-P & E Galiana, 2007; Krattinger SG & B Keller, 2016). Exceptions include three resistance genes against leaf, stem and yellow rusts of wheat: Lr67 encoding a hexose transporter (Moore JW et al., 2015); Lr34 encoding an ATP-binding cassette (ABC) transporter (Krattinger SG et al., 2009); and Yr36 encoding a chloroplast-localised kinase protein involved in detoxification of reactive oxygen species (Fu D et al., 2009, see also Develey-Rivière M-P & E Galiana, 2007 for resistances against other pathogens).

To the best of our knowledge, the role of delayed expression of plant resistance in disease management and pathogen evolution has never been investigated in simulation models (Rimbaud L et al., 2021), despite its supposed potential to promote resistance durability. Complete resistance is often assumed in modelling studies, and always considered active from the seedling stage. Yet, hosts are thought to generate different selective pressures on pathogens if they express complete, partial or delayed resistance (Stuthman DD et al., 2007; Pilet-Nayel M-L et al., 2017). While complete resistance exerts hard selection on the pathogen to restore infectivity, the pressure imposed by partial and delayed resistances (such as the one conferred by APR genes) is likely lower since they allow some ‘leaky’ levels of disease. Partial and delayed resistances can thus be seen as soft selection mechanisms that slow down the speed of pathogen evolution compared to typical major resistance genes. This slower pathogen evolution comes nonetheless at the price of weaker protection against disease, hence the potential of such resistance for disease management is still intriguing, particularly when deployed in conjunction with major gene resistance.

The aim of the present study is to investigate how resistance efficiency, time to resistance expression and target pathogenicity trait of a resistance gene jointly impact resistance durability and epidemiological disease control. Additionally, because deploying different types of resistance is likely a promising approach to benefit from their respective advantages, we also investigate the best strategies to combine a major resistance gene with an APR gene. To study these questions, we use a general simulation framework implemented in the R package *landsepi* (Rimbaud L et al., 2018b). The model is flexible enough to vary parameters related to the deployed resistance genes, and to

encompass various pathogen epidemiological traits. Thus, although this work is motivated by rust diseases of cereal crops (for which there is considerable empirical data), our broad conclusions may, to some extent, apply to numerous pathosystems.

Methods

Model overview

We used a demogenetic, spatially explicit, temporal and stochastic model developed to explore different plant resistance deployment strategies in agricultural landscapes and evaluate their epidemiological and evolutionary outcomes. A description of the mathematical model is detailed in a previous article (Rimbaud L et al., 2018c). Briefly, the model simulates the spread (by wind) and evolution (via mutation) of a spore-borne fungal pathogen in a cropping landscape where susceptible and resistant cultivars are cultivated with controlled proportions and controlled level of spatial aggregation. While the model has the capacity to simulate sexual reproduction for the pathogen, here we assume clonality. In the simulated landscape, resistance genes may be deployed in a single host cultivar as a pyramid, or in different cultivars that can be segregated in a mosaic of fields, combined within the same field as mixtures, or alternated within crop rotations. Resistance genes may target one or several pathogenicity traits (reduction of infection rate, sporulation rate or sporulation duration, lengthening of latent period duration) with complete or partial efficiency. The pathogen has the potential to adapt to each of the deployed resistance genes independently, via single or multiple mutations (leading to the emergence of new pathogen strains), possibly associated with a fitness cost on the susceptible cultivar. The pathogen is disseminated across the landscape using a power-law dispersal kernel: $g(\|z' - z\|) = \frac{(b-2)(b-1)}{2\pi a^2} \cdot \left(1 + \frac{\|z' - z\|}{a}\right)^{-b}$ with $\|z' - z\|$ the Euclidian distance between locations z and z' in fields i and i' , respectively, a the scale parameter and b a parameter linked to the width of the tail. The plant infection and immune status is modelled using a traditional SEIR ('susceptible-exposed-infectious-removed') framework. Plant harvests occur at the end of each cropping season, imposing potential bottlenecks (and thus genetic drift) on the pathogen population.

In this study, the *landsepi* model was extended to include resistance genes with a delayed expression (i.e., APR genes). Cultivars that carry an APR gene are susceptible at the beginning of the cropping season and become resistant once the gene activates. The time to resistance expression is drawn from a gamma distribution every year and for every field planted with a cultivar carrying an APR gene. For convenience, this distribution is parameterised with the expectation and variance of the time to expression. Both parameters, as well as the target pathogenicity trait and efficiency of resistance, are assumed to be genetically determined and thus characteristic of a given APR gene.

For the simulation experiments, we parameterised the model using available data from the empirical literature to represent wheat rust infection caused by a range of fungal pathogens in the genus *Puccinia* (Table 1, details on model calibration in Rimbaud L et al., 2018c), supporting information). The model is available in the R package *landsepi* version 1.1.1 (Rimbaud L et al., 2018b).

Numerical experiments

Three successive numerical experiments were carried out to explore APR. Experiment 1 is a baseline scenario destined to evaluate how the deployment of a single APR gene mitigates epidemics

in absence of pathogen evolution (i.e., here epidemics are caused by a single pathogen strain, not adapted to the APR gene). Experiment 2 reproduces the same scenario but includes pathogen evolution, to measure the durability of the APR gene and the epidemiological impact of the possible presence of adapted pathogen genotypes. Finally, Experiment 3 investigates whether APR genes and major resistance genes are competing alternatives or can be complementary to each other via appropriate spatio-temporal deployment strategies. **Table 1** summarises model parameters of interest.

In the first two experiments, the landscape (representing approximately 150 fields, total area: 2x2 km², see Fig S1 in Rimbaud L et al., 2018c) was composed of a mosaic of a susceptible (1/3 of total surface) and a resistant cultivar (2/3 of total surface) across the simulated landscape. Cultivars were randomly allocated to fields within the landscape either at low or at high degree of spatial aggregation (**Fig. 2**, left-hand column). The resistant cultivar carried a resistance targeting either infection rate, latent period duration, sporulation rate, or sporulation duration of the pathogen. Analysis of field and greenhouse trials on rust diseases of cereal crops revealed that resistance against these pathogenicity traits measured in different host genotypes can vary from 0% to 100% compared to the most susceptible cultivars (**Table S1**). Thus, in our simulations, resistance efficiency was varied from 0 to 100% with increments of 10%. The expected time to resistance expression varied from 0 to 90 days with increments of 10 days; a time to expression of 90 days (the whole epidemic season being 120 days) represents the case where the gene activates at anthesis stage. For example, if the resistance gene targets the latent period duration with an efficiency of 75% and a time to expression of 30 days, a non-adapted (i.e., 'wild type', wt) pathogen infecting a resistant cultivar will have an expected latent period of 10 days (see **Table 1**) until resistance activates, after which latent period is increased by 75% (i.e., 17.5 days) until the end of the cropping season. In the first experiment, the pathogen was not allowed to evolve, whereas in the second, it could adapt to the APR gene through mutation. In this case, the impact of fitness cost of adaptation (where fitness cost was defined in terms of loss of pathogenicity on the susceptible cultivar) was studied using three (0.00, 0.25, 0.50) different cost values. Model stochasticity includes field shape and boundaries, cultivar allocation to the different fields within the simulated landscape, time to APR gene expression, pathogen dispersal, mutation, off-season survival, and SEIR transitions. To account for this stochasticity, simulations were run on five different landscape structures and replicated 10 times, resulting in 50 replicates for every parameter combination. Thus, the complete factorial design of the first two experiments resulted in a total of 44,000 and 132,000 simulations, respectively.

In the third numerical experiment, a major resistance gene and an APR gene were jointly deployed according to one of four strategies: pyramiding, mixture, rotation or mosaic. The major resistance gene was assumed to target pathogen infection rate with complete efficiency and to be fully expressed from the beginning of the cropping season. Target pathogenicity trait, resistance efficiency and time to expression of the APR gene were varied exactly as in the first two experiments. However, for this experiment, spatial aggregation was fixed at a low value (representing a fragmented landscape), and the fitness cost of pathogen adaptation to 0.50. Indeed, results obtained in the second experiment showed that this parameterisation maximises the interaction between cultivars (in terms of pathogen dispersal and competition between pathogen genotypes) within a spatial deployment strategy. For all deployment strategies, 1/3 of the landscape was composed of the susceptible cultivar. The remaining 2/3 were occupied either by a single cultivar carrying the two genes (pyramid strategy), a mixture (in every field) of two resistant cultivars in balanced proportions (each cultivar carrying one

of the two genes; mixture strategy), a rotation of these two resistant cultivars (every year; rotation strategy), or a mosaic of the two resistant cultivars in balanced proportions (every cultivar representing 1/3 of the landscape area; mosaic strategy) (**Fig. S6**). With 50 stochastic replicates, the complete factorial design resulted in a total of 88,000 simulations.

Simulations were run for 120 time-steps per cropping season over a 30-year time period. Initially, only the wild-type pathogen (i.e., not adapted to any resistance), 'wt', was present in susceptible hosts, with a probability of any host being initially infected of $5 \cdot 10^{-4}$. The wt strain is unable to infect resistant hosts carrying an APR gene only if resistance is both complete and active. In all other situations, the wt strain is able to infect the hosts carrying an APR gene. In any case, a single mutation (with probability 10^{-4} , except in the first experiment where evolution did not occur) is required to overcome a resistance gene (should it be a major gene or an APR gene) and restore complete pathogenicity, in conformity with a gene-for-gene interaction.

Model outputs

In this work, epidemiological control is defined as the ability of a given deployment strategy to reduce disease impact on the resistant cultivar(s). Here, it is measured by the relative green leaf area (GLA), i.e., the proportion of healthy hosts relative to the total number of hosts, averaged for every cultivar across the whole simulation run. The higher the value of the GLA, the better the epidemiological control.

Evolutionary control is quantified here using resistance durability (for experiments 2 and 3), which measures the ability of a given deployment strategy to limit pathogen evolution and delay resistance breakdown (i.e., emergence of the resistance-breaking, 'rb', pathogen). Durability is evaluated using the time when the number of resistant hosts infected by the rb strain exceeds a threshold above which extinction of this strain is unlikely (fixed at 50,000, see Rimbaud L et al., 2018c), supporting Text S2 for details). To understand the contribution of the different pathogen genotypes to an epidemic, we also calculate, across the whole simulation run and for every cultivar, the proportion of infections due to each pathogen genotype relative to all infections.

229 **Table 1. Model parameter and simulation experiments.** See Text S1 in (Rimbaud L et al., 2018c) for calibration details. Parameters of interest (blue cells) were varied
230 according to a complete factorial design. Every simulation was replicated 10 times x 5 landscape structures to account for stochasticity, resulting in a total of 44,000,
231 132,000 and 88,000 simulations for the three numerical experiments, respectively.

Parameter	Experiment 1 (single APR gene, no evolution)	Experiment 2 (single APR gene)	Experiment 3 (major gene + APR gene)
Pathogen parameters			
Dispersal scale parameter (a) ^a	40		
Width of the dispersal kernel tail (b) ^a	7		
Maximal expected infection rate	0.40 spore ⁻¹		
Minimal expected latent period duration	10 days		
Variance of the latent period duration	9 days		
Maximal expected sporulation duration	24 days		
Variance of the sporulation duration	105 days		
Maximal expected sporulation rate	3.125 spores.day ⁻¹		
Initial probability of infection of susceptible hosts	5.10 ⁻⁴		
Off-season survival probability	10 ⁻⁴		
Landscape organisation ^b			
Number of fields in the landscape ^c	155; 154; 152; 153; 156		
Deployment strategy	Mosaic	Mosaic	Mosaic, mixture, rotation, pyramid
Proportion of landscape area covered by the susceptible cultivar	1/3		
Level of spatial aggregation	low; high	low; high	low
Major gene resistance			
Target pathogenicity trait	-	-	Infection rate
Resistance efficiency (ρ)	-	-	1.00
Expected time to resistance expression	-	-	0 day
Variance of the time to resistance expression	Equal to the expected time		
Adult plant resistance			
Target pathogenicity trait	Infection rate; latent period duration; sporulation rate; sporulation duration		

Resistance efficiency ^d	0.00; 0.10; 0.20; 0.30; 0.40; 0.50; 0.60; 0.70; 0.80; 0.90; 1.00		
Expected time to resistance expression ^e	0; 10; 20; 30; 40; 50; 60; 70; 80; 90 days		
Variance of the time to resistance expression ^f	Equal to the expected time		
Pathogen evolutionary ability ^g			
Mutation probability ^h	0	10 ⁻⁴	10 ⁻⁴
Fitness cost of adaptation (θ) ⁱ	-	0.00, 0.25; 0.50	0.50

^a The mean dispersal distance is given by: $\frac{2a}{(b-3)} = 20 \text{ m}$, but long-distance dispersal may also occur due to the heavy-tail shape of the power law.

^b crop cultivars are allocated using an algorithm based on latent Gaussian fields to control proportion and level of spatial aggregation of each cultivar; see Fig. S1 of (Rimbaud L et al., 2018a) for illustrations and (Rimbaud L et al., 2018c) for details on the algorithm.

^c see Fig S1 in (Rimbaud L et al., 2018c) for illustrations of landscape structures generated using a T-tesselation algorithm, and see (Papaïx J et al., 2014a) for details on the algorithm.

^d an efficiency of 0.00 is equivalent to the absence of a resistance gene.

^e a time of 90 days represents gene activation at anthesis stage.

^f when expectation and variance are 0 day, there is no variation in the time to expression.

^g same value for major gene and adult plant resistance.

^h probability of a spore changing its phenotype on a resistant cultivar carrying a resistance gene.

ⁱ paid by adapted pathogens on hosts that do not carry an active resistance – it consists of a reduction in the same pathogenicity trait as the one targeted by the resistance.

249

Results

250

251

252

253

254


255

256

257

258

Three separate numerical experiments were carried out to investigate the epidemiological and evolutionary outcomes of deployment strategies based on APR: the first two experiments were performed with an APR gene alone, and the third with a combination of an APR gene and a major resistance gene. In all these experiments, three parameters were systematically allowed to vary: resistance efficiency, time to resistance expression and target pathogenicity trait. Using this approach, we were able to explore a wide range of situations, from the absence of resistance (if resistance efficiency is 0%, **Fig. 1**) to a completely efficient major gene (if efficiency is 100% and there is no delay in resistance expression) with all possible intermediate situations (partially-efficient major gene, completely-efficient APR gene, partially-efficient APR gene).



259

260

261

262

Figure 1. Conceptual exploration of parameters associated with resistance genes: efficiency and time to expression. This formal framework encompasses a wide range of situations. MG: major gene ; APR: adult plant resistance.

263

264

Experiment 1: Deployment of a single APR gene in a susceptible landscape with no pathogen evolution

265

266

267

Disease control, measured by the Green Leaf Area averaged for every cultivar across the whole simulation run, was first evaluated when a single APR is deployed in the landscape and the pathogen does not have the possibility to overcome the resistant cultivar.

268

269

270

271

272

273


274

275

276

As expected, for the resistant cultivar, disease control increases with higher efficiency and shorter time to resistance expression (**Fig. 2**). Globally, the target pathogenicity trait offering the best level of disease control is the infection rate when resistance is expressed early in the cropping season, whereas it is the sporulation duration when resistance is expressed late (**Figs. 2 & S1**). On the susceptible cultivar, disease control is globally poor except when the level of spatial aggregation between cultivars is low and the APR carried by the resistant cultivar is almost completely efficient, expresses very early (i.e., it is roughly similar to a major gene), and targets the pathogen infection rate, sporulation rate or sporulation duration (**Fig. 2B**). This comes at the price of a slightly decreased level of control for the resistant cultivar compared to an aggregated landscape.

A. Aggregated landscape



B. Fragmented landscape




Figure 2. Simulated landscapes (examples on the left) and heatmaps (on the right) of the level of epidemiological control (i.e., disease limitation, measured by the Green Leaf Area, 'GLA') in the absence of pathogen evolution for different levels of resistance efficiency (vertical axis), time to resistance expression (horizontal axis) and target pathogenicity traits (columns), for strong (A) or weak (B) levels of spatial aggregation.

Experiment 2: Deployment of a single APR gene in a susceptible landscape with pathogen evolution

In this experiment, there were two possible pathogen genotypes: the *rb* and *wt* strains, respectively adapted and not adapted to the APR, whose performances on the different cultivars are summarised in **Table 2**.

Table 2. Plant-pathogen interaction matrix with a single resistance gene. The table shows the coefficients by which the value of the target pathogenicity trait (see **Table 1**) is multiplied (except for latent period duration: $1-p$ is replaced by $1+p$ and $1-\theta$ is replaced by $1+\theta$). The coefficients reflect the relative performance of the different pathogen genotypes on the different cultivars. p is the efficiency of the resistance gene and θ is the fitness cost of adaptation.

	Susceptible cultivar	Resistant cultivar (APR)	
		Non-active	Active
wild type pathogen (<i>wt</i>)	1	1	$1-p$
resistance-breaking pathogen (<i>rb</i>)	$1-\theta$	$1-\theta$	1

Impact of resistance efficiency and time to expression.

Regardless of the target pathogenicity trait, fitness cost and level of spatial aggregation, the results indicate that weak resistance (whether it is inefficient or delayed in expression; bottom right corner of graphics in **Figs. 3, S2, S3, S4**) is always durable (**panels A and B**), meaning that rb pathogen genotypes never emerged in the 30-year simulations (**panels E and F**). However, in this situation, resistance does not confer good epidemiological protection against the wt pathogen, as shown by the second output variable ('Disease control', **panels C and D**). In contrast, strong resistance (highly efficient and activated early in the growing season; top left corner of graphics in **Figs. 3, S2, S3, S4**) shows poor durability (**panels A and B**), indicating that the rb pathogen genotype quickly emerged and invaded the resistant host population (**panels E and F**). This again results in poor epidemiological control for the resistant cultivar (**panels C and D**). However, when fitness costs are large ($\theta=0.50$), there is a critical zone where disease control by the resistant cultivar reaches a higher level, particularly when infection rate is targeted by the APR gene. This zone corresponds to resistance efficiencies higher than 60% and time to expression between roughly 30 and 80 days (**Fig. 3CD**). **Fig. S5** illustrates examples of simulations carried out in the three contrasted scenarios described just above (weak resistance, strong resistance, critical zone).

Impact of fitness cost of adaptation.

Decreasing the loss of pathogenicity of the rb pathogen on the susceptible cultivar (effect of columns in **Figs. 3, S2, S3, S4**) tends to decrease both durability and disease control (at intermediate resistance efficiency and with delayed expression, rb genotypes emerge more often and cause more damage). In particular, when there are no fitness costs of adaptation, the critical zone previously described disappears completely.

Impact of the level of field spatial aggregation.

The strongest impact of spatial aggregation is on the genetic composition of the pathogen population and the associated epidemic damage (**Figs. 3, S2, S3, S4, panels E and F**). The susceptible cultivar is mostly infected by the wt pathogen in aggregated landscapes, leading to severe epidemics. In contrast, for strong resistance (highly efficient or activated early in the growing season) and in presence of fitness costs of adaptation, the susceptible cultivar is mostly infected by the rb pathogen in fragmented landscapes, resulting in moderate to good epidemiological control (due to the fitness penalty). Conversely, epidemiological control for the resistant cultivar seems slightly better in aggregated landscapes (especially when resistance is strong but considerably delayed in the cropping season, top right corner of heatmaps, **Fig. 3CD**). In the absence of fitness costs of adaptation or for weak resistance (inefficient or activated late in the growing season), the genetic composition of the pathogen is similar on the two cultivars, and the associated damage is high.

Impact of the target pathogenicity trait.

All the previous results hold qualitatively with the different pathogenicity traits targeted by resistance. When resistance targets sporulation rate or the duration of the sporulation period, the genetic composition of the pathogen population and the level of evolutionary control (resistance durability) are similar to what was observed for the infection rate (**Figs. S3, S4**). There are, however, quantitative changes in the epidemiological outcome, as size and location of the critical zone are slightly different depending on the target pathogenicity trait. For infection rate, as mentioned before, the critical zone of good disease control corresponds to resistance efficiencies higher than 60% and expression between 30 and 80 days. For sporulation rate (or sporulation duration), the critical zone corresponds to efficiencies higher than 80% (respectively 90%) and expression after 50 days

(respectively 80 days). Resistances increasing the duration of the latent period and having a high efficiency and a delayed expression (more than 30 days, **Fig. S2**, top right corner of graphics) are more durable than those targeting the other traits. This is a consequence of the absence of emergence of the *rb* pathogen. However, the level of epidemiological control is poor in comparison to the other target traits, and the size of the critical zone is considerably reduced (restricted to resistance efficiencies between 80 and 100% and times to expression of less than 20 days).




Figure 3. Heatmaps of the levels of evolutionary control (resistance durability as measured by the number of years before the emergence of the resistance-breaking ('*rb*') pathogen genotype; panels A and B), epidemiological control (i.e., disease limitation, measured by the Green Leaf Area ('GLA') on the susceptible ('S') and the resistant ('R') cultivars; panels C and D) and average frequency of the *rb* pathogen (panels E and F) for different levels of resistance efficiency (vertical axis), time to resistance expression (horizontal axis) and fitness cost of pathogen adaptation (columns), for strong (panels A, C, E) or weak (B, D, F) levels of spatial aggregation. The target pathogenicity trait is the infection rate. Circled numbers refer to example simulations in **Fig. S5**.

Experiment 3: Simultaneous deployment of a major resistance gene and an APR gene in a susceptible landscape

In a third numerical experiment, resistance durability and disease control were evaluated when a major resistance gene and an APR gene were simultaneously deployed across a landscape, either within the same cultivar (R_{12} , pyramiding strategy) or in two distinct cultivars (R_1 and R_2 , respectively) which could be cultivated in different fields (mosaic strategy), within the same field as mixtures, or alternated in time through crop rotations (see **Fig. S6** for examples of simulated landscapes). In this experiment, there are four possible pathogen genotypes, whose performances on the different cultivars are summarised in **Table 3**. Here, the level of spatial aggregation is fixed at a low value (fragmented landscape), and the fitness cost is 0.50.

Table 3. Plant-pathogen interaction matrix with two resistance genes, giving the coefficients by which the value of the target pathogenicity trait (see **Table 1**) is multiplied (except for latent period duration: $1-p$ is replaced by $1+p$ and $1-\theta$ is replaced by $1+\theta$). It reflects the relative performance of the wild-type (wt) and the resistance-breaking (rb_1 , rb_2 , rb_{12}) pathogen genotypes on the susceptible (S) and resistant cultivars carrying a major resistance gene (MG; cultivar R_1), an APR gene (R_2) or both (R_{12}). ρ_1 and ρ_2 are the efficiencies of the resistance genes, and θ_1 and θ_2 are the fitness costs of adaptation.

	S	R_1 (MG)	R_2 (APR)		R_{12} (MG+APR)	
			Non-active	Active	Non-active	Active
wt	1	0	1	$1-\rho_2$	0	0
rb ₁	$1-\theta_1$	1	$1-\theta_1$	$1-\rho_2$	1	$1-\rho_2$
rb ₂	$1-\theta_2$	0	$1-\theta_2$	1	0	0
rb ₁₂	$(1-\theta_1)(1-\theta_2)$	$1-\theta_2$	$(1-\theta_1)(1-\theta_2)$	$1-\theta_1$	$1-\theta_2$	1





Figure 4. Heatmaps showing the levels of A) evolutionary control (resistance durability, measured by the number of years before the emergence of resistance-breaking genotypes) and B) epidemiological control (i.e., disease limitation, measured by the Green Leaf Area, 'GLA') on a susceptible cultivar 'S', a resistant cultivar 'R1' carrying a completely efficient major gene ('MG') and a resistant cultivar 'R2' carrying an APR gene, for different levels of APR efficiency (vertical axis), time to APR expression (horizontal axis) and deployment strategies (columns; note that for pyramiding, R1 and R2 refer to the same cultivar). The target pathogenicity trait of the APR gene is the infection rate, the level of spatial aggregation is low, and the fitness cost is 0.50.



378

379 **Figure 5.** Average frequency of the different pathogen genotypes (see **Table 3** for notations) on a susceptible cultivar ‘S’, a resistant cultivar ‘R1’ carrying a completely
 380 efficient major gene and a resistant cultivar ‘R2’ carrying an APR gene, for different levels of APR efficiency (vertical axis), time to APR expression (horizontal axis)
 381 and deployment strategies (columns; note that for pyramiding, R1 and R2 refer to the same cultivar). The target pathogenicity trait of the APR gene is the infection
 382 rate, the level of spatial aggregation is low, and fitness cost is 0.50.

Impact of resistance efficiency, time to expression and deployment strategy.

Regardless of the characteristics of the APR gene (efficiency, time to expression, target pathogenicity trait), the major gene is always overcome quickly after deployment (**Figs. 4, S7, S8, S9**), except when it is pyramided with a very efficient APR gene that is activated early in the growing season (which is essentially the same as a pyramid of two major resistance genes). This rapid breakdown is mostly attributed to the emergence of the single mutant 'rb1' (except when the major gene is pyramided with a strong APR, in which case the breakdown is due to the double mutant 'rb12', **Fig. 5**). With respect to the durability of the APR gene and the level of protection it confers on the associated cultivar (R2), weak resistance (i.e., inefficient or delayed in expression) is durable (neither the 'rb2' nor the 'rb12' genotypes emerged) but offers poor protection against the 'wt' and 'rb1' genotypes (**Fig. 4 & 5**), similar to the results for Experiment 2. When resistance is strong (very efficient and activated early), it is quickly overcome (**Fig. 4**), either by 'rb2' in mosaics and mixtures, or by 'rb12' in rotations and pyramids (**Fig. 5**). In mosaics, this leads to the same critical zone previously described for Experiment 2. In contrast, in mixtures and rotations, the level of control stays high for a large range of resistance efficiencies and times to expression. In pyramids, there is a good level of control only for highly efficient resistances (**Fig. 4**). For the resistant cultivar carrying the major gene (R1), disease control shows contrasting results depending on the deployment strategy. It is globally poor in mosaics and globally good in rotations. In mixtures, it is good only when the second resistant cultivar (R2) carries a strong APR gene that is expressed early. In pyramids, it is good as long as the APR has a strong efficiency. For the susceptible cultivar, a good level of disease control can be obtained if the APR (deployed in cultivar R2) has a strong efficiency and early expression, especially if pyramided with a major gene. In this situation the susceptible cultivar is invaded by both the 'wt' and the 'rb12' pathogen genotypes (**Fig. 5**).

Impact of targeted pathogenicity trait.

The results are qualitatively the same when sporulation rate and sporulation duration are targeted by the APR gene instead of the infection rate (**Figs. S8 & S9**). When resistance conferred by the APR gene increases the length of the latent period (**Fig. S7**), it is durable for a larger range of parameter values (i.e., resistance efficiency and time to expression) compared with the other target traits. However, in this situation the level of epidemiological control for the different cultivars is poor in comparison to the other target traits.

Discussion

To the best of our knowledge, adult plant resistance (APR) has never been explored in mathematical models dealing with plant resistance deployment (Rimbaud L et al., 2021), despite its presence in numerous resistant cultivars of cereals and other crops (Burdon JJ, 1987 p56; McIntosh RA et al., 1995; Boyd LA, 2005; Chen XM, 2005; Develley-Rivière M-P & E Galiana, 2007; Chen W et al., 2014). Therefore, and because APR may affect different pathogenicity traits, in a delayed and potentially incomplete manner, we used the mathematical model implemented in the R package *landsepi* (Rimbaud L et al., 2018c) to explore three parameters associated with this type of resistance: target pathogenicity trait, efficiency and time to expression. The main objective was to evaluate the impact of these parameters on resistance durability (evolutionary pathogen control) and disease limitation (epidemiological control). We designed numerical experiments to explore three scenarios: the deployment of a single APR gene in a susceptible landscape, firstly without and secondly with pathogen evolution. The third experiment assessed the deployment of an APR gene together with a

major resistance gene according to different spatiotemporal deployment strategies (**Table 1**). Although this work was motivated by rust fungi of cereal crops, the generality of the model makes our results likely applicable to other pathosystems. Adult plant resistance has also been described in viruses (whilst rather called “mature plant resistance”). For instance, a cultivar of *Nicotiana edwardsonii*, expresses a delayed monogenic resistance against *Tobacco mosaic virus*, *Tobacco necrosis virus* and *Tobacco bushy stunt virus* (Cole AB et al., 2004). Mature plant resistance has also been demonstrated in the greenhouse against *Cucumber mosaic virus* with a complete restriction of viral movement and systemic colonisation in mature bell pepper plants (Garcia-Ruiz H & JF Murphy, 2001) and against *Potato virus Y* with a restriction of tuber infection in potato (Kumar P et al., 2022).

Favouring competition offers good epidemiological control in spite of pathogen adaptation

Globally, our results show that an APR gene is never overcome when it is inefficient with respect to reducing the target pathogenicity trait or is expressed late in the cropping season (**Figs. 3AB, 4A**). This is due to the small selection pressure applied to the pathogen population, given that the wt genotype can thrive on cultivars carrying such resistance genes almost as if they were susceptible. This is in accordance with results obtained via different simulation models (Carolan K et al., 2017; Cr  t   R et al., 2020) and confirms one of the mechanisms according to which partially efficient resistance is generally predicted to be more durable than complete resistance (Lecoq H et al., 2004; Stuthman DD et al., 2007; Zhan J et al., 2015). Such phenomena have also been described for pest adaptation to chemicals, where small application doses were shown to slow down the emergence of adapted genotypes (Hobbelen PHF et al., 2014). Partial resistance with low efficiency or delayed expression, however, results in severe epidemics and a weak level of epidemiological control (**Figs. 2, 3CD, 4B**). In contrast, when resistance strongly reduces the target pathogenicity trait of the wt pathogen, particularly when this happens early in the cropping season, it has a high potential to protect the resistant cultivar (Experiment 1, **Fig. 2**), as expected in absence of pathogen evolution and already shown in demographic models (e.g., Papa  x J et al., 2014b). However, if pathogen evolution is possible, the high selection pressure leads to the rapid emergence of a rb pathogen which invades the resistant host population, resulting in both low durability and disease control (Experiment 2, **Fig. 3**). This is similar to a scenario where a single major gene (i.e., complete resistance) is deployed in the landscape and quickly overcome (Rimbaud L et al., 2018c).

There is, however, an intermediate region of the parameter space where the APR gene is broken down but still confers a good level of epidemiological protection. This occurs in presence of pathogen evolution only (i.e., in Experiment 2 but not in Experiment 1), and when resistance is delayed in the cropping season but has sufficiently high efficiency once activated. The delay in resistance expression allows the wt genotype to infect resistant hosts early in the season, more efficiently than potential rb genotypes which suffer a fitness cost while resistance is inactive. As soon as it activates, resistance is strong enough to select for rb genotypes, but many hosts are, at this time, already infected by the wt genotype. The ensuing strong competition between the wt and rb genotypes (Experiment 2, **Fig. 3 & S5**) explains the limitation on epidemic development (Keesing F et al., 2006). In this context, a resistant crop carrying an APR may conceptually be seen as a within-season rotation between a susceptible and a resistant cultivar. The size of the critical zone (i.e., the range of parameter values leading to optimal epidemiological control for the resistant cultivar) is amplified whenever such competition between pathogen genotypes is stimulated. In our results, this is the case for high fitness costs of pathogen adaptation, which increases the penalty for rb genotypes on both susceptible hosts and hosts with still inactive APR gene and reduces the performance of these genotypes relative to the wt genotype. This corroborates other modelling studies showing that high fitness costs decrease epidemic severity

(Pietravalle S et al., 2006; Djidjou-Demasse R et al., 2017; Rimbaud L et al., 2018a; Watkinson-Powell B et al., 2020). In the absence of pathogen adaptation (i.e., when there is only one pathogen genotype, Experiment 1, **Fig. 2**) or fitness cost (Experiment 2, **Fig. 3**) this effect completely disappears.

The level of spatial aggregation of the landscape impacts interactions between cultivars

A high level of spatial aggregation between cultivars in the landscape (e.g. **Fig. 2A**) isolates cultivars and the respective pathogen genotypes that infect them. In terms of epidemiological control of a susceptible cultivar, it results in severe epidemics caused by the wt pathogen genotype (**Figs. 2A, 3C**). Conversely, in a fragmented landscape (weak level of aggregation, e.g. **Fig. 2B**), the increased connectivity between different cultivars favours pathogen migration from one cultivar to another (Taylor PD et al., 1993). This reduces epidemics on the susceptible cultivar as a result of two different mechanisms which our two first experiments help disentangle. First, there is a dilution effect (Mundt CC, 2002; Rimbaud L et al., submitted), especially in the presence of a cultivar carrying a very strong resistance expressed early in the cropping season. Indeed, in this situation, spread of the wt genotype across susceptible fields is mitigated by the intervening presence of resistant hosts (Experiment 1, **Fig. 2B**). This is similar to non-host plants that act as propagule sinks and thus decrease epidemic spread on susceptible plants (Suzuki SU & A Sasaki, 2011; Papaïx J et al., 2014b). Second, competition occurs between different pathogen genotypes when the resistant cultivar has an intermediate to strong efficiency and a delayed expression (as described previously). In this case, rb genotypes emerging in resistant fields disperse to susceptible fields (Experiment 2, **Fig. 3D**). This leads to a reduction in the damage caused to the susceptible cultivar (provided that rb genotypes suffer a fitness cost compared to the wt genotype) (Watkinson-Powell B et al., 2020). The side-effect of such a protective effect of the susceptible cultivar by the resistant cultivar is a slightly reduced level of disease control on the resistant cultivar when resistance is expressed late in the cropping season because it is more exposed to wt pathogen genotypes emerging from susceptible fields. Indeed, wt genotypes are fitter than rb genotypes on the resistant cultivar as long as resistance is inactive, due to the presence of fitness costs. Spatial aggregation has been previously demonstrated to have an ambivalent effect on disease management. In fact, earlier modelling work showed that fragmented landscapes better mitigate epidemics on susceptible crops but are more prone to resistance breakdown, compared to aggregated landscapes (Papaïx J et al., 2018; Rimbaud L et al., 2018a).

Optimal efficiency and time to expression of APR genes depend on the target pathogenicity trait

A recent opinion published by Saubin M et al. (2022) states that life history traits targeted by resistance influences its durability. In fact, in the present work, the size and location of the critical zone in parameter space depends on the pathogenicity trait targeted by the APR gene. When sporulation rate or duration of the sporulation period are targeted, the critical zone is shifted towards higher resistance efficiencies and longer times to resistance expression compared to the situation where infection rate is targeted (top right of **Figs. 3CD, S3CD, S4CD**). This shift occurs probably because sporulation takes place later than infection in the pathogen infectious cycle. Therefore, more time is required for the wt pathogen genotype to generate sporulating lesions in the resistant host population before resistance expression (which will favour rb genotypes). APR genes targeting the latent period duration seem very durable, but offer poor disease control in comparison to APR genes targeting other traits (**Fig. S2**). This is because even when resistance is fully efficient (i.e., latent period is multiplied by 2), pathogen spread is still possible, which imposes soft selection pressure in favour of rb genotype but provides weak protection against the wt pathogen. This conclusion contrasts with published literature suggesting that latent period duration should be the most influent component of pathogen

aggressiveness because it determines the number of possible infection cycles on a crop (Parlevliet JE, 1979; Leonard KJ & CC Mundt, 1984; Sandoval-Islas JS et al., 2007). Nevertheless, sensitivity analyses of models simulating epidemics of wheat leaf rust (Kulkarni RN et al., 1982) and potato late blight (Van Oijen M, 1992) have shown that latent period duration was equally or even less influential on disease spread and severity than other pathogenicity traits. These contrasted results highlight the crucial importance of the width of parameter variation ranges in numerical experiments. In our work, the range of variation for resistance efficiency was based on available data for rust fungi. Analysis of the minimal and maximal possible values of the pathogenicity traits measured on different cultivars of cereal crops (**Table S1**) showed that these traits may vary from about 0% to -100% (0% to +100% for latent period duration) relative to the most susceptible cultivar (except sporulation duration, for which there is little data). We thus allowed resistance efficiency to vary from 0 to 100% for all pathogenicity traits.

Major resistance genes and APR genes can be combined at landscape scale

The deployment of a single major resistance gene in a landscape results in rapid breakdown by the corresponding *rb1* pathogen and severe epidemics on both susceptible and resistant cultivars (the bottom line of heatmaps in **Fig. 4** shows the situation where the APR is absent, its efficiency being 0%). Combining a major gene with an APR gene in the landscape generally does not prevent the major gene from being overcome, however it may have interesting synergies in terms of epidemiological control depending on the deployment strategy (**Fig. 4**). As discussed earlier, one of the greatest benefits of APR genes is the limitation of epidemics due to competition between pathogen genotypes. Therefore, the presence of different sources of resistance in the landscape, should they be overcome, increases the number of pathogen genotypes present and thus the number of competitors. Globally, this decreases epidemic damage on all cultivars (Mikaberidze A et al., 2015).

More specifically, when a cultivar carrying a major gene is planted in mixtures (i.e., in the same field) with a cultivar carrying an APR gene, the first cultivar benefits from a dilution effect (since only *rb1* genotypes can infect it) conferred by the presence of the second one, which itself benefits from strong competition between the *wt*, *rb1* and *rb2* genotypes. While to some extent this should also occur in mosaics (i.e., different cultivars segregated in different fields), our results do not show such synergies for the mosaic strategy. This is probably because of the model assumption that the pathogen was initially present in all susceptible fields of the landscape, added to the fact that pathogen dispersal is mostly at the intra-field scale in our parameterisation (**Table 1**). The impact of landscape heterogeneity on epidemic spread via competition and dilution effects might be stronger for pathogens with different life histories (Mundt CC, 2002). Here, the best epidemiological control is obtained when crop cultivars are mixed at the finest spatial grain. Indeed, optimal disease control requires that the spatial scale of disease management matches the scale of pathogen dispersal (Gilligan CA, 2008). When the two resistant cultivars are rotated over time (rotation strategy), pathogen genotypes are confronted by an alternation of hard selection towards the *rb1* genotype (when the cultivar carrying the major gene is cultivated) and hard or soft selection towards the *rb2* genotype (when the cultivar carrying the APR gene is cultivated). If the APR is not too strong or has a delayed expression, selection towards *rb2* is soft, which allows competition between *wt*, *rb1* and *rb2* genotypes and reduces epidemics. Otherwise, selection is strong and the genotype that performs best in the system is the double mutant *rb12* (generalist genotype able to infect all cultivars). However, this genotype is penalised by severe fitness costs (**Table 3**), which reduces epidemic damage as well. This is in line with a previous modelling study comparing mosaics, mixtures, rotation and pyramids of major resistance genes: rotation had the best epidemiological outcome once all resistances had been

overcome (i.e., in the presence of rb genotypes) (Rimbaud L et al., 2018a). Finally, if the major gene and the APR gene are pyramided in the same cultivar and the efficiency of the APR gene is strong enough, the delayed action of the APR gene triggers competition between the single mutant rb1, selected for as long as the APR is inactive, and the double mutant rb12, selected for as soon as the APR expresses. This competition reduces epidemic damage on the pyramid cultivar. However, the presence of the APR gene does not prevent the major gene from being overcome, unless it is expressed very early in the cropping season. This is in agreement with previous modelling results: durability of a major gene was greater when pyramided with a quantitative resistance (expressed from the beginning of the cropping season), but only if the latter exhibited strong efficiency (Rimbaud L et al., 2018c).

General conclusions, limits and perspectives

There are several nonexclusive arguments for why APR genes are thought to be more durable than traditional major genes. Firstly, it could be inherent to the molecular mechanism of APR genes, that may be more difficult for the pathogen to overcome than classical NLR proteins frequently encoded by major genes (Oliva R & IL Quibod, 2017; Mundt CC, 2018). As described in the Introduction, the mechanisms of a few APR genes have been elucidated, such as Lr67, Lr34 and Yr36, which encode for a sugar transporter (Moore JW et al., 2015), an ATP-binding cassette transporter (Krattinger SG et al., 2009), and a detoxification protein (Fu D et al., 2009), respectively. Secondly, it could result from the fact that APR genes are rarely alone in a susceptible host genetic background but may be shielded by major genes. Finally, it could be due to the smaller selection pressure applied by APR genes on pathogens (since they allow some infection by wt pathogens by being only partially efficient and delayed in the season) (Mundt CC, 2018).

In the absence of relevant quantitative data concerning the first hypothesis, our parameterisation of the model gives the same mutation probability to overcome major genes and APR genes. Hence, the present study explores the latter two hypotheses. The possibility for APR genes to be shielded by major genes has been tested in Experiment 3 while the effect of selection pressure is highlighted by the difference between Experiments 1 and 2. The mutation probability to overcome the resistances was set at a high value, which could explain why, in our simulations, the combination of an APR gene with a major gene in a pyramided cultivar did not affect the durability of the APR gene in comparison to a cultivar that carried the APR gene only. Future work could investigate the potential of such pyramids with a lower mutation probability. On the other hand, our work emphasizes how shifts in selection pressure influence resistance durability. Indeed, APR genes were found to be very durable when they have a small efficiency and late expression. It may explain why some APR genes like Yr18, which has a small to moderate efficiency against stripe rust (Elahinia SA & JP Tewari, 2005; Qamar M et al., 2012) have shown high durability in the field (Krattinger SG et al., 2009). The efficiency of other APR genes like Lr12, Lr13, Lr22, Lr34, Lr35 and Lr37 have been measured between 80% and 90% against leaf rust (Burdon JJ, 1987 p56; McIntosh RA et al., 1995; Smale M et al., 1998). With such high efficiency, our simulations predicts that these genes could be quickly overcome. Nevertheless, depending on the time to resistance expression and the target pathogenicity trait, even if these genes were broken down, the resulting harsh competition between the different pathogen genotypes has the potential to provide some disease limitation, especially when deployed together with major resistance genes in mixture or rotation strategies. However, this conclusion strongly depends on the presence of fitness costs of pathogen adaptation to resistance. Furthermore, our results must be nuanced by the fact that we assumed that rb pathogens were penalised by a fitness cost on inactive APR genes, exactly as if the associated cultivars were susceptible. Experiments could be carried out in controlled conditions to test this hypothesis. We also assumed that APR genes switch suddenly from

being inactive to active, whereas some rare available data rather indicate a gradual expression of APR genes (Ma H & RP Singh, 1996). Finally, while in our simulations, APR genes could target only one pathogenicity trait at a time, in the real world pathogenicity traits often vary in association (Parlevliet JE, 1979; Sache I & C de Vallavieille-Pope, 1995; Leclerc M et al., 2019). For example, Lr16-Lr18 targets latent period duration as well as sporulation rate and duration (Tomerlin JR et al., 1983) and Lr34-Yr18 affects both infection rate and latent period (Qamar M et al., 2012). Regardless, our study represents a first attempt to numerically explore evolutionary and epidemiological outcomes of the deployment of adult plant resistance for the management of plant diseases.

Acknowledgements

The authors thank Marta Zaffaroni and Jean-Loup Gaussen for their contribution to improve the R package *landsepi*, Loïc Houde for computing assistance and Jeremy Burdon for stimulating discussions.

Fundings

This work benefited from GRDC grant CSP00192, ANR project “Archiv” (2019–2023, grant n°ANR-18-CE32-0004-01), AFB Ecophyto II-Leviers Territoriaux Project “Médée” (2020–2022), and the CSIRO/INRAE linkage program.

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article. Benoît Moury is recommender for PCI Evol. Biol.

Data, script and code availability

The model is available in the open-access R package *landsepi* (Rimbaud L et al., 2018b; webpage: <https://csiro-inra.pages.biosp.inrae.fr/landsepi/>). Simulations were performed on the BioSP computational cluster from INRAE (<https://biosp-cluster.mathnum.inrae.fr/>). Simulation results are available in supplementary information.

629

References

- 630 Azzimonti G, Lannou C, Sache I, Goyeau H (2013) Components of quantitative resistance to leaf rust
631 in wheat cultivars: diversity, variability and specificity. *Plant Pathology*, **62**, 970-981.
632 <https://doi.org/10.1111/ppa.12029>
- 633 Barrett LG, Heil M (2012) Unifying concepts and mechanisms in the specificity of plant–enemy
634 interactions. *Trends in Plant Science*, **17**, 282-292.
635 <https://doi.org/10.1016/j.tplants.2012.02.009>
- 636 Boyd LA (2005) Can robigus defeat an old enemy? - Yellow rust of wheat. *Journal of Agricultural*
637 *Science*, **143**, 233-243. <https://doi.org/10.1017/S0021859605005095>
- 638 Broers LHM (1997) Components of quantitative resistance to yellow rust in ten spring bread wheat
639 cultivars and their relations with field assessments. *Euphytica*, **96**, 215-223.
640 <https://doi.org/10.1023/a:1002916110347>
- 641 Broers LHM, Cuesta Subias X, López Atilano RM (1996) Field assessment of quantitative resistance to
642 yellow rust in ten spring bread wheat cultivars. *Euphytica*, **90**, 9-16.
643 <https://doi.org/10.1007/bf00025154>
- 644 Burdon JJ (1987) *Diseases and Plant Population Biology*, p56. Cambridge University Press, Cambridge.
- 645 Burdon JJ, Barrett LG, Rebetzke G, Thrall PH (2014) Guiding deployment of resistance in cereals using
646 evolutionary principles. *Evolutionary Applications*, **7**, 609-624.
647 <https://doi.org/10.1111/eva.12175>
- 648 Burdon JJ, Thrall PH (2014) What have we learned from studies of wild plant-pathogen
649 associations?—the dynamic interplay of time, space and life-history. *European Journal of*
650 *Plant Pathology*, **138**, 417-429. <https://doi.org/10.1007/s10658-013-0265-9>
- 651 Burdon JJ, Zhan J, Barrett LG, Papaix J, Thrall PH (2016) Addressing the challenges of pathogen
652 evolution on the world's arable crops. *Phytopathology*, **106**, 1117-1127.
653 <https://doi.org/10.1094/PHYTO-01-16-0036-FI>
- 654 Carolan K, Helps J, van den Berg F, Bain R, Paveley N, van den Bosch F (2017) Extending the durability
655 of cultivar resistance by limiting epidemic growth rates. *Proceedings of the Royal Society B:*
656 *Biological Sciences*, **284**, 20170828. <https://doi.org/10.1098/rspb.2017.0828>
- 657 Chen W, Wellings C, Chen X, Kang Z, Liu T (2014) Wheat stripe (yellow) rust caused by *Puccinia*
658 *striiformis* f. sp. *tritici*. *Molecular Plant Pathology*, **15**, 433-446.
659 <https://doi.org/10.1111/mpp.12116>
- 660 Chen XM (2005) Epidemiology and control of stripe rust *Puccinia striiformis* f. sp. *tritici* on wheat.
661 *Canadian Journal of Plant Pathology*, **27**, 314-337.
662 <https://doi.org/10.1080/07060660509507230>
- 663 Cole AB, Király L, Lane LC, Wiggins BE, Ross K, Schoelz JE (2004) Temporal expression of PR-1 and
664 enhanced mature plant resistance to virus infection is controlled by a single dominant gene
665 in a new *Nicotiana* hybrid. *Molecular Plant-Microbe Interactions*, **17**, 976-985.
666 <https://doi.org/10.1094/MPMI.2004.17.9.976>
- 667 Crété R, Pires RN, Barbetti MJ, Renton M (2020) Rotating and stacking genes can improve crop
668 resistance durability while potentially selecting highly virulent pathogen strains. *Scientific*
669 *Reports*, **10**, 19752. <https://doi.org/10.1038/s41598-020-76788-7>
- 670 Cromey MG (1992) Adult plant resistance to stripe rust (*Puccinia striiformis*) in some New Zealand
671 wheat cultivars. *New Zealand Journal of Crop and Horticultural Science*, **20**, 413-419.
672 <https://doi.org/10.1080/01140671.1992.10418058>
- 673 de Ronde D, Butterbach P, Kormelink R (2014) Dominant resistance against plant viruses. *Frontiers in*
674 *Plant Science*, **5**, 307. <https://doi.org/10.3389/fpls.2014.00307>
- 675 Denissen CJM (1993) Components of adult plant resistance to leaf rust in wheat. *Euphytica*, **70**, 131-
676 140. <https://doi.org/10.1007/bf00029650>

- 677 Develey-Rivière M-P, Galiana E (2007) Resistance to pathogens and host developmental stage: a
678 multifaceted relationship within the plant kingdom. *New Phytologist*, **175**, 405-416.
679 <https://doi.org/10.1111/j.1469-8137.2007.02130.x>
- 680 Djidjou-Demasse R, Moury B, Fabre F (2017) Mosaics often outperform pyramids: insights from a
681 model comparing strategies for the deployment of plant resistance genes against viruses in
682 agricultural landscapes. *New Phytologist*, **216**, 239-253. <https://doi.org/10.1111/nph.14701>
- 683 Elahinia SA, Tewari JP (2005) Assessment of two different sources of durable resistance and
684 susceptible cultivar of wheat to stripe rust (*Puccinia striiformis* f. sp. *tritici*). *Caspian Journal*
685 *of Environmental Sciences*, **3**, 117-122.
686 http://cjes.guilan.ac.ir/article_948_5e0230f27bfd99b837bd4d79b5fd8591.pdf
- 687 Flor HH (1955) Host-parasite interaction in flax rust - Its genetics and other implications.
688 *Phytopathology*, **45**, 680-685. [https://www.webofscience.com/wos/woscc/full-](https://www.webofscience.com/wos/woscc/full-record/WOS:A1955WJ02800010)
689 [record/WOS:A1955WJ02800010](https://www.webofscience.com/wos/woscc/full-record/WOS:A1955WJ02800010)
- 690 Fu D, Uauy C, Distelfeld A, Blechl A, Epstein L, Chen X, Sela H, Fahima T, Dubcovsky J (2009) A kinase-
691 START gene confers temperature-dependent resistance to wheat stripe rust. *Science*, **323**,
692 1357-1360. <https://doi.org/10.1126/science.1166289>
- 693 Gallois J-L, Moury B, German-Retana S (2018) Role of the genetic background in resistance to plant
694 viruses. *International Journal of Molecular Sciences*, **19**, 2856.
695 <https://doi.org/10.3390/ijms19102856>
- 696 García-Arenal F, McDonald BA (2003) An analysis of the durability of resistance to plant viruses.
697 *Phytopathology*, **93**, 941-952. <https://doi.org/10.1094/PHYTO.2003.93.8.941>
- 698 Garcia-Ruiz H, Murphy JF (2001) Age-related resistance in bell pepper to *Cucumber mosaic virus*.
699 *Annals of Applied Biology*, **139**, 307-317. [https://doi.org/10.1111/j.1744-](https://doi.org/10.1111/j.1744-7348.2001.tb00144.x)
700 [7348.2001.tb00144.x](https://doi.org/10.1111/j.1744-7348.2001.tb00144.x)
- 701 Gilligan CA (2008) Sustainable agriculture and plant diseases: an epidemiological perspective.
702 *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **363**, 741-
703 759. <https://doi.org/10.1098/rstb.2007.2181>
- 704 Hobbelen PHF, Paveley ND, van den Bosch F (2014) The emergence of resistance to fungicides. *PLOS*
705 *ONE*, **9**, e91910. <https://doi.org/10.1371/journal.pone.0091910>
- 706 Johnson R (1983) Genetic background of durable resistance. In: *Durable Resistance in Crops* eds
707 Lamberti F, Waller JM, & Graaff NA), pp. 5-26. Springer New York, Boston, MA.
708 https://doi.org/10.1007/978-1-4615-9305-8_2
- 709 Johnson R (1984) A critical analysis of durable resistance. *Annual Review of Phytopathology*, **22**, 309-
710 330. <https://doi.org/10.1146/annurev.py.22.090184.001521>
- 711 Keesing F, Holt RD, Ostfeld RS (2006) Effects of species diversity on disease risk. *Ecology Letters*, **9**,
712 485-498. <https://doi.org/10.1111/j.1461-0248.2006.00885.x>
- 713 Krattinger SG, Keller B (2016) Molecular genetics and evolution of disease resistance in cereals. *New*
714 *Phytologist*, **212**, 320-332. <https://doi.org/10.1111/nph.14097>
- 715 Krattinger SG, Lagudah ES, Spielmeier W, Singh RP, Huerta-Espino J, McFadden H, Bossolini E, Selter
716 LL, Keller B (2009) A putative ABC transporter confers durable resistance to multiple fungal
717 pathogens in wheat. *Science*, **323**, 1360-1363. <https://doi.org/10.1126/science.1166453>
- 718 Kulkarni RN, Chopra VL, Singh D (1982) Relative importance of components affecting the leaf rust
719 progress curve in wheat. *Theoretical and Applied Genetics*, **62**, 205-207.
720 <https://doi.org/10.1007/bf00276238>
- 721 Kumar P, Cowan GH, Squires JN, Hackett CA, Tobin AK, Torrance L, Roberts AG (2022) Phloem
722 connectivity and transport are not involved in mature plant resistance (MPR) to Potato Virus
723 Y in different potato cultivars, and MPR does not protect tubers from recombinant strains of
724 the virus. *Journal of Plant Physiology*, **275**, 153729.
725 <https://doi.org/10.1016/j.jplph.2022.153729>

726 Lannou C (2012) Variation and selection of quantitative traits in plant pathogens. *Annual Review of*
727 *Phytopathology*, **50**, 319-338. <https://doi.org/10.1146/annurev-phyto-081211-173031>

728 Leclerc M, Clément JAJ, Andrivon D, Hamelin FM (2019) Assessing the effects of quantitative host
729 resistance on the life-history traits of sporulating parasites with growing lesions. *Proceedings*
730 *of the Royal Society B: Biological Sciences*, **286**, 20191244.
731 <https://doi.org/10.1098/rspb.2019.1244>

732 Lecoq H, Moury B, Desbiez C, Palloix A, Pitrat M (2004) Durable virus resistance in plants through
733 conventional approaches: a challenge. *Virus Research*, **100**, 31-39.
734 <https://doi.org/10.1016/j.virusres.2003.12.012>

735 Lehman JS, Shaner G (1998) Genetic variation in latent period among isolates of *Puccinia recondita*
736 f.sp. *tritici* on partially resistant wheat cultivars. *Phytopathology*, **86**, 633-641.
737 <https://doi.org/10.1094/Phyto-86-633>

738 Leonard KJ, Mundt CC (1984) Methods for estimating epidemiological effects of quantitative
739 resistance to plant diseases. *Theoretical and Applied Genetics*, **67**, 219-230.
740 <https://doi.org/10.1007/bf00317041>

741 Ma H, Singh RP (1996) Expression of adult plant resistance to stripe rust at different growth stages of
742 wheat. *Plant Disease*, **80**, 375-379. <https://doi.org/10.1094/PD-80-0375>

743 McDonald BA, Linde C (2002) Pathogen population genetics, evolutionary potential, and durable
744 resistance. *Annual Review of Phytopathology*, **40**, 349-379.
745 <https://doi.org/10.1146/annurev.phyto.40.120501.101443>

746 McIntosh RA, Wellings CR, Park RF (1995) *Wheat Rusts. An Atlas of Resistance Genes*. CSIRO
747 Publications, East Melbourne, Victoria, Australia.
748 <https://ebooks.publish.csiro.au/content/wheat-rusts>

749 Mikaberidze A, McDonald BA, Bonhoeffer S (2015) Developing smarter host mixtures to control plant
750 disease. *Plant Pathology*, **64**, 996-1004. <https://doi.org/10.1111/ppa.12321>

751 Moore JW, Herrera-Foessel S, Lan C, Schnippenkoetter W, Ayliffe M, Huerta-Espino J, Lillemo M,
752 Viccars L, Milne R, Periyannan S, Kong X, Spielmeyer W, Talbot M, Bariana H, Patrick JW,
753 Dodds P, Singh R, Lagudah E (2015) A recently evolved hexose transporter variant confers
754 resistance to multiple pathogens in wheat. *Nature Genetics*, **47**, 1494-1498.
755 <https://doi.org/10.1038/ng.3439>

756 Mortensen K, Green GJ (1978) Assessment of receptivity and urediospore production as components
757 of wheat stem rust resistance. *Canadian Journal of Botany*, **56**, 1827-1839.
758 <https://doi.org/10.1139/b78-221>

759 Moury B, Fabre F, Montarry J, Janzac B, Ayme V, Palloix A (2010) L'adaptation des virus de plantes
760 aux résistances variétales. *Virologie*, **14**, 227-239. <https://doi.org/10.1684/vir.2010.0311>

761 Mundt CC (2002) Use of multiline cultivars and cultivar mixtures for disease management. *Annual*
762 *Review of Phytopathology*, **40**, 381-410.
763 <https://doi.org/10.1146/annurev.phyto.40.011402.113723>

764 Mundt CC (2018) Pyramiding for resistance durability: Theory and practice. *Phytopathology*, **108**,
765 792-802. <https://doi.org/10.1094/PHYTO-12-17-0426-RVW>

766 Niks RE, Qi X, Marcel TC (2015) Quantitative resistance to biotrophic filamentous plant pathogens:
767 Concepts, misconceptions, and mechanisms. *Annual Review of Phytopathology*, **53**, 445-470.
768 <https://doi.org/10.1146/annurev-phyto-080614-115928>

769 Oliva R, Quibod IL (2017) Immunity and starvation: new opportunities to elevate disease resistance in
770 crops. *Current Opinion in Plant Biology*, **38**, 84-91. <https://doi.org/10.1016/j.pbi.2017.04.020>

771 Papaïx J, Adamczyk-Chauvat K, Bouvier A, Kiêu K, S. T, Lannou C, Monod H (2014a) Pathogen
772 population dynamics in agricultural landscapes: The *Ddal* modelling framework. *Infection,*
773 *Genetics and Evolution*, **27**, 509-520. <https://doi.org/10.1016/j.meegid.2014.01.022>

774 Papaix J, Rimbaud L, Burdon JJ, Zhan J, Thrall PH (2018) Differential impact of landscape-scale
775 strategies for crop cultivar deployment on disease dynamics, resistance durability and long-
776 term evolutionary control. *Evolutionary Applications*, **11**, 705-717.
777 <https://doi.org/10.1111/eva.12570>

778 Papaix J, Touzeau S, Monod H, Lannou C (2014b) Can epidemic control be achieved by altering
779 landscape connectivity in agricultural systems? *Ecological Modelling*, **284**, 35-47.
780 <https://doi.org/10.1016/j.ecolmodel.2014.04.014>

781 Pariaud B, Robert C, Goyeau H, Lannou C (2009) Aggressiveness components and adaptation to a
782 host cultivar in wheat leaf rust. *Phytopathology*, **99**, 869-878.
783 <https://doi.org/10.1094/PHYTO-99-7-0869>

784 Park RF (2008) Breeding cereals for rust resistance in Australia. *Plant Pathology*, **57**, 591-602.
785 <https://doi.org/10.1111/j.1365-3059.2008.01836.x>

786 Park RF, Rees RG (1989) Expression of adult plant resistance and its effect on the development of
787 *Puccinia striiformis* f.sp. *tritici* in some Australian wheat cultivars. *Plant Pathology*, **38**, 200-
788 208. <https://doi.org/10.1111/j.1365-3059.1989.tb02134.x>

789 Parlevliet JE (1979) Components of resistance that reduce the rate of epidemic development. *Annual*
790 *Review of Phytopathology*, **17**, 203-222.
791 <https://doi.org/10.1146/annurev.py.17.090179.001223>

792 Parlevliet JE (2002) Durability of resistance against fungal, bacterial and viral pathogens; present
793 situation. *Euphytica*, **124**, 147-156. <https://doi.org/10.1023/a:1015601731446>

794 Pietravalle S, Lemarié S, van den Bosch F (2006) Durability of resistance and cost of virulence.
795 *European Journal of Plant Pathology*, **114**, 107-116. [https://doi.org/10.1007/s10658-005-](https://doi.org/10.1007/s10658-005-3479-7)
796 [3479-7](https://doi.org/10.1007/s10658-005-3479-7)

797 Pilet-Nayel M-L, Moury B, Caffier V, Montarry J, Kerlan M-C, Fournet S, Durel C-E, Delourme R (2017)
798 Quantitative resistance to plant pathogens in pyramiding strategies for durable crop
799 protection. *Frontiers in Plant Science*, **8**, 1838. <https://doi.org/10.3389/fpls.2017.01838>

800 Qamar M, Gardezi DA, Iqbal M (2012) Determination of rust resistance gene complex Lr34/Yr18 in
801 spring wheat and its effect on components of partial resistance. *Journal of Phytopathology*,
802 **160**, 628-636. <https://doi.org/10.1111/j.1439-0434.2012.01957.x>

803 Quan W, Hou G, Chen J, Du Z, Lin F, Guo Y, Liu S, Zhang Z (2013) Mapping of QTL lengthening the
804 latent period of *Puccinia striiformis* in winter wheat at the tillering growth stage. *European*
805 *Journal of Plant Pathology*, **136**, 715-727. <https://doi.org/10.1007/s10658-013-0201-z>

806 Richardson KL, Vales MI, Kling JG, Mundt CC, Hayes PM (2006) Pyramiding and dissecting disease
807 resistance QTL to barley stripe rust. *Theoretical and Applied Genetics*, **113**, 485-495.
808 <https://doi.org/10.1007/s00122-006-0314-2>

809 Rimbaud L, Fabre F, Papaix J, Moury B, Lannou C, Barrett LG, Thrall PH (2021) Models of plant
810 resistance deployment. *Annual Review of Phytopathology*, **59**, 125-152.
811 <https://doi.org/10.1146/annurev-phyto-020620-122134>

812 Rimbaud L, Fabre F, Zamberletti P, Papaix J (submitted) Revisiting dilution and barrier to better
813 disentangle their effects on epidemics in crop mixtures.

814 Rimbaud L, Papaix J, Barrett LG, Burdon JJ, Thrall PH (2018a) Mosaics, mixtures, rotations or
815 pyramiding: What is the optimal strategy to deploy major gene resistance? *Evolutionary*
816 *Applications*, **11**, 1791-1810. <https://doi.org/10.1111/eva.12681>

817 Rimbaud L, Papaix J, Rey J-F (2018b) landsepi: Landscape Epidemiology and Evolution. *R package*
818 *version 1.1.1* <https://cran.r-project.org/package=landsepi>

819 Rimbaud L, Papaix J, Rey J-F, Barrett LG, Thrall PH (2018c) Assessing the durability and efficiency of
820 landscape-based strategies to deploy plant resistance to pathogens. *Plos Computational*
821 *Biology*, **14**, e1006067. <https://doi.org/10.1371/journal.pcbi.1006067>

822 Rimé D, Robert C, Goyeau H, Lannou C (2005) Effect of host genotype on leaf rust (*Puccinia triticina*)
823 lesion development and urediniospore production in wheat seedlings. *Plant Pathology*, **54**,
824 287-298. <https://doi.org/10.1111/j.1365-3059.2005.01174.x>

825 Sache I, de Vallavieille-Pope C (1995) Classification of airborne plant pathogens based on sporulation
826 and infection characteristics. *Canadian Journal of Botany*, **73**, 1186-1195.
827 <https://doi.org/10.1139/b95-128>

828 Sandoval-Islas JS, Broers LHM, Mora-Aguilera G, Parlevliet JE, Osada-Kawasoe S, Vivar HE (2007)
829 Quantitative resistance and its components in 16 barley cultivars to yellow rust, *Puccinia*
830 *striiformis* f. sp. *hordei*. *Euphytica*, **153**, 295-308. <https://doi.org/10.1007/s10681-006-9236-y>

831 Saubin M, Louet C, Bousset L, Fabre F, Frey P, Fudal I, Grogard F, Hamelin F, Mailleret L, Stoeckel S,
832 Touzeau S, Petre B, Halkett F (2022) Improving sustainable crop protection using population
833 genetics concepts. *Molecular Ecology*, in press. <https://doi.org/10.1111/mec.16634>

834 Smale M, Singh RP, Sayre K, Pingali P, Rajaram S, Dubin HJ (1998) Estimating the economic impact of
835 breeding nonspecific resistance to leaf rust in modern bread wheats. *Plant Disease*, **82**, 1055-
836 1061. <https://doi.org/10.1094/PDIS.1998.82.9.1055>

837 Sørensen CK, Hovmøller MS, Leconte M, Dedryver F, de Vallavieille-Pope C (2014) New races of
838 *Puccinia striiformis* found in Europe reveal race specificity of long-term effective adult plant
839 resistance in wheat. *Phytopathology*, **104**, 1042-1051. <https://doi.org/10.1094/PHYTO-12-13-0337-R>

841 Stuthman DD, Leonard KJ, Miller-Garvin J (2007) Breeding crops for durable resistance to disease.
842 *Advances in Agronomy*, **95**, 319-367. [https://doi.org/10.1016/S0065-2113\(07\)95004-X](https://doi.org/10.1016/S0065-2113(07)95004-X)

843 Suzuki SU, Sasaki A (2011) How does the resistance threshold in spatially explicit epidemic dynamics
844 depend on the basic reproductive ratio and spatial correlation of crop genotypes? *Journal of*
845 *Theoretical Biology*, **276**, 117-125. <https://doi.org/10.1016/j.jtbi.2011.02.002>

846 Taylor PD, Fahrig L, Henein K, Merriam G (1993) Connectivity is a vital element of landscape
847 structure. *Oikos*, **68**, 571-573. <https://doi.org/10.2307/3544927>

848 Tomerlin JR, Eversmeyer MG, Kramer CL, Browder LE (1983) Temperature and host effects on latent
849 and infectious periods and on urediniospore production of *Puccinia recondita* f. sp. *tritici*.
850 *Phytopathology*, **73**, 414-419. <https://doi.org/10.1094/Phyto-73-414>

851 van den Bosch F, Gilligan CA (2003) Measures of durability of resistance. *Phytopathology*, **93**, 616-
852 625. <https://doi.org/10.1094/PHYTO.2003.93.5.616>

853 Van Oijen M (1992) Selection and use of a mathematical model to evaluate components of resistance
854 to *Phytophthora infestans* in potato. *Netherlands Journal of Plant Pathology*, **98**, 192-202.
855 <https://doi.org/10.1007/bf01974382>

856 Watkinson-Powell B, Gilligan CA, Cuniffe NJ (2020) When does spatial diversification usefully
857 maximize the durability of crop disease resistance? *Phytopathology*, **110**, 1808-1820.
858 <https://doi.org/10.1094/phyto-07-19-0261-r>

859 Zhan J, Thrall PH, Papaix J, Xie L, Burdon JJ (2015) Playing on a pathogen's weakness: using evolution
860 to guide sustainable plant disease control strategies. *Annual Review of Phytopathology*, **53**,
861 19-43. <https://doi.org/10.1146/annurev-phyto-080614-120040>

Supplementary information

Figure S1. Heatmaps of the optimal pathogenicity trait targeted by an APR gene.

Figure S2. Heatmaps of the levels of evolutionary and epidemiological control, and average genotype frequencies in Experiment 2 when the target pathogenicity trait is the latent period duration.

Figure S3. Heatmaps of the levels of evolutionary and epidemiological control, and average genotype frequencies in Experiment 2 when the target pathogenicity trait is the sporulation rate.

Figure S4. Heatmaps of the levels of evolutionary and epidemiological control, and average genotype frequencies in Experiment 2 when the target pathogenicity trait is the sporulation duration.

Figure S5. Epidemiological outcome and dynamics of pathogen genotype frequencies in three examples of simulations.

Figure S6. Example of simulated fragmented landscapes used in Experiment 3.


Figure S7. Heatmaps of the levels of evolutionary and epidemiological control in Experiment 3 when the target pathogenicity trait is the latent period duration.

Figure S8. Heatmaps of the levels of evolutionary and epidemiological control in Experiment 3 when the target pathogenicity trait is the sporulation rate.

Figure S9. Heatmaps of the levels of evolutionary and epidemiological control in Experiment 3 when the target pathogenicity trait is the sporulation duration.


Table S1. Observed ranges of infection rate, latent period duration, sporulation rate and sporulation duration for rust fungi.

Raw data. Dataset of simulation results used in this study.




882

883 **Figure S1.** Heatmaps indicating the optimal pathogenicity trait targeted by an APR gene with respect
884 to the level of epidemiological control (i.e., disease limitation, measured by the Green Leaf Area, 'GLA')
885 on the resistant cultivar in the absence of pathogen evolution for different levels of resistance
886 efficiency (vertical axis) and time to resistance expression (horizontal axis), for strong (top) or weak
887 (bottom) levels of spatial aggregation.




88~

Figure S2. Heatmaps of the levels of evolutionary control (resistance durability as measured by the number of years before the emergence of the resistance-breaking ('rb') pathogen genotype, panels A and B), epidemiological control (i.e. disease limitation, measured by the Green Leaf Area ('GLA') on the susceptible ('S') and the resistant ('R') cultivars, panels C and D) and average frequency of the rb pathogen (panels E and F) for different levels of resistance efficiency (vertical axis), time to resistance expression (horizontal axis) and fitness cost of pathogen adaptation (columns), for strong (panels A, C, E) or weak (B, D, F) levels of spatial aggregation. The target pathogenicity trait is the latent period duration.




85

898 **Figure S3.** Heatmaps of the levels of evolutionary control (resistance durability as measured by the
899 number of years before the emergence of the resistance-breaking ('rb') pathogen genotype, panels A
900 and B), epidemiological control (i.e. disease limitation, measured by the Green Leaf Area ('GLA') on the
901 susceptible ('S') and the resistant ('R') cultivars, panels C and D) and average frequency of the rb
902 pathogen (panels E and F) for different levels of resistance efficiency (vertical axis), time to resistance
903 expression (horizontal axis) and fitness cost of pathogen adaptation (columns), for strong (panels A, C,
904 E) or weak (B, D, F) levels of spatial aggregation. The target pathogenicity trait is the sporulation rate.




9

Figure S4. Heatmaps of the levels of evolutionary control (resistance durability as measured by the number of years before the emergence of the resistance-breaking (‘rb’) pathogen genotype, panels A and B), epidemiological control (i.e. disease limitation, measured by the Green Leaf Area (‘GLA’) on the susceptible (‘S’) and the resistant (‘R’) cultivars, panels C and D) and average frequency of the rb pathogen (panels E and F) for different levels of resistance efficiency (vertical axis), time to resistance expression (horizontal axis) and fitness cost of pathogen adaptation (columns), for strong (panels A, C, E) or weak (B, D, F) levels of spatial aggregation. The target pathogenicity trait is the sporulation duration.



914

915 **Figure S5.** Epidemiological outcome (represented by the relative Green Leaf Area, top line) and dynamics of pathogen genotype frequencies (bottom line, 'wt' refers
 916 to the wild-type and 'rb' to the resistance-breaking pathogen genotype) in three examples of simulations where a single APR is deployed in a susceptible landscape
 917 with low level of spatial aggregation. Situations 1, 2 and 3 are pointed in Figure 3. The pathogenicity trait targeted by resistance is the infection rate and the fitness
 918 cost of adaptation is $\theta=0.50$.



919

920 **Figure S6.** Example of simulated fragmented landscapes used in Experiment 3 (APR + MG). For all
 921 deployment strategies, 1/3 of the landscape was composed of the susceptible cultivar. The remaining
 922 2/3 were occupied either by: A) a single cultivar carrying the two genes (pyramid strategy); B) a mixture
 923 (in every field) of two resistant cultivars in balanced proportions (each cultivar carrying one of the two
 924 genes); C) a rotation of these two resistant cultivars (every year); or D) a mosaic of the two resistant
 925 cultivars in balanced proportions (every cultivar representing 1/3 of the landscape area).




Figure S7. Heatmaps showing the levels of A) evolutionary control (resistance durability, measured by the number of years before the emergence of resistance-breaking genotypes) and B) epidemiological control (i.e., disease limitation, measured by the Green Leaf Area, 'GLA') on a susceptible cultivar 'S', a resistant cultivar 'R1' carrying a completely efficient major gene ('MG') and a resistant cultivar 'R2' carrying an APR gene, for different levels of APR efficiency (vertical axis), time to APR expression (horizontal axis) and deployment strategies (columns; note that for pyramiding, R1 and R2 refer to the same cultivar). The target pathogenicity trait of the APR gene is the latent period duration, the level of spatial aggregation is low, and the fitness cost is 0.50.




Figure S8. Heatmaps showing the levels of A) evolutionary control (resistance durability, measured by the number of years before the emergence of resistance-breaking genotypes) and B) epidemiological control (i.e., disease limitation, measured by the Green Leaf Area, 'GLA') on a susceptible cultivar 'S', a resistant cultivar 'R1' carrying a completely efficient major gene ('MG') and a resistant cultivar 'R2' carrying an APR gene, for different levels of APR efficiency (vertical axis), time to APR expression (horizontal axis) and deployment strategies (columns; note that for pyramiding, R1 and R2 refer to the same cultivar). The target pathogenicity trait of the APR gene is the sporulation rate, the level of spatial aggregation is low, and the fitness cost is 0.50.




Figure S9. Heatmaps showing the levels of A) evolutionary control (resistance durability, measured by the number of years before the emergence of resistance-breaking genotypes) and B) epidemiological control (i.e., disease limitation, measured by the Green Leaf Area, 'GLA') on a susceptible cultivar 'S', a resistant cultivar 'R1' carrying a completely efficient major gene ('MG') and a resistant cultivar 'R2' carrying an APR gene, for different levels of APR efficiency (vertical axis), time to APR expression (horizontal axis) and deployment strategies (columns; note that for pyramiding, R1 and R2 refer to the same cultivar). The target pathogenicity trait of the APR gene is the sporulation duration, the level of spatial aggregation is low, and the fitness cost is 0.50.

Table S1. Observed ranges of infection rate, latent period duration, sporulation rate and sporulation duration for rust fungi (genus *Puccinia*) measured in different cultivars of wheat and barley. For a given study, different lines refer to different trials carried out in different conditions (time, temperature, pathogen genotype, leaf stage). Footnotes indicate when the measured variable is not exactly the same as the one used in the *landsepi* model (and of which a definition is given in the first line of the table).

Pathogen	Host	Nb of host genotypes	Nb of pathogen genotypes	Infection rate (Prop. of inoculated spores resulting in a lesion)			Latent period (Nb days from inoculation to onset of 50% of sporulating lesions)			Sporulation rate (Nb spores/lesion/day)			Sporulation duration (Nb days from end of latent period to end of sporulation)			Reference ^a
				Max	Min	Effect size	Min	Max	Effect size	Max	Min	Effect size	Max	Min	Effect size	
<i>P. hordei</i>	Barley	6	1			-60% ^b			95%			-50%			-30%	Parlevliet JE, 1979, table 4
<i>P. striiformis</i>	Wheat	10	1	5,85 ^b 8,63 ^b	0,33 ^b 0,67 ^b	-94% -92%										Broers LHM et al., 1996, Table 6
<i>P. striiformis</i>	Wheat	22	1	13,20 ^b	3,30 ^b	-75%	10,3	21,0	104%							Qamar M et al., 2012, Tables 1 & 2
				13,60 ^b	4,70 ^b	-65%	8,5	14,4	69%							
				17,00 ^b	11,90 ^b	-30%	8,2	11,9	45%							
<i>P. triticina</i>	Wheat	22	1	36,70 ^b	8,20 ^b	-78%	9,3	18,7	101%							Qamar M et al., 2012, Tables 3 & 4
				45,00 ^b	11,30 ^b	-75%	8,9	12,7	43%							
				46,70 ^b	26,30 ^b	-44%	8,2	10,0	22%							
<i>P. triticina</i>	Wheat	16	2	0,1303	0,0002	-100%	198,2 ^d	296,1 ^d	49%							Denissen CJM, 1993, Table 4
<i>P. striiformis</i>	Barley	11	1	17,50 ^b	1,20 ^b	-93%	12,1	14,9	23%	0,79 ^h	0,29 ^h	-63%				Richardson KL et al., 2006, Fig 3
<i>P. triticina</i>	Wheat	8	1	0,849	0,143	-83%	156,4 ^e	214,0 ^e	37%	15,83 ⁱ	5,12 ⁱ	-68%				Azzimonti G et al., 2013, Table 3
			1	0,696	0,101	-85%	153,9 ^e	210,9 ^e	37%	17,73 ⁱ	5,13 ⁱ	-71%				
			1	0,830	0,281	-66%	157,0 ^e	193,2 ^e	23%	20,85 ⁱ	7,72 ⁱ	-63%				
<i>P. striiformis</i>	Wheat	5	1	4,10 ^b	2,10 ^b	-49%	12,6 ^f	13,9 ^f	10%							Cromey MG, 1992, Table 1 & 2
		7	1	3,50 ^b	0,60 ^b	-83%	12,2 ^f	16,5 ^f	35%	0,51 ^h	0,25 ^h	-51%				
<i>P. striiformis</i>	Wheat	3	1	0,684 ^c	0,631 ^c	-8%	12,03 ^f	13,20 ^f	10%							Elahinia SA & JP Tewari, 2005, Tables 1 & 2
			1	0,597 ^c	0,565 ^c	-5%	14,11 ^f	17,18 ^f	22%							
				0,673 ^c	0,623 ^c	-7%	12,40 ^f	13,51 ^f	9%							
<i>P. triticina</i>	Wheat	1	3	0,593 ^c	0,558 ^c	-6%	14,27 ^f	17,40 ^f	22%							Pariaud B et al., 2009, Table 5
				18 ^b	13 ^b	-28%	147,7 ^e	156,5 ^e	6%	305 ⁱ	198 ⁱ	-35%				
				18 ^b	15 ^b	-17%				93 ⁱ	47 ⁱ	-49%				
<i>P. graminis</i>	Wheat	7	3	80 ^b	33 ^b	-59%										Mortensen K & GJ Green, 1978, Tables 3, 4, 5, 6, 7
				5,80 ^b	3,60 ^b	-38%				0,44 ⁱ	0,17 ⁱ	-61%				
				5,70 ^b	1,60 ^b	-72%				0,82 ⁱ	0,38 ⁱ	-54%				
				5,20 ^b	0,40 ^b	-92%				0,91 ⁱ	0,20 ⁱ	-78%				
				1,36 ^b	0,18 ^b	-87%				1,70 ⁱ	0,75 ⁱ	-56%				
				1,57 ^b	0,07 ^b	-96%				1,87 ⁱ	0,16 ⁱ	-91%				
				1,18 ^b	0,01 ^b	-99%				2,62 ⁱ	0,08 ⁱ	-97%				
				3,18 ^b	0,13 ^b	-96%										
<i>P. striiformis</i>	Barley	16	1	2,65 ^b	0,06 ^b	-98%										Sandoval-Islas JS et al., 2007, Tables 6, 7, 8
				2,12 ^b	0,01 ^b	-100%										
				23,10 ^b	0,01 ^b	-100%	8,6	9,5	10%							
<i>P. striiformis</i>	Wheat	10	1	14,00 ^b	0,00 ^b	-100%	15,7	24,4	55%							Broers LHM, 1997, Tables 1, 3, 4
							16,4	26,0	59%							
				23,00 ^b	0,00 ^b	-100%	10,7	13,5	26%	60 ^h	15 ^h	-75%				
				14,60 ^b	0,10 ^b	-99%	13,7	16,8	23%	93 ^h	26 ^h	-72%				
<i>P. striiformis</i>	Wheat	12	1				13,5	19,1	41%	97 ^h	13 ^h	-87%				Park RF & RG Rees, 1989, Table 5
							12,6	19,6	56%	138 ^h	18 ^h	-87%				
							11,4 ^f	13,8 ^f	21%							
<i>P. striiformis</i>	Wheat	12	1				11,2 ^f	14,5 ^f	29%							

							11,8 ^f	14,5 ^f	23%							
<i>P. striiformis</i>	Wheat	3	1				12,5 ^f	18,2 ^f	46%							Quan W et al., 2013, Table 3
							12,5 ^f	17,8 ^f	42%							
							13,2 ^f	19,2 ^f	45%							
		5	1				12,5 ^f	18,8 ^f	50%							
<i>P. striiformis</i>	Wheat	11	1				319,2 ^g	460,8 ^g	44%	64,86 ^h	19,45 ^h	-70%				Sørensen CK et al., 2014, Fig 2
			1				319,2 ^g	465,6 ^g	46%	71,35 ^h	10,81 ^h	-85%				
<i>P. striiformis</i>	Wheat	11	1				12,5 ^f	20,1 ^f	61%							Ma H & RP Singh, 1996, Table 3
							12,6 ^f	17,5 ^f	39%							
<i>P. recondita</i>	Wheat	3	1				8,3	8,8	6%	600,22 ^j	214,51 ^j	-64%	17,8	14,2	-20%	Tomerlin JR et al., 1983, Table 1 & 2
							10,1	11,6	15%	83,69 ^j	19,91 ^j	-76%	37,3	33,9	-9%	
							8,3	9,1	10%	160,78 ^j	48,85 ^j	-70%	21,4	18,3	-14%	
<i>P. triticina</i>	Wheat	4	1				149 ^d	177 ^d	19%							Rimé D et al., 2005, Fig 1
<i>P. triticina</i>	Wheat	5	7				7,75	12,60	63%							Lehman JS & G Shaner, 1998, Fig 1

^a in black a review of the literature, in green and blue experiments carried out in field and greenhouse conditions, respectively.

^b lesion density (nb lesions/surf. unit).

^c germination rate (prop. germinated spores).

^d latent period measured in hours.

^e latent period measured in degree-days.

^f latent period measured with the nb. of days until first sporulating lesions.

^g latent period measured with the nb. of hours until first sporulating lesions.

^h lesion size (surface or distance unit).

ⁱ total sporulation per lesion (mass unit of spores/lesion).

^j computed from total sporulation per lesion / sporulation duration (nb spores/lesion/day).