



HAL
open science

A NEW 665K SNP ARRAY USED FOR DETECTING SIGNATURES OF POSITIVE SELECTION IN RAINBOW TROUT (*ONCORHYNCHUS MYKISS*)

Florence Phocas, Katy Paul, Audrey Dehaullon, Maria Bernard, Gwendal Restoux, Delphine Lallias

► To cite this version:

Florence Phocas, Katy Paul, Audrey Dehaullon, Maria Bernard, Gwendal Restoux, et al.. A NEW 665K SNP ARRAY USED FOR DETECTING SIGNATURES OF POSITIVE SELECTION IN RAINBOW TROUT (*ONCORHYNCHUS MYKISS*). International Symposium on Genetics in Aquaculture, Nov 2022, Puerto Varas, Chile. hal-03778981

HAL Id: hal-03778981

<https://hal.inrae.fr/hal-03778981>

Submitted on 16 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A NEW 665K SNP ARRAY USED FOR DETECTING SIGNATURES OF POSITIVE SELECTION IN RAINBOW TROUT (ONCORHYNCHUS MYKISS)

Phocas, F.^{1*}, Paul, K.¹, Dehaullon, A.¹, Bernard, M.^{1,2}, Restoux, G.¹, and Lallias, D.¹

¹Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350 Jouy-en-Josas, France

²INRAE, SIGENAE, 78350 Jouy-en-Josas, France

e-mail: florence.phocas@inrae.fr

Native from North America, rainbow trout has been domesticated since the 1870s and introduced on all continents. Domestication and selective breeding have led to genetic changes due to inbreeding, genetic drift and selection. To address evolutionary biology questions, we used variant databases from USDA and INRAE to develop a 665K SNP array* (HD chip) for rainbow trout from a subset of the 32,372,492 SNPs. We prioritized the most variable SNPs in both databases with homogenous repartition over the genome. Of the 664,531 HD chip SNPs, 65% and 61% passed quality control and were polymorphic in samples of 288 and 175 genotyped fish from two commercial French lines (LB and LC respectively). Then, a subset of 72 LB and 67 LC genotypes of weakly related fish was pooled with the HD genotypes of 14 fish from the INRAE synthetic line (SY) and of 20 fish coming from rivers of the North West of USA (HA). These genotypes were used to reveal regions of the genome with low genetic diversity among various populations, likely to involve genes positively selected during either domestication or selection. The three French lines were moderately differentiated ($F_{st} \sim 0.10-0.12$), but strongly differed from the HA population ($F_{st} \sim 0.27-0.29$). A run of homozygosity (ROH) study revealed homozygous regions shared by all populations on Omy2, Omy12 and Omy15. These ROH islands included 26 annotated genes related to key roles in reproduction, embryonic development, genome stability and immunity.

*supported by EMFF project HypoTemp (n° PFEA470019FA1000016)