



# Most diverse, most neglected: weevils (Coleoptera: Curculionoidea) are ubiquitous specialized brood-site pollinators of tropical flora

Julien Haran, Gael J. Kergoat, Bruno A. S. de Medeiros

## ► To cite this version:

Julien Haran, Gael J. Kergoat, Bruno A. S. de Medeiros. Most diverse, most neglected: weevils (Coleoptera: Curculionoidea) are ubiquitous specialized brood-site pollinators of tropical flora. 2022. hal-03780127v3

**HAL Id: hal-03780127**

**<https://hal.inrae.fr/hal-03780127v3>**

Preprint submitted on 3 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Peer-reviewed & recommended by PCI Ecology

Recommendation: <https://doi.org/10.24072/pci.ecology.100505>

# Most diverse, most neglected: weevils (Coleoptera: Curculionoidea) are ubiquitous specialized brood-site pollinators of tropical flora

Julien Haran<sup>1</sup>, Gael J. Kergoat<sup>2</sup> & Bruno A. S. de Medeiros<sup>3</sup>

<sup>1</sup> CBGP, CIRAD, Montpellier SupAgro, INRAE, IRD, Montpellier University, Montpellier, France. ORCID: 0000-0001-9458-3785

<sup>2</sup> CBGP, INRAE, CIRAD, IRD, Montpellier Institut Agro, Univ. Montpellier, Montpellier, France. ORCID: 0000-0002-8284-6215

<sup>3</sup> Field Museum of Natural History, Chicago-IL, USA and Smithsonian Tropical Research Institute, Panama City, Panama. ORCID: 0000-0003-1663-668X

## Abstract

In tropical environments, and especially tropical rainforests, a major part of pollination services is provided by diverse insect lineages. Unbeknownst to most, beetles, and more specifically hyperdiverse weevils (Coleoptera: Curculionoidea), play a substantial role there as specialist mutualist brood pollinators. The latter contrasts with a common view where they are only regarded as plant antagonists. This study aims at giving a comprehensive understanding of what is known on plant-weevil brood-site mutualistic interactions, through a review of the known behavioral, morphological and physiological features found in these systems, and the identification of potential knowledge gaps to fill. To date, plant-weevil associations have been described or indicated in no less than 600 instances. Representatives of major plant lineages are involved in these interactions, which have emerged independently at least a dozen times. Strikingly, these mutualistic interactions are associated with a range of convergent traits in plants and weevils. Plants engaged in weevil-mediated pollination are generally of typical cantharophilous type exhibiting large, white and fragrant flowers or inflorescences and they also show specific structures to host the larval stages of their specialist pollinators. Another characteristic feature is that flowers often perform thermogenesis and exhibit a range of strategies to separate sexual phases, either spatially or temporally. Conversely, lineages of brood-site weevil pollinators present numerous shared behavioral and physiological traits, and often form multispecific assemblages of closely related species on a single host; recent studies also revealed that they generally display a high degree of phylogenetic niche conservatism. This pollination mutualism occurs in all tropical regions, and the contrasts between the known and expected diversity of these systems suggests that a wide range of interactions remain to be described globally. Our early estimates of the species richness of the corresponding weevil clades and the marked pattern of phylogenetic niche conservatism of host use further suggest that weevil-based pollination far exceeds the diversity of other brood-site mutualistic systems, which are generally restricted to one or a few groups of plants. As such, weevil pollinators constitute a relevant model to explore the emergence and evolution of specialized brood-site pollination systems in the tropics.

## Keywords

Cantharophily, entomophily, mutualism, nursery pollination, plant-insect interactions, tropical rainforests

67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

**Table of contents**

<b>Abstract</b>	2
<b>Table of contents</b>	3
<b>Introduction</b>	4
<b>1. A wide spectrum of mutualistic interactions</b>	5
<b>2. Reciprocal traits related to BSPM by weevils</b>	10
2.1 Plant traits	10
2.2 Weevil morphology and behavior	15
2.3 Evolutionary trends	21
2.3.1 Larval host specificity	21
2.3.2 Sympatric species assemblages	22
2.3.3 Cryptic and closely related weevil species	23
2.3.4 Trade offs in cost-benefit of interactions	24
2.3.5 Evolutionary dynamics of host use	25
<b>3. Economic importance and conservation</b>	26
<b>4. Oversights and estimates of global richness</b>	27
4.1 The inconspicuousness of plant-weevils pollination interaction	27
4.2 An underestimated diversity of species and interactions	28
<b>5. Concluding remarks: a road map for future research</b>	30
5.1 Searching for new plant-weevil BSPM systems	30
5.2 Priority research axes	32
5.2.1 A plea for description of systems	32
5.2.2 Evolutionary trajectories : when, why and how?	32
<b>Acknowledgements</b>	33
<b>Conflict of interest disclosure</b>	33
<b>References</b>	34
<b>Supplementary information</b>	52

## Introduction

Most of the extant biodiversity in tropical regions is concentrated in tropical rainforests, which were the first biomes to be coined as 'biodiversity hotspots' (Myers 1988). The outstanding diversity of pollinating systems in these ecosystems offers unique opportunities to study the evolution of reciprocally beneficial interactions among organisms. Pollination ecology in tropical rainforests indeed presents unique characteristics due to specific constraints. For instance, anemophily (wind pollination) is considered to be infrequent and potentially limited because of a combination of marked spatial constraints (evergreen canopy, structural heterogeneity and complexity) associated with frequent periods of high rainfall preventing pollen dispersal (Regal 1982; Williams & Adam 1994). Anemophilous plants are also presumably disadvantaged whenever they are over-dispersed (Regal 1982; see also below). Hence, pollination by animal vectors is predominant and can concern up to 100% of the plant species in some tropical rainforests (Bawa 1990; Ollerton et al. 2011; Rech et al. 2016). The high level of plant diversity that characterizes tropical rainforests is also associated with a spatial trend where the distribution of conspecific individuals is usually patchy (Bawa 1990; Williams & Adam 1994). The latter limits the efficiency of generalist pollinators, as they are not constrained to actively seeking out specific plants (Whitehead 1968; Bawa 1990; but see Wolowski et al. 2014). To cope with these major constraints, the reliance on more intricate pollination mechanisms, involving specialist pollinators (especially insects), is a hallmark of tropical rainforests (Bawa 1990; Renner & Feil 1993; Vizentin-Bugoni et al. 2018) and also widespread in all tropical or subtropical biomes (e.g., Gottsberger 1986; Ramirez 2004; Maruyama et al. 2013; Guy et al. 2021).

A mode of specialized pollination that particularly stands out is brood-site pollination (or nursery pollination; hereafter called BSPM for brood-site pollination mutualism), a system where immature stages of a pollinator develop within tissues (either flowers, ovules or pollen; Sakai 2002) of a plant as a reward for its pollination. It has independently evolved several times, mostly in tropical and subtropical biomes (Sakai 2002; Dufaÿ & Anstett 2003). These specialized mutualistic systems have been unequivocally documented in at least a dozen plant families and five insect orders (Coleoptera, Diptera, Hymenoptera, Lepidoptera and Thysanoptera) (Sakai 2002; Dufaÿ & Anstett 2003; Kawakita & Kato 2009; Hembry & Althoff 2016). Textbook examples of brood-site pollination are traditionally exemplified by the highly specialized fig/fig-wasp and yucca/yucca-moth models (Janzen 1979; Wiebes 1979; Pellmyr 2003; Herre et al. 2008); studies on these models have yielded important discoveries on the origin and maintenance of mutualisms and on the degree of co-evolution, with obligatory interactions possibly leading to evolutionary co-diversifications (e.g., Hembry & Althoff 2016). However, the number of insect and plant lineages engaging in brood-site pollination likely goes well beyond the few iconic interactions typically studied. Here we advocate the need to consider other systems to obtain a broader and more complete view of the ecology and evolution of biological interactions in general, and pollination in particular.

Among the neglected insect lineages involved in BSPM, the weevils (Coleoptera: Curculionoidea) are probably the most overlooked. Weevils contain over 62,000 described species and a colossal amount of unknown diversity (Oberprieler et al. 2007). This clade of beetle is exclusively phytophagous and is the most speciose lineage of phytophagous insects, usually only depicted as antagonistic to plants (Anderson 1995). Indeed, brood-site

pollination by weevils is generally considered anecdotal and is often overlooked in reviews on brood-site pollination (Dufayé et al. 2003; Hembry & Althoff 2016; Toon et al. 2020). Weevils are not even mentioned as one of the most relevant groups of beetle pollinators in some studies (e.g., Kevan & Baker 1983; Wardhaugh 2015; IPBES 2016). However, in recent decades, a number of plant-weevil interactions involving brood-site pollination have been progressively documented and described in detail (Franz & Valente 2005, Caldara et al. 2014; Nunes et al. 2018; de Medeiros et al. 2019; Saunders 2020; Toon et al. 2020). Moreover, recent plant-focused studies on tropical pollinators have recovered weevils as one of the most important pollinator groups (Krimse & Chaboo 2020; Paz et al. 2021). Taken altogether, all these studies suggest that these associations may be more widespread than previously thought (Caldara et al. 2014); it also contradicts the common view that weevils are not pollinators, and suggests that much of the global insect pollinator diversity is currently left unnoticed. For example, the combined species diversity of the three weevil families containing brood-site pollinator lineages (ca. 350, 4400 and 51,000 species, for Belidae, Brentidae and Curculionidae, respectively; Marvaldi & Ferrer 2014; Oberprieler 2014a, 2014b) is at least twice as large as the diversity of bees (ca. 20,000 species; Zattara & Aizen 2021), which are usually considered to be the most important group of insect pollinators (Potts et al. 2016). As we detail in this review, multiple and diverse weevil lineages commonly visit flowers, and an ever-increasing number of them can be considered true pollinators. This broad reassessment of the potential role of weevils as pollinators is highly relevant for the understanding of ecosystem functioning (including for agroecosystems), especially in tropical regions where bees are much less diverse than weevils (Oberprieler et al. 2007; Orr et al. 2021; Freitas et al. 2022).

The aim of this review is to provide a synthetic overview of all plant-weevil brood-site pollination mutualisms described to date. We begin by aggregating available data about weevil flower visitors and detail the extent of plant and weevil lineages known to be engaged in such interactions. We further summarize known reciprocal adaptations or traits and evolutionary trends to highlight both common patterns and specificities within these interactions. Finally, we investigate possible reasons why weevils have been previously overlooked in the pollination literature, estimate the extent of potential undescribed interactions, and conclude with a road map for future research on plant-weevil pollination mutualisms.

## 1. A wide spectrum of mutualistic interactions

Weevils are the most diverse group of insects that visit flowers and develop in reproductive structures (Oberprieler et al. 2007; Krimse & Chaboo 2020). This close interaction with the reproductive structures of plants probably preceded the colonization and diversification of weevils on angiosperms. The oldest weevil fossils (such as †*Belonotaris punctatissimus*) were found in the Karabastau Formation (ranging from the Oxfordian to Callovian stages, 166.1 to 157.3 million years ago (Mya); Walker et al. 2018); at that time they were likely associated with gymnosperm-dominated floras (e.g., see Oberprieler & Oberprieler 2012 for slightly younger weevil fossils from the Tithonian stage, 150.8 to 145.5 Mya). Their origin therefore clearly predates the rise to dominance of angiosperms during the Cretaceous Terrestrial Revolution, from 125 to 80 Mya (Dilcher 2000; Magallón & Castillo 2009; Benton 2010). In addition, several early diverging weevil lineages, such as Caridae or

Nemonychidae, typically breed on gymnosperm male or female reproductive structures (Oberprieler et al. 2007). Associations with angiosperm flowers and seeds evolved both within these early-diverging lineages (Ferrer et al. 2011; Kuschel & Leschen 2011) but also in more derived lineages belonging to the two most speciose weevil families (Brentidae and Curculionidae). Weevils in these clades generally visit flowers when feeding on pollen and ovipositing in buds, ovaries and fruits where larval development occurs (Oberprieler et al. 2007; Caldara et al. 2014). When adults fly between flowers to mate or oviposit on one or on a fairly narrow range of hosts, they can carry pollen between conspecific plant species, thereby pollinating them. Therefore, the flower-visiting and plant oviposition behaviors of several weevil lineages create the context for the evolution of BSPM from antagonistic to mutualistic interactions, or conversely (Figure 1). Interestingly, the ever-growing body of accumulated knowledge on plant-weevil interactions suggests that increasingly diverse and complex specialized plant-weevil BSPM interactions are to be expected (Franz & Valente 2005; Franz 2006; Saunders 2020; Toon et al. 2020).

The first report of weevil pollination activity refers to a palm-weevil interaction and dates back to the 19<sup>th</sup> century (Von Martius 1823). Since then, most efforts to describe and document weevil pollination have focused on palm-weevil and cycad-weevil mutualisms (Franz & Valente 2005; Toon et al. 2020). This focus on cycads and palms can be explained by their characteristic conspicuous inflorescences which attract massive gatherings of weevils, so much so that they have triggered specific research. To date, weevil BSPM has been reported for at least 87 palm species (Arecaceae) belonging to 26 genera (Table 1). Weevils pollinating palms consist of about 200 species from 17 genera, mostly belonging to the pantropical tribes Acalyptini and Derelomini (Curculionidae: Curculioninae) *sensu* Caldara et al. (2014), hereafter called Derelomini or derelomine weevils; see also Franz (2006) and Alonso-Zarazaga (2007) for nomenclatural uncertainties. Among the cycads, weevil BSPM has been reported for 91 cycad species belonging to seven genera from all three extant cycad families (Cycadaceae, Stangeriaceae and Zamiaceae). Weevils pollinating cycads belong to a diverse range of lineages; they are predominantly found within Curculioninae and Molytinae among the Curculionidae (ca. 50 species in six Paleotropical genera), but also within Belidae (24 species in six Neotropical genera) and Brentidae (12 species in two Afrotropical genera).

The study of these interactions, their mechanisms and their patterns of associations with palms and cycads has stimulated the discovery of similar cases among several monocot families of angiosperms. For instance, in Neotropical *Anthurium* (Araceae), 10 species are engaged in BSPM with *Cyclanthura* weevils (Derelomini, seven species; Franz 2003, 2006). In the same region, 34 species from eight genera of cyclanths (Cyclanthaceae) are pollinated by 27 weevil species belonging to six genera of Derelomini (Franz 2007a, 2008). In Neotropical Orchidaceae, three weevil species belonging to two genera of Baridinae (Curculionidae, alternatively classified as the supertribe Bariditae among Conoderinae) ensure their pollination (Nunes et al. 2018). More anecdotally, *Strelitzia nicolai* (Strelitziaceae) from southern Africa appears to be co-pollinated by two derelomine species belonging to two distinct genera (Haran et al. 2022a; Haran unpublished). Importantly, the majority of weevils engaged in BSPM show a marked pattern of phylogenetic niche conservatism (PNC) at the genus level (Table 1, Table S1 & Appendix 1; Franz & Valente 2005; Toon et al. 2020; Haran et al. 2021, 2022b, see section 2.3.5 for details). Such a pattern is far from unusual among groups of internal feeding insects, such as seed beetles




(Kergoat et al. 2007, 2015) or noctuid stem borers (Kergoat et al. 2018). Therefore, we postulate that, in the absence of direct experimental evidence for a weevil species, knowledge about other species in a genus may provide an indication of pollination activity. This generalization can produce hypotheses to be tested and also provide a first assessment of the extent of plant-weevil BSPM in overlooked tropical biomes (Toon et al. 2020).

Several dicot families have been also increasingly reported to be engaged in BSPM with weevils. In the Australasian region, some species of *Eupomatia* (Eupomatiaceae), *Myristica* (Myristicaceae) and *Exospermum* (Winteraceae) are pollinated by specific weevil lineages (*Elleschodes* spp., indet. genus (see Caldara et al. 2014) and *Palontus* spp. respectively; Armstrong & Irvine 1989a, 1989b, 1990; Pellmyr et al. 1990; Armstrong 1997). In African Ebenaceae (*Euclea*), Lecythidaceae (*Napoleonaea*) and Malvaceae (*Thespesia*) weevil pollination can be inferred on the basis of PNC since congeneric weevil species associated with other plant groups are brood pollinators (Haran et al. 2022a, 2022b; Haran pers. obs.). In the Neotropics, most species of *Cecropia* (Urticaceae) surveyed to date are associated with weevils belonging to the genus *Udeus* (Curculioninae: Eugnomini), which visit male inflorescences in all species (Lira et al. unpublished) and female inflorescences in at least one case (Mendonça 2004). In the speciose pantropical family Annonaceae, at least 22 species in 13 genera are associated with 30 species of weevils pollinators belonging to two Ochyromerini genera (Curculionidae: Curculioninae) and other unidentified weevil groups (Momose et al. 1998; Ratnayake et al. 2006; Lau et al. 2017; Saunders 2020; Dao et al. 2023; B. de Medeiros unpublished).


All in all, the extensive literature search conducted in this review underlines that no less than 600 instances of plant-weevils species-species BSPM interactions have been recorded to date or suggested based on the marked trend of PNC for brood pollination (Table 1; see Table S1 and Appendix 1 for details). Overall, BSPM with weevils has been demonstrated experimentally in 12 of the 22 plant families involved in this mutualistic system, with the remaining 10 families suggested based on both PNC of weevil genera and traits of floral structures associated with cantharophily. Representatives of all major plant lineages, gymnosperms, monocots and dicots, are engaged in BSPM with weevils; collectively representing ca. 250 species belonging to 72 genera (Figure 1) associated with about 300 species of weevils. In this regard, weevils are by far the most ecologically successful group of insects engaged in BSPM interactions with the plant kingdom. Our review shows that weevil-based BSPM has emerged several times independently with plants involved in these interactions typically belonging to groups considered to be ancient lineages (Pant 1987; Franz 2004; Saunders 2020; Baker & Couvreur 2013). We also show that this type of interaction is currently predominantly found in tropical and subtropical regions of the eastern and western hemisphere, with only a few cases under Mediterranean climates (Figure 2).





**Figure 1.** Summary of known and expected brood-site pollination mutualisms (BSPM) between plants and weevils (Coleoptera: Curculionoidea). The boxes in blue and green refer respectively to the weevil and plant lineages involved (the size of boxes is correlated to the number of species involved; see also the approximate scale at the bottom left). The phylogenetic tree on the left presents a synthetic view of phylogenetic relationships among weevils based on the phylogenomic study of Shin et al. (2018). The relationships among the so-called CCCMS clade (for Baridinae, Cossoninae, Curculioninae, Molytinae and Scolytinae) are unresolved; several subfamilies belonging to this clade and not involved in BSPM (e.g., Scolytinae, Lixinae) are not shown. Most of the plant families reported here contain at least one experimentally verified case of BSPM; however, for the plant families highlighted with an “\*”, BSPM interactions are postulated based on the phylogenetic niche conservatism of the weevil genera involved (see Methods in Appendix 1). Some minor interactions between several lineages of Curculioninae and dicots are not shown for clarity; see Table S1 for details. Pictures of selected examples of inflorescences of plant lineages involved: **A.** *Encephalartos* sp. (Zamiaceae) pollinated by weevils from the Brentidae family and Curculionidae Molytinae. **B.**

*Elaeis guineensis* (Arecaceae) pollinated by Derelomini. **C.** *Syagrus coronata* (Arecaceae) pollinated by Derelomini. **D.** *Carludovica palmata* (Cyclanthaceae) pollinated by Derelomini. **E.** *Euclea racemosa* (Ebenaceae) associated with Derelomini. **F.** *Annona senegalensis* (Annonaceae) pollinated by Ochyromerini. **G.** *Cecropia peltata* (Urticaceae), a genus including species pollinated by Eugnomini. (credits: A-B, E: J. Haran; C-D, G: B. de Medeiros, F: M. Gueye).



**Figure 2.** Geographical distribution of brood-site pollination mutualism (BSPM) involving weevil lineages and estimates of knowledge gaps. Distribution data collated from [Alonso-Zarazaga & Lyal \(1999\)](#), articles listed in this review, and unpublished observations by the authors (note that the distribution of an unidentified weevil genus associated with Myristicaceae is not shown). Bars show the relative number of known and estimated BSPM interactions for a selection of plant-weevil interactions. Beige: number of plant species involved in verified interactions (involving brood-site and/or synchronization of weevil and flower phenology and/or pollen transfer observed). Light brown: number of plant species involved estimated based on floral visitors, including weevils and PNC of weevils involved. Dark brown: global diversity of interaction estimated from the diversity of plant lineages engaged in BSPM with weevils (Derelomini-Carludoviceae; [Franz & Valente 2005](#); [Franz 2006](#); see details on Methods in Appendix 1) and the diversity of weevil genera engaged in these interactions (other interactions; Table S1, GBIF data and unpublished authors observations, see section 4.2). y-axis: species number.

[illegible]

**Table 1.** Synthetic list of known plant-weevils BSPM systems. See Table S1 for details on species-specific interactions, species authorship, tissues for larval development, presence of cryptic lineages and corresponding reference list. See Figure 2 for information on the geographic distribution of weevil lineages involved in BSPM.

## 2. Traits related to BSPM by weevils

### 2.1 Plant traits

Flowers and inflorescences of plants engaged in BSPM with weevils generally display typical traits related to cantharophily (beetle pollination) and remarkable convergent morphologies, physiologies, phenology and reproductive strategies. Some of the traits in these flowers are associated with beetle pollination more generally and not specifically with brood pollination. Disentangling the role of each trait in the interaction is a task that remains to be done in most cases. Here we point out features that have been proposed to increase the attractiveness of flowers to weevils engaged in BSPM, provide them with food rewards and sometimes protect them from predators. They also may improve pollen transfer to weevil integuments and play a role in weevil pollinator selection and retention, thereby improving pollination efficiency.

A first major characteristic for weevil-pollinated plant lineages is that they tend to display **large flowers or inflorescences** (Figure 1). In cycads and most monocot lineages involved in BSPM (Araceae, Cyclanthaceae, some Arecaceae), flowers or female ovules are clustered in large compact cones or inflorescences (Franz & Valente 2005; Toon et al. 2020). Individual flowers can also reach substantial sizes in Annonaceae or Lecythidaceae (Prance & Jongkind 2015; Saunders 2020). In contrast, other plant lineages producing smaller individual flowers form either dense inflorescences (most Arecaceae) or

synchronized blooms (Ebenaceae – *Euclea*, Orchidaceae – *Dichaea*, *Grobya* etc). Field observations and experiments on various Annonaceae flowers and Cyclanthaceae inflorescences show that larger floral structures attract greater number of pollinators (Gottsberger 1999; Franz 2007a), revealing that this floral trait has been independently selected to increase weevil attraction and pollination.

The main characteristic of weevil-based cantharophily is the transformation of floral structures to host the weevil larval stages. Two main types of floral morphology are typical of weevil-pollinated plants: (i) **clustered inflorescences** (in cycads and all monocots involved in BSPM except Orchidaceae and Strelitziaceae), and (ii) **individual floral chambers, bowls or claws** (in most dicots involved in BSPM; Gottsberger 1999; Prance & Jongkind 2015; Saunders 2020). Clustered inflorescences provide fleshy and nutrient-rich tissues for the development of weevil larvae, generally in male inflorescences (flower capsules, microsporophylls and rachises; Anstett 1999; Downie et al. 2008; Teichert et al. 2018; Haran et al. 2020). Within individual floral chambers, tissues for larval development include anthers, or fleshy petals (Saunders 2020; Haran et al. 2022a; Dao et al. 2023). In clustered inflorescences as in floral chambers, the modified structures effectively filter access to specific pollinators and sometimes further retain and protect them. For instance, in Cyclanthaceae, the inflorescences of *Asplundia*, *Evodianthus* and *Ludivia* exhibit narrow interfloral entrances allowing only a specific weevil pollinator to reach the hidden floral reproductive structures (Franz 2007a; Teichert et al. 2018), while larger, non-pollinating species are prevented from reaching them. Narrow interfloral openings can also promote the transfer of pollen from anthers to the integuments of pollinators as they exit the floral chambers (Teichert et al. 2018). The Annonaceae provide the most striking examples of enclosed floral chambers, which are used to retain and protect *Endaeus* weevil pollinators and also ward reproductive tissues from predation (Gottsberger 1999; Ratnayake et al. 2006; Lau et al. 2017; Saunders 2020). Other structures associated with weevil pollination include staminodes (in Cyclanthaceae and Eupomatiaceae), which are stamen-like structures that do not produce pollen but rather sticky secretions or odoriferous oils (elongated projections in Fig. 1D). They increase attraction, provide food, shelter or protection for weevil pollinators (Armstrong & Irvine 1990; Franz 2007a; Teichert et al. 2018). In Annonaceae sticky pollenkit and short pollen-connecting threads create large aggregates of pollen grains that are considered specific adaptations to improve the efficiency of pollen transfer onto weevils integuments (Ratnayake et al. 2006). The fleshy petals and other floral tissues of plants implicated in BSPM with weevils typically show extensive feeding damages (Ratnayake et al. 2006; Saunder 2020; Haran et al. 2022a; Dao et al. 2023; Figure 3F), indicating that these structures are also food sources for the adult pollinators.

The flowers of plant lineages engaged in BSPM with weevils are **predominantly white or creamy white** (Figure 1), but pale red, pale yellow and pale green flowers are occasionally encountered (Franz & Valente 2005; Ratnayake et al. 2006; Saunders 2020; Toon et al. 2020). In Cyclanthaceae, weevils show a preference for natural white inflorescences in contrast to those experimentally shaded by an opaque fabric (but letting the fragrance to be emitted; Franz 2007a). It should be noted that many weevils lineages involved in BSPM have nocturnal or crepuscular activity (Franz 2007a; Auffray et al. 2017; Lau et al. 2017; Auffray et al. 2022; Haran et al. 2022a), where light intensity is low. Under these conditions, the pale colors were hypothesized to provide better contrast with the surrounding environment in a way that likely enhances their visual recognition by weevils



(Franz 2007a; Saunders 2020). This suggests that visual recognition of inflorescences is an important dimension of weevil attraction, at least as important as olfaction (see below), and that white or pale colors are more attractive to weevils.

All major plant lineages engaged in BSPM with weevils include representatives with **thermogenic** activity, allowing inflorescence or flower temperatures to rise several degrees above ambient temperature (Franz 2007a; Teichert et al. 2018; Saunders 2020; Toon et al. 2020). This phenomenon is postulated to act as an attractant for weevil and other beetle pollinators, by increasing the emission of volatiles or infrared radiation itself. It has also been hypothesized to provide an energy reward to individuals reaching floral tissues (Rands & Whitney 2008), but this is likely not the case for weevils since thermogenic activity is usually correlated with short phases of attraction and is not maintained during an ‘interim’ phase when individuals stand on flowers or are enclosed in floral chambers (Teichert et al. 2018; Toon et al. 2020). In Annonaceae, thermogenesis is identified as typical of flowers pollinated by beetles, and weevils in particular (Gottsberger 1999). The temperature reached appears to be positively correlated with inflorescence size (Franz 2007a), which may explain why large flowers and inflorescences evolved convergently in weevil-pollinated plants. In several systems (Carludovicoideae-Derelomini; Zamiaceae-Belidae), the first peak of thermogenesis associated with the attraction phase of the weevils is followed, after an ‘interim’ phase of several hours, by a second peak where weevils leave the inflorescences (Franz 2007a; Teichert et al. 2018; Salzman et al. 2020). These second peaks of thermogenesis coincide with the staminate phase and probably act as repellent or as an aid for improved take-off that favors weevil departure once they are covered with pollen (Teichert et al. 2018). This kind of push-pull mechanism where weevils are successively attracted and then repelled during the circadian flowering cycle promotes pollinator movement between plant congeners and thus cross-pollination (Salzman et al. 2020).


The **fragrances** produced by weevil-pollinated flowers during anthesis are strong and detectable by human olfactory senses, and are commonly associated with fruity or floral fragrances. In most genera of weevil-pollinated Cyclanthaceae (*Aspludia*, *Carludovica*, *Chorigyne*, *Dicranopygium* and *Evodianthus*), production of aromas reminiscent of various fruits and flowers has been reported (Franz 2007a). Similarly, in *Polyalthia* (Annonaceae), a fruity odor is produced during anthesis (Ratnayake et al. 2006). In *Elaeis guineensis* (Arecaceae), anthesis is correlated with the production of estragol (Lajis et al. 1985), while in *Attalea phalerata* (Arecaceae), weevils and other beetle pollinators are attracted by a strong emission of methyl acetate (Maia et al. 2021). In *Macrozamia* and *Zamia* (Zamiaceae) the emission of linalool or methyl salicylate has been recorded (Toon et al. 2020; Salzman et al. 2021). Overall, the emission in large amount of one or a few compounds that are specific to the interaction seems to be a signature of chemical communication in BSPM involving weevils (Ervik et al. 1999; Teichert et al. 2018; Salzman et al. 2021; T. Auffray pers. com.). Notable exceptions of floral odor emission are found in *Chamaerops humilis* (Arecaceae), which attracts *Derelomus* species via the volatiles emitted from the leaves (Dufay et al. 2003), and in *Syagrus coronata* (Arecaceae), where the volatiles are given off by a persistent peduncular bract (Barbosa et al. 2020).

The morphological and physiological features described above are generally organized into **timed phenological sequences** that coincide with **phases of flower and inflorescence receptivity**. In most plant lineages producing large inflorescences (e.g.,

Araceae, Arecaceae, Zamiaceae), anthesis occurs over several days and is accompanied by circadian thermogenic cycles and emission of volatiles (Franz 2007a; Auffray et al. 2017; Toon et al. 2020). When flowers are isolated (in Annonaceae and Eupomatiaceae), but also in Cyclanthaceae, each individual flower or inflorescence goes through one cycle of anthesis with the following stages: (i) a phase of thermogenesis/attraction associated with the receptivity of female reproductive structures (pistillate phase), (ii) an interim period when weevils can be trapped in the floral chamber for a variable time, and (iii) a phase when pollinators are released with exposure to male reproductive structures (stamen and pollen, staminate phase), which can be achieved by abscission of the floral chamber petals and/or by a second peak of thermogenesis (Armstrong & Irvine 1990; Franz 2007a; Lau et al. 2017; Teichert et al. 2018; Saunders 2020). In the first case, the anthesis lasts between several days and two weeks (Toon et al. 2020), while in the second case, the complete cycle of anthesis is short and usually lasts at most 48h (Franz 2007a; Saunders 2020). Interestingly, the pistillate and staminate phase cycles found in some Cyclanthaceae (*Asplundia*; Franz 2007a) are very similar to those found in some Annonaceae (Saunders 2020), indicating a remarkable convergent evolution of plant-weevil pollination systems. With the notable exception of cycads, orchids and some palms, in all plant-weevil BSPM systems, the **pistillate and staminate phases coincide with peaks of crepuscular or nocturnal peaks in pollinator activity** (Armstrong & Irvine 1990; K  chmeister et al. 1998; Franz 2007a; Auffray et al. 2017; Saunders 2020; Auffray et al. 2022). These converging nocturnal phenology suggest that nocturnal conditions improve pollination efficiency, possibly through increased attraction of weevil pollinators. It has been also hypothesized that the movement of pollinators and the dispersal of floral fragrances are facilitated at night because there is less wind; lower temperatures also potentially improve the detection of thermogenic flowers (Borges et al. 2016; Auffray et al. 2022).

A remarkable feature of plant lineages engaged in BSPM with weevils is the **separation of sexes**. About 95% of the species recorded in this review show physical or functional dioecy, a rare feature in the plant kingdom in general. In cycads, all species are dioecious, with male plants producing pollen cones and female plants ovulate cones (Toon et al. 2020). Weevil-pollinated palms may be dioecious or monoecious, but anthesis of male and female flowers are always separated in time in the latter case (Barfod et al. 2011). One example is the monoecious palm genus *Elaeis* (Arecaceae), in which individuals cyclically produce functionally unisexual male and female inflorescences, resulting in an allogamous mode of reproduction called temporal dioecy (Adam et al. 2011). In other genera, such as *Acrocomia* and *Bactris* the pistillate and staminate phases of the **protogynous inflorescences** are separated by a short interim phase resulting in circadian dioecy (Henderson et al. 2000; Carre  o-Barrera et al. 2021). Similar patterns are observed in other monoecious plant lineages such as Annonaceae and Cyclanthaceae (Franz 2007a, 2007b; Saunders 2020). In Annonaceae in particular, several other types of monoecy that may result in functional dioecy have been reported (i.e. separate pistillate and staminate flowers on the same plant, with a few flowers per individual at a time; Saunders 2020). Finally, a number of palms exhibit **protandrous inflorescences** with very extended interim phases, also resulting in functional dioecy (Silberbauer-Gottsberger et al, 2013; N    ez-Avellaneda et al, 2008, 2015; de Medeiros et al. 2019). Another striking feature of systems involving Arecaceae, Annonaceae and Cyclanthaceae is the small number of flowering inflorescences per individual or the small number of flowering individuals at time, a strategy that also likely promotes cross-pollination (Saunders 2020). The resulting low number of propagules

available at a certain time may be balanced by the long flowering periods of the plants engaged in these BSPM interactions, for instance up to seven-months in *Asplundia* (Cyclanthaceae; Franz 2007a) and many species of *Syagrus* (Arecaceae; Noblick 2017). These long flowering periods have also been proposed as a strategy for constant breeding of specialized brood pollinators (Carreño-Barrera et al. 2020).



**Figure 3.** Examples of weevils and plants engaged in brood-site pollination mutualisms. **A.** *Anchylorhynchus trapezicollis* (Derelomini) on pistillate flowers of *Syagrus coronata* (Arecaceae) in South America. **B.** *Derelomus* sp. (*costiger* group; Derelomini) pollinator of *Phoenix reclinata* (Arecaceae) in southern Africa. **C.** *Perelleschus evelynae* (Derelomini) on pistillate flowers of *Carludovica palmata* (Cyclanthaceae) in Central America. **D.** *Tranes lyterioides* (Molytinae) on sporophylls of *Macrozamia communis* (Zamiaceae) in Australia. **E.** *Udeus eugnomoides* (Eugnomini) on pistillate flowers of *Cecropia peltata* (Urticaceae) in Central America. **F.** Flowers of the African custard apple (*Annona senegalensis*; Annonaceae) with aggregation of *Endaeus* spp. pollinators (Ochyromerini) in tropical Africa. (credits: A/C/E: B, E: B. de Medeiros, B: J. Haran, D: R. Oberprieler, F: Z. Dao).




## 2.2 Weevil morphology and behavior

Like plants involved in BSPM, weevil pollinators display a series of morphological, physiological and behavioral features that may constitute adaptations to pollination mutualisms. Weevil pollinators have been less studied than the plants they pollinate, so not only their shared morphological features have been scarcely reported, but also the function and adaptive significance of these features are generally unexplored. This section provides a summary of what is known to date, with the aim of identifying general trends associated with BSPM pollination, as well as pointing out more peculiar cases.

Known weevil species involved in BSPM are **small-sized beetles**, with body size ranging from a few millimeters to about one centimeter (1.1 mm, *Staminodeus inermis* (Franz 2001); 8.6 mm in *Anchylorhynchus burmeisteri* (de Medeiros & Vanin 2020), rostrum excluded). Body shape is often consistent with floral morphology; for example, the body of weevils associated with cycads in Belidae and Brentidae is very flat and allows individuals to enter the cracks of megasporophylls. In most species, the body colors are **pale yellow, reddish or brown**; they usually match the color or shape of their floral substrates, probably to limit predation (Figures 3, 4). The matching colors are either obtained due to the color of the integument itself (e.g., in *Andranthobius*, *Cotithene*, *Derelomus*, *Ebenacobius*, *Elaeidobius*, *Prosoestus*) or due to the colors of the integument and scales covering it (e.g., in *Anchylorhynchus*; Valente & da Silva 2014; de Medeiros & Vanin 2020; Haran et al. 2020, 2022a). In some species, the shape of the elytra forms an angled surface mimicking the shape of the petals of the host plant (e.g., *Derelomus subcostatus*, *D. costiger* and *D. piriformis*; see Figure 4E). Overall, there seems to be a selective pressure to mimic floral substrates as suggested by the various examples of strongly converging phenotypes among unrelated species associated with the same host plant (e.g., *Derelomus pallidus* and *Ebenacobius rectirostris* on *Euclea racemosa*; Haran et al. 2022b; Figure 4A, B), but whether these phenotypes really provide an advantage in escaping predation has not been formally explored. Interestingly, patterns of mimicry are generally found in species active during the day on inflorescences, with little or no space to hide (i.e. most Derelomini associated with Arecaceae or Ebenaceae). Conversely, in weevil lineages that remain hidden in floral chambers or inflorescences during the day, the integuments are usually pale brown or reddish in color, and they do not specifically match floral substrates (e.g., in *Cyclanthura*; Ochyromerini; Figure 4C, F). Notable exceptions to this pattern include *Montella* weevils (Baridinae), which exhibit a dark body integument contrasting with the color of its host's flowers (Nunes et al. 2018) and some species of *Anchylorhynchus* weevils with extreme polymorphism showing disruptive patterns such as black and yellow stripes (de Medeiros & Vanin 2020).


A series of putative pollen-carrying morphologies are found in weevils engaged in BSPM. Most of these lineages have species with **erect setae** on their body surface (Haran et al. 2020, 2022a; Dao et al. 2023). Hair or scale cover on the integument are widespread in weevils, so these may represent instances of exaptation. However, their peculiar size or arrangement in some lineages acting as brood-site pollinators led to suggestions that they may improve pollen transport (Syed et al. 1982). The erect setae can cover the entire body surface (in Ochyromerini or in some *Ebenacobius* and *Celetes*; Valente 2005; Ratnayake et al. 2006; Haran et al. 2022b; Figure 4C, F), or form combs on the edge of the elytra (in some *Elaeidobius*; Figure 4D), or on the tibiae (in some *Derelomus*); setae may also be found on

the ventral region (in some *Elaeidobius* and *Anchylorhynchus*). In *Anchylorhynchus*, a genus that breeds on female flowers and thus may benefit directly from pollination, these ventral setae may be highly branched in some species, reminiscent of setae found in bee corbiculae (de Medeiros et al. 2019; de Medeiros & Vanin 2020). Interestingly, in some weevil species, the presence of setae is a dimorphic feature, with only males exhibiting long setae, suggesting that they may play a differential role in pollen transport (Haran et al. 2020; Hsiao & Oberprieler 2022). That said, even weevil species without specific setae (e.g., *Derelomus chamaeropsis*, *Andranthobius* spp., Baridinae and Belidae) are known to be effective pollinators (Anstett 1999; Nunes et al. 2018; Salzman et al. 2020; Carreño-Barrera et al. 2021), suggesting that integuments alone can efficiently transport pollen. Several unrelated lineages of weevils engaged in BSPM also exhibit prosternal processes or tubercles (in *Andranthobius*, *Diplothemiobius*, *Ebenacobius*, *Elaeidobius*, *Miltotranes*, *Porthetes*, *Prosoestus*, *Tranes*; Bondar 1941; Valente & da Silva 2014; Haran et al. 2020, 2022a; Hsiao & Oberprieler 2022; Haran pers. obs.) consisting of cuticular apophyses located near the procoxae of males. In some Baridinae not involved in BSPM, such structures grow allometrically and have been shown to be used in interspecific fights to dislodge competing males (Eberhard & Garcia 2000; Eberhard et al. 2000; Davis & Engel 2010). Since fights between males have been described in Derelomini (Franz 2003, 2006), it can be postulated that these prosternal processes or tubercles probably have a similar function for the aforementioned genera. Finally, in some weevil lineages, structural complexity and dense cover of setae in these structures have been hypothesized to possibly improve pollen transport (see Hsiao & Oberprieler 2022). It is not known why the weevils evolved these pollen-carrying structures, except in the rare cases where the weevils directly benefit from the pollination of their hosts. Indeed, there are only two known cases where weevils breed on tissues that result from their direct pollination: in the genus *Montella* (Nunes et al. 2018) and in the genus *Anchylorhynchus* (de Medeiros & Vanin 2020). *Montella* weevils display active pollination behavior (carrying orchid pollinaries; Nunes et al. (2018), which is consistent with other cases of specialized pollinating seed predators such as fig-wasps (Jousselin et al. 2003) and yucca moths (Pellmyr 1997). For *Anchylorhynchus* the evidence is more ambiguous; females rub their bodies against female flowers after oviposition, but it remains to be shown whether this behavior serves to mark oviposition and avoid competition, increases the likelihood of pollen deposition, or both (de Medeiros et al. 2014).



**Figure 4.** Dorsal habitus of weevils involved in BSPM. **A-B.** *Derelomus pallidus* and *Ebenacobius rectirostris* (Derelomini), two species associated with inflorescences of *Euclea* bushes (Ebenaceae) showing a marked morphological convergence. **C.** *Ebenacobius curvistetis* (Derelomini), a species with body surface covered with erect setae, thought to be an adaptation to pollen transportation. **D.** *Elaeidobius kamerunicus* (Derelomini) pollinator of *Elaeis guineensis* (Arecaceae) with combs of setae on edge of elytra in males. **E.** *Derelomus costiger* (Derelomini) with shape of elytra mimicking the structure of petals of its host *Phoenix reclinata* (Arecaceae). **F.** *Endaeus floralis* (Ochyromerini) pollinator of *Annona senegalensis* (Annonaceae) with body surface covered by setae. (credits: A-F: J. Haran).





**Figure 5.** Breeding sites of weevils engaged in BSPM. **A.** Pupa of *Elaeidobius plagiatus* (Derelomini) in flower capsules of male inflorescences of *Elaeis guineensis* (Arecaceae). **B.** Egg of *Phytotribus platyrhinus* (Derelomini) inside rachis of inflorescence of *Syagrus coronata* (Arecaceae). **C.** Larva of *Andranthobius bondari* (Derelomini) in aborted staminate flower of *Syagrus coronata* (Arecaceae). **D.** Larva of *Celetes impar* (Derelomini) in a spathe of *Syagrus coronata* (Arecaceae). **E.** Larvae of *Anchylorhynchus bicarinatus* (Derelomini) inside a pistillate flower petal of *Oenocarpus mapora* (Arecaceae). **F.** Larva of *Derelomus chamaeropsis* (Derelomini) inside a rachis of male inflorescence of *Chamaerops humilis* (Arecaceae). (credits: A, F: J. Haran, B-E: B. de Medeiros).



**Figure 6.** Example of sympatric species assemblage of weevils of a host including species from other beetle lineages. **A.** Inflorescence of *Syagrus coronata* (Arecaceae) with *Anchylorhynchus trapezicollis* (Derelomini, top right), *Andranthobius bondari* (Derelomini, top left), *Microstrates ypsilon* (Baridinae, black weevils) and sap beetles (Nitidulidae, *Mystrops palmarum*). **B.** Inflorescences in anthesis of *Elaeis guineensis* (Areaceae) with multiple specific pollinators including weevils (*Elaeidobius plagiatus* (left), *E. piliventris* (right) and *E. subvittatus* (bottom left), Derelomini). (credits: A: B. de Medeiros, B: J. Haran).

Weevils engaged in BSPM likely have good **flight abilities**, which allow them to fly actively between inflorescences of distant hosts ([Auffray et al. 2017](#); [Lau et al. 2017](#);

Saunders 2020; Toon et al. 2020). This is supported by experimental data in other small weevils, which are capable to fly up to kilometers in flight mills (McKibben et al. 1998; Evenden et al. 2014) and by our own experience in collecting specimens on isolated hosts, suggesting that they can move over long distances (J. Haran and B. de Medeiros pers. obs.). Moreover, unlike social insects such as bees, the home range of pollinator weevils is not constrained by the location of a nest. Adult weevils feed on pollen or nectar produced by flowers (de Medeiros et al. 2014; Lau et al. 2017; de Medeiros et al. 2019) but also often on floral structures themselves (*Elaeidobius*, *Ebenacobius*, *Endaeus*; Haran et al. 2022a; Dao et al. 2023). The active behavior of adults is not surprising, since weevils **respond well to host specific floral volatiles**. This has, for example, been demonstrated in several cycad specialists, with electroantennography experiments and pheromone traps (Salzman et al. 2021). Several weevil lineages have also developed specific olfactory globules in their antennae which are considered to be adaptations for the detection of these volatiles (Crowson 1991; Oberprieler 2004). Following olfactory stimuli, weevils can reach or leave the flowers and inflorescences of their hosts, according to circadian cycles remarkably synchronized with flowering cycles (Franz 2007a; Auffray et al. 2017; Mendeléz-Jácome et al. 2019; Saunders 2020; Salzman et al. 2021). The weevil's reaction to a volatile can be concentration-dependent, with the same molecule acting as either an attractant or a repellent depending on the amount emitted from the floral structures. This change in behavior in turn creates a push-pull effect enhancing their movement between flowers and therefore pollination (Franz 2007a; Salzmann et al. 2021). When floral structures do not produce peaks of volatiles (the so-called 'interim' phase'; Saunder 2020), weevils may simply remain on the inflorescences (in cycads and palms; Auffray et al. 2017; de Medeiros et al. 2019; Toon et al. 2020), be hidden or trapped there (in Araceae and Annonaceae; Franz 2007b; Saunders 2020; Dao et al. 2023), or leave inflorescences (Ratnayake et al. 2006; de Medeiros et al. 2019). In the genus *Ebenacobius*, many species have only been collected from the leaf litter during the day, which suggests that they hide there between their crepuscular and night phases of movements (Haran et al. 2022a). Weevils are fairly long-lived insects, with an adult lifespan of up to 79 days (mean 31 days) in *Elaeidobius subvittatus* (Syed et al. 1982). As the anthesis of the host inflorescence lasts only a few hours or a few days (five days in the case of *Elaeis guineensis*), the same individual can visit many flowers or inflorescences during its lifespan.

The **larval stages** of weevils **develop on the various types of substrates** provided by their hosts. In all the weevil lineages engaged in BSPM, the larvae are generally endophagous, developing either in live tissues or as detritivores on decaying plant material, in more or less woody tissues (Figure 5; Anstett 1999; Franz 2007a,b; Nunes et al. 2018; Toon et al. 2020; Dao et al. 2023). In Ebenaceae, which have small individual flowers, larvae of pollinators are ectophagous and feed at the base of the anthers (Haran et al. 2022a). In *Anchylorhynchus*, eggs are deposited externally between petals (Figure 5E) and first instars drill into female flowers to continue their development internally (de Medeiros et al. 2014). When larvae feed on limited spaces such as individual flowers or fruits, some species have developed a cannibalistic behavior (Figure 5E) and morphology to cope with a potential overpopulation in the brood site (de Medeiros et al. 2014). For most plant-weevil BSPM, the interplay between the presence of secondary compounds in plant tissues and a potential adaptation of associated weevil lineages has not been yet investigated, although they may drive specialization patterns (Wang et al. 2021). For example in weevils associated with



cycads (Belidae), the larvae of some species have been found associated with specific bacteria involved in the control of toxic compounds from their host (Salzman et al. 2018).

## 2.3 Evolutionary trends

### 2.3.1 Larval host specificity

The vast majority of weevils engaged in brood-site pollination appear to be strictly monophagous as larvae (Franz & Valente 2005; de Medeiros & Núñez-Avellaneda 2013; Valente & de Medeiros 2013; Toon et al. 2020; de Medeiros & Farrell 2020; de Medeiros & Vanin 2020; Haran et al. 2021; Hsiao & Oberprieler 2022). Such a level of specialization is rare for weevils and phytophagous insects in general, even when considering other groups of endophagous feeders, which are known to be generally highly specialized (e.g., Gaston et al. 1992; Anderson 1993; Bernays & Chapman 1994; Kergoat et al. 2008, 2018). One of the most striking case reflecting this host specificity is the pollination ecology of sympatric palms belonging to the genus *Oenocarpus* in the Amazon (Núñez-Avellaneda et al. 2015). The three species studied have a very similar floral structure and phenology, but each has a specific assemblage of weevil pollinators, only a few of which are able to visit several *Oenocarpus* species. Also, in *Montella* weevils actively pollinating *Dichaea cogniauxiana*, adults visit the flowers of another orchid species (*D. pendula*), but can only complete their life cycle in the former due differences in self-compatibility of the reproductive system (Nunes et al. 2016). Cases of oligophagy or polyphagy in weevil pollinators have been suggested in several systems based on general observations of inflorescences visitors, but these have never been formally tested (Franz & Valente 2005; Ratnayake et al. 2006; Saunders 2020). In the Annonaceae – Ochyromerini system, a species of *Endaeus* has been identified as the pollinator of two congeneric and sympatric *Polyalthia* species (Ratnayake et al. 2006). In the Araceae – Derelomini and Cyclanthaceae – Derelomini systems, a few intriguing cases of oligophagy or even polyphagy have been reported for weevil species acting as effective pollinators (Franz & O'brien 2001a,b; Franz 2006, 2007); however, most of these accounts should be viewed with caution as they are based on observations of adult stages only. Weevils can be slightly eclectic in the selection of flowers visited at the adult stages (Haran et al. 2022a) and over-interpretation of these visits can lead to erroneous conclusions about larval host specificity. The complexity of the host plant taxonomy can also make it difficult to interpret historical records and blur patterns of host-associations (de Medeiros & Vanin 2020). All of this highlights that understanding the level of species-specificity in these systems is primarily hampered by the lack of actual host plant data for the larval stages (Franz 2004). More generally, integrative and detailed studies are needed to accurately assess plant-weevil BSPM interactions. All detailed studies of host plant associations using molecular data have found a higher than previously thought weevil diversity and a very high degree of specialization on their hosts (Downie & Williams 2009; Brookes et al. 2015; de Medeiros & Farrell 2020; Haran et al. 2021).



### 2.3.2 Sympatric species assemblages

While most weevils engaged in pollination mutualism are specialized and dependent on a single host, the reciprocal condition does not apply to plants, which are usually pollinated by various insect guilds. This imbalanced pattern, also referred to as mutualism asymmetry, is a well-known trend in mutualistic associations as a whole, and it has been hypothesized that it could potentially determine the stability of associations over time and their level of specificity (Chomicki et al. 2020).

With respect to weevil specialists, plants involved in BSPM are often pollinated by diverse multi-species assemblages of weevil pollinators living in sympatry (Figures 3F, 6; Hotchkiss 1958; Ervik et al. 1999; Franz & Valente 2005; Núñez-Avellaneda et al. 2015; de Medeiros et al. 2019; Saunders 2020; Toon et al. 2020; Haran et al. 2021; 2022b; Auffray et al. 2022; Dao et al. 2023). A remarkable aspect of these assemblages is their stability over the range of their hosts (Núñez-Avellaneda et al. 2015; de Medeiros et al. 2019; Haran et al. 2020), in stark contrast to other brood-site pollination systems (Jousselin et al. 2008; McLeish & van Noort 2012). Sympatric weevil assemblages may coexist on the same host as they often develop on distinct ecological niches such as male or female inflorescences, as observed in the genera *Elaeidobius* and *Prosoestus* on *Elaeis guineensis* (Syed 1979; Alibert 1938) or in several cycad-associated weevils (Toon et al. 2020). Niche partitioning can also be observed within monoecious inflorescences, either on different tissues (de Medeiros et al. 2019; Figure 6) or in relation to distinct phenological anthesis stages (Haran et al. in prep.). Unexpectedly, these sympatric multi-species assemblages have been described in all plant-weevil mutualism systems, suggesting that they may be the rule rather than the exception. The prevalence of these assemblages seems specific to weevil pollinators; in other BSPM systems such as the fig/fig-wasp model, instances of multispecific assemblages of pollinators are only reported in about 10% of associations (Cruaud et al. 2012), even though usage of molecular data for wasp species delimitation has been increasing this number (Satler et al. 2022; Su et al. 2022). In comparison with other brood-site pollinator systems, weevils also present the greatest diversity of congeneric species associated with the same host: this is the case of the African oil Palm (*Elaeis guineensis*), which hosts no less than eight species of *Elaeidobius* on its male inflorescences (Compton et al. 2009; Haran et al. 2020a, 2021; Figure 6B). The exact role of these assemblages is not yet elucidated and several potential evolutionary processes have been proposed to explain their emergence. A first process is the lineage duplication induced by the fragmentation of the host range in relation to past environmental changes (such as climate oscillations), followed by the reconnection of host populations. This case has been highlighted in the *Elaeis*-derelomine system where past cycles of fragmentation of *Elaeis guineensis* populations into the upper and lower Guinean forest blocks would have led to the diversification of *Elaeidobius* and *Prosoestus* species (Haran et al. 2021). When populations reconnect after isolation, differences in generation times between the host plant and its pollinators appear to allow palm populations to exchange genes again while weevil populations are already too divergent and already correspond to distinct species. A second process results from independent past colonizations (historical host shifts) of a host by multiple lineages of weevils. This process is suggested by the structure of sympatric assemblages themselves, where weevil species in an assemblage belong to distinct and phylogenetically unrelated genera (Franz 2007a; de Medeiros et al. 2019; Toon et al. 2020;

Haran et al. 2022b). Interestingly these two processes are not mutually exclusive, as underlined by the example of the weevil community associated with the inflorescences of *Phoenix reclinata* (Arecaceae). Here, lineage duplication and independent colonizations led to the emergence of a sympatric assemblage of six species of brood-site pollinators (Haran et al. 2022b).

In plant-weevil BSPM, multi-species assemblages are not limited to weevils but also include a range of insects of various orders that develop on host inflorescences (Desmier de Chenon 1981; Búrquez et al. 1987; Ervik et al. 1999; Gottsberger 1999; Núñez-Avellaneda et al. 2015; de Medeiros et al. 2019; Toon et al. 2020; Auffray et al. 2022). It is very common, for example, that weevil-pollinated plants are also visited by sap beetles (Nitidulidae, Figure 6A). In Annonaceae they belong to the genus *Carpophilus*, in Arecaceae the most common genera are *Carpophilus*, *Microporum* and *Mystrops*, in cycads they belong to the genus *Carpophilus*, and in Cyclanthaceae they belong to the genus *Mystrops* (Desmier de Chenon 1981; Knudsen et al. 2001; Núñez-Avellaneda et al. 2005; Ratnayake et al. 2006; Franz 2007a; Núñez-Avellaneda et al. 2015; Toon et al. 2020). Sap beetles range from the most important pollinators in some plants to negligible in others (Ratnayake et al. 2006; Núñez-Avellaneda et al. 2015; Lau et al. 2017; de Medeiros et al. 2019; Toon et al. 2020; Carreño-Barrera et al. 2021). Indeed, the patterns observed in cycad-associated brood-site pollinators show that sap beetles, but also certain lineages belonging to other families of beetles (Boganiidae, Erotylidae and Tenebrionidae) or even other orders of insects (Lepidoptera and Thysanoptera) may have become brood-site pollinators of cycads (Toon et al. 2020). It is therefore not surprising that species of various groups of insects have independently colonized the same host and co-exist on it. In palms, multi-species assemblages sometimes even include generalist insects that have also been identified as pollinators. In some cases, bees are the most important pollinators with weevils being accessory (Núñez & Carreño, 2017; Bezerra et al. 2020), but more commonly generalist pollinators play a minor role (Desmier de Chenon 1981; Núñez-Avellaneda et al. 2015; de Medeiros et al. 2019). The latter echoes the fact that, whenever weevils are involved in these diverse assemblages, they generally play a predominant role in host pollination (Desmier de Chenon 1981; Ratnayake et al. 2006; Núñez-Avellaneda et al. 2015; Toon et al. 2020). The precise role of these secondary pollinators (brood site or not) is not clear, but a reduction of extinction risk by limiting dependence on a specialized mutualist has been suggested (Silberbauer-Gottsberger 1990; Chomicki et al. 2020).

### 2.3.3 Cryptic and closely related weevil species

The evolution of plant-weevil BSPM is a dynamic system with many instances of recent or ongoing speciation processes. This trend was first suggested thanks to the results of several morphological studies, which revealed that BSPM weevil genera often consist of diverse closely related species (Oberprieler 1996; Oberprieler 2004; Valente 2005; Franz & Valente 2005; de Medeiros & Vanin 2020; Hsiao & Oberprieler 2022). This trend has been confirmed by molecular studies, which have revealed additional layers of hidden diversity in the form of cryptic and closely related species often co-occurring on a single host (Downie & Williams 2009; Brookes et al. 2015; Nunes et al. 2018; de Medeiros & Farrell 2020; Haran et al. 2021, 2022b,b). All of these cryptic and closely related weevil species (Table S1) likely reflect recent speciation events. For example, dating analyses from two recent molecular studies have recovered numerous instances of recent speciation events (1.2 to 0.2 Mya old), which

may be associated with recent changes in paleoenvironments (Haran et al. 2021, 2022b). Morphological features enabling to distinguish these closely related species include subtle differences such as the location and size of certain setae on the male ventrites, the arrangement of setae on the elytra or the ratios on the male genitalia. The repeated occurrence of these cases of cryptic and closely related species highlight the need for detailed taxonomic studies to properly estimate the levels of specificity of plant-weevil BSPM systems.

#### 2.3.4 Trade offs in cost-benefit of interactions

The balance between costs and benefits in plant-weevil BSPM interactions has been investigated in several systems (e.g., Dufay & Anstett 2004; Franz 2004; Franz & Valente 2005; Nunes et al. 2018; Saunders 2020; Toon et al. 2020). Plants engaged in BSPM with weevils require cross-pollination for a large fruit set. Therefore, a major pressure they have experienced is pollen limitation, linked to inadequate pollen quality and/or quantity (Franz 2007a; Nunes et al. 2018), a widely observed condition for tropical flora (Tremblay et al. 2005; Wolowski et al. 2014). Under this pressure, the advantage of having specialized pollinators is believed to be high (Silberbauer-Gottsberger 1990), especially in the understory where small plants do not compete well for generalist pollinators because their floral displays are less conspicuous (Turner 2001). The benefit to the plant is clear when weevils develop only on male flowers or on non-reproductive tissues (e.g., live or decaying stems, bracts and petals), and therefore do not affect propagule production (Dufay & Anstett 2003; Dao et al. 2023). This case is widely encountered in “pollination by deceit” systems (Dufay & Anstett 2003) described in many palms (Arecaceae), where adult weevils search for male inflorescences where development of larval stages take place but are misled by female inflorescences producing similar attractants but no rewards. Such a benefit is, however, less evident when the flower buds or seeds are destroyed during the larval development of the weevils, and sometimes it even results in a sharp reduction in the fitness of the host. For example, brentids in the genus *Antliarhinus* are ovule parasites that can destroy up to 80% of the seeds of their cycad host *Encephalartos*. Despite this damage, it has been shown that adults contribute up to 10% to the cross-pollination of their host (Donaldson 1997). Similarly, the main pollinator of the palm *Syagrus coronata* is a specialist weevil of the genus *Anchylorhynchus* (Derelomini), which are known to be seed predators (de Medeiros et al. 2019). However, it has been recently shown that there is variation in the genus, with some species only being able to develop on aborted female flowers and therefore harmless to the plant (de Medeiros 2022 in press). Studies on these peculiar cases can reveal why such costly pollination systems are maintained when less costly ones, involving other specialist or generalist pollinators, are potentially available and how interactions evolve along the mutualism-antagonism spectrum (Fenster et al. 2004; de Medeiros et al. 2019).

In some cases, the detrimental effect of the larval development of weevil pollinators is mitigated by specific plant adaptations. In the dwarf palm (*Chamaerops humilis*), the process associated with fruit development leads to the termination of larval development on female inflorescences of *Derelomus chamaeropis*. Thus, the maintenance of the populations of this obligate pollinator relies solely on male inflorescence tissues, with no impact on seed sets (Dufay & Anstett 2004). Control of the detrimental effects of larval development may also include other organisms. For example, in the orchid *Dichaea cogniauxiana*, pollination

depends heavily on the activity of a baridine weevil developing as an ovule parasite. Normally, about 20% of fruits are lost to weevil larvae, but the activity of a parasitic wasp killing weevil larvae at an early stage can significantly reduce the fraction of seeds lost to the weevil (Nunes et al. 2018). Such tripartite interactions involving a plant, a pollinator and a parasitoid are probably overlooked, although they potentially play an important role in weevil-based BSPM systems.

Entirely detrimental interactions for the plants have also emerged within weevil lineages generally recognized as pollinators. The species involved have been labeled as ‘cheaters’ because they develop in the tissues of their hosts without providing pollination service (Franz, 2003, 2004; Franz & Valente, 2005; de Medeiros et al., 2019; de Medeiros & Farrell, 2020). In several plant-weevil BSPM, beneficial and detrimental weevil lineages coexist on the host inflorescence. For example, in the cyclanths (genus *Carludovica*) – weevil BSPM system, pollination is ensured by *Ganglionus* weevils (Derelomini) developing in the male flowers of the monoecious inflorescences. However, they coexist with another genus of derelomine weevils (genus *Systemotelus*), which predates *Carludovica* seeds while providing no pollination service. Since most derelomine weevils are brood pollinators, this cheater habit has therefore been inferred as a transformation from a beneficial to a detrimental one (Franz & Valente 2005). Such transitions of beneficial to detrimental life habits are a common feature of brood-pollinators system, and have been extensively documented in fig/fig-wasp and yucca/yucca-moth systems (e.g., Pellmyr et al. 1996; West et al. 1996; Segraves et al. 2005).

### 2.3.5 Evolutionary dynamics of host use

Since most of the plant groups actually associated with weevil brood pollinators are categorized as ancient tropical lineages (Annonaceae, Araceae, Arecaceae, cycads, Cyclanthaceae, Ebenaceae and Strelitziaceae; Franz & Valente 2005; Kress & Specht 2006; Toon et al. 2020), extant plant-weevil associations have sometimes been assumed to be ancient and possibly concurrent with the origin of plant lineages (Brenner et al. 2003; Oberprieler 2004; Franz 2006). However, such a pattern has not been backed up by any solid evidence, as all studies integrating knowledge of the fossil record and the results of molecular dating analyses support the hypothesis that weevil brood pollinator lineages began to diversify long after the origin of the plant lineages on which they specialize. For example, the estimated origin of palm-specialists from the Derelomini tribe (ca. 40 Mya; Haran et al. 2022b) significantly postdate the well-documented appearance of palms ca. 100 Mya (Baker & Couvreur 2013). Likewise, the phylogenetically diverse lineages of weevils that shifted on cycads (Oberprieler 2004) colonized them relatively recently (Downie et al. 2008; Hsiao & Oberprieler 2022), well after their origin more than 250 Mya (Condamine et al. 2015).

Weevils show a marked pattern of phylogenetic niche conservatism for the specific plant lineages they pollinate. Consistency of association with plant clades is observed at the tribe, genus or species-group level (Franz & Valente 2005; de Medeiros & Vanin 2020; Haran et al. 2021, 2022b; Figure 1). This consistency is particularly strong at the genus level for BSPM interactions (Table S1). In other words, when a BSPM interaction is established for a weevil species, it is very likely that congeneric species are also involved in such an interaction. An exception to this pattern has been reported in the genus *Cotithene*



(Derelomini) associated with *Evodianthus* (Cyclanthaceae) in Amazonia: only one weevil species in this genus is a brood site pollinator., Congeners visiting this cyclanth are not engaged in such relationship due to a size mismatch with the host flower morphology (Valente et al. 2019).

Unlike other brood-pollinators, weevils display a remarkable ability to shift to unrelated plant lineages at various evolutionary scales. A striking example of this pattern is seen in eastern hemisphere Derelomini with two parallel instances of secondary shifts from palms (Arecaceae) to Ebenaceae (genus *Euclea*; Haran et al. 2022b). Interestingly, such secondary shifts did not necessarily constitute evolutionary dead ends since they are sometimes followed by lineage diversification and even additional secondary shifts (Haran et al. 2022b). Other notable examples of host shifts include the parallel and independent colonization of the closely related palms *Elaeis guineensis* and *Elaeis oleifera* by two distinct lineages of derelomine weevils (Haran et al. 2021) and the independent colonization of the legume tree *Baikiaea insignis* (Fabaceae) by both Ochyromerini (two species of *Endaeus*) and Derelomini (*Lomederus ghesquierei*) (Marshall 1932, 1933). Overall, the evolutionary pattern of host use in weevils engaged in BSPM is much more dynamic than what is reported in other brood pollinators systems (Pellmyr 2003; Herre et al. 2008) and thus reinforces the interest of using weevils as an additional model system to investigate evolutionary dynamics of pollination mutualism in the tropics (Haran et al. 2022b).

### 3. Economic importance and conservation

Many plants engaged in BSPM with weevils have economic, cultural or ornamental values, features that have sometimes triggered the discovery and study of these mutualistic systems. Thanks to a better understanding of the plant-weevil BSPM systems, it was possible to achieve practical objectives. For example, a classic application in agriculture is the introduction of the derelomine weevil *Elaeidobius kamerunicus* from Africa to Asia, to improve pollination of cultivated African oil palm (*Elaeis guineensis*; Syed et al. 1982). The successful establishment of this weevil, also known as the ‘million dollar weevil’, has resulted in a dramatic increase in pollination rates and yields (Krantz & Poinar 2004; Li et al. 2019).

Many weevil pollinated palms are an important source of food, goods and building materials (e.g., *Acrocomia*, *Cocos*, *Euterpe*, *Oenocarpus*, *Phytelephas*, *Syagrus*; Núñez-Avellaneda et al. 2015; de Medeiros et al. 2019; Bezerra et al 2020; Carreño-Barrera et al 2021; Auffray et al. 2022). In tropical and subtropical regions, wild fruits and leaves of Annonaceae (e.g., *Annona senegalensis*), Clusiaceae (e.g., *Pentadesma butyracea*) and Fabaceae (e.g., *Piliostigma reticulatum*) for which pollination by weevils is indicated based on PNC are commonly consumed by local populations or have a cosmetic or medical use (e.g., Babajide et al. 2008; Traoré et al. 2011; Dao et al. 2023). In cycad lineages primarily pollinated by weevils (Toon et al. 2020), the high ornamental and subsequent economic value of many cycad species has also led to over-collection (Mankga & Yessoufou 2017). In these groups, reproductive failure, among habitat loss and medicinal use, has been identified as one of the main drivers of population decline, highlighting the importance of detailed investigations of their pollination ecology. More generally, while plant-weevil BSPM probably emerged to balance some pollination constraints in tropical biomes, it is also a risky evolutionary path if the specialized pollinators become rare or extinct. Yet too little is known

about the impact of pollinator population dynamics on the reproductive success and long-term survival of the plants involved (Toon et al. 2020).

## 4. Oversights and estimates of global richness

### 4.1 The inconspicuousness of plant-weevils pollination interaction

One may wonder why such widespread mutualistic interactions have taken so long to be described and recognized. Pollination ecology is a dynamic discipline and pollination syndromes have been explored for a significant range of tropical flora (Momose et al. 1998; Schleuning et al. 2012; Vizentin-Bugoni et al. 2018), several causes may be invoked for the delays in description of plant-weevil BSPM systems.

First, it can be explained by common misconceptions about plant and weevil biology. For example, in the cycad/molytine and palm/derelomine systems, massive aggregations of weevils on inflorescences are highly suggestive of pollination activity. Although these associations have been known for a long time (e.g., von Martius 1823), the common misconception that palms and cycads are primarily anemophilous groups has hampered research on brood-site mutualism in these systems for decades (Von Martius 1823; Lepesme 1947; Bondar 1943; Henderson 1986; Dufay & Anstett 2003; Barfod et al 2011; Toon et al. 2020). These flower visitors have long been considered parasites, providing no pollination service (Bondar 1943; Lepesme 1947), consistent with antagonistic interactions generally reported in weevils (Oberprieler et al. 2007). In a striking example of how prior misconceptions can prevent discovery, a study characterizing floral visitors of *Butia catarinensis* by collecting only during the day and using sweep nets failed to find a relevant number of beetles in female flowers (Rosa 2000). However, B. de Medeiros visited the same locality less than 10 years later and found hundreds of individuals of *Anchylorhynchus eriospathae* hiding at the base of inflorescences during the day (de Medeiros & Vanin 2020). These weevils breed on female flowers (de Medeiros et al. 2014) and adequate collection and observation methods could have recorded them visiting active female flowers for oviposition. To date, the cycad and palm-weevil associations remain by far the best studied plant-weevil BSPM systems, in particular because the abundance of these insects on the inflorescences and their specialized nature greatly facilitates the identification of the interactions (Franz & Valente 2005).

Second, the inconspicuous nature of many interactions hampered and slowed down the discovery of several plant-weevil BSPM interactions. For example, in the case of the Annonaceae – *Endaeus* mutualisms, the flowers are isolated and therefore the interactions are thus far less conspicuous. In these systems, weevils typically reach flowers in very low numbers, sometimes at a very specific time of the circadian flowering cycle (early in the morning or late in the evening), and may be hidden or “enclosed” in floral chambers during most of the day (Ratnayake et al. 2006; Lau et al. 2017; Saunders 2020) or present on the flower only at night and notably absent during the day (Ratnayake et al. 2006). When not enclosed in flowers, *Endaeus* weevils and related weevil genera are usually collected while flying over the canopy (Kojima & Morimoto 1995; Sprick and Florent 2018), using light traps

(Kojima & Morimoto 1995; Haran et al. 2022a) or by beating plants only used as shelters, generally in low numbers (Kojima & Morimoto 1995). Therefore, unraveling BSPM interactions between weevils and their host plants is usually a difficult task. To illustrate this point, a striking example of inconspicuous interactions is found in the *Anthurium* (Araceae) / Derelomini mutualism. In this system, only a few individuals of the derelomine *Cyclanthura carinata* reach *Anthurium* inflorescences at night during the 1-2 weeks of anthesis, and further remain hidden in areas protected by the spathe, leaving the inflorescences apparently empty. This inconspicuousness probably explains why this obligate pollination mutualistic system has been overlooked for decades (Franz and Valente 2005; Franz 2007b).

## 4.2 An underestimated diversity of species and interactions

How much of the diversity of weevil-based pollination is still unknown? A basic calculation of the discrepancies between the number of species whose biology is reported and the total richness of their clade suggests that only a small portion of the interactions are at least superficially known (Figure 2). There is also a strong geographic and taxonomic bias in the description of plant-weevil BSPM systems.

For example, host plant identity information is available for most Neotropical derelomine genera (Franz & Valente 2005). By comparison, host records are only available for five of the 13 Paleotropical genera of Derelomini (Alonso-Zarazaga & Lyal 1999; Franz & Valente 2005). The estimated number of derelomine species overall (ca. 300 species; Caldara et al. 2014; de Medeiros & Vanin 2020; Haran et al. 2020, 2022a) pales in comparison with the diversity estimates of their main hosts, because the species diversity in the palm family (Arecaceae) is estimated at about 2500 species (Dransfield et al. 2008; Baker & Couvreur 2013). Since pollination ecology and pollinators have only been identified for a small fraction of extant palm species, we can hypothesize that dozens – if not hundreds – of palm-weevil BSPM systems remain to be discovered. Additionally, even though Derelomini are best known for their associations with palms, recent research has unraveled that some of them are also associated with plants other than palms. For example, in the Southern African derelomine fauna, the recently described genus *Ebenacobius* has been found associated with Ebenaceae and other basal angiosperms. In this genus, no less than 14 new species have been recently described (Haran et al. 2022a) and at least four others are awaiting description. Moreover, based on the inferred pattern of PNC in BSPM weevils, Franz & Valente (2005) and Franz (2006) suggested that most of the ca. 230 species of Carludovicoideae (Cyclanthaceae) and a significant part of the ca. 800 known *Anthurium* (Araceae) species in the Neotropical region are pollinated by derelomine weevils. In cycad-weevil BSPM systems, the range of interactions reported here only reflects our current state of knowledge, and many cycads species have been insufficiently studied and many interactions remain to be described (Toon et al. 2020).

The general lack of knowledge on plant-weevil BSPM systems can be exemplified by our understanding of Ochyromerini. Pollination activity in this tribe has been only reported between a few species of *Endaenidius* and *Endaeus* and their Annonaceae hosts and BSPM has been only formerly identified in one system (*Annona senegalensis* - *Endaeus* spp.; Lau



et al. 2017; Saunders 2020; Dao et al. 2023). Importantly, Ochyromerini has been determined to be the dominant weevil guild circulating above the canopy in a Southeast Asian dipterocarp forest (Sprick & Floren 2018), with over 130 undescribed ochyromerine species recorded on Borneo island alone. Similar species richness was also observed in Cameroon using light traps (Haran unpublished). Finally, a multitude of undescribed Ochyromerini species (mostly belonging to the genus *Endaeus*) have been reported by several authors in Asia and Africa (Hustache 1924; Marshall 1926; Oberprieler 1993; Momose et al. 1998), and it can be suspected that this group could play an important role in the pollination of Paleotropical rainforests.

In the Annonaceae, a pantropical clade with more than 2400 species, cantharophily seems to be the most common pollination syndrome (Momose et al. 1998; Gottsberger 1999; Saunders 2020). The Neotropical species in this plant family sum up to more than 950 species (Maas et al. 2011), and it is in this region that cantharophilous flowers involving weevil pollination are the most widespread (Gottsberger 1999; Silberbauer-Gottsberger et al. 2003; Saravy et al. 2021). This set of interactions requires further investigations in this region since the weevil lineages involved remain unidentified and possibly include undescribed lineages (the pollinators of Annonaceae reported (Ochyromerini) are exclusively paleotropical; Alonso-Zarazaga & Lyal 1999).

Our general overview of weevil-based pollination shows that several additional pantropical families of angiosperms are – or at least can be considered as partially – pollinated by weevils (Table S1), with only a small fraction of these interactions having been uncovered so far. For example, in Clusiaceae, Ebenaceae, Lecythidaceae, Myristicaceae and Sapotaceae, BSPM by weevils have been only indicated for a small number of species; however there are likely many more of such interactions because cantharophilous flowers are known in several genera of these families and the associated weevil lineages also contain a significant number of undescribed species (Oberprieler 1993; Caldara et al. 2014; Haran et al. 2022b). In some cases, the weevil diversity also exceeds that of the known potential hosts; for example, 25 species of the Australian weevil genus *Elleschodes* are known (Armstrong & Irvine 1990; Caldara et al. 2014; Pullen et al. 2014), of which several species are known to pollinate the only three species of *Eupomatia* (Eupomoatiaceae) so it is likely that this genus also pollinates other plant families. Going further, the Orchidaceae-Baridinae system highlights the fact that even weevils lineages that are not known to be pollinators and that lack the associated typical morphological features can also be engaged in BSPM (Nunes et al. 2018), with a potential rich array of interactions (van der Cingel 2001). From the host plant perspective, the orchid species studied (*Dichaea cogniauxiana*) also lacks features associated with weevil pollination in other groups (i.e., it has a standard orchid floral morphology and phenology; Nunes et al. 2018). Such examples greatly expand the potential spectrum of brood-site mutualistic interactions between weevils and plants in tropical biomes and call for a complete shift in our perception of the role of these beetles in the functioning of tropical ecosystems.

A preliminary estimate of the expected diversity of plant-weevil BSPM systems can be made based on the discrepancies between the number of systems described and inferred based on PNC (Figure 2; see Appendix 1). The calculation of the ratios for a selection of systems (see figure 2) shows that the expected total diversity of species involved is on average eight times richer than the number of species inferred from PNC and about 17 times

richer than the number of species with experimentally-verified BSPM. Based on the species counts obtained in this review (see section 1), it is estimated that ca. 2000 plant species could be engaged in BSPM with weevils globally. Similarly, the species richness of weevils involved in these interactions could be around 2500 species. Thus, plant-weevil interactions probably represent the most speciose case of brood-site pollination mutualism, exceeding the diversity of other BSPM species-rich groups (e.g., [Lopez-Vaamonde et al. 2009](#); [Cruaud et al. 2010](#)).

## 5. Concluding remarks: a road map for future research

### 5.1 Searching for new plant-weevil BSPM systems

Accumulating evidence for the role of weevils as specialized brood-pollinators of plants is changing our perspective on the role of this super-diverse group of beetles in tropical ecosystems and beyond. As such, it is time to expand our understanding of pollination ecology and to research and test this mutualism in a range of plant and weevil groups that have been overlooked. We show in this review that some common features associated with BSPM between plants and weevils can help identify novel systems. That said, we also show that these mutualistic interactions can exist between lineages that do not exhibit these characteristics at all. In this regard, we should also change our perception of the structure of these interactions themselves. In light of this reasoning, a wide range of plant-weevil interactions should be re-evaluated. A few selected “priority” examples include the potential role of several ancient weevil groups (Belidae, Caridae, Cimberididae and Nemonychidae) that visit and breed in gymnosperm cones. With the exception of cycads, they are not considered pollinators ([Oberprieler et al. 2007](#)), but these cases have never been verified in detail. In the Neotropics, *Oxycorynus* species (Belidae, Oxycoryninae) are known to visit the male and female flowers of parasitic plants of the genus *Lophophytum* (Balanophoraceae), the larvae developing in the inflorescences ([Ferrer et al. 2011](#)).

Plant-weevil BSPM systems can also be expected in temperate regions, where the predominance of anemophily and entomophily based on generalist pollinators might have precluded investigations of more specialized systems. In some Western Palearctic willows (Salicaceae), flowering cycles show intriguing spikes of volatiles which are not correlated with the visits of pollinators considered as efficient ([Jürgens et al. 2014](#)). Interestingly, a part of these compounds (4-1 dimethoxybenzene) is also produced by several South American palms for the specific attraction of beetles, including weevils ([Knudsen et al. 2001](#); [Núñez-Avellaneda et al. 2005](#)). This case deserves special attention since the male and female inflorescences of these dioecious trees also harbor the larval stages of several typical genera of flower weevils (*Acalyptus*, *Dorytomus*, *Ellescus*; [Hoffmann 1958](#)).

While this review focuses specifically on BSPM, it is also important to highlight that weevils may be involved in the pollination of generalist plants, and may not necessarily use breeding sites as rewards. One group within Curculionidae that deserves attention is the subfamily Baridinae. Here we found only one example of a demonstrated effective pollinator (*Montella*, a brood-site pollinator of orchids; [Nunes et al. 2018](#)), but this mainly Neotropical

taxon harbors an extreme diversity with more than 4,000 described species, many of which are found in flowers as adults (Prena et al. 2014; B. de Medeiros pers. obs.). Due to their unresolved phylogeny, complex taxonomy and very high number of undescribed taxa (Davis 2011), they are rarely identified beyond the subfamily or tribe level in pollination studies, preventing accumulation of knowledge. A great diversity of baridine pollinators, with or without brood sites as rewards, remains to be discovered. Interactions may also occur in even more unexpected groups of flower-associated weevils. In orchids, a species endemic to the Cayman Islands (*Myrmecophila thomsoniana*) is apparently pollinated by the weevil *Lachnopus vanessablockae* (Curculionidae: Entiminae; Girón et al. 2018) with pollinia transferred by contact of the elytra on the stigmatic surface of flowers. As typical for entimines, the larvae of this species reproduce in the soil on the root systems of plants (Girón et al. 2018), so it is unlikely to be a case of BSPM. These cases demonstrate the need for a better understanding of the evolution of pollinators in weevils more broadly, and the contexts leading to BSPM more specifically.

## 5.2 *Priority research axes*

### 5.2.1 A plea for description of systems

A better understanding of the diversity and functioning of plant-weevil BSPM systems requires detailed studies of the corresponding interactions ([Fenster et al. 2004](#)), but also of the identity and boundaries of the species involved. To date, too many plant-weevil BSPM studies have a focus on plants and poorly account for pollinator identity. This is partly due to the complexity of the taxonomy in these groups (species complexes, small sized species) but also to taxonomic shortfalls. As [Franz & Engel \(2010\)](#) pointed out, there are no shortcuts to understanding the evolution and diversification of plant-weevil interactions and only detailed case studies can provide relevant insights. Taxonomic descriptions are essential prerequisites for any study of pollination in the highly specialized plant-weevil BSPM systems, and this effort must be made with the relevant expertise of trained taxonomists using integrative approaches based on morphology and molecular tools ([Toon et al. 2020](#); [Engel et al. 2021](#)). When expertise is not available, we strongly recommend producing at least one DNA barcode sequence for each weevil morphospecies involved in a system and depositing it on international databases (e.g., [de Medeiros et al. 2019](#)), and using images in addition to names to document weevil flower visitors (e.g., [Núñez-Avellaneda & Rojas-Robles, 2008](#); [Núñez et al. 2015](#); [Guerrero-Olaya 2017](#); [de Medeiros et al. 2019](#)). The associated voucher specimens should also be deposited in permanent collections of museums or institutions. These two procedures will make it possible to considerably improve species description rates, but also the knowledge on their ecology and distribution.

### 5.2.2 Evolutionary trajectories : when, why and how?

Although preliminary insights on the emergence and maintenance of plant-weevil BSPM are readily available, a significant number of evolutionary aspects of these interactions remain to be investigated. A first aspect lies in the age of these interactions. Reliable fossils of weevil lineages engaged in BSPM are noticeably absent, precluding making precise inferences about the ages of these mutualisms. The reasons why so many ancient and more derived plant lineages have engaged in BSPM with weevils can be elucidated if their emergence coincides with specific past biogeographical or climatic events. More generally, plant-weevil BSPM systems offer a rich and diverse range of systems considered as potential models for understanding the conditions favoring: (i) the shift from detrimental to beneficial interactions, and the role of tripartite interactions in driving these interactions, (ii) transitions between pollination syndromes such as between BSPM, other kinds of entomophily, anemophily, intermediate or ambophilous systems, and (iii) whether there is a general trend towards higher frequency of specialized pollinators engaging in BSPM in tropical biomes. Interestingly, almost all plant lineages exhibiting cantharophilous and thermogenic flowers are widely associated with weevils. It remains to be determined to what extent the association with weevils specifically was a key factor in the emergence of cantharophily in general (Bernhardt 2000). Other poorly documented aspects of these interactions include the potential association with specific microorganisms for colonization of breeding sites and whether plants alter their physiology to favor the development of the hosted pollinators. Finally, an unexplored aspect of these specialized interactions lies in their evolution in the context of current human-induced ecosystem disruption. The present multifaceted changes affecting the biosphere and insects in particular (Potts et al. 2016; Vanderplanck et al. 2021; Wagner et al. 2021) provide a context for an assessment of the resilience or vulnerability of these systems.

## Acknowledgements

We thank the editor and the four reviewers for their very detailed and constructive comments on an earlier version of this study. We would like to thank all the researchers who have contributed to the recognition of the importance of plant-weevil pollination mutualism through detailed and time-consuming descriptions of the taxonomy, ecology and physiology of these systems. We thank Z. Dao (Joseph Ki-Zerbo University, Ouagadougou, Burkina Faso), M. Gueye (UCAD, Dakar, Senegal) and R. G. Oberprieler (CSIRO, Canberra, Australia) for providing images of weevils and host plants. This work was supported by recurring funding from CIRAD (Julien Haran) and INRAE (Gael J. Kergoat). Bruno de Medeiros was funded by a STRI postdoctoral fellowship while writing this manuscript.

## Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article. Gael J. Kergoat is one of the *PCI Ecology* recommenders.

## References

- Adam H, Collin M, Richaud F, Beulé T, Cros D, Omoré A, Nodichao L, Nouy B & Tregear JW. 2011. Environmental regulation of sex determination in oil palm: current knowledge and insights from other species. *Annals of Botany* 108(8): 1529–1537. <https://doi.org/10.1093/aob/mcr151>
- Alibert H. 1938. Étude sur les insectes parasites du palmier à huile au Dahomey. *Journal d'agriculture traditionnelle et de botanique appliquée* 207: 745–773.
- Alonso-Zarazaga MA & Lyal CHC. 1999. *A World Catalogue of Families and Genera of Curculionoidea (Insecta: Coleoptera) (Excepting Scolytidae and Platypodidae)*. Entomopraxis S.C.P., Barcelona, pp. 1–315.
- Alonso-Zarazaga MA. 2007. Acalyptini C. G. Thomson 1859, correct name for a tribe in Curculioninae (Coleoptera: Curculionidae) and a comment on nomenclatural terminology. *The Coleopterists Bulletin* 61(4): 559. [https://doi.org/10.1649/0010-065X\(2007\)61\[559:ACGTCN\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2007)61[559:ACGTCN]2.0.CO;2)
- Anderson RS. 1993. Weevils and plants: phylogenetic versus ecological mediation of evolution of host plant associations in Curculioninae (Coleoptera: Curculionidae). *Memoirs of the Entomological Society of Canada* 165: 197–232. <https://doi.org/10.4039/entm125165197-1>
- Armstrong JE & Irvine AK. 1989a. Flowering, sex ratios, pollen-ovule ratios, fruit set, and reproductive effort of a dioecious tree, *Myristica insipida* (Myristicaceae), in two different rain forest communities. *American Journal of Botany* 76(1): 74–85. <https://doi.org/10.2307/2444776>
- Armstrong JE & Irvine AK. 1989b. Floral biology of *Myristica insipida* (Myristicaceae), a distinctive beetle pollination syndrome. *American Journal of Botany* 76(1): 86–94. <https://doi.org/10.2307/2444777>
- Armstrong JE & Irvine AK. 1990. Functions of staminodia in the beetle-pollinated flowers of *Eupomatia laurina*. *Biotropica* 22(4): 429–431. <https://doi.org/10.2307/2388563>
- Armstrong JE. 1997. Pollination by deceit in nutmeg (*Myristica insipida*, Myristicaceae): floral displays and beetle activity at male and female trees. *American Journal of Botany* 84(9): 1266–1274. <https://doi.org/10.2307/2446051>
- Anderson R. 1995. An evolutionary perspective on diversity in Curculionoidea. *Memoirs of the Entomological Society of Washington* 14: 130–114.
- Anstett M-C. 1999. An experimental study of the interaction between the dwarf palm (*Chamaerops humilis*) and its floral visitor *Derelomus chamaeropsis* throughout the life cycle of the weevil. *Acta Oecologica* 20(5): 551–558. [https://doi.org/10.1016/S1146-609X\(00\)86622-9](https://doi.org/10.1016/S1146-609X(00)86622-9)



1200 Auffray T, Frérot B, Poveda R, Louise C & Beaudoin-Ollivier L. 2017. Diel patterns of activity  
 1201 for insect pollinators of two oil palm species (Arecaceae). *Journal of Insect*  
 1202 *Science* 17(2): 1–6. <https://doi.org/10.1093/jisesa/iex018>  
 1203

1204 Auffray T, Montufar R, Uquillas SXP, Barragan A, Pincebourde S, Gibernau M & Dangles O.  
 1205 2022. Fine-scale temporal dynamics of flower visitors sheds light on insect-assemblage  
 1206 overlap between sexes in a dioecious Ecuadorian palm. *Biotropica* 55(1): 256–267.  
 1207 <https://doi.org/10.1111/btp.13182>  
 1208

1209 Babajide OJ, Babajide OO, Daramola AO & Mabusela WT. 2008. Flavonols and an  
 1210 oxychromonol from *Piliostigma reticulatum*. *Phytochemistry* 69(11): 2245–2250.  
 1211 <https://doi.org/10.1016/j.phytochem.2008.05.003>  
 1212

1213 Baker WJ, Couvreur TL. 2013. Global biogeography and diversification of palms sheds light  
 1214 on the evolution of tropical lineages. I. Historical biogeography. *Journal of Biogeography*  
 1215 40(2): 274–285. <https://doi.org/10.1111/j.1365-2699.2012.02795.x>  
 1216

1217 Barbosa CM, Maia ACD, Martel C, Regueira JCS, Navarro AF, Raguso RA, Milet-Pinheiro P  
 1218 & Machado IC. 2020. Reproductive biology of *Syagrus coronata* (Arecaceae): sex-biased  
 1219 insect visitation and the unusual case of scent emission by peduncular bracts. *Plant Biology*  
 1220 23(1): 100–110. <https://doi.org/10.1111/plb.13162>  
 1221

1222 Barfod AS, Hagen M & Borchsenius F. 2011. Twenty-five years of progress in understanding  
 1223 pollination mechanisms in palms (Arecaceae). *Annals of Botany* 108(8): 1503–1516.  
 1224 <https://doi.org/10.1093/aob/mcr192>  
 1225

1226 Bawa KS. 1990. Plant-pollinator interactions in tropical rain forests. *Annual Review of*  
 1227 *Ecology and Systematics* 21: 399–422.  
 1228 <https://doi.org/10.1146/annurev.es.21.110190.002151>  
 1229

1230 Benton MJ. 2010. The origins of modern biodiversity on land. *Philosophical Transactions of*  
 1231 *the Royal Society B Biological Sciences* 365(1558): 3667–3679.  
 1232 <https://doi.org/10.1098/rstb.2010.0269>  
 1233

1234 Bernhardt P. 2000. Convergent evolution and adaptive radiation of beetle-pollinated  
 1235 angiosperms. *Plant Systematics and Evolution* 222: 293–320.  
 1236 <https://doi.org/10.1007/BF00984108>  
 1237

1238 Bernays EA & Chapman RE. 1994. *Host selection by phytophagous insects*. Chapman &  
 1239 Hall, New York, pp. 1–199.  
 1240

1241 Bezerra LA, Campbell AJ, Brito TF, Menezes C & Maués MM. 2020. Pollen loads of flower  
 1242 visitors to açai palm (*Euterpe oleracea*) and implications for management of pollination  
 1243 services. *Neotropical Entomology* 49:482–490. <https://doi.org/10.1007/s13744-020-00790-x>  
 1244

1245 Bondar GG. 1941. Notas Entomológicas da Baía. VII. *Revista de Entomologia* 12: 268–303.  
 1246

1247 Bondar GG. 1943. Notas Entomológicas da Baía. XIII. *Revista de Entomologia* 14: 337–388.



1248  
1249 Borges RM, Somanathan H & Kelber A. 2016. Patterns and processes in nocturnal and  
1250 crepuscular pollination services. *The Quarterly Review of Biology* 91(4): 389–418.  
1251 <https://doi.org/10.1086/689481>  
1252  
1253 Brenner ED, Stevenson DW & Twigg RW. 2003. Cycads: evolutionary innovations and the  
1254 role of plant-derived neurotoxins. *Trends in Plant Science* 8(9): 446–452.  
1255 [https://doi.org/10.1016/S1360-1385\(03\)00190-0](https://doi.org/10.1016/S1360-1385(03)00190-0)  
1256  
1257 Brookes DR, Hereward JP, Terry LI & Walter GH. 2015. Evolutionary dynamics of a cycad  
1258 obligate pollination mutualism – pattern and process in extant *Macrozamia* cycads and their  
1259 specialist thrips pollinators. *Molecular Phylogenetics and Evolution* 93: 83–93.  
1260 <https://doi.org/10.1016/j.ympev.2015.07.003>  
  
1261 Búrquez A, Sarukhán KJ & Pedroza AL. 1987. Floral biology of a primary rain forest palm,  
1262 *Astrocaryum mexicanum* Liebm, *Botanical Journal of the Linnean Society* 94(4): 407–419.  
1263 <https://doi.org/10.1111/j.1095-8339.1987.tb01058.x>  
  
1264 Caldara R, Franz NM, Oberprieler RG. 2014. *Curculioninae latreille, 1802*. In: Leschen RAB  
1265 & Beutel RG, eds. Handbook of Zoology. Coleoptera, Beetles – Morphology and  
1266 Systematics. De Gruyter, Berlin, pp. 589–628.  
1267  
1268 Carreño-Barrera J, Maia ACD, Colombo CA & Núñez-Avellaneda LA. 2021. Co-pollination,  
1269 constancy, and efficiency over time: small beetles and the reproductive success of  
1270 *Acrocomia aculeata* (Arecaceae) in the Colombian Orinoquia. *Botany Letters* 168(3): 395–  
1271 407. <https://doi.org/10.1080/23818107.2021.1893215>  
1272  
1273 Carreño-Barrera J, Núñez-Avellaneda LA, Sanín MJ & Maia ACD. 2020. Orchestrated  
1274 flowering and interspecific facilitation: Key factors in the maintenance of the main pollinator  
1275 of coexisting threatened species of andean wax palms (*Ceroxylon* spp.). *Annals of the*  
1276 *Missouri Botanical Garden* 105(3): 281–299. <https://doi.org/10.3417/2020590>  
1277  
1278 Chomicki G, Kiers ET & Renner SS. 2020. The evolution of mutualistic dependence. *Annual*  
1279 *Review of Ecology, Evolution, and Systematics* 51: 409–432.  
1280 <https://doi.org/10.1146/annurev-ecolsys-110218-024629>  
1281  
1282 Compton SG, Grehan K & van Noort S. 2009. A fig pollinated by three or more species of  
1283 agaonid fig wasps. *African Entomology* 17(2): 215–222.  
1284 <https://doi.org/10.4001/003.017.0212>  
1285  
1286 Condamine FL, Nagalingum NS, Marshall CR & Morlon H. 2015. Origin and diversification of  
1287 living cycads: a cautionary tale on the impact of the branching process prior in Bayesian  
1288 molecular dating. *BMC Evolutionary Biology* 15: 65. [https://doi.org/10.1186/s12862-015-](https://doi.org/10.1186/s12862-015-0347-8)  
1289 [0347-8](https://doi.org/10.1186/s12862-015-0347-8)  
1290  
1291 Crowson RH. 1991. *Relationships to cycads*. In: Zunino M, Belles X & Blas M, eds.  
1292 *Advances in Coleopterology*. European Association of Coleopterology, pp. 13–28.  
1293

1294 Cruaud A, Jabbour-Zahab R, Genson G, Cruaud C, Couloux A, Kjellberg F, Van Noort S &  
 1295 Rasplus J-Y. 2010. Laying the foundations for a new classification of Agaonidae  
 1296 (Hymenoptera: Chalcidoidea), a multilocus phylogenetic approach. *Cladistics* 26(4): 359–  
 1297 387. <https://doi.org/10.1111/j.1096-0031.2009.00291.x>  
 1298  
 1299 Cruaud A, Cook J, Da-Rong Y, Genson G, Jabbour-Zahab R, Kjellberg F, Pereira RAS,  
 1300 Rønsted N, Santos-Mattos O, Savolainen V, Ubaidillah R, van Noort S, Yan-Qiong P &  
 1301 Rasplus J-Y. 2012. *Fig-fig wasps mutualism: the fall of the strict cospeciation paradigm?* In:  
 1302 Patiny S, ed. *Evolution of Plant-Pollinator Interactions*. Cambridge University Press, pp. 68–  
 1303 102.  
 1304  
 1305 Dao ZA, Romba R, Jaloux B, Haran J, Ouedraogo A & Gnankine O. 2023. Pollination  
 1306 syndrome of the African custard apple (*Annona senegalensis* Pers.) reveals reliance on  
 1307 specialized brood-site weevil pollinators in Annonaceae. Preprint.  
 1308 <https://doi.org/10.21203/rs.3.rs-1933322/v1>  
 1309  
 1310 Davis SR. 2011. Delimiting baridine weevil evolution (Coleoptera: Curculionidae: Baridinae).  
 1311 *Zoological Journal of the Linnean Society* 161(1): 88–156. [https://doi.org/10.1111/j.1096-](https://doi.org/10.1111/j.1096-3642.2010.00619.x)  
 1312 [3642.2010.00619.x](https://doi.org/10.1111/j.1096-3642.2010.00619.x)  
 1313  
 1314 Davis SR & Engel MS. 2010. Antiquity and evolution of prosternal horns in baridine weevils  
 1315 (Coleoptera: Curculionidae). *Journal of Paleontology* 84(5): 918–926.  
 1316 <https://doi.org/10.1666/09-160.1>  
 1317  
 1318 de Medeiros BAS. 2022. *The natural history of conflict and cooperation in a community of*  
 1319 *palm flower insects*. In: XX, XX, eds. Barro Colorado Island Centennial Book. Barro  
 1320 Colorado Island Centennial Book. In press, pp. XX–XX. *in press*  
 1321  
 1322 de Medeiros BAS & Farrell BD. 2020. Evaluating insect-host interactions as a driver of  
 1323 species divergence in palm flower weevils. *Communications Biology* 3: 749.  
 1324 <https://doi.org/10.1038/s42003-020-01482-3>  
 1325  
 1326 de Medeiros BAS, de Cássia Bená D & Vanin SA. 2014. *Curculio Curculis lupus*: biology,  
 1327 behavior and morphology of immatures of the cannibal weevil *Anchylorhynchus eriospathae*  
 1328 G.G. Bondar, 1943. *PeerJ* 2:e502. <https://doi.org/10.7717/peerj.502>  
 1329  
 1330 de Medeiros BAS & Núñez-Avellaneda LA 2013. Three new species of *Anchylorhynchus*  
 1331 Schoenherr, 1836 from Colombia (Coleoptera: Curculionidae; Curculioninae; Acalyptini).  
 1332 *Zootaxa* 3636(2): 394–400. <https://doi.org/10.11646/zootaxa.3636.2.10>  
 1333  
 1334 de Medeiros BAS, Núñez-Avellaneda LA, Hernandez AM & Farrell BD. 2019. Flower visitors  
 1335 of the licuri palm (*Syagrus coronata*): brood pollinators coexist with a diverse community of  
 1336 antagonists and mutualists. *Biological Journal of the Linnean Society* 126(4): 666–687.  
 1337 <https://doi.org/10.1093/biolinnean/blz008>  
 1338  
 1339 de Medeiros BAS & Vanin SA. 2020. Systematic revision and morphological phylogenetic  
 1340 analysis of *Anchylorhynchus* Schoenherr, 1836 (Coleoptera, Curculionidae: Derelomini).  
 1341 *Zootaxa* 4839(1): 1–98. <https://doi.org/10.11646/zootaxa.4839.1.1>

- Desmier de Chenon R. 1981. *New data on the entomophil pollination of oil palm in West Africa*. Technical report, Socfindo, Medan, Indonesia, pp. 1–47.
- Dilcher D. 2000. Toward a new synthesis: major evolutionary trends in the angiosperm fossil record. *Proceedings of the National Academy of Sciences of the U.S.A.* 97(13): 7030–7036. <https://doi.org/10.1073/pnas.97.13.7030>
- Donaldson JS. 1997. Is there a floral parasite mutualism in cycad pollination? The pollination biology of *Encephalartos villosus* (Zamiaceae). *American Journal of Botany* 84(10): 1398–406. <https://doi.org/10.2307/2446138>
- Downie DA, Donaldson JS & Oberprieler RG. 2008. Molecular systematics and evolution in an African cycad-weevil interaction: Amorphocerini (Coleoptera: Curculionidae: Molytinae) weevils on *Encephalartos*. *Molecular Phylogenetics and Evolution* 47(1): 102–116. <https://doi.org/10.1016/j.ympev.2008.01.023>
- Downie DA & Williams JG. 2009. Population structure of *Porthetes hispidus* (Coleoptera: Curculionidae), a pollinator of the African cycad *Encephalartos friderici-guilielmi*. *Annals of the Entomological Society of America* 102(6): 1126–1134. <https://doi.org/10.1603/008.102.0622>
- Dransfield J, Uhl NW, Asmussen CB, Baker WJ, Harley M.M. & Lewis CE. 2008. *Genera Palmarum: The evolution and classification of palms*. Kew Publishing, Kew, pp. 1–732.
- Dufaÿ M & Anstett MC. 2003. Conflicts between plants and pollinators that reproduce within inflorescences: evolutionary variations on a theme. *Oikos* 100(1): 3–14. <https://doi.org/10.1034/j.1600-0706.2003.12053.x>
- Dufaÿ M & Anstett MC. 2004. Cheating is not always punished: killer female plants and pollination by deceit in the dwarf palm *Chamaerops humilis*. *Journal of Evolutionary Biology* 17(4): 862–868. <https://doi.org/10.1111/j.1420-9101.2004.00714.x>
- Dufaÿ M, Hossaert-McKey M & Anstett MC. 2003. When leaves act like flowers: how dwarf palms attract their pollinators. *Ecology Letters* 6(1): 28–34. <https://doi.org/10.1046/j.1461-0248.2003.00382.x>
- Eberhard WG & Garcia-C JM. 2000. Ritual jousting by horned *Parisoschoenus expositus* weevils (Coleoptera, Curculionidae, Baridinae). *Psyche: A Journal of Entomology* 103: 55–84. <https://doi.org/10.1155/2000/16361>
- Eberhard WG, Garcia JM & Lobo J. 2000. Size-specific defensive structures in a horned weevil confirm a classic battle plan: avoid fights with larger opponents. *Proceedings of the Royal Society B: Biological Sciences* 267(1448): 1129–1134. <https://doi.org/10.1098/rspb.2000.1118>
- Engel MS, Ceríaco LMP, Daniel GM, Dellapé PM, Löbl I, Marinov M, Reis RE, Young MT, Dubois A, Agarwal I, Lehmann P, Alvarado M, Alvarez N, Andreone F, Araujo-Vieira K,

Ascher JS, Baêta D, Baldo D, Bandeira SA, Barden P, Barrasso DA, Bendifallah L, Bockmann FA, Böhme W, Borkent A, Brandão CRF, Busack SD, Bybee SM, Channing A, Chatzimanolis S, Christenhusz MJM, Crisci JV, D'elfia G, Da Costa LM, Davis SR, De Lucena CAS, Deuve T, Elizalde SF, Faivovich J, Farooq H, Ferguson AW, Gippoliti S, Gonçalves FMP, Gonzalez VH, Greenbaum E, Hinojosa-Díaz IA, Ineich I, Jiang J, Kahono S, Kury AB, Lucinda PHF, Lynch JD, Malécot V, Marques MP, Marris JWM, Mckellar RC, Mendes LF, Nihei SS, Nishikawa K, Ohler A, Orrico VGD, Ota H, Paiva J, Parrinha D, Pauwels OSG, Pereyra MO, Pestana LB, Pinheiro PDP, Prendini L, Prokop J, Rasmussen C, Rödel M-O, Rodrigues MT, Rodríguez SM, Salasnaya H, Sampaio Í, Sánchez-García A, Shebl MA, Santos BS, Solórzano-Kraemer MM, Sousa ACA, Stoev P, Teta P, Trape J-F, Dos Santos CV-D, Vasudevan K, Vink CJ, Vogel G, Wagner P, Wappler T, Ware JL, Wedmann S & Zacharie CK. 2021. The taxonomic impediment: a shortage of taxonomists, not the lack of technical approaches. *Zoological Journal of the Linnean Society* 193(2): 381–387. <https://doi.org/10.1093/zoolinlean/zlab072>

Ervik F, Tollsten L & Knudsen JT. 1999. Floral scent chemistry and pollination ecology in phytelephantoid palms (Arecaceae). *Plant Systematics and Evolution* 217: 279–297. <https://doi.org/10.1007/BF00984371>

Evenden ML, Whitehouse CM & Sykes J. 2014. Factors influencing flight capacity of the mountain pine beetle (Coleoptera: Curculionidae: Scolytinae). *Environmental Entomology* 43(1):187–96. <https://doi.org/10.1603/EN13244>

Fenster CB, Armbruster SW, Wilson P, Dudash MR & Thomson JD. 2004. Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution, and Systematics* 35: 375–403. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132347>

Ferrer MS, Marvaldi AE, Sato HA & Gonzalez AM. 2011. Biological notes on two species of *Oxycorynus* (Coleoptera: Belidae) associated with parasitic plants of the genus *Lophophytum* (Balanophoraceae), and new distribution records in Argentina. *Revista de la Sociedad Entomológica Argentina* 70(3–4): 351–355.

Franz NM. 2001. Description and phylogeny of *Staminodeus*, a new genus of Derelomini (Coleoptera: Curculionidae) associated with Cyclanthaceae. *The Coleopterists Bulletin* 55(4): 411–432.

Franz NM. 2003. Systematics of *Cyclanthura* gen. n., a new genus of Derelomini (Coleoptera: Curculionidae). *Insect Systematics and Evolution* 34(2): 153–198. <https://doi.org/10.1163/187631203788964818>

Franz NM. 2004. Analysing the history of the derelomine flower weevil–*Carludovica* association (Coleoptera: Curculionidae; Cyclanthaceae). *Biological Journal of the Linnean Society* 8(4): 483–517. <https://doi.org/10.1111/j.1095-8312.2003.00293.x>

Franz NM. 2006. Towards a phylogenetic system of derelomine flower weevils (Coleoptera: Curculionidae). *Systematic Entomology* 31(2): 220–287. <https://doi.org/10.1111/j.1365-3113.2005.00308.x>

1438 Franz NM. 2007a. Reproductive trade-offs in a specialized plant-pollinator system involving  
 1439 *Asplundia uncinata* (Cyclanthaceae) and a derelomine flower weevil (Coleoptera:  
 1440 Curculionidae). *Plant Systematics and Evolution* 269: 183–201.  
 1441 <https://doi.org/10.1007/s00606-007-0595-1>  
 1442  
 1443 Franz NM. 2007b. Pollination of *Anthurium* by derelomine flower weevils (Coleoptera:  
 1444 Curculionidae). *Revista de Biología Tropical* 55(1): 269–271.  
 1445 <https://doi.org/10.15517/RBT.V55I1.6079>  
 1446  
 1447 Franz NM. 2008. Revision, phylogeny and natural history of *Cotithene* Voss (Coleoptera:  
 1448 Curculionidae). *Zootaxa* 1782(1): 1–33. <https://doi.org/10.11646/zootaxa.1782.1.1>  
 1449  
 1450 Franz NM & Engel MS. 2010. Can higher-level phylogenies of weevils explain their  
 1451 evolutionary success? A critical review. *Systematic Entomology* 35(4): 597–606.  
 1452 <https://doi.org/10.1111/j.1365-3113.2010.00534.x>  
 1453  
 1454 Franz NM & O'Brien CW. 2001a. *Ganglionus*, a new genus of Derelomini (Coleoptera:  
 1455 Curculionidae) associated with *Carludovica* (Cyclanthaceae). *Annals of the Entomological*  
 1456 *Society of America* 74(6): 835–850.  
 1457 [https://doi.org/10.1603/0013-8746\(2001\)094\[0835:GANGOD\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2001)094[0835:GANGOD]2.0.CO;2)  
 1458  
 1459 Franz NM & O'Brien CW. 2001b. Revision and Phylogeny of *Perelleschus* (Coleoptera:  
 1460 Curculionidae), with notes on its association with *Carludovica* (Cyclanthaceae). *Transactions*  
 1461 *of the American Entomological Society* 127(2): 255–287. <https://doi.org/10.2307/25078744>  
 1462  
 1463 Franz NM & Valente RM. 2005. Evolutionary trends in derelomine flower weevils  
 1464 (Coleoptera: Curculionidae): from associations to homology. *Invertebrate Systematics* 19(6):  
 1465 499–530. <https://doi.org/10.1071/IS05026>  
 1466  
 1467 Freitas FV, Branstetter MG, Casali DM, Aguiar AJC, Griswold T & Almeida EAB. 2022.  
 1468 Phylogenomic dating and Bayesian biogeography illuminate an antitropical pattern for  
 1469 eucerine bees. *Journal of Biogeography* 49(6): 1034–1047.  
 1470 <https://doi.org/10.1111/jbi.14359>  
 1471  
 1472 Gaston KJ, Reavy D & Valladares GR. 1992. Intimacy and fidelity: Internal and external  
 1473 feeding by the British microlepidoptera. *Ecological Entomology* 17: 86–88.  
 1474 <https://doi.org/10.1111/j.1365-2311.1992.tb01044.x>  
 1475  
 1476 Girón JC, O'Brien CW & Rose-Smyth MC. 2018. On the West Indian weevil genus  
 1477 *Lachnopus* Schönherr, 1840 (Coleoptera: Curculionidae: Entiminae): descriptions of six new  
 1478 species, a proposal for species-groups, and an annotated checklist. *Zootaxa* 4423(1): 1–85.  
 1479 <https://doi.org/10.11646/zootaxa.4423.1.1>  
 1480  
 1481 Gottsberger G. 1986. Some pollination strategies in neotropical savannas and forests. *Plant*  
 1482 *Systematics and Evolution* 152: 29–45. <https://doi.org/10.1007/BF00985349>  
 1483  
 1484 Gottsberger G. 1999. Pollination and evolution in neotropical Annonaceae. *Plant Species*  
 1485 *Biology* 14(2): 143–152. <https://doi.org/10.1046/j.1442-1984.1999.00018.x>



1486 Guerrero-Olaya NY & Núñez-Avellaneda LA. 2017. Ecología de la polinización de *Syagrus*  
 1487 *smithii* (Arecaceae), una palma cantarofila de la Amazonia Colombiana. *Revista Peruana de*  
 1488 *Biología*. 24: 43–54. <http://doi.org/10.15381/rpb.v24i1.13102>

1489 Guy TJ, Hutchinson MC, Baldock KCR, Kayser E, Baiser B, Staniczenko PPA, Goheen JR,  
 1490 Pringle RM & Palmer TM. 2021. Large herbivores transform plant-pollinator networks in an  
 1491 African savanna. *Current Biology* 31(13): 2964–2971.  
 1492 <https://doi.org/10.1016/j.cub.2021.04.051>

1493 Haran J, Beaudoin-Ollivier L, Benoit L & Kergoat GJ. 2021. The origin of an extreme case of  
 1494 sister-species sympatry in a palm-pollinator mutualistic system. *Journal of Biogeography*  
 1495 48(12): 3158–3169. <https://doi.org/10.1111/jbi.14273>

1496 Haran J, Beaudoin-Ollivier L, Benoit L & Kuschel G. 2020. Revision of the palm-pollinating  
 1497 weevil genus *Elaeidobius* Kuschel, 1952 (Curculionidae, Curculioninae, Derelomini) with  
 1498 descriptions of two new species. *European Journal of Taxonomy* 684: 1–32.  
 1499 <https://doi.org/10.5852/ejt.2020.684>

1500 Haran J, Benoit L, Procheş Ş & Kergoat GJ. 2022a. *Ebenacobius* Haran, a new southern  
 1501 African genus of flower weevils (Coleoptera: Curculioninae: Derelomini) associated with  
 1502 dicotyledonous plants. *European Journal of Taxonomy* 818: 1–54.  
 1503 <https://doi.org/10.5852/ejt.2022.818.1771>

1504 Haran J, Procheş Ş, Benoit L & Kergoat GJ. 2022b. From monocots to dicots: host shifts in  
 1505 Afrotropical derelomine weevils shed light on the evolution of non-obligatory brood  
 1506 pollination mutualism. *Biological Journal of the Linnean Society* in press.  
 1507 <https://doi.org/10.1093/biolinnean/blac069>

1508 Hembry DH & Althoff DM. 2016. Diversification and coevolution in brood pollination  
 1509 mutualisms: Windows into the role of biotic interactions in generating biological diversity.  
 1510 *American Journal of Botany* 103: 1783–1792.  
 1511 <https://doi.org/10.3732/ajb.1600056>

1512 Henderson A. 1986. A review of pollination studies in the Palmae. *The Botanical Review* 52:  
 1513 221–259. <http://doi.org/10.1007/BF02860996>

1514 Henderson A, Pardini R, Dos Santos Rebello JF, Vanin S & Almeida D. 2000. Pollination of  
 1515 *Bactris* (Palmae) in an Amazon forest. *Brittonia* 52: 160–171.  
 1516 <https://doi.org/10.2307/2666507>

1517 Herre AE, Jandér KC & Machado CA. 2008. Evolutionary ecology of figs and their  
 1518 associates: recent progress and outstanding puzzles. *Annual Review of Ecology, Evolution*  
 1519 *and Systematics* 39: 439–458. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110232>

1520 Hoffmann A. 1958. *Faune de France*. 62. Coléoptères Curculionides, vol. 3. Lechevalier,  
 1521 Paris, pp. 1–545.

- Hotchkiss AT. 1958. Pollen and pollination in the Eupomatiaceae. *Proceedings of the Linnean Society of New South Wales* 83: 86–91.
- Hsiao Y & Oberprieler RG. 2022. Taxonomic revision of the genus *Miltotranes* Zimmerman, 1994 (Coleoptera: Curculionidae: Molytinae), the *Bowenia*-pollinating cycad weevils in Australia, with description of a new species and implications for the systematics of *Bowenia*. *Insects* 13(5): 456. <https://doi.org/10.3390/insects13050456>
- Hustache A. 1924. Synopsis des Curculionides de la faune malgache. *Bulletin de l'Académie Malgache* 7: 1–582.
- IPBES. 2016. *The assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production*. In: Potts EG, Imperatriz-Fonseca VL & Ngo HT, eds. Secretariat of the intergovernmental science-policy platform on biodiversity and ecosystem services. Bonn, Germany, pp. 1–552. <https://doi.org/10.5281/zenodo.3402856>
- Janzen DH. 1979. How to be a Fig. *Annual Review of Ecology and Systematics* 10: 13–51. <https://doi.org/10.1146/annurev.es.10.110179.000305>
- Jousselin E, Hossaert-McKey M, Herre EA & Kjellberg F. 2003. Why do fig wasps actively pollinate monoecious figs? *Oecologia* 134: 381–387. <https://doi.org/10.1007/s00442-002-1116-0>
- Jousselin, E, van Noort S, Rasplus J-Y, Rønsted N, Erasmus C & Greeff J. 2008. One tree to bind them all: host conservatism in a fig wasp community unraveled by cospeciation analyses among pollinating and non pollinating fig wasps. *Evolution* 62(7): 1777–1797. <https://doi.org/10.1111/j.1558-5646.2008.00406.x>
- Jürgens A, Glück U, Aas G & Dötterl S. 2014. Diel fragrance pattern correlates with olfactory preferences of diurnal and nocturnal flower visitors in *Salix caprea* (Salicaceae). *Botanical Journal of the Linnean Society* 175(4): 624–640. <https://doi.org/10.1111/boj.12183>
- Kawakita A & Kato M. 2009. Repeated independent evolution of obligate pollination mutualism in the Phyllanthae–*Epicephala* association. *Proceedings of the Royal Society B: Biological Sciences* 276(1656): 417–426. <https://doi.org/10.1098/rspb.2008.1226>
- Kergoat GJ, Condamine FL, Toussaint EFA., Capdevielle-Dulac C, Clamens A-L, Barbut J, Goldstein PZ & Le Ru BP. 2018. Opposite macroevolutionary responses to environmental changes in grasses and insects during the Neogene grassland expansion. *Nature Communications* 9: 5089. <https://doi.org/10.1038/s41467-018-07537-8>
- Kergoat GJ, Delobel A, Le Ru B & Silvain J-F. 2008. *Seed-beetles in the age of the molecule: recent advances on systematics and host-plant association patterns*. In: Jolivet P, Santiago-Blay J & Schmitt M, eds. *Researches on Chrysomelidae Volume 1*. Brill, Leiden, the Netherlands, pp. 59–86.

1577 Kergoat GJ, Le Ru BP, Sadeghi SE, Tuda M, Reid CAM, György Z, Genson G, Ribeiro-  
 1578 Costa CS & Delobel A. 2015. Evolution of *Spermophagus* seed beetles (Coleoptera,  
 1579 Bruchinae, Amblycerini) indicates both synchronous and delayed colonizations of host  
 1580 plants. *Molecular Phylogenetics and Evolution* 89: 91–103.  
 1581 <https://doi.org/10.1016/j.ympev.2015.04.014>  
 1582  
 1583 Kergoat GJ, Silvain J-F, Delobel A, Tuda M & Anton K-W. 2007. Defining the limits of  
 1584 taxonomic conservatism in host–plant use for phytophagous insects: Molecular systematics  
 1585 and evolution of host–plant associations in the seed-beetle genus *Bruchus* Linnaeus  
 1586 (Coleoptera: Chrysomelidae: Bruchinae). *Molecular Phylogenetics and Evolution* 43(1): 251–  
 1587 269. <https://doi.org/10.1016/j.ympev.2006.11.026>  
 1588  
 1589 Kevan PG & Baker HG. 1983. Insects as flower visitors and pollinators. *Annual Review of*  
 1590 *Entomology* 28: 407–453. <https://doi.org/10.1146/annurev.en.28.010183.002203>  
 1591  
 1592 Kirmse S & Chaboo CS. 2020. Flowers are essential to maintain high beetle diversity  
 1593 (Coleoptera) in a neotropical rainforest canopy. *Journal of Natural History* 54(25–26): 1661–  
 1594 1696. <https://doi.org/10.1080/00222933.2020.1811414>.  
 1595  
 1596 Knudsen JT, Tollsten L & Ervik F. 2001. Flower scent and pollination in selected neotropical  
 1597 palms. *Plant Biology* 3: 642–653. <https://doi.org/10.1055/s-2001-19366>  
 1598  
 1599 Kojima H & Morimoto K. 1995. Study on the tribe Ochyromerini (Coleoptera, Curculionidae)  
 1600 from East Asia I, descriptions of new species of the genera *Endaeus* and *Endaenidius*.  
 1601 *ESAKIA* 35: 63–102. <https://doi.org/10.5109/2582>  
 1602  
 1603 Krantz GW & Poinar, Jr GO. 2004. Mites, nematodes and the multimillion dollar weevil.  
 1604 *Journal of Natural History* 38(2): 135–141. <https://doi.org/10.1080/0022293021000041734>  
 1605  
 1606 Kress WJ & Specht CD. 2006. The evolutionary and biogeographic origin and diversification  
 1607 of the tropical monocot order Zingiberales. *Aliso: A Journal of Systematic and Floristic*  
 1608 *Botany* 22(1): 49. <https://scholarship.claremont.edu/aliso/vol22/iss1/49>  
 1609  
 1610 Küchmeister H, Webber AC, Silberbauer-Gottsberger I & Gottsberger G. 1998. A polinização  
 1611 e sua relação com a termogênese em espécies de Arecaceae e Annonaceae da Amazônia  
 1612 central. *Acta Amazônica* 28: 217–245.  
 1613 <https://doi.org/10.1590/1809-43921998283245>  
 1614  
 1615 Kuschel G & Leschen RAB. 2011. Phylogeny and taxonomy of the Rhinorhynchinae  
 1616 (Coleoptera: Nemonychidae). *Invertebrate Systematics* 24(6): 573–615.  
 1617 <https://doi.org/10.1071/IS09027>  
 1618  
 1619 Lajis MN, Hussein Y & Toia RF. 1985. Extraction and identification of the main compound  
 1620 present in *Elaeis guineensis* flower volatiles. *Pertanika* 8: 105–105.  
 1621  
 1622 Lau JYY, Guo X, Pang C-C, Tang CC, Thomas DC, Saunders RMK. 2017. Time-dependent  
 1623 trapping of pollinators driven by the alignment of floral phenology with insect circadian  
 1624 rhythms. *Frontiers in Plant Science* 8: 1119. <https://doi.org/10.3389/fpls.2017.01119>

1625  
1626 Lepesme P. 1947. *Les insectes des palmiers*. Paul Lechevalier, Paris, pp. 1–903.  
1627  
1628 Li K, Tschardt T, Saintes B, Buchori D & Grass I. 2019. Critical factors limiting pollination  
1629 success in oil palm: a systematic review. *Agriculture, Ecosystem, Environment* 280: 152–  
1630 160. <https://doi.org/10.1016/j.agee.2019.05.001>  
1631  
1632 Lopez-Vaamonde C, Wikström N, Kjer KM, Weiblen GD, Rasplus JY, Machado CA & Cook  
1633 JM. 2009. Molecular dating and biogeography of fig-pollinating wasps. *Molecular*  
1634 *Phylogenetics and Evolution* 52(3): 715–726. <https://doi.org/10.1016/j.ympev.2009.05.028>  
1635  
1636 Maas PJM, Westra LYT, Rainer H, Lobão AQ, Erkens RHJ. 2011. An updated index to  
1637 genera, species, and infraspecific taxa of Neotropical Annonaceae. *Nordic Journal of Botany*  
1638 29(3): 257–356. <https://doi.org/10.1111/j.1756-1051.2011.01092.x>  
1639  
1640 Magallón S & Castillo A. 2009. Angiosperm diversification through time. *American Journal of*  
1641 *Botany* 96: 349–365. <https://doi.org/10.3732/ajb.0800060>  
1642  
1643 Maia ACD, do Amaral Ferraz Navarro DM, Núñez-Avellaneda LA, Carreño-Barrera J,  
1644 Iannuzzi L, Cardona-Duque & J Nantes WAG. 2021. Methyl acetate, a highly volatile floral  
1645 semiochemical mediating specialized plant-beetle interactions. *The Science of Nature* 108:  
1646 21. <https://doi.org/10.1007/s00114-021-01731-3>  
1647  
1648 Mankga LT & Yessoufou K. 2017. Factors driving the global decline of cycad diversity. *AoB*  
1649 *Plants*. 9(4): plx022. <https://doi.org/10.1093/aobpla/plx022>  
1650  
1651 Marshall GAK. 1926. II. – On new Curculionidae from the oriental region (Coleoptera).  
1652 *Journal of Natural History Series* 9 17(100): 353–371.  
1653 <http://dx.doi.org/10.1080/00222932608633428>  
1654  
1655 Marshall GAK. 1932. XXVII. – New Curculionidae (Coleoptera) from tropical Africa. *Journal*  
1656 *of Natural History Series* 10 10(57): 217–230. <http://dx.doi.org/10.1080/00222933208673569>  
1657  
1658 Marshall GAK. 1933. I. – New Curculionidae (Coleoptera) from the Belgian Congo. *Journal*  
1659 *of Natural History Series* 10 11(61): 1–16. <http://dx.doi.org/10.1080/00222933308673619>  
1660  
1661 Maruyama PK, Oliveira GM, Ferreira C, Dalsgaard B & Oliveira PE. 2013. Pollination  
1662 syndromes ignored: importance of non-ornithophilous flowers to Neotropical savanna  
1663 hummingbirds. *Naturwissenschaften* 100: 1061–1068. [https://doi.org/10.1007/s00114-013-](https://doi.org/10.1007/s00114-013-1111-9)  
1664 [1111-9](https://doi.org/10.1007/s00114-013-1111-9)  
1665  
1666 Marvaldi AE & Ferrer MS. 2014. 3.3. *Belidae Schoenherr, 1826*. In: Leschen RAB & Beutel  
1667 RG, eds. *Morphology and Systematics (Phytophaga) Volume 3*. Berlin, München, Boston,  
1668 De Gruyter, pp. 316–328. <https://doi.org/10.1515/9783110274462.316>  
1669  
1670 McKibben GH, Grodowitz MJ & Villavaso EJ. 1988. Comparison of Flight Ability of Native  
1671 and Two Laboratory-Reared Strains of Boll Weevils (Coleoptera: Curculionidae) on a Flight  
1672 Mill. *Environmental Entomology* 17(5), 852–854. <https://doi.org/10.1093/ee/17.5.852>

1673  
1674 McLeish MJ & van Noort S. 2012. Codivergence and multiple host species use by fig wasp  
1675 populations of the *Ficus* pollination mutualism. *BMC Evolutionary Biology* 12: 1.  
1676 <https://doi.org/10.1186/1471-2148-12-1>  
1677  
1678 Meléndez-Jácome MR, Racines-Oliva MA, Galvis AA, Dávila AS & Ponce WP. 2019. Oil  
1679 palm pollinator dynamics and their behavior on flowers of different oil palm species *Elaeis*  
1680 *guineensis*, *Elaeis oleifera* and the *oleifera* x *guineensis* hybrid in Ecuador. *Pertanika Journal*  
1681 *of Tropical Agricultural Science* 42: 1295–1320.  
1682  
1683 Mendonça EN. 2004. Aspectos da autoecologia de *Cecropia glaziovii* Snethl.  
1684 (Cecropiaceae), fundamentos para o manejo e conservação de populações naturais da  
1685 espécie. Florianópolis, SC, pp. 1–78. Dissertação – Programa de Pós-Graduação em  
1686 Biologia Vegetal. Departamento de Botânica. Centro de Ciências Biológicas. Universidade  
1687 Federal de Santa Catarina. (available from ;  
1688 [https://repositorio.ufsc.br/xmlui/bitstream/handle/123456789/86682/1+autoecologia+de+cecr](https://repositorio.ufsc.br/xmlui/bitstream/handle/123456789/86682/1+autoecologia+de+cecropia+glaziovii.pdf;jsessionid=5A187EAF090193737CCFE4CF1BEA8A3F?sequence=1)  
1689 [opia+glaziovii.pdf;jsessionid=5A187EAF090193737CCFE4CF1BEA8A3F?sequence=1](https://repositorio.ufsc.br/xmlui/bitstream/handle/123456789/86682/1+autoecologia+de+cecropia+glaziovii.pdf;jsessionid=5A187EAF090193737CCFE4CF1BEA8A3F?sequence=1))  
1690  
1691 Momose K, Yumoto T, Nagamitsu T, Kato M, Nagamasu H, Sakai S, Harrison R, Itioka T,  
1692 Hamid A & Inoue T. 1998. Pollination biology in a lowland dipterocarp forest in Sarawak,  
1693 Malaysia. I. Characteristics of the plant-pollinator community in a lowland dipterocarp forest.  
1694 *American Journal of Botany* 85(10): 1477–501. <https://doi.org/10.2307/2446404>  
1695  
1696 Myers N. 1988. Threatened biotas: ‘hot-spots’ in tropical forests. *The Environmentalist* 8:  
1697 187–208. <https://doi.org/10.1007/BF02240252>  
1698  
1699 Noblick LR. 2017. A revision of the genus *Syagrus* (Arecaceae). *Phytotaxa* 294(1): 1–262.  
1700 <https://doi.org/10.11646/phytotaxa.294.1.1>  
1701  
1702 Núñez-Avellaneda LA, R. Bernal R & Knudsen JT. 2005. Diurnal palm pollination by  
1703 mystropine beetles: is it weather-related? *Plant Systematics and Evolution* 254: 149–171.  
1704 <https://doi.org/10.1007/s00606-005-0340-6>  
1705  
1706 Núñez-Avellaneda LA & Rojas-Robles R. 2008. Biología reproductiva y ecología de la  
1707 polinización de la palma milpesos *Oenocarpus bataua* en los Andes colombianos. *Caldasia*  
1708 30: 101–125.  
1709  
1710 Núñez-Avellaneda LA, Isaza C & Galeano G. 2015. Ecología de la polinización de tres  
1711 especies de *Oenocarpus* (Arecaceae) simpátricas en la Amazonia Colombiana. *Revista De*  
1712 *Biología Tropical* 63: 35–55. <https://doi.org/10.15517/rbt.v63i1.13030>  
1713  
1714 Núñez-Avellaneda LA & Carreño JI. 2017. Polinización por abejas en *Syagrus orinocensis*  
1715 (Arecaceae) en la Orinoquia colombiana. *Acta Biológica Colombiana* 22: 221–233.  
<https://doi.org/10.15446/abc.v22n2.58925>



1716 euglossine-pollinated orchid *Dichaea pendula*. *Oecologia* 182(4): 933–946.  
 1717 <https://doi.org/10.1007/s00442-016-3703-5>  
 1718  
 1719 Nunes PCE, Maruyama PK, Azevedo-Silva M & Sazima M. 2018. Parasitoids turn  
 1720 herbivores into mutualists in a nursery system involving active pollination. *Current Biology*  
 1721 28(6): 980–986.. <https://doi.org/10.1016/j.cub.2018.02.013>  
 1722  
 1723 Oberprieler RG. 1993. New taxa of Ochyromerina from Africa, with comments on the  
 1724 subtribe and description of immature stages (Coleoptera: Curculionidae: Curculioninae:  
 1725 Tychiini). *Journal of African Entomology* 107: 217–252.  
 1726  
 1727 Oberprieler RG. 1996. Systematics and evolution of the tribe Amorphocerini, with a review of  
 1728 the cycad weevils of the world. Ph.D. dissertation, University of the Free State,  
 1729 Bloemfontein, South Africa.  
 1730  
 1731 Oberprieler RG. 2004. “*Evil weevils*” – the key to cycad survival and diversification? In:  
 1732 Lindstrom AJ, ed. The Biology, Structure and Systematics of the Cycadales. Proceedings of  
 1733 the 6th International Conference on Cycad Biology. Nong Nooch Tropical Botanical Garden,  
 1734 Chonburi, pp. 170–194.  
 1735  
 1736 Oberprieler RG. 2014a. 3.3. *Belidae Schoenherr, 1826*. In: Leschen RAB & Beutel RG, eds.  
 1737 Morphology and Systematics (Phytophaga), Volume 3. Berlin, München, Boston, De  
 1738 Gruyter, pp. 316–328. <https://doi.org/10.1515/9783110274462.316>  
 1739  
 1740 Oberprieler RG. 2014b. 3.7. *Curculionidae Latreille, 1802*. In: Leschen RAB & Beutel RG,  
 1741 eds. Morphology and Systematics (Phytophaga), Volume 3. Berlin, München, Boston, De  
 1742 Gruyter, pp. 423–648. <https://doi.org/10.1515/9783110274462.423>  
 1743  
 1744 Oberprieler RG, Marvaldi AE & Anderson RS. 2007. Weevils, weevils, weevils everywhere.  
 1745 *Zootaxa* 520: 5326–5326. <https://doi.org/10.11646/zootaxa.1668.1.24>  
 1746  
 1747 Oberprieler RG & Oberprieler SK. 2012. *Talbragarus averyi* gen. et sp. n., the first Jurassic  
 1748 weevil from the southern hemisphere (Coleoptera: Curculionoidea: Nemomychidae). *Zootaxa*  
 1749 266: 256–266. <https://doi.org/10.11646/zootaxa.3478.1.25>  
 1750  
 1751 Ollerton J, Winfree R & Tarrant S. 2011. How many flowering plants are pollinated by  
 1752 animals? *Oikos* 120: 321–326. <https://doi.org/10.1111/j.1600-0706.2010.18644.x>  
 1753  
 1754 Orr MC, Hughes AC, Chesters D, Pickering J, Zhu CD & Ascher JS. 2021. Global patterns  
 1755 and drivers of bee distribution. *Current Biology* 31(3): 451–458.  
 1756 <https://doi.org/10.1016/j.cub.2020.10.053>  
 1757  
 1758 Pant DD. 1987. The fossil history and phylogeny of the cycadales. *Geophytology* 17(2): 125–  
 1759 162.  
 1760  
 1761 Paz FS, Pinto CE, de Brito RM, Imperatriz-Fonseca VL & Giannini TC. 2021. Edible fruit  
 1762 plant species in the Amazon forest rely mostly on bees and beetles as pollinators. *Journal of*  
 1763 *Economic Entomology* 114 (2): 710–722. <https://doi.org/10.1093/jee/toaa284>

1764  
 1765 Pellmyr O. 1997. Pollinating seed eaters: Why is active pollination so rare? *Ecology* 78:  
 1766 1655. <https://doi.org/10.2307/2266090>  
 1767  
 1768 Pellmyr O. 2003. Yuccas, yucca moths, and coevolution: a review. *Annals of the Missouri*  
 1769 *Botanical Garden* 90: 35–55. <https://doi.org/10.2307/3298524>  
 1770  
 1771 Pellmyr O, Thien LB, Bergström G & Groth I. 1990. Pollination of New Caledonian  
 1772 Winteraceae: opportunistic shifts or parallel radiation with their pollinators? *Plant*  
 1773 *Systematics and Evolution* 173(3):143–157. <https://doi.org/10.1007/BF00940859>  
 1774  
 1775 Pellmyr O, Thompson JN, Brown JM & Harrison RG. 1996. Evolution of pollination and  
 1776 mutualism in the yucca moth lineage. *The American Naturalist* 196 148(5): 827–847.  
 1777 <https://doi.org/10.1086/285958>  
 1778  
 1779 Potts SG, Imperatriz-Fonseca V, Ngo H, Aizen MA, Biesmeijer JC, Breeze TD, Dicks LV,  
 1780 Garibaldi LA, Hill R, Settele J & Vanbergen AJ. 2016. Safeguarding pollinators and their  
 1781 values to human well-being. *Nature* 540: 220–229. <https://doi.org/10.1038/nature20588>  
 1782  
 1783 Prance GT & Jongkind CCH. 2015. A revision of African Lecythidaceae. *Kew Bulletin* 70(1):  
 1784 1–68. <https://doi.org/10.1007/s12225-014-9547-4>  
 1785  
 1786 Prena J, Colonnelli E & Hespenheide HA 2014. 3.7.9 *Conoderinae* Schoenherr, 1833. In:  
 1787 Leschen RAB & Beutel RG, eds. Morphology and Systematics (Phytophaga), Volume 3.  
 1788 Berlin, München, Boston, De Gruyter, pp. 577–589.  
 1789  
 1790 Pullen KR, Jennings D & Oberprieler RG. 2014. Annotated catalogue of Australian weevils  
 1791 (Coleoptera: Curculionoidea). *Zootaxa* 3896: 1–481.  
 1792 <https://doi.org/10.11646/zootaxa.3896.1.1>  
 1793  
 1794 Ramírez N. 2004. Ecology of pollination in a tropical venezuelan savanna. *Plant Ecology*  
 1795 173(2): 171–189. <http://www.jstor.org/stable/20146634>  
 1796  
 1797 Rands SA & Whitney HM. 2008. Floral temperature and optimal foraging: Is heat a feasible  
 1798 floral reward for pollinators? *PLoS One* 3: e2007.  
 1799 <https://doi.org/10.1371/journal.pone.0002007>  
 1800  
 1801 Ratnayake RMCS, Gunatilleke IAUN, Wijesundara DSA & Saunders RMK. 2006.  
 1802 Reproductive biology of two sympatric species of *Polyalthia* (Annonaceae) in Sri Lanka. I.  
 1803 Pollination by Curculionid Beetles. *International Journal of Plant Sciences* 167(3): 483–493.  
 1804 <https://doi.org/10.1086/502715>  
 1805  
 1806 Rech AR, Dalsgaard B, Sandel B, Sonne J, Svenning J-C, Holmes N & Ollerton J. 2016. The  
 1807 macroecology of animal versus wind pollination: ecological factors are more important than  
 1808 historical climate stability. *Plant Ecology & Diversity* 9(3): 253–262.  
 1809 <https://doi.org/10.1080/17550874.2016.1207722>  
 1810

1811 Regal PJ. 1982. Pollination by wind and animals: ecology of geographic patterns. *Annual*  
 1812 *Review of Ecology and Systematics* 13(1): 497–524.  
 1813 <https://doi.org/10.1146/annurev.es.13.110182.002433>  
 1814  
 1815 Renner SS & Feil JP. 1993. Pollinators of tropical dioecious angiosperms. *American Journal*  
 1816 *of Botany* 80: 1100–1107. <https://doi.org/10.1002/j.1537-2197.1993.tb15337.x>  
 1817  
 1818 Rosa L. 2000. Ecologia da polinização de *Butia capitata* (Martius) Beccari var. *odorata*  
 1819 (Palmae), no sul do Brasil. MSc Thesis. Universidade Federal de Santa Catarina.  
 1820  
 1821 Sakai S. 2002. A review of brood-site pollination mutualism: plants providing breeding sites  
 1822 for their pollinators. *Journal of Plant Research* 115: 161–168.  
 1823 <https://doi.org/10.1007/s102650200021>  
 1824  
 1825 Salzman S, Crook D, Calonje M, Stevenson DW, Pierce NE & Hopkins R. 2021. Cycad-  
 1826 weevil pollination symbiosis is characterized by rapidly evolving and highly specific plant-  
 1827 insect chemical communication. *Frontiers in Plant Science* 12: 639368.  
 1828 <https://doi.org/10.3389/fpls.2021.639368>  
 1829  
 1830 Salzman S, Crook D, Crall JD, Hopkins R & Pierce NE. 2020. An ancient push-pull  
 1831 pollination mechanism in cycads. *Science Advances* 6: eaay6169.  
 1832 <https://doi.org/10.1126/sciadv.aay6169>  
 1833  
 1834 Salzman S, Whitaker M & Pierce NE. 2018. Cycad-feeding insects share a core gut  
 1835 microbiome. *Biological Journal of the Linnean Society* 123(4): 728–738.  
 1836 <https://doi.org/10.1093/biolinnean/bly017>  
 1837  
 1838 Saravy FP, Marques MI & Schuchmann KL. 2021. Coleopteran pollinators of Annonaceae in  
 1839 the Brazilian Cerrado – a review. *Diversity* 13(9): 438. <https://doi.org/10.3390/d13090438>  
 1840  
 1841 Satler JD, Herre EA, Heath TA, Machado CA, Zúñiga AG & Nason JD. 2022. Genome-wide  
 1842 sequence data show no evidence of hybridization and introgression among pollinator wasps  
 1843 associated with a community of Panamanian strangler figs. *Molecular Ecology* 31(7): 2106–  
 1844 2123. <https://doi.org/10.1111/mec.16373>  
 1845  
 1846 Saunders RMK. 2020. The evolution of key functional floral traits in the early divergent  
 1847 angiosperm family Annonaceae. *Journal of Systematics and Evolution* 58(4): 369–392.  
 1848 <https://doi.org/10.1111/jse.12645>  
 1849  
 1850 Schleuning M, Fründ J, Klein AM, Abrahamczyk S, Alarcón R, Albrecht M, Andersson GKS,  
 1851 Bazarian S, Böhning-Gaese K, Bommarco R, Dalsgaard B, Dehling DM, Gotlieb A, Hagen  
 1852 M, Hickler T, Holzschuh A, Kaiser-Bunbury CN, Kreft H, Morris RJ, Sandel B, Sutherland  
 1853 WJ, Svenning JC, Tschamntke T, Watts S, Weiner CN, Werner M, Williams NM, Winqvist C,  
 1854 Dormann CF & Blüthgen N. 2012. Specialization of mutualistic interaction networks  
 1855 decreases towards tropical latitudes. *Current Biology* 22: 1925–1931.  
 1856 <https://doi.org/10.1016/j.cub.2012.08.015>  
 1857

1858 Segraves KA, Althoff DM & Pellmyr O. 2005. Limiting cheaters in mutualism: evidence from  
 1859 hybridization between mutualist and cheater yucca moths. *Proceedings of the Royal Society*  
 1860 *B Biological Sciences* 272(1577): 2195–2201.  
 1861 <http://doi.org/10.1098/rspb.2005.3201>  
 1862

1863 Shin S, Clarke DJ, Lemmon AR, Lemmon EM, Aitken AL, Haddad S, Farrell BD, Marvaldi  
 1864 AE, Oberprieler RG, McKenna DD. 2018. Phylogenomic data yield new and robust insights  
 1865 into the phylogeny and evolution of weevils. *Molecular Biology and Evolution* 35: 823–836.  
 1866 <http://doi.org/10.1093/molbev/msx324>  
 1867

1868 Silberbauer-Gottsberger I. 1990. Pollination and evolution in palms. *Phyton* 30(2): 213–233.  
 1869

1870 Silberbauer-Gottsberger I, Gottsberger G & Webber AC. 2003. Morphological and functional  
 1871 flower characteristics of New and Old World Annonaceae with respect to their mode of  
 1872 pollination. *Taxon* 52: 701–718. <http://doi.org/10.2307/3647345>

1873 Silberbauer-Gottsberger I, Vanin SA & Gottsberger G. 2013. Interactions of the Cerrado  
 1874 palms *Butia paraguayensis* and *Syagrus petraea* with parasitic and pollinating insects.  
 1875 *Sociobiology* 60: 306–316. <http://doi.org/10.13102/sociobiology.v60i3.306-316>

1876 Sprick P & Floren A. 2018. Diversity of Curculionoidea in humid rain forest canopies of  
 1877 Borneo: a taxonomic blank spot. *Diversity* 10(4): 116. <https://doi.org/10.3390/d10040116>  
 1878

1879 Su ZH, Sasaki A., Kusumi J., Chou A-P, Tzeng H-Y, Li H-Q & Yu H. 2022. Pollinator sharing,  
 1880 copollination, and speciation by host shifting among six closely related dioecious fig species.  
 1881 *Communications Biology* 5: 284. <https://doi.org/10.1038/s42003-022-03223-0>  
 1882

1883 Syed RA. 1979. Studies on oil palm pollination by insects. *Bulletin of Entomological*  
 1884 *Research* 69: 213–224. <http://doi.org/10.1017/S000748530001767>  
 1885

1886 Syed RA, Law IH & Corley RHV. 1982. Insect pollination of oil palm: introduction,  
 1887 establishment and pollinating efficiency of *Elaeidobius kamerunicus* in Malaysia. *Planter* 58:  
 1888 547–561.  
 1889

1890 Teichert H, Dötterl S & Gottsberger G. 2018. Scent emissions and floral nutrients of  
 1891 Carludovicoideae (Cyclanthaceae) and their importance for associated beetles. *Plant*  
 1892 *Systematics and Evolution* 304: 831–839. <https://doi.org/10.1007/s00606-018-1513-4>  
 1893

1894 Toon A, Terry LI, Tang W, Walter GH & Cook LG. 2020. Insect pollination of cycads. *Austral*  
 1895 *Ecology* 45: 1033–1058. <https://doi.org/10.1111/aec.12925>  
 1896

1897 Traoré L, Ouédraogo I, Ouédraogo A & Thiombiano A. 2011. Perceptions, usages et  
 1898 vulnérabilité des ressources végétales ligneuses dans le Sud-Ouest du Burkina Faso.  
 1899 *International Journal of Biological and Chemical Sciences* 5(1): 258–278.  
 1900 <https://doi.org/10.4314/ijbcs.v5i1.68103>  
 1901

1902 Tremblay RL, Ackerman JD, Zimmerman JK & Calvo RN. 2005. Variation in sexual  
 1903 reproduction in orchids and its evolutionary consequences: A spasmodic journey to

1904 diversification. *Biological Journal of the Linnean Society* 84: 1–54.  
 1905 <https://doi.org/10.1111/j.1095-8312.2004.00400.x>  
 1906  
 1907 Turner IM. 2001. *The ecology of trees in the tropical rain forest*. Cambridge Tropical Biology  
 1908 Series, Cambridge University Press, Cambridge, pp. 1–298.  
 1909  
 1910 Valente RM. 2005. O gênero *Celetes* Schoenherr, 1836 (Curculionidae, Eirrhiniinae):  
 1911 filogenia, sistemática e evolução com palmeiras hospedeiras. Doctoral thesis, University of  
 1912 São Paulo, São Paulo, Brazil.  
 1913  
 1914 Valente RM & de Medeiros BAS. 2013. A new species of *Anchylorhynchus* Schoenherr  
 1915 (Coleoptera: Curculionidae) from the Amazon, with a record of a new host palm for the  
 1916 genus. *Zootaxa* 3709(4): 304–400. <https://doi.org/10.11646/zootaxa.3709.4.6>  
 1917  
 1918 Valente RM & Da Silva PA. 2014. The first Amazonian species of *Andranthobius* Kuschel  
 1919 (Coleoptera: Curculionidae), with records of new host palms for the genus. *Zootaxa* 3786(4):  
 1920 458–468. <https://doi.org/10.11646/zootaxa.3786.4.4>  
 1921  
 1922 Valente RM, DA Silva PAL & de Medeiros BAS. 2019. The first species of *Cotithene* Voss  
 1923 (Coleoptera: Curculionidae: Curculioninae) from Amazonian Brazil, with notes on its role as  
 1924 a pollinator of *Evodianthus funifer* (Poit.) Lindm. (Cyclanthaceae). *Zootaxa* 4576.3.3.  
 1925 <https://doi.org/10.11646/zootaxa.4576.3.3>  
 1926  
 1927 van der Cingel NA. 2001. *An atlas of orchid pollination: America, Africa, Asia and Australia*  
 1928 CRC Press, Rotterdam, pp. 1–296.  
 1929  
 1930 Vanderplanck M, Lapeyre B, Brondani M, Opsommer M, Dufay M, Hossaert-McKey M &  
 1931 Proffit M. 2021. Ozone pollution alters olfaction and behavior of pollinators. *Antioxidants*  
 1932 10(5): 636. <https://doi.org/10.3390/antiox10050636>  
 1933  
 1934 Vizentin-Bugoni J, Maruyama PK, de Souza CS, Ollerton J, Rech AR & Sazima M. 2018.  
 1935 *Plant-pollinator networks in the Tropics: A review*. In: Dáttilo W & Rico-Gray V, eds.  
 1936 Ecological networks in the tropics. Springer, Cham, Switzerland, pp. 73–91.  
 1937 [https://doi.org/10.1007/978-3-319-68228-0\\_6](https://doi.org/10.1007/978-3-319-68228-0_6)  
 1938  
 1939 Von Martius KFP. 1823. '*Historia naturalis palmarum. Opus tripatitum, cuius volumen*  
 1940 *primum palmas generatim tractat; volumen secundum Brasiliae palmas singulatim*  
 1941 *descriptione et icone illustrat; volumen tertium ordinis, familiarum, generum characteres*  
 1942 *recenset species selectas describit et figuris adumbrat adiecta omnium synopsi*, Vol. 1.'  
 1943 (T.O. Weigel: Lipsiae).  
 1944  
 1945 Wagner DL, Grames EM, Forister ML, Berenbaum MR & Stopak D. 2021. Insect decline in  
 1946 the Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of*  
 1947 *Sciences of the U.S.A.* 118(2): e2023989118 <https://doi.org/10.1073/pnas.2023989118>  
 1948  
 1949 Walker JD, Geissman JW, Bowring SA, Babcock LE. 2018. Geologic Time Scale v. 5.0:  
 1950 Geological Society of America.  
 1951 <https://www.geosociety.org/documents/gsa/timescale/timescl.pdf>



1952

1953 Wang R, Yang Y, Jing Y, Segar ST, Zhang Y, Wang G, Chen J, Liu Q-F, Chen S, Chen Y,

1954 Cruaud A, Ding Y-Y, Dunn DW, Gao Q, Gilmartin PM, Jiang K, Kjellberg F, Li H-Q, Li Y-Y,

1955 Liu J-Q, Liu J, Machado CA, Ming R, Rasplus J-Y, Tong X, Wen P, Yang H-M, Yang J-J, Yin

1956 Y, Zhang XT, Zhang Y-Y, Yu H, Yue Z, Compton SG & Chen XY. 2021. Molecular

1957 mechanisms of mutualistic and antagonistic interactions in a plant–pollinator association.

1958 *Nature Ecology and Evolution* 5: 974–986. <https://doi.org/10.1038/s41559-021-01469-1>

1959

1960 Wardhaugh CW. 2015. How many species of arthropods visit flowers? *Arthropod-Plant*

1961 *Interactions* 9(6): 547–65. <https://doi.org/10.1007/s11829-015-9398-4>

1962

1963 West SA, Herre EA, Windsor DM & Green PRS. 1996. The ecology and evolution of the New

1964 World non-pollinating fig wasp communities. *Journal of Biogeography* 23: 447–458.

1965 <https://doi.org/10.1111/j.1365-2699.1996.tb00006.x>

1966

1967 Whitehead DR. 1968. Wind pollination in the angiosperms: evolutionary and environmental

1968 considerations. *Evolution* 23: 28–35. <https://doi.org/10.2307/2406479>

1969

1970 Wiebes JT. 1979. Co-evolution of figs and their insect pollinators. *Annual Review of Ecology*

1971 *and Systematics* 10: 1–12. <https://doi.org/10.1146/annurev.es.10.110179.000245>

1972

1973 Williams G & Adam P. 1994. A review of rainforest pollination and plant-pollinator

1974 interactions with particular reference to Australian subtropical rainforests. *Australian*

1975 *Zoologist* 29 (3–4): 177–212. <https://doi.org/10.7882/AZ.1994.006>

1976

1977 Wolowski M, Ashman TL & Freitas L. 2014. Meta-analysis of pollen limitation reveals the

1978 relevance of pollination generalization in the Atlantic forest of Brazil. *PLoS ONE* 9(2):

1979 e89498. <https://doi.org/10.1371/journal.pone.0089498>

1980

1981 Zattara EE & Aizen MA. 2021. Worldwide occurrence records suggest a global decline in

1982 bee species richness. *One Earth* 4(1): 114–123.

1983 <https://doi.org/10.1016/j.oneear.2020.12.005>

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

## Supplementary information

**Table S1.** List of plant-weevil brood-site pollination mutualism obtained through literature review completed by unpublished observations. Weevil names and authorship follows the Weevil taxa database (accessed January 2023) otherwise stated. Plant names have been updated according to the World Flora Online (<http://www.worldfloraonline.org/>, accessed August 2022).

## Appendix 1

### Material and Methods

The literature review was conducted directly through search of keywords on Web of Science and indirectly through reference cited in review papers on specific plant-weevil pollination systems. About 110 articles, including five review papers were screened to search for species-specific plant weevil interactions, but also the ecological, phenological and physiological traits associated with these interactions. These published data were completed by unpublished field observations provided by colleagues or the authors themselves. For each species-species interaction, we compiled an indication of classification (family rank), the location of the breeding site of larval stages and the known evidence of pollination mutualism. The experimental demonstration of an effective brood-site pollination mutualism requires the observation of breeding sites of larval stage of the putative pollinator and the investigation of pollination efficacy of the adults by a pollinator exclusion method. This time consuming process is rarely conducted, except for a few model systems. In order to integrate a consistent number of interactions without risk of overestimation, we followed the following rationale: BSPM interactions were reported only when, 1) the weevil species is identified at least at the tribal level, 2) BSPM interactions has been shown in a congeneric plant or weevil species (phylogenetic niche conservatism, see sections 1 and 2.3.5) and/or 3) plants and/or weevils show typical traits associated with BSPM (see sections 1 and 2). We explicitly excluded cases of flower visitors reported in one publication that were demonstrated not to be relevant pollinators in another one. This strategy has limitations but reflects the current state of knowledge on these systems and should be seen as a first assessment of their extent in overlooked tropical biomes. When relevant, details or a discussion on knowledge gaps were added (see column "notes" in Table S1).

A subset of plant-weevil interactions was used to estimate the extent of undescribed interactions (Derelomini-angiosperms dicots/palms/Carludovicoideae, Ochyromerini-angiosperms dicots, Stoerini-Eupomiataceae). All these types of interactions have in common (except in the Derelomini-angiosperms dicots): effective BSPM has been reported at least in one system; highly similar interactions are known in closely related genera and species; and an estimate of the global diversity of the weevil of plant clade is available (published records or author unpublished observations, see section 4 and Table S1). For these five groups, we reported the number of effective BSPM interactions reported, the expected number of BSPM expected based on phylogenetic niche conservatism (see section 1 and above) and an estimate of the full diversity (of the weevil except for the Carludovicoideae) of the clade involved in the interactions (see statistics in figure 2). Based on these values obtained from the literature we computed the ratios between experimentally-verified BSPM and expected BSPM based on PNC, and also between experimentally-

2043 verified BSPM and the full diversity of the lineages included (ratio of 8 and 17 respectively).  
2044 Considering the species counts obtained in this review (see section 1) and these ratios, we  
2045 estimated that about 2000 plants (250 x 8) and around 2500 weevil species (300 x 8) could  
2046 be engaged in BSPM globally. We chose the lower range estimates to avoid any  
2047 overstatement.

2048

2049