



HAL
open science

Unifying ammonia inhibitory limits in anaerobic digestion: link with operational conditions and microbial communities

Gabriel Capson-Tojo, Roman Moscoviz, Ángel Robles, Sergi Astals,
Jean-Philippe Steyer

► To cite this version:

Gabriel Capson-Tojo, Roman Moscoviz, Ángel Robles, Sergi Astals, Jean-Philippe Steyer. Unifying ammonia inhibitory limits in anaerobic digestion: link with operational conditions and microbial communities. 16th IWA World Conference on Anaerobic Digestion, Jun 2019, Delft, Netherlands. 1p. hal-03782803

HAL Id: hal-03782803

<https://hal.inrae.fr/hal-03782803v1>

Submitted on 21 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Unifying ammonia inhibitory limits in anaerobic digestion: link with operational conditions and microbial communities

16TH IWA World conference on Anaerobic Digestion
23-27th June, Delft, The Netherlands



G. Capson-Tojo ^{a,b}, R. Moscoviz ^a, A. Robles ^c, S. Astals ^{b,d}, J.-P. Steyer ^e

^a Suez, CIRSEE, 38 rue du Président Wilson, 78230, Le Pecq, France

^b Advanced Water Management Centre, The University of Queensland, Brisbane, QLD 4072, Australia

^c Departament d'Enginyeria Química, ETSE-UV, Universitat de València, Avinguda de la Universitat s/n, 46100 Burjassot, València, Spain

^d Department of Chemical Engineering and Analytical Chemistry, University of Barcelona, C/Martí i Franquès 1, 08028 Barcelona, Spain


^e LBE, Univ. Montpellier, INRA, 102 avenue des Etangs, 11100, Narbonne, France



Introduction

- Free ammonia nitrogen (FAN) is a major inhibitor in anaerobic digestion processes
- There is no current agreement in the literature regarding inhibitory limits
- Data from literature was used to unify inhibitory limits and link inhibition resilience to operational parameters and archaeal populations

Main results



Reduction of the specific methanogenic activities (SMA) at different FAN concentrations. The set of inhibition curves and IC_{50} values resulting from the clustering approach are also shown.


- Six different inhibition curves were obtained, forming **six data clusters with increasing resilience to FAN** (higher IC_{50} values)
- A **sensitivity analysis** verified that the cluster distribution was optimal according to defined criteria
- The median **temperatures and pH** of each cluster **increased with the IC_{50}**
- The total ammonia nitrogen (TAN) and the NH_4^+ concentrations were **not significantly different between clusters**

Take home message

- Resilience to FAN inhibition is affected by both the operational parameters and the archaeal composition
- The working pH and temperature, rather than the TAN content itself, are the main factors affecting FAN resilience
- Methanosaeta*-dominated reactors presented lower inhibition limits when compared to mixotrophic and hydrogenotrophic systems
- More research must be carried out to study the microbial communities in nitrogen-rich systems at high FAN levels

Materials and methods

- A database containing 1,590 points was generated (from 50 articles)
- FAN concentrations were recalculated using a modified Davies equation to account for the ionic strength in N-rich media (Capson-Tojo *et al.*)
- The inhibitory constants (IC_{50}) were calculated using a threshold inhibition function (Astals *et al.*, 2018)
- An iterative clustering approach based on regression analysis over the inhibition function was used to obtain different IC_{50} values



Repartition of the main archaeal genus after anaerobic digestion for each resulting cluster. Only the predominant genus (that with the highest percentage in relative abundance) was considered for the counting. "Non-reported" corresponds to studies in which the microbial communities were not analysed.

- Clusters with **low IC_{50} values** were dominated by **acetoclastic** methanogens (mainly *Methanosaeta*)
- Methanosarcina* (mixotroph) dominated at **low-intermediate IC_{50} values**
- Hydrogenotrophic** methanogens showed the **highest IC_{50}**

References

Astals *et al.*, (2018), Characterising and modelling free ammonia and ammonium inhibition in anaerobic systems. *Water Res.* 143:127–35

Capson-Tojo *et al.*, Unraveling the literature chaos around free ammonia inhibition in anaerobic digestion. Submitted to *Renewable & Sustainable Energy Reviews*