HAL
open science

Methanosarcina sp. as key archaea to avoid acidification in dry anaerobic digestion of food waste

Gabriel Capson-Tojo, Maxime Rouez, Marion Crest, Eric Trably, Jean-Philippe Steyer, Jean Philippe Delgenes, Renaud Escudié

To cite this version:

Gabriel Capson-Tojo, Maxime Rouez, Marion Crest, Eric Trably, Jean-Philippe Steyer, et al.. Methanosarcina sp. as key archaea to avoid acidification in dry anaerobic digestion of food waste. 15th IWA World Conference On Anaerobic Digestion, Oct 2017, Beijing, China. , 1p. hal-03782805

HAL Id: hal-03782805

https://hal.inrae.fr/hal-03782805v1

Submitted on 21 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

METHANOSARCINA SP. AS KEY ARCHAEA TO AVOID ACIDIFICATION IN DRY ANAEROBIC DIGESTION OF FOOD WASTE

15TH IWA WORLD CONFERENCE ON ANAEROBIC DIGESTION

18-20th October. Beijing (China)

G. Capson-Tojo*,, M. Rouez**, M. Crest**, E. Trably*, J.-P. Steyer*, J.-P. Delgenès*, R. Escudié***

* LBE, INRA, Univ. Montpellier, 102 avenue des Etangs, 11100, Narbonne, France

** Suez, CIRSEE, 38 rue du Président Wilson, 78230, Le Pecq, France

INTRODUCTION

- **Food waste** (FW) is a highly biodegradable substrate rich in nitrogen [1]. These characteristics often lead to accumulation of volatile fatty acids (**VFAs**) and free ammonia nitrogen (**FAN**), causing low methane yields and even AD failure [2,3]. It is then critical to have well-adapted resilient **archaeal communities** in the reactors, resistant to these common inhibitors (such as mixotrophic archaea [4])
- **Objective**: assess influence of the composition of the archaeal communities of different inocula on the methane production from FW and cardboard (CB) at different working conditions (*i.e.* **total solid** (TS) contents, **co-digestion proportions** and **substrate loads**)

MATERIALS & METHODS

- Model FW (VALORGAS report [5]) and compact CB as substrates
- Three different inocula: UASB1 (high TS, low TAN), UASB2 (high TS, low TAN), OW (low TS, high TAN)
- 5 pairs of comparable experiments at different co-digestion proportions (1.00-4.00), substrate to inoculum (S/X) ratios (0.25-1.00 g VS·g VS^{-1}) and initial TS contents (20-30 %)
- Follow-up of methane production and analysis of concentration of TAN, VFAs and archaeal composition in inocula and digestates

Operational conditions of the batch experiments and obtained methane yields

Pair	Inoculum	Substrate	S/X (g VS·g VS^{-1})	Co-dig. ratio (g TS FW·g TS CB^{-1})	Initial TS (%)	Methane yield (ml CH_4·g VS^{-1})
A	UASB2	FW+CB	0.25	1.00	27.5	393±9.0
	UASB1	FW+CB	0.25	1.00	30.0	7.9±1.9
B	UASB2	FW+CB	0.25	1.86	27.5	409±11
	UASB1	FW+CB	0.25	1.75	30.0	11±2.7
C	UASB2	FW+CB	0.25	1.00	35.0	401±16
	UASB1	FW+CB	0.25	1.00	35.0	17±2.3
D	OW	FW	0.25	-	20.0	464±14
	UASB1	FW	0.25	-	20.0	0.7±0.9*
E	OW	FW	1.00	-	20.0	375±17
	UASB2	FW+CB	1.00	4.00	27.5	0±0

MAIN RESULTS



Composition of the archaeal communities in the inocula (left) and the digestates (right)

- High TAN and transient VFA concentrations in the reactors (up to 5 g·l^{-1} and 23 g COD·l^{-1}, respectively)
- One reactor of each pair produced methane efficiently and the other was acidified due to VFA accumulation at the beginning of the batch AD process
- No methane with UASB1; with UABS2, only at an S/X ratio of 0.25 g VS·g VS^{-1}; OW worked at high loads
- In UASB1: no ***Methanosarcina*** sp. initially; in UASB2, minority initially; in OW: predominant methanogen
- In all reactors producing methane: ***Methanosarcina*** sp. as main methanogen
- No *Methanosaeta* sp. in digestates due to inhibition

Take-Home Message

- While the tests started with inocula rich in ***Methanosarcina*** sp. led to efficient methane production, VFAs accumulated in the reactors when inocula were poor in this archaea and no methane was produced
- Higher proportions of *Methanosarcina* sp. in the inocula allowed greater substrate loads
- Regardless of the inoculum used, ***Methanosarcina*** sp. was the dominant methanogen in the **methane-producing** reactors
- The initial archaeal composition of the inoculum is crucial during reactor start-up to achieve stable anaerobic digestion

REFERENCES

[1] Capson-Tojo G., et al., *Food waste valorization via anaerobic processes: a review.* Reviews in Environmental Science and Bio/Technology, 2016. **15**: p. 499–547.
[2] Capson-Tojo G., et al., *Dry anaerobic digestion of food waste and cardboard at different substrate loads, solid contents and co-digestion proportions.* Bioresource Technology, 2017. **233**: p. 166–175.
[3] Capson-Tojo G., et al., *Accumulation of propionic acid during consecutive batch anaerobic digestion of commercial food waste*. Bioresource Technology, 2017. **245**, Part A: p. 724-733.
[4] De Vrieze J., et al., Methanosarcina: The rediscovered methanogen for heavy duty biomethanation. Bioresource Technology, 2012. **112**: p. 1–9.
[5] VALORGAS. *D2.1: Compositional analysis of food waste from study sites in geographically distinct regions of Europe*. 2010.

INSTITUT NATIONAL DE LA RECHERCHE AGRONOMIQUE
Laboratoire de Biotechnologie de l'Environnement
102 Avenue des Etangs • Narbonne • F-11100 • France
Tel : + 33 6 38 12 79 57
E-mail: gabriel.capson-tojo@supagro.inra.fr
www.montpellier.inra.fr/narbonne