



Cyclic di-AMP Oversight of Counter-Ion Osmolyte Pools Impacts Intrinsic Cefuroxime Resistance in *Lactococcus lactis*

Huong Thi Pham, Wen Shi, Yuwei Xiang, Su Yi Foo, Manuel Plan, Pascal Courtin, Marie-Pierre Chapot-Chartier, Eddy Smid, Zhao-Xun Liang, Esteban Marcellin, et al.

► To cite this version:

Huong Thi Pham, Wen Shi, Yuwei Xiang, Su Yi Foo, Manuel Plan, et al.. Cyclic di-AMP Oversight of Counter-Ion Osmolyte Pools Impacts Intrinsic Cefuroxime Resistance in *Lactococcus lactis*. *mBio*, 2021, 12 (2), pp.e00324-21. 10.1128/mBio.00324-21 . hal-03786864

HAL Id: hal-03786864

<https://hal.inrae.fr/hal-03786864v1>

Submitted on 10 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cyclic di-AMP oversight of counter-ion osmolyte pools impacts intrinsic cefuroxime resistance in *Lactococcus lactis*

Pham, Huong Thi; Shi, Wen; Xiang, Yuwei; Foo, Su Yi; Plan, Manuel R.; Courtin, Pascal; Chapot-Chartier, Marie-Pierre; Smid, Eddy J.; Liang, Zhao-Xun; Marcellin, Esteban; Turner, Mark S.

2021

Pham, H. T., Shi, W., Xiang, Y., Foo, S. Y., Plan, M. R., Courtin, P., Chapot-Chartier, M., Smid, E. J., Liang, Z., Marcellin, E. & Turner, M. S. (2021). Cyclic di-AMP oversight of counter-ion osmolyte pools impacts intrinsic cefuroxime resistance in *Lactococcus lactis*. *MBio*, 12(2), e00324-21-. <https://dx.doi.org/10.1128/mBio.00324-21>

<https://hdl.handle.net/10356/154055>

<https://doi.org/10.1128/mBio.00324-21>

© 2021 Pham et al. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license

Downloaded on 10 Sep 2024 22:38:38 SGT



Cyclic di-AMP Oversight of Counter-Ion Osmolyte Pools Impacts Intrinsic Cefuroxime Resistance in *Lactococcus lactis*

Huong Thi Pham,^{a,b} Wen Shi,^a Yuwei Xiang,^a Su Yi Foo,^a Manuel R. Plan,^{c,d} Pascal Courtin,^e  Marie-Pierre Chapot-Chartier,^e Eddy J. Smid,^f  Zhao-Xun Liang,^g Esteban Marcellin,^c  Mark S. Turner^{a,h}

^aSchool of Agriculture and Food Sciences, University of Queensland, Brisbane, Queensland, Australia

^bThe University of Danang, University of Science and Technology, Da Nang, Vietnam

^cAustralian Institute for Bioengineering and Nanotechnology, University of Queensland, Brisbane, Queensland, Australia

^dMetabolomics Australia, AIBN, The University of Queensland, Brisbane, Queensland, Australia

^eUniversité Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, Jouy-en-Josas, France

^fLaboratory of Food Microbiology, Wageningen University and Research, Wageningen, The Netherlands

^gSchool of Biological Sciences, Nanyang Technological University, Singapore

^hQueensland Alliance for Agriculture and Food Innovation, University of Queensland, Brisbane, Queensland, Australia

ABSTRACT The broadly conserved cyclic di-AMP (c-di-AMP) is a conditionally essential bacterial second messenger. The pool of c-di-AMP is fine-tuned through diadenylate cyclase and phosphodiesterase activities, and direct binding of c-di-AMP to proteins and riboswitches allows the regulation of a broad spectrum of cellular processes. c-di-AMP has a significant impact on intrinsic β -lactam antibiotic resistance in Gram-positive bacteria; however, the reason for this is currently unclear. In this work, genetic studies revealed that suppressor mutations that decrease the activity of the potassium (K^+) importer KupB or the glutamine importer GlnPQ restore cefuroxime (CEF) resistance in diadenylate cyclase (*cdaA*) mutants of *Lactococcus lactis*. Metabolite analyses showed that glutamine is imported by GlnPQ and then rapidly converted to glutamate, and GlnPQ mutations or c-di-AMP negatively affects the pools of the most abundant free amino acids (glutamate and aspartate) during growth. In a high-c-di-AMP mutant, GlnPQ activity could be increased by raising the internal K^+ level through the overexpression of a c-di-AMP-insensitive KupB variant. These results demonstrate that c-di-AMP reduces GlnPQ activity and, therefore, the level of the major free anions in *L. lactis* through its inhibition of K^+ import. Excessive ion accumulation in *cdaA* mutants results in greater spontaneous cell lysis under hypotonic conditions, while CEF-resistant suppressors exhibit reduced cell lysis and lower osmoresistance. This work demonstrates that the overaccumulation of major counter-ion osmolyte pools in c-di-AMP-defective mutants of *L. lactis* causes cefuroxime sensitivity.

IMPORTANCE The bacterial second messenger cyclic di-AMP (c-di-AMP) is a global regulator of potassium homeostasis and compatible solute uptake in many Gram-positive bacteria, making it essential for osmoregulation. The role that c-di-AMP plays in β -lactam resistance, however, is unclear despite being first identified a decade ago. Here, we demonstrate that the overaccumulation of potassium or free amino acids leads to cefuroxime sensitivity in *Lactococcus lactis* mutants partially defective in c-di-AMP synthesis. It was shown that c-di-AMP negatively affects the levels of the most abundant free amino acids (glutamate and aspartate) in *L. lactis*. Regulation of these major free anions was found to occur via the glutamine transporter GlnPQ, whose activity increased in response to intracellular potassium levels, which are under c-di-AMP control. Evidence is also presented showing that they are major osmolytes that enhance osmoresistance and cell lysis. The regulatory reach of c-di-AMP can be extended to include the main free anions in bacteria.

Citation Pham HT, Shi W, Xiang Y, Foo SY, Plan MR, Courtin P, Chapot-Chartier M-P, Smid EJ, Liang Z-X, Marcellin E, Turner MS. 2021. Cyclic di-AMP oversight of counter-ion osmolyte pools impacts intrinsic cefuroxime resistance in *Lactococcus lactis*. mBio 12:e00324-21. <https://doi.org/10.1128/mBio.00324-21>.

Invited Editor Maria L. Marco, University of California, Davis

Editor Kimberly A. Kline, Nanyang Technological University

Copyright © 2021 Pham et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mark S. Turner, m.turner2@uq.edu.au.

Received 9 February 2021

Accepted 8 March 2021

Published 8 April 2021

KEYWORDS cyclic di-AMP, antibiotic resistance, osmolyte, regulation, *Lactococcus*, lactic acid bacteria, osmolytes, osmoregulation, second messenger

Cyclic di-AMP (c-di-AMP) is a class of diffusible nucleotide second messengers that transmit signals based on their intracellular concentration (1). Once a c-di-AMP concentration threshold is reached, it is able to bind to protein and riboswitch receptors to alter their activity (2). Most *Firmicutes* contain one diadenylate cyclase (DAC) (named either CdaA or DacA) and one or two phosphodiesterases (PDEs) (GdpP or PgpH), which fine-tune the intracellular c-di-AMP level (3). Despite a significant number of different c-di-AMP receptors being identified, most play a role in the regulation of K⁺ or compatible solute (e.g., carnitine and glycine-betaine) accumulation (2). In several bacterial genera, a common role for c-di-AMP in adaptation to environmental osmolarity has been identified and aligns well with the receptors to which c-di-AMP binds (4). PDE mutants with elevated c-di-AMP grow poorly in high-osmolarity media (5). In contrast, a lack of c-di-AMP can restrict bacterial growth and result in lysis of cells in rich media due to the overaccumulation of a range of osmolytes, including peptides, K⁺, and glycine-betaine (6–10). Growth of *cdaA* mutants can be rescued in high-osmolarity growth media (6, 11) or through the omission of major osmolytes that are imported by bacteria present in growth media (6–8), suggesting that the internal osmotic pressure is high in *cdaA* mutants.

Another common phenotype observed in bacterial mutants with altered c-di-AMP levels is varying resistance to cell wall-acting antibiotics of the β -lactam family. β -Lactam antibiotics inhibit the transpeptidase activity of penicillin-binding proteins needed to cross-link the glycan strands in peptidoglycan. β -Lactam treatment results in decreased peptidoglycan cross-linking and cell lysis (12). In different species, *gdpP* mutants have shown elevated β -lactam resistance (13–15), and *cdaA* (*dacA*) mutants exhibited β -lactam sensitivity (6, 16, 17). Clinical isolates of *Staphylococcus aureus* that have developed intrinsic β -lactam resistance have also been found to contain destructive mutations in the *gdpP* gene (18–21). The mechanism by which c-di-AMP regulates β -lactam resistance is, however, unclear despite the first evidence of a connection being reported a decade ago. Two hypotheses have been proposed to explain how c-di-AMP might regulate β -lactam resistance. The first is that c-di-AMP regulates cell wall peptidoglycan synthesis. This could be possible in part via the phosphoglucosamine mutase enzyme GlmM, which is involved in peptidoglycan precursor biosynthesis. The *glmM* gene is colocated in an operon with the most common c-di-AMP synthesis gene, and GlmM regulates c-di-AMP synthesis through direct binding to CdaA/DacA (22–24). The second hypothesis is that a lack of c-di-AMP results in the overaccumulation of intracellular osmolytes that increase internal osmotic pressure to such an extent that the β -lactam-weakened cell wall is unable to prevent lysis (5, 6). This is supported by results showing that β -lactam resistance in c-di-AMP-depleted mutants can be rescued in high-osmolarity media containing sucrose or NaCl (6, 14).

In this study, we identified and characterized genes that contribute to sensitivity to the β -lactam antibiotic cefuroxime (CEF) in c-di-AMP synthesis mutants of the model lactic acid bacterium *Lactococcus lactis*. CEF was chosen since the depletion of c-di-AMP in *Bacillus subtilis* and *Listeria monocytogenes* results in strong CEF hypersensitivity (6, 14, 16). We found that the overaccumulation of K⁺ and the major anionic amino acids (Glu and aspartate [Asp]) contributes to CEF sensitivity in *L. lactis* *cdaA* mutants. CEF-resistant suppressor mutants became sensitive to osmotic stress and lysed significantly less in a hypotonic environment, suggesting that mutants with reduced osmotic pressure have greater stability. We also identified the mechanism by which c-di-AMP regulates the dominant anionic amino acid levels in *L. lactis*.

RESULTS

CEF-resistant suppressors of *cdaA* mutants commonly contain mutations in a K⁺ or amino acid uptake system. Previously, we identified a large number of independent mutations in *cdaA* in a suppressor screen of different high-c-di-AMP *gdpP*




FIG 1 Isolation of cefuroxime-resistant suppressors from mutants defective in *cdaA*. (A) Osmoresistance and CEF resistance of the wild type, a *gdpP* mutant (*gdpP-1*), and a variety of osmoresistant *cdaA* suppressor mutants (labeled with green writing) obtained from parent *gdpP-1* and *gdpP-2* strains. For this and all other dilution spot plates below, spots are 10 µl of 10-fold serial dilutions of mid-log-phase cultures starting from a 10⁻¹ dilution at the top. (B) CEF resistance of strains using a disk diffusion assay with 0.15 µg CEF per disk. (C) Confirmation of CEF-resistant suppressors (labeled with blue writing) obtained from the *cdaA-1* and *cdaA-2* strains.

mutants under elevated NaCl concentrations (22, 25). Most of the *cdaA* mutants, which were all more osmoresistant than their *gdpP* mutant parent, contained single amino acid changes in CdaA, while some possessed frameshift mutations or mutations affecting the *cdaA* ribosome-binding site (Fig. 1A). Here, we tested a range of different *cdaA*




FIG 2 Loss-of-function mutations in *kupB* restore CEF resistance in the *cdaA-2* strain. (A) Location of mutations in KupB in CEF-resistant suppressors of the *cdaA-2* strain. (B) Promoter activity of wild-type and *kupB-2* mutant promoters in the WT strain background. Means \pm standard deviations (SD) from three independent biological replicate experiments are shown. ***, $P < 0.001$ using an unpaired *t* test. (C) CEF resistance of strains, including the *kupB-2* strain containing a WT copy of *kupB* introduced on the pGh9 plasmid.

suppressor mutants for their CEF resistance. Out of 11 *cdaA* mutants, 6 were CEF sensitive (Fig. 1A). We found that in a disk diffusion assay with CEF, strongly growing colonies frequently formed within the inhibition zone of the *cdaA-1* (*cdaA* mutant 1) strain but not its *gdpP-1* parent strain (Fig. 1B). Purification of these colonies and retesting using the disk diffusion assay showed that they underwent one or more mutations to become resistant to CEF (see an example of the *glnP-3* strain in Fig. 1B). To better understand why *L. lactis* *cdaA* mutant strains are sensitive to CEF, we characterized suppressor mutations that restored CEF resistance.

CEF-sensitive *cdaA-1* and *cdaA-2* strains were chosen for further study. They both have lower c-di-AMP levels than a *gdpP* mutant strain (see Fig. S1 in the supplemental material). They were plated with inhibitory levels of CEF, and 12 suppressor mutants were obtained and confirmed (Fig. 1C). Whole-genome sequencing (WGS) of these suppressors revealed that distinct mutations occurred in suppressors from the two different parent strains *cdaA-1* and *cdaA-2* (Table S1).

For the *cdaA-2* strain, 4 out of 5 CEF-resistant suppressors possessed mutations in *kupB* (Table S1). Additional isolation of CEF-resistant suppressors and sequencing of *kupB* revealed three more independent mutations in this gene (G28S, P526L, and F550L) (Fig. 2A). KupB is a K⁺ importer and has been previously identified as a c-di-AMP receptor protein in *L. lactis* (26), and gain-of-function mutations in *kupB* restored osmoresistance in a high-c-di-AMP *gdpP* mutant (25). Interestingly, one suppressor strain (*kupB-4*) also contained a transposon insertion upstream of *busAA* (Table S1). Analysis of this IS905 insertion revealed that it is not oriented in a direction that would provide activation of downstream genes, like that described previously (27, 28). Therefore, the insertion likely disrupts transcription initiated from the native *busAA* promoter located 135 bp upstream.

Interestingly, the KupB amino acid changes identified in our CEF-resistant suppressor screen are mostly located in transmembrane helices within the proximity of residues that interact with K⁺ or protons in KimA from *B. subtilis* (29) (Fig. S2). To determine if the mutations in *kupB* caused a gain or loss of function, we examined the effect of the mutation 27 bp upstream of *kupB* found in the *kupB-2* strain on gene expression using a *lacZ* reporter. It was found that the G \rightarrow T mutation reduced expression from the *kupB* promoter by 30% (Fig. 2B). Inspection of the upstream region did not reveal any obvious changes in -10 or -35 sigma factor recognition boxes, so the reason for this downregulation is unclear at this stage. Next, we introduced a wild-type (WT) copy

of *kupB* into the *kupB-2* suppressor, which restored CEF sensitivity (Fig. 2C). Taken together, these results demonstrate that reduced K⁺ uptake in the *kupB* suppressor mutants increases CEF resistance.

For the *cdaA-1* strain, 6 out of 7 CEF-resistant suppressors contained mutations in the amino acid ATP-binding cassette (ABC) transport system GlnPQ (Table S1). Additional isolation of CEF-resistant suppressors and sequencing of *glnPQ* revealed three more independent mutations in these genes (G569W and S621N in GlnP and H199Q in GlnQ) (Fig. 3A). GlnP is composed of a fusion of two substrate-binding domains (SBDs) to the transmembrane permease domain, and GlnQ is an ATPase (30–33). The primary substrate of GlnPQ is glutamine (Gln); however, other amino acids can be imported through this transporter with various affinities. To determine if the mutations in *glnPQ* caused a gain or loss of function, we introduced the wild-type *glnP* gene in the *glnP-1* strain, which lowered CEF resistance (Fig. 3B). Curing of the *glnP* expression plasmid from this strain resulted in the restoration of CEF resistance (Fig. 3B). Next, we compared the resistances of strains to the toxic Gln analog L-5-N-hydroxyglutamine. It was found that 2 of the 3 *glnPQ* suppressor mutants (*glnP-1* and *glnQ-1*) grew well in the presence of the toxic analog (Fig. 3C). It is likely that the *glnP-1* and *glnQ-1* strains contain more destructive *glnPQ* mutations than the *glnP-3* strain, which grew poorly at this concentration of analog tested. Next, we compared the growths of strains in chemically defined medium (CDM) with various Gln levels. The *glnQ-1* strain grew poorly compared with its *cdaA-1* parent strain in low-Gln medium (Fig. 3D). Interestingly, the *cdaA-1* and *gdpP-1* strains grew better and worse, respectively, than the WT at low Gln concentrations (Fig. 3D), suggesting that c-di-AMP may negatively influence the Gln uptake ability.

To obtain a c-di-AMP synthesis-defective strain completely defective in GlnPQ activity, we plated the *cdaA-1* strain onto agar containing an inhibitory concentration of the toxic analog L-5-N-hydroxyglutamine. Two analog-resistant suppressors were obtained (Fig. S3), and analysis of *glnP* revealed single nucleotide changes that introduced a TAA stop codon at codon 44 (*glnP-7*) or codon 442 (*glnP-6*). Both the *glnP-6* and *glnP-7* strains were more CEF resistant than their parent *cdaA-1* strain (Fig. 3E). They were also unable to fully grow in CDM unless very high levels of Gln were provided (Fig. 3F), showing that the Gln acquisition ability of the *glnP-6* and *glnP-7* strains is severely impaired. Taken together, these results demonstrate that destructive *glnPQ* mutations lead to reduced uptake of Gln (and possibly other amino acids), which results in CEF resistance in a c-di-AMP synthase-defective strain.

CEF-resistant suppressors possess an osmosensitive phenotype. We hypothesized that the CEF-resistant suppressors have lower concentrations of intracellular osmolytes (K⁺ or free amino acid pool), which either directly or indirectly results in reduced internal osmotic pressure leading to greater cell stability during CEF-induced cell wall weakening. A lower level of intracellular osmolytes would also be expected to reduce osmoresistance. The CEF-resistant suppressors tested were all found to be more sensitive to osmotic stress than their parent strains (*cdaA-1* or *cdaA-2*) (Fig. S4A). Some CEF-resistant strains (*kupB-1*, *kupB-4*, *glnP-1*, and *glnQ-1*) were found to be highly NaCl sensitive, suggesting that their mutations are more severe. The expression of *glnP* rescued NaCl resistance in the *glnP-1* strain (Fig. 3B), and the L-5-N-hydroxyglutamine-selected suppressors *glnP-6* and *glnP-7* were highly NaCl sensitive (Fig. 3E), confirming that in *cdaA* mutants, GlnPQ activity is required for growth under high osmolarity. We next determined if higher growth medium osmolarity could rescue the CEF resistance of the *cdaA-1* and *cdaA-2* strains. It was found that the addition of increased NaCl enhanced the growth of these strains on CEF-containing agar (Fig. S4B). Together, these data show that mutations that allow for CEF resistance lower the osmotic pressure within the cell, and CEF-sensitive *cdaA* mutant cells can be stabilized by elevated external osmolarity.

CEF-resistant suppressors exhibit reduced cell lysis. In c-di-AMP-depleted mutants of *B. subtilis* and *L. monocytogenes*, elevated cell lysis occurs during growth in rich media (14, 16). We examined if *cdaA* mutants of *L. lactis* would exhibit greater lysis




FIG 3 Loss-of-function mutations in GlnPQ restore CEF resistance in the *cdaA-1* strain. (A) Location of GlnPQ mutations in CEF-resistant suppressors obtained from the *cdaA-1* strain. Membrane-spanning regions were predicted using TOPCONS. Note that GlnPQ functions as a homodimer but is shown here as a monomer. (B) CEF and NaCl resistance of strains, including the *glnP-1* strain containing a WT copy of *glnP* introduced on the pGh9 plasmid. A strain in which the plasmid was cured from the *glnP-1-pGh9-glnP* strain was also included. (C) Growth of strains with the toxic glutamine analog L-5-N-hydroxyglutamine. (D) Growth (after 24 h) of strains in chemically defined medium with various glutamine concentrations. (E) CEF and NaCl resistance of strains, including the L-5-N-hydroxyglutamine-resistant suppressors obtained from the *cdaA-1* strain (*glnP-6* and *glnP-7*). (F) Growth (after 24 h) of strains in chemically defined medium with various glutamine concentrations. In panels C, D, and F, means \pm SD from three independent biological replicate experiments are shown.

during growth with CEF. Culture supernatants were examined for the presence of DNA and RNA by gel electrophoresis as an indicator of cell lysis. Both the *cdaA-1* and *cdaA-2* strains were found to lyse significantly when cultured with CEF, while CEF-resistant suppressors, the WT, and *gdpP* mutants showed no or minimal lysis (Fig. 4A). *L. lactis cdaA-1* was also found to undergo some lysis during growth in medium without CEF (Fig. 4A). To explore the role of osmotic pressure in the stability of strains, we compared the amounts of spontaneous lysis of washed cells grown to mid-log-phase, which were suspended in pure water (a hypotonic solution). *L. lactis cdaA-1* cells grown in GM17 and heart infusion (HI) media were relatively stable when suspended in water; however, cells grown in glucose-yeast (GY) broth were much more prone to lysis. It was found that both the *cdaA-1* and *cdaA-2* strains lysed more in water than CEF resistance suppressors, the WT, and *gdpP* mutants (Fig. 4B). To lower the internal osmotic pressure of cells, *cdaA* mutants were pregrown in GY medium with increasing concentrations of NaCl. This resulted in greater stability of cells following their resuspension in water, most likely due to a lowering of cell turgor pressure (Fig. 4C). These results suggest that c-di-AMP synthesis mutants possess high internal osmotic pressure, which leads to reduced cell stability.

We next examined if there are cell wall peptidoglycan changes in c-di-AMP-defective strains that may be contributing to CEF sensitivity and reduced cell stability. Remarkably, the *cdaA-1* strain, which is CEF sensitive and exhibited less stability than the other strains, had a cell wall that was the same thickness as that of the WT and thicker than the cell walls of the *gdpP-1* and *glnP-1* strains (Fig. S5A). This suggests that a defect in c-di-AMP synthesis in *L. lactis* does not negatively affect cell wall biosynthesis. In previous work, we found that an *L. lactis gdpP* mutant contained elevated peptidoglycan precursor UDP-N-acetylglucosamine (UDP-NAG) levels, which were lowered upon mutation of the phosphoglucosamine mutase gene *glmM* (22). Here, we measured UDP-NAG levels and found that they negatively correlated with cell wall thickness. UDP-NAG levels were higher in mutants with thinner cell walls, indicating that slower cell wall biosynthesis may result in the accumulation of peptidoglycan precursors (Fig. S5B). Peptidoglycan muropeptides and the peptidoglycan cross-linking index of strains were also analyzed (Fig. S5C). Since CEF inhibits peptidoglycan cross-linking (34), cells with reduced cross-linking may exhibit greater sensitivity. However, a small but statistically significant increase in cross-linking was observed in the CEF-sensitive *cdaA-1* mutant compared to its parent *gdpP-1* strain (Fig. S5C). Taken together, cell wall analyses did not provide an explanation for the variations in CEF resistance and cell integrity observed in the strains examined here.

Mutations in GlnPQ lower intracellular Gln, Glu, and Asp levels. While the roles of K⁺ and the c-di-AMP-binding receptor KupB have been studied previously in the osmoresistance of *L. lactis* (25, 26), the role of GlnPQ in c-di-AMP-regulated processes has not been reported. GlnPQ has been found to bind and transport 3 different amino acids (Gln, Glu, and Asn) with different affinities and rates (30, 32, 33). GlnPQ transports only the protonated form of Glu, not the anion, which is the dominant species at physiological pH, and its capacity to transport Asn is lower than its capacity to transport Gln (33). It is the sole transport system in *L. lactis* for the essential amino acids Gln and Glu (32). This work points to Gln being the primary target of GlnPQ.

The CEF-sensitive *cdaA-1* strain contained a higher level of Gln than its parent *gdpP-1* strain, which had a low level like the *glnPQ* suppressors (Fig. 5A). Quantitation of all free amino acid levels in WT *L. lactis* revealed that Asp and Glu were by far the most abundant, present at ~100-fold-higher concentrations than Gln (Fig. 5B). Interestingly, their levels also varied significantly between the strains tested. The *cdaA-1* strain contained significantly higher Glu and Asp levels than its CEF-resistant *glnPQ* suppressors (Fig. 5A). In addition, the c-di-AMP level had a negative effect on Glu and Asp levels, with *cdaA-1* and *gdpP-1* mutants having significantly higher and lower levels, respectively, than their parents (Fig. 5A). From this, we hypothesized that following import by GlnPQ, Gln is converted to Glu and Asp, which together form a major anionic solute




FIG 4 CEF-resistant suppressors release less DNA and RNA during growth with CEF or following resuspension in water. (A) Agarose gel electrophoresis of supernatants from strains grown for various times in HI broth with various CEF concentrations. The locations of genomic DNA (gDNA) and rRNA are indicated. (B) Agarose gel electrophoresis analysis of the supernatant following resuspension of cells (grown to mid-log phase in GY medium) in water. (C) Agarose gel electrophoresis analysis of the supernatant following resuspension of cells (grown to mid-log phase in GY medium with different NaCl concentrations) in water. The gradient from left to right for each strain indicates cells grown in GY medium with 0, 0.05, 0.1, 0.15, and 0.2 M NaCl. The M lane indicates the DNA ladder.




FIG 5 GlnPQ mutations lower the levels of Gln, Glu, and Asp in CEF-resistant suppressors. (A) Intracellular levels of Gln, Glu, and Asp in the indicated strains. Data are means \pm SD from three independent biological replicate experiments. ***, $P < 0.001$ using one-way analysis of variance (ANOVA) followed by Tukey's test for multiple comparisons. (B) All intracellular free amino acid levels in WT *L. lactis*. The inset shows amino acids of low abundance. Data are means \pm SD from three independent biological replicate experiments. (C) Proposed pathway for the conversion of Gln to Glu and Asp in *L. lactis*. Note that several other enzymes in *L. lactis* can convert Gln to Glu during nucleotide, nucleotide-sugar, and amino acid biosynthesis. Pyr, pyruvate; OA, oxaloacetate; α -KG, α -ketoglutarate; GltB and GltD, glutamate synthase large and small subunits, respectively; AspC, aspartate aminotransferase. The red circles depict c-di-AMP binding to pyruvate carboxylase (PC).

pool. A proposed pathway showing the conversion of Gln to Glu and Asp in *L. lactis* is shown in Fig. 5C. The pathway also includes the c-di-AMP receptor pyruvate carboxylase (PC), which synthesizes the Asp precursor oxaloacetate. From these results, it was of interest to investigate how c-di-AMP levels might also affect GlnPQ activity and conversion of Gln to Glu and Asp.

Gln uptake by GlnPQ is activated by increased intracellular ionic strength and K^+ , which provides high levels of the counter-ion Glu. The experiments described above were carried out using cells grown in rich complex media, so intracellular amino acids can derive from both imported peptides as well as free extracellular amino acids. To verify that the changes in Glu and Asp levels seen in strains were specifically due to altered import of Gln by GlnPQ, we carried out Gln feeding assays with resting (nongrowing) cells in buffer with glucose (Fig. 6A). In the WT, rapid and large increases in intracellular Gln and Glu levels were observed; however, no increase in Asp was observed even after 60 min with Gln (Fig. 6B; Fig. S6). This suggests that Gln/Glu is not used to generate Asp in resting cells. We therefore used intracellular Glu as a marker for GlnPQ activity since its level continued to rise over 60 min, while the level of Gln fell due to its conversion to Glu. This confirms that following ATP-dependent import of Gln, rapid conversion to Glu occurs. The high-c-di-AMP *gdpP* mutant strain *gdpP-1* was unable to increase its Glu level to high levels, and its levels remained around 7-fold lower than that of the WT after 5 min (Fig. 6B). The restricted increase in Glu in the *gdpP-1* strain was the result of reduced activity of GlnPQ since intracellular Gln levels measured after 5 min of feeding with Gln were also much lower (~8-fold) than those of the WT. The *cdaA-1* strain contained high initial levels of Glu but further increased this level following Gln feeding to remain higher than that of the WT. The *glnP* suppressor mutant *glnP-1* had a low starting level of Glu and following Gln addition remained low, as expected, since the GlnPQ transporter is defective in this strain.

The finding that the *gdpP* mutant strain *gdpP-1* is unable to strongly increase Glu levels in this assay suggests that GlnPQ activity is inhibited in this strain. The *gdpP-1* strain contains a high level of c-di-AMP (22), which leads to inhibition of K^+ import through direct binding to KupB (25, 26). We hypothesized that GlnPQ activity is affected by ionic strength in *L. lactis* cells and that the low K^+ level observed in the *gdpP* mutant prevents the activation of GlnPQ. Exposing cells to elevated external osmotic conditions, which we predicted would increase the intracellular ionic strength, led to significantly higher intracellular Glu levels (Fig. 6C).

Next, we examined if an increase in intracellular K^+ in a high-c-di-AMP *gdpP* mutant could activate GlnPQ. The *gdpP-2* strain containing either a single copy or multiple copies of a constitutively active *kupB^{A618V}* gene variant accumulated significantly higher Glu levels following Gln feeding (Fig. 6D). Levels of Glu and Asp in the *gdpP-2* strain expressing *kupB^{A618V}* were also significantly higher in cells during growth in rich media (Fig. 6E). Therefore, the level of intracellular K^+ , which is inhibited by c-di-AMP binding to KupB, positively affects the accumulation of the major counter-ion amino acids Glu and Asp in *L. lactis*.

Finally, we were interested in determining if overfeeding of Gln to *cdaA-1* cells would trigger greater cell lysis in nongrowing cells. We grew *cdaA-1* cells in CDM with low Gln levels to decrease the Glu pool and reduce cell lysis. Cells were then incubated with either glucose only or glucose and Gln (Fig. 6F), washed, and then resuspended in water. It was found that *cdaA-1* cells incubated with glucose and Gln lysed more than the same batch of cells incubated with only glucose (Fig. 6G). Since this assay was performed using nongrowing cells, this confirms that lysis is the direct result of Gln uptake (and Glu overaccumulation) and is unrelated to a change in a metabolic process (e.g., cell wall biosynthesis). Gln feeding to the WT did not result in any observable increase in lysis, which suggests that its osmotic pressure is still lower than that in the *cdaA* mutant. This is likely due to the intact c-di-AMP system in the WT lowering the levels of other osmolytes within the cell, unlike the *cdaA* mutant.




FIG 6 GlnPQ activity controls intracellular Glu in response to intracellular K⁺, and overaccumulation of Glu leads to greater cell instability in the *cdaA-1* strain. (A) Pathway showing ATP-dependent Gln uptake by GlnPQ and conversion to Glu. (B) Intracellular Glu levels in resting cells following 5 min of incubation with glucose with or without Gln. The means from three independent biological replicate experiments are shown as horizontal bars. (C) Effect of increased osmolarity on Gln uptake in resting cells of the WT after 20 min in buffer. Levels are relative to those in cells in buffer with no additional osmolyte added. *, $P < 0.001$ using one-way ANOVA followed by Tukey's test for multiple comparisons. (D) Effect of expression of the c-di-AMP-insensitive KupB variant KupB^{A618V} in a single copy (*gdpP-2-kupB^{A618V}*) and multiple copies (*gdpP-2-pGh9-kupB^{A618V}*) on Gln uptake by resting cells of the high-c-di-AMP *gdpP-2* mutant in buffer. ***, $P < 0.001$ using one-way ANOVA followed by Tukey's test for multiple comparisons. (E) Intracellular Glu and Asp levels in *gdpP-2* cells expressing the c-di-AMP-insensitive K⁺ importer KupB^{A618V} in (Continued on next page)

DISCUSSION

c-di-AMP has emerged as a global osmoregulatory signal in numerous bacterial genera (35). Evidence has indicated that the link between c-di-AMP and β -lactam antibiotic resistance is an additional consequence of its regulation of intracellular osmolyte levels (5, 6). This hypothesis proposes that cells possessing higher internal osmotic pressure (i.e., *cdaA* mutants) will be more susceptible to osmotic lysis, especially when the stress-bearing peptidoglycan layer is compromised upon CEF exposure. Here, in a screen for suppressors that rescued the CEF resistance of partially defective *cdaA* *L. lactis* mutants, mutations that lowered the levels of major inorganic or organic ions (K^+ , Glu, and Asp) were found. GlnPQ and KupB suppressors grew poorly on media with elevated salt, demonstrating that these ions play important roles in osmoregulation in *L. lactis*. In nongrowing cells, Gln uptake (and Glu accumulation) triggered greater lysis of an *L. lactis* *cdaA* mutant, providing further evidence for osmotic pressure being an important contributor to cell instability in a strain with defective c-di-AMP synthesis. Reduced ion accumulation has also been observed in CEF-resistant suppressors of an *L. monocytogenes* *cdaA* mutant (6). Mutations that restricted PC activity led to lower levels of the citrate anion. With respect to a direct role of c-di-AMP in cell wall homeostasis, differences in peptidoglycan cross-linking or precursor synthesis have been identified previously in high-c-di-AMP mutants (13, 22, 36). Our analysis of peptidoglycan thickness, cross-linking, and precursor levels, however, did not reveal alterations that would appear to render the *L. lactis* *cdaA* mutant peptidoglycan more susceptible to CEF. Indeed, the high-c-di-AMP *gdpP* mutant, which had the thinnest cell wall and the least cross-linked peptidoglycan, was CEF resistant. A role for more subtle cell wall changes in CEF resistance, however, cannot be excluded. Therefore, our findings provide additional support for a model whereby *cdaA* mutants accumulate unhealthy levels of several different osmolytes, which, when combined, lead to a critical internal osmotic pressure that a normally structured cell wall is unable to fully withstand.

In support of this theory, we identified that the *cdaA-1* mutant contains a *busAA-AB* promoter mutation that destroys the transcription of the transporter for the compatible solute glycine-betaine (see Table S2 and Fig. S7 in the supplemental material). This mutation likely permitted the growth of this suppressor, which contains a severe *cdaA* frameshift mutation, on normal media. This is similar to that seen in a *Streptococcus agalactiae* *cdaA* mutant, which was viable only after the inactivation of its glycine-betaine transporter (8). Although viable, the *L. lactis* *cdaA-1* mutant is still sensitive to CEF, and a subsequent lowering of additional osmolytes through the inactivation of GlnPQ is necessary to restore CEF resistance. Therefore, c-di-AMP-controlled osmolytes likely have an additive effect on internal osmotic pressure whereby moderate osmolyte overaccumulation permits growth but generates a CEF-sensitive phenotype, while extreme osmolyte overaccumulation results in a complete loss of viability. It is well established that the turgor pressure of Gram-positive bacteria is up to 10-fold higher than that of Gram-negative bacteria and therefore needs to be tightly controlled (37). Our findings and those of others (5–9, 11, 25, 38) suggest that phenotypes affecting the growth and cell integrity of low- and high-c-di-AMP mutants can in most part be explained by variations in the levels of turgor-inducing internal osmolytes.

In *L. lactis*, c-di-AMP negatively regulates Asp, K^+ , and glycine-betaine levels through direct binding to PC, Kup homologs, BusR, and BusAA/OpuAA (25, 26, 39, 40). Its regulatory reach can now be extended to the control of Glu (and Asp) levels through indirect regulation of GlnPQ. Glu is the major anion in most bacteria, and its accumulation allows the cell to balance the charge of significant levels of K^+ (41, 42). In the *L. lactis* WT, Glu is present at a high level, but the other anionic amino acid Asp

FIG 6 Legend (Continued)

cells grown in GM17 medium. (F) Model for enhanced lysis in Glu-loaded cells. (G) DNA and RNA release from cells resuspended in water after being grown in CDM with a low glutamine concentration (100 μ M) and then incubated for 30 min with glucose with or without Gln in buffer. Quantitative data are presented as means \pm SD from three independent biological replicate experiments.

is also present at a similar concentration. Together, these two amino acids represent 55% of the total free amino acids in WT *L. lactis* (Fig. 5B). In *L. lactis*, Glu (and Gln) is unable to be synthesized due to an incomplete tricarboxylic acid (TCA) cycle and needs to be sourced from peptides or amino acids external to the cell (33). The GlnPQ transporter in *L. lactis* plays an important role through its ability to efficiently transport Gln with both high and low affinities (33). Studies of *L. lactis* growing in defined media revealed that out of 20 amino acids provided to cells in defined media, Gln was the most consumed amino acid, accounting for up to 50% of the total nitrogen imported (43). Therefore, Gln uptake and conversion are significant processes in *L. lactis* and allow the cell to generate large osmolyte pools of Glu and Asp. Here, we provide evidence that in *L. lactis*, c-di-AMP has a significant influence on GlnPQ transporter activity through its control of K⁺ levels via direct binding to KupB (25, 26, 44). c-di-AMP can therefore modulate the levels of the major K⁺ counter-ions Glu and Asp. The presence of structurally and functionally similar GlnPQ transport systems in streptococci and enterococci (30) suggests that this regulation mechanism likely extends beyond *L. lactis*. Based on our work and others, a model for c-di-AMP-regulated homeostasis of the major intracellular ions and their effect on CEF resistance in *L. lactis* can be proposed (Fig. 7).

Recent work in *S. aureus* and *B. subtilis* identified suppressor mutations in Gln and Glu transporters, respectively, which rescued the growth of mutants devoid of c-di-AMP (9, 38). These results align well with those found in *L. lactis* and suggest that cells defective in c-di-AMP production are unable to regulate intracellular levels of major osmolyte amino acids. The mode of regulation of the mutated transporters (AlsT, AimA, and YfkC) is not known; however, it would be of interest to explore if they are controlled indirectly by c-di-AMP, like GlnPQ in *L. lactis*. In support of our findings, a previous *in vitro* characterization of *L. lactis* GlnPQ found that Gln uptake in reconstituted proteoliposomes was activated ~4-fold by an increased salt concentration in the lumen (32). In other work, the ATPase component (GlnQ) of *S. agalactiae* was found not to bind c-di-AMP (8), therefore making it unlikely that c-di-AMP-regulated Gln uptake occurs via a direct interaction. Indeed, when we overexpressed constitutively active KupB^{A618V} in the *gdpP-2* strain, which triggers an accumulation of c-di-AMP (25), GlnPQ was found to be more active due to elevated K⁺ import (Fig. 6D). Recent work in *B. subtilis* has found that Glu availability affects K⁺ import activity by KtrCD (45), further confirming the need for alignment of ionic osmolyte levels. Coordination of K⁺ and anion levels ensures a balancing of the charge within the cell, and by controlling both through a common orchestrator (c-di-AMP), it can occur efficiently and rapidly.

MATERIALS AND METHODS

Strains, media, and chemicals used. *L. lactis* strains (see Table S2 in the supplemental material) were routinely grown in nonshaking tubes at 30°C in M17 medium (Difco, USA) supplemented with 0.5% (wt/vol) glucose (GM17). When needed, 3 µg/ml erythromycin (Em) was added to the media for *L. lactis*. DNA release assays were carried out on cells grown in either heart infusion (HI) medium (Oxoid) or glucose-yeast (GY) medium (1.33% yeast extract [Oxoid], 1.33% glucose, and 1% 0.1 M K₂HPO₄). To characterize the growth of *L. lactis* under different concentrations of glutamine (Gln), chemically defined minimal medium (CDM) using two types of Dulbecco's modified Eagle's medium (DMEM) (Merck) was used as the base (39). The first type of DMEM (catalog no. 5671) contains 4.5 g/liter glucose and sodium bicarbonate, while the second type of DMEM (catalog no. 5546) contains 1 g/liter glucose and 0.11 g/liter pyruvic acid. Both media are devoid of Gln and Glu. These media were supplemented with histidine at 0.13 mg/ml, arginine at 0.72 mg/ml, leucine at 1 mg/ml, valine at 0.6 mg/ml, glucose at 0.15%, sodium acetate at 0.75 mg/ml, morpholinepropanesulfonic acid (MOPS) at 13 mg/ml, guanine at 0.05 mg/ml, xanthine at 0.05 mg/ml, FeSO₄ at 0.1 mg/ml, ZnSO₄ at 0.1 mg/ml, and adenine at 0.2 mg/ml. Unless stated otherwise, cells were washed and resuspended in KPM buffer (0.1 M K₂HPO₄ adjusted with H₃PO₄ acid to pH 6.5 and supplemented with 10 mM MgSO₄) (46). *Escherichia coli* NEB-5α cells containing pGh9 derivatives were grown in HI medium containing 150 µg/ml Em at 30°C with aeration at 250 rpm.

c-di-AMP extraction and quantification. c-di-AMP from *L. Lactis* was extracted as previously described (25). c-di-AMP was detected and quantified by liquid chromatography-coupled tandem mass spectrometry (LCMS-8060; Shimadzu, Japan). Chromatographic separation was performed on an ultra-high-pressure liquid chromatography (UHPLC) Nextera X2 instrument using a Shim-pack Velox SP-C₁₈ column (1.8 µm, 2.1 by 150 mm; Shimadzu, Japan). Eluents A and B consisted of 0.05% (vol/vol) formic acid in water and acetonitrile (Merck), respectively. The sample volume was 10 µl with a flow rate of




FIG 7 Proposed model for changes leading to CEF sensitivity in the *cdaA-1* strain and CEF resistance in *glnPQ* suppressor mutant strains. (A) The WT normally contains low c-di-AMP levels, resulting in high K⁺ uptake by KupB. High intracellular K⁺ levels lead to active Gln import by GlnPQ and conversion to high levels of anionic Glu and Asp. PC is also uninhibited in this strain and contributes to the Asp pool. WT cells can modulate their c-di-AMP level in response to environmental stressors. (B) Inactivation of GdpP in the *gdpP-1* strain results in a high level of c-di-AMP, which binds and reduces K⁺ uptake by KupB. Lower intracellular K⁺ concentrations result in less activation of GlnPQ and smaller Glu and Asp pools. PC is also inhibited in this strain by c-di-AMP. This strain is highly salt sensitive due to low levels of intracellular osmolytes. (C) Selection for salt-resistant mutants of the *gdpP-1* strain resulted in partial CdaA inactivation and low-level c-di-AMP synthesis, leading to very high K⁺ uptake by an uninhibited KupB. A high level of K⁺ strongly activates Gln uptake by GlnPQ, and high Glu and Asp pools form. In this mutant, PC is also not inhibited by c-di-AMP and increases the production of the Asp precursor oxaloacetate. This mutant is CEF sensitive due to osmolyte overaccumulation. (D) Selection of CEF-resistant suppressors of the *cdaA-1* strain results in partial inactivation of GlnPQ, which lowers Gln uptake and the major pools of Glu and Asp. This reduces major intracellular anions. PC is still uninhibited in this strain, but the lower levels of Glu result in lower levels of Asp. c-di-AMP is depicted as two red circles, and its breakdown product pApA is 5'-phosphadenylyl-adenosine. Black or gray arrows show transport or enzymatic conversion steps. Thick arrows indicate greater flux, while dashed arrows indicate reduced flux. Green arrows indicate activation of GlnPQ by K⁺. Red lines indicate binding and inhibition of KupB and PC by c-di-AMP.

0.3 ml min⁻¹. Eluent A (95%) was used from 0 to 1 min, followed by a linear gradient from 95% to 50% eluent A until 10 min. The column was then washed with 90% eluent B for 3 min and then reequilibrated with 95% eluent A for 2 min prior to reinjection. The internal standard of azidothymidine (AZT) (Sigma) was used. c-di-AMP was detected with a triple-quadrupole mass spectrometer equipped with an electrospray ionization

source using multiple reaction monitoring transitions of m/z 657→124 in negative ionization mode. Data obtained were curated using LabSolutions Insight version 3.2 SP1 and LabSolutions Postrun/QuantBrowser version 5.95 (Shimadzu Corporation).

Isolation of CEF-resistant suppressors and WGS. The *cdaA* mutant strains *cdaA-1* and *cdaA-2* were streaked or spread either from mid-log-phase cultures, from broth cultures grown overnight, or directly from frozen glycerol stocks (40% glycerol) onto GM17 agar containing $\geq 0.08 \mu\text{g/ml}$ CEF (Merck) and incubated for 2 days at 30°C. Colonies were picked and restreaked on agar containing the same concentration of CEF, from where they were obtained to ensure purity. CEF resistance confirmation was carried out by serial dilution of mid-log-phase cultures onto GM17 agar with CEF.

CEF disk diffusion assays were carried out by mixing 5 μl of a mid-log-phase culture (optical density [OD] of ~ 0.7) with 7 ml of 0.75% GM17 agar and pouring the culture onto a 15-ml 1.5% GM17 agar base. Following drying, a sterile 8-mm disk was placed on the top agar, and a 10- μl solution of CEF was added (0.15 μg). Following incubation overnight, zones of inhibition were observed. CEF-resistant suppressors were checked for *cdaA* back-mutations using PCR (Table S3), as described previously (22), before being analyzed by WGS. Genomic DNA extractions were performed as described previously (47). Sequencing was performed using the Illumina NovaSeq 6000 platform (Macrogen, South Korea). Single nucleotide polymorphisms (SNPs) were analyzed using Geneious Prime (Biomatters Ltd., New Zealand) (22, 25).

Genetic manipulation of strains. Plasmids and primers used in this study are shown in Tables S2 and S3, respectively. Electroporation of *L. lactis* strains was done as previously described (48), with minor changes for some strains. Following electroporation of pGh9-*kupB* and pGh9-*glnP* into the *kupB-2* and *glnP-1* strains, respectively, cells were plated onto GM17 agar with 3 $\mu\text{g/ml}$ Em supplemented with 0.1 M NaCl. The activities of promoters were determined using pTCV-lac (49). The WT promoter of *kupB* and the mutated variant from the *kupB-2* strain were amplified and cloned into pTCV-lac and assayed for activity in the *L. lactis* WT using a β -galactosidase assay.

Isolation of L-5-N-hydroxyglutamine-resistant suppressors. The *cdaA* mutant strain *cdaA-1* was streaked onto GM17 agar containing 1 mM the toxic glutamine analog L-5-N-hydroxyglutamine (Merck) and incubated for 2 days at 30°C. Colonies were picked and restreaked on agar containing the same concentration of the analog to ensure purity. Resistance confirmation was carried out by the serial dilution drop plate method as described above.

Cell wall thickness analysis. Cells grown to mid-log phase were fixed with 2.5% glutaraldehyde (ProSciTech) in phosphate-buffered saline (pH 7.4), and after washing in buffer, they were postfixed in 1% osmium tetroxide. They were then gradually dehydrated in ethanol (30 to 100%), infiltrated with a gradual increase in the concentration of Epon resin, and then polymerized for 2 days at 60°C. Ultrathin (60-nm) sections were collected onto 200-mesh copper grids and stained with uranyl acetate and lead citrate. Grids were examined using a Hitachi HT7700 electron microscope (Hitachi, Japan) operated at 80 kV. Images were acquired with a complementary metal oxide semiconductor (CMOS) camera (Advanced Microscopy Techniques), and peptidoglycan thickness was measured on micrographs.

UDP-NAG quantitation. Mid-log-phase (OD at 600 nm [OD₆₀₀] of ~ 0.6) cells were collected by centrifugation at $5,000 \times g$ for 10 min at 4°C and washed 2 times with 1/10 KPM buffer. UDP-NAG was extracted and quantified as previously described (22).

Peptidoglycan muropeptide and cross-linking analysis. Peptidoglycan was extracted from exponential-phase cells (OD₆₀₀ of ~ 0.8) and then digested with mutanolysin as described previously (50). The resulting soluble muropeptides were reduced with sodium borohydride and separated by reverse-phase UHPLC (RP-UHPLC) with a 1290 chromatography system (Agilent Technologies) and a Zorbax Eclipse Plus C₁₈ Rapid Resolution High Definition column (100 by 2.1 mm with a particle size of 1.8 μm ; Agilent Technologies) at 50°C using ammonium phosphate buffer and a methanol linear gradient as described previously (51). Muropeptides were identified according to their retention times by comparison with an *L. lactis* muropeptide reference chromatogram (51). The different muropeptides were quantified by integration of the peak areas, and the percentage of each peak was calculated as the ratio of its area over the sum of all peak areas. The peptidoglycan cross-linking index was calculated according to methods described previously (52), as follows: $(1/2 \sum \text{dimers} + 2/3 \sum \text{trimers} + 3/4 \sum \text{tetramers}) / \sum \text{all muropeptides}$.

Quantification of amino acid pools in growing *L. lactis* cells. Cells were grown in 30 ml GM17 medium until an OD₆₀₀ of ~ 0.7 was reached, collected by centrifugation at $5,000 \times g$ for 10 min at 4°C, and washed twice in 1/10 KPM buffer. After resuspension in 1.8 ml of 50% acetonitrile, cells were lysed using a Precellys 24 homogenizer (Bertin Technologies) with a 0.5-ml equivalent of 0.1-mm zirconia/silica beads (6,000 rpm for 30 s and repeated 3 times, with chilling on ice between repeats). Following centrifugation at $17,000 \times g$ for 15 min at 4°C, the supernatant was mixed 1:1 with an internal standard of sarcosine and 2-aminobutanoic acid. Amino acids were derivatized and analyzed with an Agilent 1200-SL HPLC system with a fluorescence detector (FLD) (catalog no. G1321A; Agilent) as described previously (53).

Extraction and quantification of intracellular Glu in the Gln uptake assay. Cells were grown in 50 ml of CDM containing a low level of Gln (200 μM) until an OD₆₀₀ of ~ 0.4 was reached. Following centrifugation at $5,000 \times g$ for 10 min at 25°C, cells were washed twice in KPM buffer and then resuspended in 3 ml of 1/10 KPM buffer. Uptake assays were performed using 0.5 ml of cells. Cells were energized with 20 mM glucose (final concentration) first before adding Gln (1 mM final concentration). The uptake assay mixture was incubated at 30°C for 5 min. Control reaction mixtures without glucose were included. For osmotic treatments, NaCl, KCl, and sucrose were added before Gln and incubated at 30°C for 20 min. After incubation, samples were centrifuged at $17,000 \times g$ for 1 min at 4°C and washed twice with KPM buffer. Thereafter, Glu was extracted from cells using acetonitrile-methanol-H₂O at a ratio of 2:2:1 using the same method as that described previously for c-di-AMP extraction (25). The supernatant (600 μl) was dried in an RVC 2-18 CDplus rotation vacuum concentrator (Christ). Glu and Gln were measured using

the Glu assay kit (catalog no. MAK004-1KT; Merck) or the Gln and Glu determination kit (catalog no. GLN1-1KT; Merck), with minor adjustments to the protocols. For the Glu assay kit, the dried samples were resuspended in 50 μ l of Glu assay buffer, vortexed well, and centrifuged at $16,000 \times g$ for 5 min at 25°C. A portion (2 μ l) was added into the kit master mix before incubation and reading of the OD₄₅₀ using a NanoDrop One instrument (Thermo Fisher Scientific). For quantification, Glu standards were prepared at 1,000 to 31.25 μ M in serial 2-fold dilutions.

DNA/RNA release (lysis) assays during growth and under hypotonic conditions. Strains were grown in HI broth until an OD₆₀₀ of ~0.2 was reached. HI broth was chosen instead of GM17 medium for this experiment as the latter produced fluorescent smears in the gels, making it less sensitive. The culture was split into 10-ml volumes, where CEF was added at different concentrations. Samples (100 μ l) were collected every 2 h and then centrifuged at $17,000 \times g$ for 3 min. The supernatant (20 μ l) was taken and analyzed for the presence of genomic DNA and RNA by agarose gel electrophoresis using SYBR Safe stain (Invitrogen) and the 1-kb Plus DNA ladder (Thermo Fisher Scientific).

Lysis was also tested for strains suspended in hypotonic liquid. *L. lactis* strains were grown in GY broth with or without NaCl until an OD₆₀₀ of ~0.5 was reached. Cells (1.5 ml) were collected by centrifugation at $12,000 \times g$ for 3 min, and the pellets were then washed with KPM followed by 1/10 KPM and centrifuged at $12,000 \times g$ for 1 min. Cells were resuspended in 100 μ l of MilliQ (Merck)-treated deionized water and centrifuged within 1 min at $17,000 \times g$ for 3 min. The supernatant (20 μ l) was taken and analyzed by agarose gel electrophoresis as described above.

The effect of Gln uptake on cell lysis in hypotonic liquid was also examined. WT and *cdaA-1* strains (15 ml) were grown to an OD₆₀₀ of ~0.6 in CDM broth containing a low level of Gln (100 μ M), washed twice with KPM buffer before being resuspended in 1 ml of KPM buffer, and then divided into 2 500- μ l aliquots for the sample and control. To obtain high internal Glu levels, 30 mM glucose and 10 mM Gln were added to cell suspensions and incubated at 30°C for 30 min. The control samples contained glucose but no Gln. After incubation, cells were centrifuged and washed with KPM buffer and then 1/10 KPM buffer at $12,000 \times g$ for 1 min. The cell pellet was resuspended in 200 μ l of MilliQ-treated deionized water and centrifuged within 1 min at $17,000 \times g$ for 3 min. The supernatant (20 μ l) was taken and analyzed by agarose gel electrophoresis as described above.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

FIG S1, DOCX file, 0.02 MB.

FIG S2, DOCX file, 0.8 MB.

FIG S3, DOCX file, 1.7 MB.

FIG S4, DOCX file, 1.5 MB.

FIG S5, DOCX file, 0.04 MB.

FIG S6, DOCX file, 0.02 MB.

FIG S7, DOCX file, 2.1 MB.

TABLE S1, DOCX file, 0.01 MB.

TABLE S2, DOCX file, 0.03 MB.

TABLE S3, DOCX file, 0.01 MB.

ACKNOWLEDGMENTS

We acknowledge the work of Wanqing Zhang and Andrea Yeap for their assistance with experiments.

Research funding for this project awarded to M.S.T., E.M., and Z.-X.L. is from the Australian Research Council (grant DP190100827). Elements of this research used equipment from the Queensland node of Metabolomics Australia funded by Bioplatforms Australia, an NCRIS-funded initiative.

REFERENCES

- Corrigan RM, Grundling A. 2013. Cyclic di-AMP: another second messenger enters the fray. *Nat Rev Microbiol* 11:513–524. <https://doi.org/10.1038/nrmicro3069>.
- He J, Yin W, Galperin MY, Chou SH. 2020. Cyclic di-AMP, a second messenger of primary importance: tertiary structures and binding mechanisms. *Nucleic Acids Res* 48:2807–2829. <https://doi.org/10.1093/nar/gkaa112>.
- Commichau FM, Heidemann JL, Ficner R, Stulke J. 2019. Making and breaking of an essential poison: the cyclases and phosphodiesterases that produce and degrade the essential second messenger cyclic di-AMP in bacteria. *J Bacteriol* 201:e00462–18. <https://doi.org/10.1128/JB.00462-18>.
- Turner MS, Vu TNM, Marcellin E, Liang Z-X, Pham HT. 2020. Osmoregulation via cyclic di-AMP signaling, p 177–189. In Chou S-H, Guilian N, Lee VT, Römling U (ed), *Microbial cyclic di-nucleotide signaling*. Springer, Cham, Switzerland.
- Commichau FM, Gibhardt J, Halbedel S, Gundlach J, Stulke J. 2018. A delicate connection: c-di-AMP affects cell integrity by controlling osmolyte transport. *Trends Microbiol* 26:175–185. <https://doi.org/10.1016/j.tim.2017.09.003>.
- Whiteley AT, Garelis NE, Peterson BN, Choi PH, Tong L, Woodward JJ, Portnoy DA. 2017. c-di-AMP modulates *Listeria monocytogenes* central metabolism to regulate growth, antibiotic resistance and osmoregulation. *Mol Microbiol* 104:212–233. <https://doi.org/10.1111/mmi.13622>.
- Gundlach J, Herzberg C, Kaever V, Gunka K, Hoffmann T, Weiß M, Gibhardt J, Thürmer A, Hertel D, Daniel R, Bremer E, Commichau FM, Stülke J. 2017. Control of potassium homeostasis is an essential function

- of the second messenger cyclic di-AMP in *Bacillus subtilis*. *Sci Signal* 10: eaal3011. <https://doi.org/10.1126/scisignal.aal3011>.
8. Devaux L, Sleiman D, Mazzuoli MV, Gominet M, Lanotte P, Trieu-Cuot P, Kaminski PA, Firon A. 2018. Cyclic di-AMP regulation of osmotic homeostasis is essential in group B *Streptococcus*. *PLoS Genet* 14:e1007342. <https://doi.org/10.1371/journal.pgen.1007342>.
 9. Zeden MS, Kviatkovski I, Schuster CF, Thomas VC, Fey PD, Grundling A. 2020. Identification of the main glutamine and glutamate transporters in *Staphylococcus aureus* and their impact on c-di-AMP production. *Mol Microbiol* 113:1085–1100. <https://doi.org/10.1111/mmi.14479>.
 10. Whiteley AT, Pollock AJ, Portnoy DA. 2015. The PAMP c-di-AMP is essential for *Listeria monocytogenes* growth in rich but not minimal media due to a toxic increase in (p)ppGpp. *Cell Host Microbe* 17:788–798. <https://doi.org/10.1016/j.chom.2015.05.006>.
 11. Zeden MS, Schuster CF, Bowman L, Zhong Q, Williams HD, Grundling A. 2018. Cyclic-di-adenosine monophosphate (c-di-AMP) is required for osmotic regulation in *Staphylococcus aureus* but dispensable for viability in anaerobic conditions. *J Biol Chem* 293:3180–3200. <https://doi.org/10.1074/jbc.M117.818716>.
 12. Kohanski MA, Dwyer DJ, Collins JJ. 2010. How antibiotics kill bacteria: from targets to networks. *Nat Rev Microbiol* 8:423–435. <https://doi.org/10.1038/nrmicro2333>.
 13. Corrigan RM, Abbott JC, Burhenne H, Kaever V, Grundling A. 2011. c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress. *PLoS Pathog* 7:e1002217. <https://doi.org/10.1371/journal.ppat.1002217>.
 14. Luo Y, Helmann JD. 2012. Analysis of the role of *Bacillus subtilis* s^M in beta-lactam resistance reveals an essential role for c-di-AMP in peptidoglycan homeostasis. *Mol Microbiol* 83:623–639. <https://doi.org/10.1111/j.1365-2958.2011.07953.x>.
 15. Griffiths JM, O'Neill AJ. 2012. Loss of function of the *gdpP* protein leads to joint beta-lactam/glycopeptide tolerance in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 56:579–581. <https://doi.org/10.1128/AAC.05148-11>.
 16. Witte CE, Whiteley AT, Burke TP, Sauer JD, Portnoy DA, Woodward JJ. 2013. Cyclic di-AMP is critical for *Listeria monocytogenes* growth, cell wall homeostasis, and establishment of infection. *mBio* 4:e00282-13. <https://doi.org/10.1128/mBio.00282-13>.
 17. Rismondo J, Gibhardt J, Rosenberg J, Kaever V, Halbedel S, Commichau FM. 2016. Phenotypes associated with the essential diadenylate cyclase CdaA and its potential regulator CdaR in the human pathogen *Listeria monocytogenes*. *J Bacteriol* 198:416–426. <https://doi.org/10.1128/JB.00845-15>.
 18. Ba X, Kalmar L, Hadjirin NF, Kerschner H, Apfalter P, Morgan FJ, Paterson GK, Girvan SL, Zhou R, Harrison EM, Holmes MA. 2019. Truncation of GdpP mediates beta-lactam resistance in clinical isolates of *Staphylococcus aureus*. *J Antimicrob Chemother* 74:1182–1191. <https://doi.org/10.1093/jac/dkz013>.
 19. Banerjee R, Gretes M, Harlem C, Basuino L, Chambers HF. 2010. A *mecA*-negative strain of methicillin-resistant *Staphylococcus aureus* with high-level beta-lactam resistance contains mutations in three genes. *Antimicrob Agents Chemother* 54:4900–4902. <https://doi.org/10.1128/AAC.00594-10>.
 20. Argudin MA, Roisin S, Nienhaus L, Dodemont M, de Mendonca R, Nonhoff C, Deplano A, Denis O. 2018. Genetic diversity among *Staphylococcus aureus* isolates showing oxacillin and/or cefoxitin resistance not linked to the presence of *mec* genes. *Antimicrob Agents Chemother* 62:e00091-18. <https://doi.org/10.1128/AAC.00091-18>.
 21. Giulieri SG, Guerillot R, Kwong JC, Monk IR, Hayes AS, Daniel D, Baines S, Sherry NL, Holmes NE, Ward P, Gao W, Seemann T, Stinear TP, Howden BP. 2020. Comprehensive genomic investigation of adaptive mutations driving the low-level oxacillin resistance phenotype in *Staphylococcus aureus*. *mBio* 11:e02882-20. <https://doi.org/10.1128/mBio.02882-20>.
 22. Zhu Y, Pham TH, Nhiep THN, Vu NMT, Marcellin E, Chakraborti A, Wang Y, Waanders J, Lo R, Huston WM, Bansal N, Nielsen LK, Liang Z-X, Turner MS. 2016. Cyclic-di-AMP synthesis by the diadenylate cyclase CdaA is modulated by the peptidoglycan biosynthesis enzyme GlmM in *Lactococcus lactis*. *Mol Microbiol* 99:1015–1027. <https://doi.org/10.1111/mmi.13281>.
 23. Tosi T, Hoshiga F, Millership C, Singh R, Eldrid C, Patin D, Mengin-Lecreulx D, Thalassinou K, Freemont P, Grundling A. 2019. Inhibition of the *Staphylococcus aureus* c-di-AMP cyclase DacA by direct interaction with the phosphoglucosamine mutase GlmM. *PLoS Pathog* 15:e1007537. <https://doi.org/10.1371/journal.ppat.1007537>.
 24. Gibhardt J, Heidemann JJ, Bremskamp R, Rosenberg J, Seifert R, Kaever V, Ficner R, Commichau FM. 2020. An extracytoplasmic protein and a moonlighting enzyme modulate synthesis of c-di-AMP in *Listeria monocytogenes*. *Environ Microbiol* 22:2771–2791. <https://doi.org/10.1111/1462-2920.15008>.
 25. Pham HT, Nhiep NTH, Vu TNM, Huynh TN, Zhu Y, Huynh ALD, Chakraborti A, Marcellin E, Lo R, Howard CB, Bansal N, Woodward JJ, Liang ZX, Turner MS. 2018. Enhanced uptake of potassium or glycine betaine or export of cyclic-di-AMP restores osmoresistance in a high cyclic-di-AMP *Lactococcus lactis* mutant. *PLoS Genet* 14:e1007574. <https://doi.org/10.1371/journal.pgen.1007574>.
 26. Quintana IM, Gibhardt J, Turdieu A, Hammer E, Commichau FM, Lee VT, Magni C, Stülke J. 2019. The KupA and KupB proteins of *Lactococcus lactis* IL1403 are novel c-di-AMP receptor proteins responsible for potassium uptake. *J Bacteriol* 201:e00028-19. <https://doi.org/10.1128/JB.00028-19>.
 27. Bachmann H, Fischlechner M, Rabbers I, Barfa N, Branco dos Santos F, Molenaar D, Teusink B. 2013. Availability of public goods shapes the evolution of competing metabolic strategies. *Proc Natl Acad Sci U S A* 110:14302–14307. <https://doi.org/10.1073/pnas.1308523110>.
 28. Dodd HM, Horn N, Gasson MJ. 1994. Characterization of IS905, a new multicopy insertion sequence identified in lactococci. *J Bacteriol* 176:3393–3396. <https://doi.org/10.1128/jb.176.11.3393-3396.1994>.
 29. Tascon I, Sousa JS, Corey RA, Mills DJ, Griwatz D, Aumuller N, Mikusevic V, Stansfeld PJ, Vonck J, Hanelt I. 2020. Structural basis of proton-coupled potassium transport in the KUP family. *Nat Commun* 11:626. <https://doi.org/10.1038/s41467-020-14441-7>.
 30. Fulyani F, Schuurman-Wolters GK, Zagar AV, Guskov A, Slotboom DJ, Poolman B. 2013. Functional diversity of tandem substrate-binding domains in ABC transporters from pathogenic bacteria. *Structure* 21:1879–1888. <https://doi.org/10.1016/j.str.2013.07.020>.
 31. Poolman B, Smid EJ, Konings WN. 1987. Kinetic properties of a phosphate-bond-driven glutamate-glutamine transport system in *Streptococcus lactis* and *Streptococcus cremoris*. *J Bacteriol* 169:2755–2761. <https://doi.org/10.1128/jb.169.6.2755-2761.1987>.
 32. Schuurman-Wolters GK, Poolman B. 2005. Substrate specificity and ionic regulation of GlnPQ from *Lactococcus lactis*. An ATP-binding cassette transporter with four extracytoplasmic substrate-binding domains. *J Biol Chem* 280:23785–23790. <https://doi.org/10.1074/jbc.M500522200>.
 33. Fulyani F, Schuurman-Wolters GK, Slotboom DJ, Poolman B. 2016. Relative rates of amino acid import via the ABC transporter GlnPQ determine the growth performance of *Lactococcus lactis*. *J Bacteriol* 198:477–485. <https://doi.org/10.1128/JB.00685-15>.
 34. Curtis NA, Hughes JM, Ross GW. 1976. Inhibition of peptidoglycan cross-linking in growing cells of *Escherichia coli* by penicillins and cephalosporins, and its prevention by R factor-mediated beta-lactamase. *Antimicrob Agents Chemother* 9:208–213. <https://doi.org/10.1128/aac.9.2.208>.
 35. Stülke J, Krüger L. 2020. Cyclic di-AMP signaling in bacteria. *Annu Rev Microbiol* 74:159–179. <https://doi.org/10.1146/annurev-micro-020518-115943>.
 36. Massa SM, Sharma AD, Siletti C, Tu Z, Godfrey JJ, Gutheil WG, Huynh TN. 2020. c-di-AMP accumulation impairs mureptide synthesis in *Listeria monocytogenes*. *J Bacteriol* 202:e00307-20. <https://doi.org/10.1128/JB.00307-20>.
 37. Rojas ER, Huang KC. 2018. Regulation of microbial growth by turgor pressure. *Curr Opin Microbiol* 42:62–70. <https://doi.org/10.1016/j.mib.2017.10.015>.
 38. Krüger L, Herzberg C, Rath H, Pedreira T, Ischebeck T, Poehlein A, Gundlach J, Daniel R, Volker U, Mader U, Stulke J. 2021. Essentiality of c-di-AMP in *Bacillus subtilis*: bypassing mutations converge in potassium and glutamate homeostasis. *PLoS Genet* 17:e1009092. <https://doi.org/10.1371/journal.pgen.1009092>.
 39. Choi PH, Vu TMN, Pham HT, Woodward JJ, Turner MS, Tong L. 2017. Structural and functional studies of pyruvate carboxylase regulation by cyclic di-AMP in lactic acid bacteria. *Proc Natl Acad Sci U S A* 114:E7226–E7235. <https://doi.org/10.1073/pnas.1704756114>.
 40. Sikkema HR, van den Noort M, Rheinberger J, de Boer M, Krepel ST, Schuurman-Wolters GK, Paulino C, Poolman B. 2020. Gating by ionic strength and safety check by cyclic-di-AMP in the ABC transporter OpuA. *Sci Adv* 6:eabd7697. <https://doi.org/10.1126/sciadv.abd7697>.
 41. McLaggan D, Naprstek J, Buurman ET, Epstein W. 1994. Interdependence of K⁺ and glutamate accumulation during osmotic adaptation of *Escherichia coli*. *J Biol Chem* 269:1911–1917. [https://doi.org/10.1016/S0021-9258\(17\)42113-2](https://doi.org/10.1016/S0021-9258(17)42113-2).
 42. Gundlach J, Commichau FM, Stulke J. 2018. Of ions and messengers: an intricate link between potassium, glutamate, and cyclic di-AMP. *Curr Genet* 64:191–195. <https://doi.org/10.1007/s00294-017-0734-3>.
 43. Lahtvee PJ, Adamberg K, Arike L, Nahku R, Aller K, Vilu R. 2011. Multi-omics approach to study the growth efficiency and amino acid metabolism in *Lactococcus lactis* at various specific growth rates. *Microb Cell Fact* 10:12. <https://doi.org/10.1186/1475-2859-10-12>.

44. Pham HT, Turner MS. 2019. Onward and [K⁺]upward: a new potassium importer under the spell of cyclic di-AMP. *J Bacteriol* 201:e00150-19. <https://doi.org/10.1128/JB.00150-19>.
45. Krüger L, Herzberg C, Warneke R, Poehlein A, Stautz J, Weiß M, Daniel R, Hänelt I, Stülke J. 2020. Two ways to convert a low-affinity potassium channel to high affinity: control of *Bacillus subtilis* KtrCD by glutamate. *J Bacteriol* 202:e00138-20. <https://doi.org/10.1128/JB.00138-20>.
46. Turner MS, Woodberry T, Hafner LM, Giffard PM. 1999. The *bspA* locus of *Lactobacillus fermentum* BR11 encodes an L-cystine uptake system. *J Bacteriol* 181:2192–2198. <https://doi.org/10.1128/JB.181.7.2192-2198.1999>.
47. Prasad P, Turner MS. 2011. What bacteria are living in my food? An open-ended practical series involving identification of unknown foodborne bacteria using molecular techniques. *Biochem Mol Biol Educ* 39:384–390. <https://doi.org/10.1002/bmb.20532>.
48. Wells JM, Wilson PW, Le Page RW. 1993. Improved cloning vectors and transformation procedure for *Lactococcus lactis*. *J Appl Bacteriol* 74:629–636. <https://doi.org/10.1111/j.1365-2672.1993.tb05195.x>.
49. Poyart C, Trieu-Cuot P. 1997. A broad-host-range mobilizable shuttle vector for the construction of transcriptional fusions to beta-galactosidase in Gram-positive bacteria. *FEMS Microbiol Lett* 156:193–198. <https://doi.org/10.1111/j.1574-6968.1997.tb12726.x>.
50. Meyrand M, Boughammoura A, Courtin P, Mezange C, Guillot A, Chapot-Chartier M-P. 2007. Peptidoglycan N-acetylglucosamine deacetylation decreases autolysis in *Lactococcus lactis*. *Microbiology (Reading)* 153:3275–3285. <https://doi.org/10.1099/mic.0.2007/005835-0>.
51. Courtin P, Miranda G, Guillot A, Wessner F, Mezange C, Domakova E, Kulakauskas S, Chapot-Chartier M-P. 2006. Peptidoglycan structure analysis of *Lactococcus lactis* reveals the presence of an L,D-carboxypeptidase involved in peptidoglycan maturation. *J Bacteriol* 188:5293–5298. <https://doi.org/10.1128/JB.00285-06>.
52. Glauner B. 1988. Separation and quantification of muropeptides with high-performance liquid chromatography. *Anal Biochem* 172:451–464. [https://doi.org/10.1016/0003-2697\(88\)90468-x](https://doi.org/10.1016/0003-2697(88)90468-x).
53. Valgepea K, Loi KQ, Behrendorff JB, Lemgruber RSP, Plan M, Hodson MP, Kopke M, Nielsen LK, Marcellin E. 2017. Arginine deiminase pathway provides ATP and boosts growth of the gas-fermenting acetogen *Clostridium autoethanogenum*. *Metab Eng* 41:202–211. <https://doi.org/10.1016/j.ymben.2017.04.007>.