



HAL
open science

Fisheries management in spatially structured Atlantic salmon populations

Amaia Lamarins, Etienne Prévost, Stephanie Carlson, Mathieu Buoro

► **To cite this version:**

Amaia Lamarins, Etienne Prévost, Stephanie Carlson, Mathieu Buoro. Fisheries management in spatially structured Atlantic salmon populations. NoWPaS 2022, Mar 2022, Saint Etienne de Baigorry, France. hal-03790558

HAL Id: hal-03790558

<https://hal.inrae.fr/hal-03790558>

Submitted on 28 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fisheries management in spatially structured Atlantic salmon populations

Amaïa Lamarins, Etienne Prévost, Stephanie Carlson, Mathieu Buoro

UMR ECOBIOP INRAE-UPPA, ESPM Dpt, University of California Berkeley

Keywords: dispersal, fisheries, management, modeling

Abstract: Despite extensive research on the causes and consequences of spatial dispersal, the implications of connectivity for conservation and management are poorly appreciated, especially for species for which dispersal rates are ignored or considered negligible such as salmonids. For such species, exploitation by fishing usually does not consider populations connectivity, and fisheries management rather focuses at population scale. To date, prior work already warned about the danger of ignoring spatial structure and connectivity of populations but a very few studies have explicitly compared different strategies of exploitation while also considering the spatial structure of populations. We use a spatially explicit demo-genetic agent-based model as a virtual laboratory to mimic a realistic Atlantic salmon populations network and compare several spatialized fishing strategies (e.g. fishing all populations, sink, or source only). We assess their consequences at the demographic, phenotypic and genotypic levels. We show different effects of spatialized exploitation strategies on metapopulation size, stability and synchrony, as well as life history strategies and genetic evolution of traits. Importantly, we show that the consequences of spatialized fishing strategies depend on the exploitation pressure acting on the local populations and on the metapopulation as a whole. Altogether, we argue that it is critical to account for metapopulation structure in defining fisheries management because spatialized exploitation of connected populations can lead to various demographic outcomes but also complex evolutionary trajectories.