



**HAL**  
open science

# Evolutionary-driven management of brown trout (*Salmo trutta*. L) intraspecific diversity

Dorinda Folio, Jacques Labonne, Arnaud Caudron

► **To cite this version:**

Dorinda Folio, Jacques Labonne, Arnaud Caudron. Evolutionary-driven management of brown trout (*Salmo trutta*. L) intraspecific diversity. NoWPaS 2022, Mar 2022, Saint Etienne de Baigorry, France. hal-03790849

**HAL Id: hal-03790849**

**<https://hal.inrae.fr/hal-03790849>**

Submitted on 28 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Evolutionary-driven management of brown trout (*Salmo trutta*. L) intraspecific diversity

Dorinda Folio, Jacques Labonne, Arnaud Caudron

<sup>1</sup>Université de Pau et des Pays de l'Adour, UMR INRAE-UPPA, Ecobiop, FR-64310  
Saint-Pée-sur-Nivelle, France

<sup>2</sup>SCIMABIO INTERFACE, FR-74200, Thonon-les-Bains, France

**Keywords:** Management, Intraspecific diversity, Eco-evolution, Modelling

**Abstract:** Human interventions have often led to the admixture of previously allopatric lineages, notably for salmonid species. In France, intensive stocking of Atlantic brown trout has led to a mosaic of outcomes regarding hybridization and introgression of Mediterranean alleles by Atlantic ones. To understand and predict this diversity dynamic, we developed an individual-based demo-genetic model, which implements identified reproductive isolation mechanisms – female preference and gene-by-environment embryonic survival – between both lineages. The model aims at representing the evolution of intraspecific diversity dynamics in various environmental contexts and under different management practices, by including eco-evolutionary mechanisms as well as their genetic basis. Simulations, considering pre- and post-zygotic isolation, indicate that evolutionary mechanisms help at predicting introgression dynamics, depending on the environmental context. Managers interested in native or intraspecific diversity conservation should therefore consider the role of both, reproductive isolation mechanisms and environmental factors in their decision-making regarding conservation action.