



**HAL**  
open science

## European catfish *Silurus glanis* behaviour in response to a strong summer hypoxic event in a shallow lake

Samuel Westrelin, Stéphanie Boulêtreau, Frédéric Santoul

### ► To cite this version:

Samuel Westrelin, Stéphanie Boulêtreau, Frédéric Santoul. European catfish *Silurus glanis* behaviour in response to a strong summer hypoxic event in a shallow lake. *Aquatic Ecology*, 2022, 809 (1), pp.121-139. 10.1007/s10452-022-09952-y . hal-03791179

**HAL Id: hal-03791179**

**<https://hal.inrae.fr/hal-03791179v1>**

Submitted on 19 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **How to cite this article:**

2 Westrelin S, Boulêtreau S, Santoul F (2022) European catfish *Silurus glanis* behaviour in  
3 response to a strong summer hypoxic event in a shallow lake. *Aquatic Ecology*.  
4 doi:10.1007/s10452-022-09952-y

5  
6 You can fully access to the **journal online version** at :

7 <https://link.springer.com/article/10.1007/s10452-022-09952-y>

8  
9 You can **publicly read** the paper at the following link :

10 <https://rdcu.be/cluiF>

11  
12  
13 **European catfish *Silurus glanis* behaviour in response to a strong summer**  
14 **hypoxic event in a shallow lake**

15  
16 Westrelin Samuel<sup>(1)</sup>, Boulêtreau Stéphanie<sup>(2)</sup>, Santoul Frédéric<sup>(3)</sup>

17  
18 (1) INRAE, Aix Marseille Univ, Pôle R&D ECLA, RECOVER, 3275 Route de Cézanne - CS  
19 40061, F-13182 Aix-en-Provence Cedex 5, FRANCE.

20 <https://orcid.org/0000-0002-0169-1363>

21 (2) Laboratoire Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS,  
22 Toulouse, FRANCE.

23 <https://orcid.org/0000-0002-0094-0196>

24 (3) Laboratoire Evolution & Diversité Biologique, Université Paul Sabatier, CNRS, ENFA,

25 UMR5174 EDB, 118 route de Narbonne, 31062 Toulouse, FRANCE. <https://orcid.org/0000->

26 [0002-2932-2172](https://orcid.org/0000-0002-2932-2172)

27  
28 **Corresponding author**

29 Samuel Westrelin

30 INRAE

31 3275 Route de Cézanne - CS 40061

32 F-13182 Aix-en-Provence Cedex 5

33 France

34 samuel.westrelin@inrae.fr

35 +33 (0)4 42 66 69 71

36

## 37 Acknowledgements

38 We are very grateful to Virginie Diouloufet, Julien Dublon, Dorian Milesi and Ange Molina for their  
39 investment in the field work and data entry and to Yann Le Coarer and Nathalie Reynaud for dGPS  
40 treatments including fieldwork support. We would like to thank Mailove Benoliel, Léa Voisin and  
41 numerous other people who occasionally helped in the field as well as Coralie Garron for English  
42 correction of this manuscript. We also thank Lionel Allègre, Stéphanie Bertrand and Dominique  
43 Ghione for their warm welcome and logistical support on the study site. Finally, we appreciated the  
44 positive and relevant comments of the two reviewers, Johann Mourier and an anonymous reader, that  
45 significantly improved the finished document. This project was in part funded by the « Département  
46 des Bouches-du-Rhône ».

47

48

49

## 50 Abstract

51 Hypoxic events have always naturally occurred in freshwater ecosystems but are worsening due to  
52 anthropogenic activities. Hypoxia tolerance greatly varies among fish species and is difficult to  
53 quantify in nature in large fish species. We analysed the movements of 40 subadult and adult  
54 European catfish *Silurus glanis* ([727; 2150] mm) exposed to a natural summer hypoxic event in a  
55 shallow lake of southeastern France. Catfish could withstand very low dissolved oxygen  
56 concentrations (DOC), down to 1.3 mg/L in the upper half of the water column (corresponding to a  
57 mean dissolved oxygen saturation rate of 16%), when their preferred benthic habitats were anoxic.  
58 While hypoxia was becoming more severe, individuals significantly increased their activity and the  
59 surface area they visited, whatever their size. This led them to a refuge zone where they aggregated or  
60 stayed in close vicinity, very little mobile, over one and a half day during the overall anoxia of the  
61 lake. This zone, located very close to the well oxygenated water inflow, was probably one of the most  
62 oxygenated accessible zone. During this aggregation, the smallest individuals were however more  
63 active than the largest ones. This was probably because they more often needed to move to better  
64 oxygenated places within the gathering area, compared to larger dominant fish that occupied the best  
65 places. The ability of catfish to withstand very low DOC, along with its high optimum temperature  
66 range, could give it a competitive advantage over other predatory species in the context of global  
67 change.

## 68 Key Words

69 Aggregation, escape response, fish, hypoxia, lake

70

## 71 Introduction

72 Low dissolved oxygen conditions, so called hypoxic events, occur in a wide range of marine and  
73 freshwater ecosystems (Diaz & Rosenberg, 2011). They happen when oxygen consumption, primarily  
74 by decomposing organic matter, exceeds oxygen supply by photosynthetic production and diffusion  
75 from the atmosphere. Hypoxic events have always naturally occurred in aquatic ecosystems but the  
76 gradual rise in nutrient and organic enrichment due to human activities (sewage, industrial and land

77 runoff) has resulted in the increase in their frequency and seriousness, sometimes leading to anoxia  
78 (Druon et al., 2004; Hagy et al., 2004). Increasing hypoxia is now recognized as an environmental  
79 issue of global importance for fresh, coastal and oceanic waters (Breitburg et al., 2009; Diaz &  
80 Rosenberg, 2011; Jenny et al., 2016). In lentic and lotic freshwater systems, hypoxia varies in  
81 seasonality, frequency and persistence, depending on many factors, including eutrophication, inflow of  
82 industrial waste, reduced mixing due to depth or wind conditions, thermal variations and ice cover  
83 (Poff et al., 2002; Ficke et al., 2007).

84         Compared with most birds and mammals, ectothermic vertebrates (fish, amphibian, and  
85 reptile) are tolerant of variable oxygen availability (Bickler & Buck, 2007). In fish, hypoxia tolerance  
86 greatly varies among the 20,000 species. Species such as trout (*Oncorhynchus mykiss*) and tuna  
87 (*Katsuwonus pelamis*, *Thunnus albacares*, *Thunnus obesus*), that extensively depend on aerobic  
88 metabolism for rapid and sustained swimming, are moderately to extremely sensitive to hypoxia  
89 (Gesser, 1977; Bushnell et al., 1990; Gamperl & Driedzic, 2009). Carp (*Cyprinus carpio*), eel  
90 (*Anguilla anguilla*), catfish (*Silurus glanis*) and hagfish (*Myxine glutinosa*) can manage with low  
91 oxygen concentrations (Weber et al., 1976; Gesser, 1977; Axelsson et al., 1990; Massabuau & Forgeue,  
92 1995). At the extreme, the crucian carp (*Carassius carassius*) is able to endure months of hypoxia at  
93 low temperature (Nilsson & Renshaw, 2004; Stecyk et al., 2004). Such hypoxia tolerance involves  
94 metabolic adjustments, including metabolism depression, tolerance of metabolic products during  
95 anaerobiosis, and strategies for avoiding or repairing cellular injuries during reoxygenation (Brauner et  
96 al., 2004; Wells, 2009).

97         Behavioural responses can provide additional flexibility to mitigate exposure to hypoxic  
98 stress. Changes in spontaneous swimming activity have been described in a wide range of fish species  
99 when exposed to hypoxia (Schurmann & Steffensen, 1994; Chapman & McKenzie, 2009). Domenici  
100 et al. (2000) found that the lower was the spontaneous speed in Atlantic herring (*Clupeus harengus*) in  
101 normoxia, the greater was the increase in speed in hypoxia. Some fish up-regulate their speed,  
102 performing an escape response, defined as a type of fast start characterized by a brief and sudden  
103 acceleration (Domenici & Blake, 1997). Alternatively, others down-regulate their speed, they show  
104 freezing behaviour by adopting a fixed and immobile posture through which they become less

105 susceptible to detection by predators. It has also been suggested that species that reduce their activity  
106 in hypoxia tend to be demersal or benthopelagic, with a relatively sedentary lifestyle during which  
107 they may often encounter hypoxia in their habitat; whereas species that increase activity in case of  
108 hypoxia tend to be active pelagic schooling fishes (Domenici et al., 2000; Herbert & Steffensen, 2005;  
109 Herbert & Steffensen, 2006). Therefore, changes in swimming activity as behavioural responses to  
110 mitigate exposure to hypoxia are difficult to predict since they largely depend on fish species and  
111 context (Chapman & McKenzie, 2009). Moreover, such behaviours are difficult to describe and  
112 quantify in nature, and also in laboratory conditions for very large species.

113         The European catfish (*Silurus glanis*) is the largest fish species inhabiting European  
114 freshwaters (up to 2.7 m in body length and 130 kg of weight, Boulêtreau & Santoul, 2016). The  
115 species is native from Eastern Europe and has been introduced in Southwestern Europe during the 19<sup>th</sup>  
116 Century for sport fishing and aquaculture (Copp et al., 2009). It has successfully established in most of  
117 the large Southwest European watersheds (Boulêtreau et al., 2020). Several features could explain its  
118 colonization and expansion success. The range of temperatures within which adults do not show any  
119 sign of abnormal behaviour is quite large and falls between 12 °C and 28 °C, but reproduction  
120 optimally occurs above 20 °C (Souchon & Tissot, 2012). It is also tolerant to water pollution, partly  
121 due to low oxygen requirements as little as 1-1.5 mg/L depending on the temperature (Massabuau &  
122 Forgue, 1995). The species is considered to use oxygen very efficiently, partly thanks to a high  
123 haematocrit (35-38%, Mihalik, 1995). Nevertheless, such physiological capacities have only been  
124 measured on young individuals in laboratory conditions. In natural conditions, one study has reported  
125 the displacements of 19 juvenile catfish (total length less than 400 mm) in response to a winter  
126 hypoxia in one oxbow lake of the river Elbe (Czech Republic). Fish were shown to exhibit unexpected  
127 high activity and displacements (Daněk et al., 2014). But no behavioural response to hypoxia has ever  
128 been studied on adult European catfish in natural conditions.

129         In an experiment set up to assess the space use and activity cycles of the European catfish, 40  
130 large individuals (subadults and adults whose body length ranged in 727 - 2150 mm) have been  
131 tracked by acoustic telemetry for three years in a 104-ha shallow eutrophic lake located in South-  
132 Eastern France. Within this period, the lake experienced a severe hypoxia, leading to the mortality of

133 many fish from different species, including carp, eel and some small European catfish. However,  
134 fourteen months later, our telemetry tracking data revealed that all the 40 tagged catfish were still  
135 alive. Therefore, we aimed to analyse how large European catfish individuals could have resisted to  
136 hypoxia in natural conditions. More specifically, we examined how telemetry data could provide  
137 valuable information to highlight catfish individual tolerance and behaviour changes in response to  
138 hypoxia depending on catfish body size.

139

## 140 **Material and methods**

### 141 **Study site**

142 “Etang des Aulnes” is a shallow natural lake, mean depth 3.8 m, maximum depth 6 m, 104 ha area,  
143 located in South-Eastern France in a protected natural area. A primary canal and a secondary one  
144 collect irrigation waters that feed the lake. The lake then outflows in another canal (Figure 1). The lake  
145 water residence time is 300 days. The outflow is regulated to get high water levels in winter and low  
146 water levels in summer (maximal difference of 0.6 m).  
147 The fish assemblage, determined by fyke nets, fishing traps and electro fishing in October 2017, 2018  
148 and 2019 was composed of 16 species. The most dominant species were bream (*Abramis brama*,  
149 relative abundance 65%), European perch (*Perca fluviatilis*, 13%), pumpkinseed (*Lepomis gibbosus*,  
150 8%), tench (*Tinca tinca*, 4%), pikeperch (*Sander lucioperca*, 4%), European catfish (*Silurus glanis*,  
151 3%) and Northern pike (*Esox lucius*, 2%). In addition, two crayfishes were present: *Procambarus*  
152 *clarkii* and *Faxonius limosus*. Fishing is allowed but only during daytime from the eastern bank of the  
153 lake and no other activity is authorised.

154

### 155 **Physical and chemical lake conditions**

156 Hourly vertical profiles of water temperature (0.5, 3 and 5 m above the bottom) and dissolved oxygen  
157 concentration (DOC) (0.5, 1.5, 2.5 and 3.5 m above the bottom) were recorded at the deepest point in  
158 the lake (location 42 on Figure 1). HOBO data loggers U22 were used for temperature and U26 for  
159 DOC. The dissolved oxygen saturation rate (DOS) was calculated from DOC and temperature values.  
160 At 0.5 m above the bottom, there were no records from 09-02 03:00 to 09-04 12:00, because of a

161 sensor failure. 0.5-1.5 m, 2.5-3 m and 3.5-5 m heights of measurements above the bottom are named  
162 bottom, middle and surface, respectively, in the following text. In addition, vertical profiles of  
163 temperature and DOC were recorded on 09-04 with a YSI Exo2 multiparameter sonde at different  
164 locations in the lake to get a spatial picture of what happened all over the lake at different depths  
165 (Figure 1).

166 The mean hourly wind speed at a standard height of 10 meters above ground (10-m wind) was  
167 measured at the meteorological station of Salon-de-Provence, located 24 km east of the lake and  
168 representative of the weather conditions on the lake. The meteorological data were provided by  
169 Météo-France, the French meteorological institute, and available from the INRAE CLIMATIK  
170 platform (<https://intranet.inrae.fr/climatik/>, in French) managed by the AgroClim laboratory of  
171 Avignon, France.

172

### 173 **Fish tagging**

174 A total of 40 European catfish were caught by fyke nets, angling or electrofishing over two sampling  
175 campaigns: 10 in October 2017 and 30 in October 2018. Different techniques were used to sample the  
176 whole range of sizes among subadults and adults and individual behaviours (Harkonen et al., 2016).  
177 Once captured, catfish were stocked for a few hours in large aerated basins (2x1.25x0.5 m<sup>3</sup>) filled with  
178 regularly changed lake water to check their condition. Fish were then individually anaesthetised, which  
179 took 5-6 minutes, by immersion in a smooth and smaller tank 1.8x0.5x0.7 m<sup>3</sup> containing an aerated  
180 solution of benzocaine (80 mg/L). Once the fish had lost its balance (ventral side up), did not respond  
181 to stimuli anymore and had a very slow and steady operculum rate, it was weighed, measured and  
182 placed ventral side up in an identical tank containing an aerated solution of benzocaine (40 mg/L) to  
183 irrigate the gills during surgery. A 15 to 20 mm long incision was made with a scalpel in the middle of  
184 an imaginary line that would join the basis of the pectoral fin to the pelvic fin. An acoustic transmitter  
185 sterilised in surgical spirit and rinsed with physiological liquid was inserted into the peritoneal cavity.  
186 Vemco V13-1L acoustic transmitters (30.5 mm long, 9.2 g in the air, mean battery life 1825 days, 180  
187 s - range 120-240s - mean burst interval for the 12 used in 2017 and 320 s - range 260-380s - in 2018)  
188 were used. The transmitter weight in the air did not exceed 2% of the fish body weight (Winter, 1996;



189 Snobl et al., 2015). The incision was closed using 2 to 3 simple surgical sutures (3-0 Polydioxanon  
190 resorbable monofilament) placed 5 mm apart. An antiseptic and antibiotic dressing was applied on the  
191 incision wound to help healing and limit the risk of infection. Two surgeons took turn every four fish  
192 to operate, one fish surgery taking 5-6 min. Fish were then put in large, aerated recovery basins  
193 ( $2 \times 1.25 \times 0.5 \text{ m}^3$ ), where they were continually observed until they recovered normal opercular activity,  
194 swimming ability, balance and behavioural response to stimuli, usually after 10 min. Fish spent 3 to 6  
195 hours in this recovery basin before being released to their capture site. All individuals could be  
196 released in good shape.

197 At the time they were tagged, total length of the 40 tracked European catfish ranged in [727; 2 150]  
198 mm (mean 1 033 mm) and weight in [2 301; 64 380] g (mean 9 658 g) (Table 1).

199

## 200 [Fish tracking](#)

201 An array of 52 underwater omnidirectional Vemco acoustic receivers (20 VR2W 69kHz and 32  
202 VR2Tx 69kHz) with their associated synchronisation tag (additional V16-1L transmitter for VR2W  
203 and built in V16-like transmitter for VR2Tx, 500-700 s, used to correct for receiver internal clock  
204 drift) were anchored to the bottom throughout the lake from December 2017 (Figure 1). Seven  
205 reference tags (V13-1L, 840-960s) were added to detect anomalies in the tracking system. On average,  
206 neighbouring receivers were positioned 155 m from each other (range, 100-209 m), in 3.9 m water  
207 depth (range, 1.5 - 6 m), 0.5 m above the bottom. Receivers were removed roughly every 6 months to  
208 download fish detections. From these detections, fish 2D positions were calculated with the Vemco  
209 Positioning System (VPS) (Smith, 2013). The horizontal position error, a dimensionless parameter  
210 calculated by the VPS for each position, gives information on the quality of the position estimate, and  
211 was used to filter the data set (Espinoza et al., 2011). Here, we retained only positions with horizontal  
212 position error not exceeding 100; this limit represented a good compromise between the mean position  
213 error (7.4 m) and the percentage of positions kept (87%).

214

## 215 **Space use metrics**

216 Fish have been continuously tracked from December 2017 but, here, we focused on ten days  
217 throughout August-September of 2019, when a severe hypoxic event occurred. Individual raw  
218 positions were interpolated using the R package *trajr* (McLean & Skowron Volponi, 2018) for each  
219 quarter hour between the first and the last position to get synchronised individual tracks. Interpolated  
220 positions from gaps in raw positions longer than 1 hour were discarded. These tracks were all together  
221 plotted on the lake map to create a video of the catfish displacements, useful to get insights on catfish  
222 space use (Online Resource 1). Distances between consecutive positions were computed and set to  
223 zero if less than the telemetry system mean position error (7.4 m). To represent fish swimming  
224 activity, for each individual and each hour, the mean speed (m/h) was calculated. A mean daily speed  
225 (m/h) was calculated for each individual if at least seven hourly mean speeds were available in a day.  
226 The mean distance of each individual to all others was calculated using the R package *spatstat*  
227 (Baddeley et al., 2015). The distance to shore of each individual was calculated using the R package  
228 *rgeos* (Bivand & Rundel, 2019). These distances were calculated for every quarter hour and averaged  
229 over every hour and every day. The individual daily home ranges were estimated with an  
230 Epanechnikov kernel as the utilization distribution with probability levels of 95% and 50%; the home  
231 range 50% is often referred to as the core area (Powell, 2000). These both metrics were also estimated  
232 for all pooled individuals over different periods of the ten days to map the areas used by the fish.  
233 These spatial analyses were conducted using the R package *adehabitatHR* (Calenge, 2006).

234 The video (Online Resource 1) showed an attractive location where individuals gathered. To identify  
235 possible differences in individual behaviours during the aggregation dynamics, we analysed input-  
236 output movements of every fish when they could be identified on video recording. We could extract  
237 the time when the fish reached the aggregation and stayed inside for 34 among 40 fish and the time  
238 when it definitively left (i.e. it did never swim back to the aggregation location) for 30 among 40 fish.

239

## 240 **Statistical analyses**

241 We applied an algorithm to detect possible breakpoints corresponding to structural changes in the 2-  
242 month time series of DOC and of the average hourly speed of all 40 individuals (Zeileis et al., 2003).

243 This was done with the *strucchange* R package (Zeileis et al., 2002). Then, we compared the  
244 statistically detected breakpoints in each time series to identify potential concomitance between  
245 changes in fish swimming activity and DOC dynamics. Among these periods bounded by the  
246 breakpoints, one comprised the severe hypoxia. The same algorithm was re-run on the mean hourly  
247 speed within this period to detect possible different levels of catfish activity depending on the hypoxia  
248 severity. DOC was compared among these different periods of activity by Kruskal-Wallis tests and  
249 pairwise comparisons by Fisher's least significant difference (R package *agricolae*, de Mendiburu,  
250 2020).

251 To assess relationships between fish behaviour (characterised by speed, distance to shore, distance to  
252 others and home ranges) and DOC, generalized linear mixed-effects models (Zuur et al., 2009) were  
253 used by focusing on this hypoxic period. To get rid of temporal correlation that impeded the model  
254 robustness, daily means of the different variables were used. After preliminary trials, fish size class  
255 and day gave from far the best model adjustments compared to size in mm and DOC, very probably  
256 because of threshold effects of DOC on the behaviour. Fish size was defined from body length  
257 measured during fish tagging, i.e. up to 22 months before the anoxic period, and was classified into  
258 three classes: “small”, “large” and “medium” corresponding to total length < 850 mm, ≥1100 mm and  
259 in-between, respectively (Table 1). The two extreme classes were considered in order to maximise the  
260 chances that the size of individuals from these both classes still differed in August-September 2019.  
261 The fish identity was considered as a random effect to explicitly account for individual variability. To  
262 take into account the skewed distribution of mean individual daily speed or distance to shore towards  
263 zero, a Tweedie family function with a log-link was used (Gilman et al., 2012).

264 The model could be written as follows:

$$265 \quad \log(\overline{\text{METRICS}}_{\text{ind}}) = \alpha + \text{SIZE} * \text{DAY} + s(\text{ind}) + \varepsilon$$

266

267 where  $\overline{\text{METRICS}}_{\text{ind}}$  is the expected daily mean individual speed, distance to the shore or home range,  $\alpha$   
268 is the overall intercept, SIZE is the size class (Table 1),  $s(\text{ind})$  is a smoothing function modeling the  
269 individual effects (Wood, 2008), and " $\varepsilon$ " is the error term following a normal distribution with zero

270 mean. Means among days and among sizes by day were pairwise compared by using the *emmeans* R  
271 package (Lenth, 2016). The model fitting was assessed with regards to the homogeneity and normality  
272 of the residuals (Zuur et al., 2009) and to the percentage of explained variance (Hastie & Tibshirani,  
273 1990). For mean individual distance to others, as we could not find a reliable model, we compared the  
274 distributions between days and between size classes among days by using Kruskal-Wallis tests and, if  
275 significantly different, we made pairwise comparisons by Fisher's least significant difference (R  
276 package *agricolae*, de Mendiburu, 2020).  
277 All statistical analyses were made with R version 3.6.3 (R Core Team, 2020).

278

## 279 Results

### 280 DOC dynamics

281 The surface DOC showed large variations over August-September 2019 (Figure 2). A striking event  
282 occurred at the end of August when the surface DOC collapsed and, during several days, reached very  
283 low values (mean surface DOC was 1.2 mg/L over 08-29 – 09-04), even becoming null for 52 hours.  
284 This corresponded to a full anoxia which first appeared at the bottom on 08-26 and propagated at the  
285 surface to make the whole water column anoxic on 08-30 03:00 (Online Resource 2). At this time,  
286 water temperatures were stratified and ranged from 22.8 °C at the bottom to 26.7 °C at the surface  
287 (Online Resource 2). The anoxia ended on 09-02 16:00 when surface DOC raised up to 1.6 mg/L and  
288 the oxygenation was homogeneous over the whole water column (middle and bottom DOC  
289 respectively at 1.6, mg/L and 1.5 mg/L). One hour before, DOC was 0, 0.2 and 0.2 mg/L, at the  
290 surface, in the middle and at the bottom respectively. In parallel, water temperatures started to  
291 homogenize from 09-01 14:00 to be mixed on 09-02 04:00 (24.7 °C at the bottom and 24.8 °C at the  
292 surface). This was very linked to the wind that strengthened from 09-01 09:00 (greater than 3 m/s,  
293 Online Resource 2).

294

### 295 DOC dynamics and fish activity

296 Several structural change points were found in the hourly mean speed and DOC time series (Figure 2).  
297 The corresponding dates for 2-month speed time series matched well those of surface DOC. Worthy of

298 note, they did not match with dates of changes of deeper DOC (Online Resource 3). In the next, we  
299 focus on surface DOC.

300 The 08-27 23:00 to 09-06 06:00 period, that comprises the full anoxia, shows very large variations of  
301 hourly speed that appear different from the diel cycle that could be observed outside of this period  
302 (Figure 2b). Within this period, four subperiods were detected: at the beginning, fish activity appeared  
303 quite similar to the previous period; then, speed sharply increased for 32 hours before fish suddenly  
304 stopped and performed very few movements during one and a half day, after which they started to  
305 progressively move again. The surface DOC was different between all four speed subperiods  
306 (Kruskal-Wallis test,  $\chi^2=94.010$ , 3 d.f.,  $p<0.001$ ): the surface DOC was 1.1 mg/L in average (DOS  
307 13%) during the highest activity subperiod and 0.1 mg/L (DOS 1%) during the lowest. The daily  
308 analysis also showed that the speed increase (on 08-30) and decrease (on 08-31 and 09-01) were  
309 significant (Table 2a, Figure 3b). This enhanced activity corresponded to higher individual home  
310 ranges (Tables 2c, 2d, Figures 3e, 3f) and to larger areas visited by the pool of individuals (Figure 4b).  
311 Conversely, the home range 95% as well as the core area were considerably reduced when the activity  
312 was the lowest (Figures 3e, 3f) and most of individuals gathered in a tiny area (Figure 4c) where they  
313 aggregated (Figure 3d). On 09-02, when fish recovered their activity, the areas they travelled over  
314 were still reduced (Figure 3f) and nearby the area where they had aggregated (Figure 4d).  
315 The highest activity subperiod started when surface DOC dropped down to 1.3 mg/L (DOS 16%) and  
316 ceased when the whole water column became anoxic. After the reduced activity subperiod, fish started  
317 to progressively move again when surface DOC raised up to 0.7 mg/L (DOS 8%) whereas the half  
318 lower of the water column was still anoxic (Online Resource 2).

319

### 320 [Aggregation location and dynamics](#)

321 From 08-31 to 09-01, catfish were almost inactive, closer to each other, closer to the bank and  
322 gathered in a same tiny area (Figures 3b, 3c, 3d, 4c). They aggregated where the main tributary flows  
323 into the pond. This was among the places that exhibited the highest DOC values recorded on 09-04 in  
324 the whole lake, a few days after the full anoxia (Figure 5). At the main inflow location (points 91 and

325 93), 1-m deep, DOC was near 8 mg/L (DOS 85%) and temperatures much cooler, 18°C against 23 -  
326 24°C everywhere else (not shown).

327 The time span between the first fish to join the aggregation location and the last one was 26 hours 15  
328 min (from 08-30 16:45 to 08-31 19:00), but 27 individuals joined the location in a short time, 4 hours  
329 45 min (from 08-30 20:30 to 08-31 01:15). The dates at which individuals definitively left the  
330 aggregation location spanned over a much longer period: 3 days 6 hours 45 min (from 09-01 14:15 to  
331 09-04 21:00). 26 individuals left it in 1 day 17 hours 45 min (from 09-02 02:15 to 09-03 20:00).

332

### 333 Fish behaviour and size dependence

334 The swimming activity, proxied by the mean daily speed, was significantly different among the days  
335 during the hypoxic event, and, some days, size class behave differently (Table 2a). The distance to  
336 shore was dependent on fish size and also different among days (Table 2b). In details on Figure 3b,  
337 whatever their size, all fish significantly increased their activity on 08-30 when the bottom anoxia was  
338 propagating to the surface. This corresponded to higher individual core areas for all fish whatever their  
339 size and, to a lesser extent, home ranges 95% (Figures 3e, 3f). On both following days, when the  
340 whole water column was anoxic and fish swimming activity the lowest, the larger individuals were  
341 even less active than the smaller ones (Table 2a, Figure 3b) and occupied a more reduced area on 09-  
342 01 (Tables 2c, 2d, Figures 3e, 3f); the smaller individuals were further to others than the larger ones  
343 (Figure 3d). In general, the smaller fish were further to the shore (Table 2b, Figure 3c, p-values of  
344 pairwise comparisons between Small and Large and Small and Medium <0.001) and had larger home  
345 ranges 95% (Table 2c, Figure 3e, p-values of pairwise comparisons between Small and Large and  
346 Small and Medium <0.05).

347

## 348 Discussion

349 Our dataset gave an excellent context to analyse the *in situ* response of a species to an environmental  
350 stress. The high resolution and high frequency of the positions collected by telemetry and the number  
351 of tagged individuals allowed to detect changes in behaviour at individual level and at timesteps  
352 suitable to be confronted with DOC variations (Bauer & Schlott, 2006; Daněk et al., 2014).

353

#### 354 **Catfish tolerance to low DOC**

355 All the 40 tagged catfish survived the severe hypoxic event that lasted more than two days and led to  
356 the death of lots of other fish, including species known to manage very well with low oxygen  
357 conditions such as carp and eel (Weber et al., 1976; Gesser, 1977). Subadult and adult European  
358 catfish could go through very low oxygen conditions during the summer when water temperatures  
359 ranged in 23-27 °C. Their behaviour was impacted only when, at the deepest point in the lake, the  
360 lower half of the water column was anoxic and when DOC dropped down to 1.3 mg/L (corresponding  
361 to a mean DOS of 16%) in the upper half. This DOC value was in the lower range of what Daněk et al.  
362 (2014) found on juvenile catfish, 1.3-2.4 mg/L in winter conditions (water temperature around 5 °C).  
363 It was in the range of Massabuau & Forgue (1995) laboratory results which concluded that very young  
364 catfish (weighing 100-150 g) could maintain dioxygen homeostasis in 1-1.5 mg/L DOC range at 13°C  
365 and even very probably in a 10-23 °C temperature range. However, increased temperature lessens  
366 oxygen solubility and thus reduces oxygen supply for ichtyofauna; it also elevates basal oxygen  
367 demand (Rogers et al., 2016). With comparable critical DOC but higher temperature, we can thus  
368 reasonably hypothesise that oxygen supply was more critical in our study. The possible lower critical  
369 oxygen threshold in our study could be explained by large differences in catfish body weights between  
370 studies. Large fish could have an advantage thanks to their lower mass-specific metabolic rate  
371 (Nilsson & Östlund-Nilsson, 2008). These tolerance values were in all cases much lower than the limit  
372 of 3-3.5 mg/L reported by Mihalik (1995).

373

#### 374 **Catfish behaviour in response to the hypoxia extent**

375 Horizontal catfish movements were not altered when anoxia was limited to the half lower part of the  
376 lake. However, these extreme conditions are likely to considerably reduce the suitable habitats  
377 regarding oxygen conditions in this shallow lake, by the way compressing fish habitat (Kraus et al.,  
378 2015). The tags did not record the pressure and thus could not provide information on fish depth.  
379 Nevertheless, we could suppose that catfish rose to the surface layer to find tolerable DOC, while they  
380 are known to mainly occupy benthic habitats (Bruton, 1996; Cucherousset et al., 2018). This remains

381 questioning. One day before complete anoxia at the deepest point of the lake, catfish exhibited a  
382 sudden higher level of activity and unusual large displacements over greater areas for about one day,  
383 all fish sizes alike. Many fish species change their spontaneous swimming activity when exposed to  
384 hypoxia, reducing or increasing their activity (Chapman & McKenzie, 2009). Usually, sedentary  
385 species decrease their swimming speed to save energy (Domenici et al., 2013). Crucian carp  
386 (*Carassius carassius*) that can endure anoxia for several hours and even days (at 9 °C) reduces its  
387 activity by 50% (Nilsson et al., 1993). The increased activity observed in the tagged catfish was  
388 unexpected for such a large species that tries to reduce its energy costs (Slavík et al., 2014) and usually  
389 performs few movements (Carol et al., 2007; Capra et al., 2018). The stress caused by the resource  
390 unavailability can significantly increase catfish movement activity (Slavík et al., 2016). Intense  
391 agitation of fish in deep hypoxia could also be interpreted as an avoidance response that helps to find a  
392 more suitable place (Domenici et al., 2000; Herbert & Steffensen, 2006; Chapman & McKenzie,  
393 2009) as catfish not only became faster but also explored extended areas. Such an increase in activity  
394 was also observed on juvenile catfish facing dissolved oxygen deficiency before they found a refuge  
395 (Daněk et al., 2014) or on school of Atlantic herrings (*Clupeus harengus*) whose speed peaked during  
396 severe hypoxia before decreasing until the school disrupted (Domenici et al., 2000). The reasons for  
397 such an agitation need further investigations.

398

### 399 [Aggregation and refuge location](#)

400 Most of catfish finally converged and gathered where the main canal flows into the lake. They stayed  
401 there or in close vicinity by considerably reducing their swimming activity for one and a half day.  
402 With the inflow canal, this gathering place was likely the most oxygenated in the lake, able to fulfill  
403 their oxygen requirements. In this summer period, irrigation led to quite a strong current in the main  
404 canal that continuously brought cool and well oxygenated water flowing through the gathering place.  
405 This place and places very close to the bank outside the receiver network were in general at the edge  
406 of the telemetry coverage area so that individuals were less often located during the anoxia (Smith,  
407 2013), which could sometimes be visible on the video (Online Resource 1). Overall unsuitable  
408 environmental conditions temporarily forced catfish to share very limited space and resources. As a



409 consequence, competition between individuals likely increased. European catfish have been reported  
410 to actively defend their access to resources (Cucherousset et al., 2018) and have been shown to expend  
411 more energy when in contact with conspecifics in preferred areas of habitat (Slavík & Horký, 2009).  
412 This could give advantage to the biggest individuals. High body mass was also shown to decrease  
413 stress from limited availability of resources (Slavík et al., 2016) which could explain why the smallest  
414 individuals were more active and further from the shore than the biggest during the anoxic event.  
415 Catfish aggregation, that can be compared to a school, led to a high oxygen consumption. According  
416 to the position in this school, in front of the inflow current or in the rear of the school, DOC can vary a  
417 lot so that some individuals need to change position leading to a reshuffle (Domenici et al., 2002;  
418 Herbert & Steffensen, 2006). The largest catfish, dominant, could occupy the most suitable positions,  
419 while the smallest individuals would be left with the least favourable ones and would then move much  
420 more often to change position toward better-oxygenated areas in the aggregation.  
421 Remarkably, the synchronisation they showed to join the refuge place contrasted with the time needed  
422 for all individuals to definitively leave the location. This would also need further investigations.

423

424 Hypoxic conditions are likely to become more frequent and severe with temperature rising and  
425 increasing eutrophication of ecosystems due to human activities. Aside temperature, DOC is a key  
426 environmental parameter driving space use by fish population. The ability of catfish to withstand very  
427 low DOC, along with its high optimum temperature range, could lead to an extent of the suitable  
428 geographical range for this species in the future. This is important to account for when engaged in  
429 conservation or fisheries management.

430

### 431 **Figure captions**

432 **Fig1** Bathymetric map of « Etang des Aulnes » at the water level of 11.14 m above sea level and  
433 experimental setup. Acoustic receivers and their associated synchronising tags are represented by  
434 white dots. Reference tags are symbolised by white squares. Monthly (and also hourly for location 42)  
435 temperature and dissolved oxygen concentration profiles are located by crosses; location 42 is the

436 deepest in the lake. One primary canal flows into the lake on the eastern bank at location 91 and a  
437 secondary one on the northern bank close to location 0. The lake outflows in a canal at its extreme  
438 south-west

439

440 **Fig2** Catfish mean speed and surface DOC during the summer period (from 1<sup>st</sup> August to 30  
441 September 2019). Panel a: hourly surface DOC (in mg/L at 3.5 m above the bottom at the deepest  
442 point of the lake, 5.5 m depth). Panel b: mean hourly speed over all individuals (in m/h). The dates of  
443 structural changes over the 2-month time series and their 95% confidence interval are labelled on the  
444 x-axis and represented by vertical dotted lines and interval at their basis. Hereafter, dates are given in  
445 the format mm-dd hh. These dates and associated 95% confidence interval are 08-27 23 [08-16 10 ;  
446 08-28 09], 09-06 06 [09-06 02 ; 09-08 06] and 09-18 17 [09-17 08 ; 09-21 04] for mean speed, and 08-  
447 12 00 [08-11 05 ; 08-12 19], 08-28 03 [08-28 01 ; 08-28 04], 09-06 06 [09-06 00 ; 09-06 10], 09-18 16  
448 [09-18 07 ; 09-19 04] for surface DOC. In addition, the dates of structural changes of mean speed time  
449 series within the period 08-27 23 to 09-06 06, comprising the anoxia, are labelled above the panel and  
450 represented by dashed vertical lines and associated 95% confidence interval at their basis

451

452 **Fig3** Catfish daily space use during the hypoxic period (from 28 August to 6 September 2019). Panel  
453 a: mean daily DOC (at 1.5, 2.5 and 3.5 m above the bottom in dotted, dashed and solid line  
454 respectively) at the deepest point in the lake, 5.5 m deep. In panels b, c and d, the boxplots represent  
455 the minimum, the first quartile, the median, the third quartile and the maximum of the distribution. In  
456 panels b, c and d, the dotted, dashed and solid lines represent the daily means over small, medium and  
457 large catfish respectively. Panel b: distribution of mean daily individual speeds (m/h). Panel c:  
458 distribution of mean individual daily distances to all others. Panel d: distribution of mean individual  
459 daily distances to shore (m). Panel e: distribution of individual home ranges 95% (hectares). Panel f:  
460 distribution of individual home ranges 50% (hectares). Letters above the boxplots stand for post-hoc  
461 comparisons between days: days which share a same letter have distributions which do not  
462 significantly differ (at the 5% significance level). Letters below the boxplots stand for comparisons  
463 between sizes within a day; no letter means the three sizes do not differ; sizes that share the same letter

464 do not significantly differ (within a day, letters from left to right correspond to small, medium and  
465 large catfish, respectively). As the interaction between size and day was not significant for distance to  
466 shore, the comparison between sizes within a day has not been performed

467

468 **Fig4** Catfish home ranges during the hypoxic period (from 28 August to 6 September 2019). The  
469 home range 95% and the core area (home range 50%) are mapped in pale grey with a dotted contour  
470 and in grey with a solid contour, respectively. They have been calculated with all pooled individuals  
471 over different grouping days brought out from figures 3e and 3f. Panel a stands for days 08-28 and 08-  
472 29, b for day 08-30 when the activity was the highest, c for days 08-31 and 09-01 when the activity  
473 was the lowest, d for day 09-02 when fish recovered their activity and e for days 09-03 to 09-06. The  
474 corresponding areas (in hectares) are given in each panel

475

476 **Fig5** Vertical profile of DOC at different locations in the pond on 09-04. The labels of the legend  
477 correspond to the different locations labelled on the lake map and symbolised by a cross. Data go from  
478 the surface to approximately 0.5 m above the bottom. Points 91 and 93 are located near the main  
479 inflow and point 11 is the closest to the outflow

480

## 481 [Supplementary Information](#)

482 **Online Resource 1** Video of catfish movements over 25 August - 6 September 2019

483 **Online Resource 2** Environmental conditions during the summer period (from 1<sup>st</sup> August to 30  
484 September 2019)

485 **Online Resource 3** Catfish mean speed and DOC during the summer period (from 1<sup>st</sup> August to 30  
486 September 2019)

487

## 488 [Author contributions](#)

489 S.W. designed the study and analysed the data. S.W. and S.B. wrote the paper. F.S. designed the paper  
490 with S.W. and S.B., carefully read it and made substantial improvements.

491

## 492 **Significance statement**

493 Hypoxic events have always naturally occurred in freshwater ecosystems but are worsening due to  
494 anthropogenic activities. Fish response to hypoxia are difficult to observe and quantify in nature for  
495 very large species. We observed that subadult and adult European catfish (40 individuals, 0.7 – 2.3m  
496 length) could resist to such a stress without any alteration in their behaviour down to 1.3 mg/L of  
497 dissolved oxygen concentration in summer conditions. Below this threshold, they enhanced their  
498 speed and the surface area they visited which led them to aggregate in a more oxygenated refuge zone.  
499 They all survived this severe event which strenghtens their capacity to cope with global change.

500

## 501 **Declarations**

### 502 **Conflicts of interests**

503 The authors declare they have no conflicts of interest.

### 504 **Funding**

505 This project was in part funded by the « Département des Bouches-du-Rhône ».

### 506 **Ethics**

507 The care and use of experimental animals complied with French animal welfare laws, guidelines and  
508 policies as approved by the French Ministry of Research through the authorisation number  
509 APAFIS#11294-2017091809143058 v2.

510

## 511 **Data availability**

512 The datasets analysed during the current study are available from the corresponding author on  
513 reasonable request.

## 514 **References**

515 Axelsson, M., A. P. Farrell & S. Nilsson, 1990. Effects of Hypoxia and Drugs on the  
516 Cardiovascular Dynamics of the Atlantic Hagfish *Myxine Glutinosa*. *Journal of*  
517 *Experimental Biology* 151:297-316.  
518 Baddeley, A., E. Rubak & R. Turner, 2015. *Spatial point patterns: methodology and*  
519 *applications with R*. Chapman and Hall/CRC Press, London.

520 Bauer, C. & G. Schlott, 2006. Reaction of common carp (*Cyprinus carpio*, L.) to oxygen  
521 deficiency in winter as an example for the suitability of radio telemetry for  
522 monitoring the reaction of fish to stress factors in pond aquaculture. *Aquaculture*  
523 *Research* 37:248-254 doi:<https://doi.org/10.1111/j.1365-2109.2005.01426.x>.

524 Bickler, P. E. & L. T. Buck, 2007. Hypoxia tolerance in reptiles, amphibians, and fishes: life  
525 with variable oxygen availability. *Annu Rev Physiol* 69:145-70  
526 doi:<https://doi.org/10.1146/annurev.physiol.69.031905.162529>.

527 Bivand, R. & C. Rundel, 2019. rgeos: Interface to Geometry Engine - Open Source ('GEOS'). R  
528 package version:0.5-2.

529 Boulêtreau, S., T. Fauvel, M. Laventure, R. Delacour, W. Bouyssonnié, F. Azémar & F. Santoul,  
530 2020. "The giants' feast": predation of the large introduced European catfish on  
531 spawning migrating allis shads. *Aquatic Ecology* doi:[https://doi.org/10.1007/s10452-](https://doi.org/10.1007/s10452-020-09811-8)  
532 [020-09811-8](https://doi.org/10.1007/s10452-020-09811-8).

533 Boulêtreau, S. & F. Santoul, 2016. The end of the mythical giant catfish. *Ecosphere* 7  
534 doi:<https://doi.org/10.1002/ecs2.1606>.

535 Brauner, C. J., V. Matey, J. M. Wilson, N. J. Bernier & A. L. Val, 2004. Transition in organ  
536 function during the evolution of air-breathing; insights from *Arapaima gigas*, an  
537 obligate air-breathing teleost from the Amazon. *Journal of Experimental Biology*  
538 207:1433-1438 doi:<https://doi.org/10.1242/jeb.00887>.

539 Breitburg, D. L., D. W. Hondorp, L. A. Davias & R. J. Diaz, 2009. Hypoxia, Nitrogen, and  
540 Fisheries: Integrating Effects Across Local and Global Landscapes. *Annual Review of*  
541 *Marine Science* 1:329-349  
542 doi:<https://doi.org/10.1146/annurev.marine.010908.163754>.

543 Bruton, M. N., 1996. Alternative life-history strategies of catfishes. *Aquatic Living Resources*  
544 9:35-41.

545 Bushnell, P. G., R. W. Brill & R. E. Bourke, 1990. Cardiorespiratory responses of skipjack tuna  
546 (*Katsuwonus pelamis*), yellowfin tuna (*Thunnus albacares*), and bigeye tuna (*Thunnus*  
547 *obesus*) to acute reductions of ambient oxygen. *Canadian Journal of Zoology*  
548 68:1857-1865.

549 Calenge, C., 2006. The package "adehabitat" for the R software: A tool for the analysis of  
550 space and habitat use by animals. *Ecological Modelling* 197:516-519  
551 doi:10.1016/j.ecolmodel.2006.03.017.

552 Capra, H., H. Pella & M. Ovidio, 2018. Individual movements, home ranges and habitat use  
553 by native rheophilic cyprinids and non-native catfish in a large regulated river.  
554 *Fisheries Management and Ecology* 25:136-149  
555 doi:<https://doi.org/10.1111/fme.12272>.

556 Carol, J., L. Zamora & E. García-Berthou, 2007. Preliminary telemetry data on the movement  
557 patterns and habitat use of European catfish (*Silurus glanis*) in a reservoir of the River  
558 Ebro, Spain. *Ecology of Freshwater Fish* 16:450-456  
559 doi:<https://doi.org/10.1111/j.1600-0633.2007.00225.x>.

560 Chapman, L. J. & D. J. McKenzie, 2009. Chapter 2 Behavioral Responses and Ecological  
561 Consequences. In Richards, J. G., A. P. Farrell & C. J. Brauner (eds) *Fish Physiology*. vol  
562 27. Academic Press, 25-77.

563 Copp, G. H., J. Robert Britton, J. Cucherousset, E. García-Berthou, R. Kirk, E. Peeler & S.  
564 Stakenas, 2009. Voracious invader or benign feline? A review of the environmental  
565 biology of European catfish *silurus glanis* in its native and introduced ranges. *Fish and*  
566 *Fisheries* 10:252-282 doi:<https://doi.org/10.1111/j.1467-2979.2008.00321.x>.

567 Cucherousset, J., P. Horky, O. Slavik, M. Ovidio, R. Arlinghaus, S. Bouletreau, R. Britton, E.  
568 Garcia-Berthou & F. Santoul, 2018. Ecology, behaviour and management of the  
569 European catfish. *Reviews in Fish Biology and Fisheries* 28:177-190  
570 doi:<https://doi.org/10.1007/s11160-017-9507-9>.

571 Daněk, T., L. Kalous, M. Petrtýl & P. Horký, 2014. Move or die: change in European catfish  
572 (*Silurus glanis* L.) behaviour caused by oxygen deficiency. *Knowl Managt Aquatic*  
573 *Ecosyst*:08.

574 de Mendiburu, F., 2020. agricolae: Statistical Procedures for Agricultural Research. R package  
575 version:1.3-2.

576 Diaz, R. J. & R. Rosenberg, 2011. Introduction to environmental and economic consequences  
577 of hypoxia. *International Journal of Water Resources Development* 27:71-82.

578 Domenici, P. & R. Blake, 1997. The kinematics and performance of fish fast-start swimming.  
579 *The Journal of Experimental Biology* 200:1165-1178.

580 Domenici, P., N. A. Herbert, C. Lefrançois, J. F. Steffensen & D. J. McKenzie, 2013. The effect  
581 of hypoxia on fish swimming performance and behaviour *Swimming physiology of*  
582 *fish*. Springer, 129-159.

583 Domenici, P., R. Silvana Ferrari, J. F. Steffensen & R. S. Batty, 2002. The effect of progressive  
584 hypoxia on school structure and dynamics in Atlantic herring *Clupea harengus*.  
585 *Proceedings of the Royal Society of London Series B: Biological Sciences* 269:2103-  
586 2111.

587 Domenici, P., J. F. Steffensen & R. S. Batty, 2000. The effect of progressive hypoxia on  
588 swimming activity and schooling in Atlantic herring. *Journal of Fish Biology* 57:1526-  
589 1538 doi:<https://doi.org/10.1111/j.1095-8649.2000.tb02229.x>.

590 Druon, J.-N., W. Schrimpf, S. Dobricic & A. Stips, 2004. Comparative assessment of large-  
591 scale marine eutrophication: North Sea area and Adriatic Sea as case studies. *Marine*  
592 *Ecology Progress Series* 272:1-23.

593 Espinoza, M., T. J. Farrugia, D. M. Webber, F. Smith & C. G. Lowe, 2011. Testing a new  
594 acoustic telemetry technique to quantify long-term, fine-scale movements of aquatic  
595 animals. *Fisheries Research* 108:364-371  
596 doi:<https://doi.org/10.1016/j.fishres.2011.01.011>.

597 Ficke, A. D., C. A. Myrick & L. J. Hansen, 2007. Potential impacts of global climate change on  
598 freshwater fisheries. *Reviews in Fish Biology and Fisheries* 17:581-613.

599 Gamperl, A. K. & W. R. Driedzic, 2009. Chapter 7 Cardiovascular Function and Cardiac  
600 Metabolism. In Richards, J. G., A. P. Farrell & C. J. Brauner (eds) *Fish Physiology*. vol  
601 27. Academic Press, 301-360.

602 Gesser, H., 1977. The effects of hypoxia and reoxygenation on force development in  
603 myocardia of carp and rainbow trout: protective effects of CO<sub>2</sub>/HCO<sub>3</sub>. *The Journal of*  
604 *Experimental Biology* 69:199-206.

605 Hagy, J. D., W. R. Boynton, C. W. Keefe & K. V. Wood, 2004. Hypoxia in Chesapeake Bay,  
606 1950–2001: Long-term change in relation to nutrient loading and river flow. *Estuaries*  
607 27:634-658 doi:<https://doi.org/10.1007/BF02907650>.

608 Harkonen, L., P. Hyvarinen, P. T. Niemela & A. Vainikka, 2016. Behavioural variation in  
609 Eurasian perch populations with respect to relative catchability. *Acta Ethologica*  
610 19:21-31 doi:<https://doi.org/10.1007/s10211-015-0219-7>.

611 Herbert, N. A. & J. F. Steffensen, 2005. The response of Atlantic cod, *Gadus morhua*, to  
612 progressive hypoxia: fish swimming speed and physiological stress. *Marine Biology*  
613 147:1403-1412 doi:<https://doi.org/10.1007/s00227-005-0003-8>.

614 Herbert, N. A. & J. F. Steffensen, 2006. Hypoxia increases the behavioural activity of  
615 schooling herring: a response to physiological stress or respiratory distress? Marine  
616 Biology 149:1217-1225 doi:<https://doi.org/10.1007/s00227-006-0284-6>.

617 Jenny, J.-P., P. Francus, A. Normandeau, F. Lapointe, M.-E. Perga, A. Ojala, A. Schimmelmann  
618 & B. Zolitschka, 2016. Global spread of hypoxia in freshwater ecosystems during the  
619 last three centuries is caused by rising local human pressure. Global Change Biology  
620 22:1481-1489 doi:<https://doi.org/10.1111/gcb.13193>.

621 Kraus, R. T., C. T. Knight, T. M. Farmer, A. M. Gorman, P. D. Collingsworth, G. J. Warren, P. M.  
622 Kocovsky & J. D. Conroy, 2015. Dynamic hypoxic zones in Lake Erie compress fish  
623 habitat, altering vulnerability to fishing gears. Canadian Journal of Fisheries and  
624 Aquatic Sciences 72:797-806 doi:<https://doi.org/10.1139/cjfas-2014-0517>.

625 Lenth, R. V., 2016. Least-Squares Means: The R Package lsmeans. Journal of Statistical  
626 Software 69:1-33 doi:10.18637/jss.v069.i01.

627 Massabuau, J.-C. & J. Forgeue, 1995. Les capacités d'adaptation du silure glane en hypoxie :  
628 un cas exemplaire d'homéostasie du milieu intérieur. Aquatic Living Resources 8:423-  
629 430 doi:<https://doi.org/10.1051/alr:1995050>.

630 McLean, D. J. & M. A. Skowron Volponi, 2018. trajr: An R package for characterisation of  
631 animal trajectories. Ethology 124:440-448.

632 Mihalik, J., 1995. Der Wels, 2nd edn. Westarp Wissenschaften, Magdeburg.

633 Nilsson, G. E. & S. Östlund-Nilsson, 2008. Does size matter for hypoxia tolerance in fish?  
634 Biological Reviews 83:173-189 doi:[https://doi.org/10.1111/j.1469-  
635 185X.2008.00038.x](https://doi.org/10.1111/j.1469-185X.2008.00038.x).

636 Nilsson, G. E. & G. M. C. Renshaw, 2004. Hypoxic survival strategies in two fishes: extreme  
637 anoxia tolerance in the North European crucian carp and natural hypoxic  
638 preconditioning in a coral-reef shark. Journal of Experimental Biology 207:3131-3139  
639 doi:<https://doi.org/10.1242/jeb.00979>.

640 Nilsson, G. E., P. R. Rosen & D. Johansson, 1993. Anoxic depression of spontaneous  
641 locomotor activity in Crucian carp quantified by a computerized imaging technique.  
642 The Journal of Experimental Biology 180:153-162.

643 Poff, N., M. Brinson & J. Day, 2002. Potential impacts on inland freshwater and coastal  
644 wetland ecosystems in the United States. Aquatic Ecosystem and Global Climate  
645 Change.

646 Powell, R., 2000. Animal home ranges and territories and home range estimators In: Boitani  
647 L, Fuller T, editors. Research techniques in animal ecology: controversies and  
648 consequences. New York: Columbia University Press.

649 R Core Team, 2020. R: A language and environment for statistical computing. . R Foundation  
650 for Statistical Computing, Vienna, Austria.

651 Rogers, N. J., M. A. Urbina, E. E. Reardon, D. J. McKenzie & R. W. Wilson, 2016. A new  
652 analysis of hypoxia tolerance in fishes using a database of critical oxygen level (Pcrit).  
653 Conservation physiology 4 doi:<https://doi.org/10.1093/conphys/cow012>.

654 Schurmann, H. & J. Steffensen, 1994. Spontaneous swimming activity of atlantic Cod *gadus*  
655 *morhua* exposed to graded hypoxia at three temperatures. The Journal of  
656 Experimental Biology 197:129-142.

657 Slavík, O. & P. Horký, 2009. When fish meet fish as determined by physiological sensors.  
658 Ecology of Freshwater Fish 18:501-506 doi:[https://doi.org/10.1111/j.1600-  
659 0633.2009.00370.x](https://doi.org/10.1111/j.1600-0633.2009.00370.x).

660 Slavík, O., P. Horký, M. Maciak & M. Wackermannová, 2016. Familiarity, prior residency,  
661 resource availability and body mass as predictors of the movement activity of the  
662 European catfish. *Journal of Ethology* 34:23-30 doi:[https://doi.org/10.1007/s10164-](https://doi.org/10.1007/s10164-015-0441-9)  
663 [015-0441-9](https://doi.org/10.1007/s10164-015-0441-9).

664 Slavík, O., P. Horký & L. Závorka, 2014. Energy Costs of Catfish Space Use as Determined by  
665 Biotelemetry. *Plos One* 9:e98997 doi:<https://doi.org/10.1371/journal.pone.0098997>.

666 Smith, F., 2013. Understanding HPE in the VPS Telemetry System. VEMCO Tutorials.

667 Snobl, Z. R., R. P. Koenigs, R. M. Bruch & F. P. Binkowski, 2015. Do Tags Exceeding 2% of Total  
668 Body Weight Impair Lake Sturgeon Movement? *North American Journal of Fisheries*  
669 *Management* 35:880-884 doi:<https://doi.org/10.1080/02755947.2015.1069425>.

670 Souchon, Y. & L. Tissot, 2012. Synthesis of thermal tolerances of the common freshwater fish  
671 species in large Western Europe rivers. *Knowledge and Management of Aquatic*  
672 *Ecosystems*:03.

673 Stecyk, J. A. W., K.-O. Stensløkken, A. P. Farrell & G. E. Nilsson, 2004. Maintained Cardiac  
674 Pumping in Anoxic Crucian Carp. *Science* 306:77-77  
675 doi:<https://doi.org/10.1126/science.1100763>.

676 Weber, R. E., G. Lykkeboe & K. Johansen, 1976. Physiological properties of eel haemoglobin:  
677 hypoxic acclimation, phosphate effects and multiplicity. *The Journal of Experimental*  
678 *Biology* 64:75-88.

679 Wells, R. M. G., 2009. Chapter 6 Blood-Gas Transport and Hemoglobin Function: Adaptations  
680 for Functional and Environmental Hypoxia. In Richards, J. G., A. P. Farrell & C. J.  
681 Brauner (eds) *Fish Physiology*. vol 27. Academic Press, 255-299.

682 Winter, J., 1996. Advances in underwater biotelemetry. *Fisheries techniques*, 2nd edition  
683 American Fisheries Society, Bethesda, Maryland:555-590.

684 Zeileis, A., C. Kleiber, W. Krämer & K. Hornik, 2003. Testing and dating of structural changes  
685 in practice. *Computational Statistics & Data Analysis* 44:109-123  
686 doi:[https://doi.org/10.1016/S0167-9473\(03\)00030-6](https://doi.org/10.1016/S0167-9473(03)00030-6).

687 Zeileis, A., F. Leisch, K. Hornik & C. Kleiber, 2002. strucchange: An R Package for Testing for  
688 Structural Change in Linear Regression Models. 2002 7:38  
689 doi:<https://doi.org/10.18637/jss.v007.i02>.

690 Zuur, A. F., E. N. Ieno, N. J. Walker, A. A. Saveliev & G. M. Smith, 2009. Mixed effects models  
691 and extensions in ecology with R; Gail M, Krickeberg K, Samet JM, Tsiatis A, Wong W,  
692 editors. New York, NY: Spring Science and Business Media.

693

## 694 Tables

695

696 **Table 1** Total length (in mm, mean, sd in italics and range) and weight (in g, mean, sd in  
697 italics and range) of all 40 tracked catfish, 9 smallest, 24 medium and 7 largest ones.

	Total length (mm)	Weight (g)
All 40 individuals	1 033 ( <i>328</i> ) [727 ; 2 150]	9 658 ( <i>13 389</i> ) [2 301 ; 64 380]



9 smallest individuals	812 (41) [727 ; 847]	3 370 (688) [2 301 ; 4 180]
24 medium individuals	944 (63) [855 ; 1 060]	5 563 (1 065) [3 760 ; 7 680]
7 largest individuals	1 623 (423) [1 100 ; 2 150]	31 786 (22 494) [8 380 ; 64 380]

699 **Table 2** Numeric results from the Generalized Linear Mixed-Effects Model that tested the  
700 fixed effects of fish size, day and their interactions on mean individual daily speed (panel a),  
701 mean individual distance to shore (panel b), individual home ranges 95% (panel c) and  
702 individual core areas (panel d). Fish identity was used as a random effect.

Daily speed <b>a</b>			
	d.f.	F	p-value
Size	2	1.301	0.274
Day	9	12.083	< 0.001
Size: Day	18	3.897	< 0.001
Individual	16.26	0.79	0.004
Explained variance (%)	56.6		
Distance to shore <b>b</b>			
	d.f.	F	p-value
Size	2	3.163	0.043
Day	9	29.599	< 0.001
Size: Day	18	1.199	0.260
Individual	24.1	1.88	< 0.001
Explained variance (%)	59.6		
Home range 95% <b>c</b>			
	d.f.	F	p-value
Size	2	4.689	0.010
Day	9	11.069	< 0.001
Size: Day	18	2.517	< 0.001
Individual	6.887	1.88	0.207
Explained variance (%)	53.7		
Core area <b>d</b>			
	d.f.	F	p-value
Size	2	2.824	0.061
Day	9	11.984	< 0.001
Size: Day	18	2.963	< 0.001

---

Individual	15.67	0.712	0.001
Explained variance (%)	49.9		

---

703

704