



HAL
open science

LIPOMECE: towards a better understanding of ruminant milk lipolysis through an integrative biology approach in milk and mammary gland

Catherine Hurtaud, Laurence Bernard, P. Trossat, Marine Gele, S. Meurisse, Anne Barbat, Didier Boichard, H el ene Larroque, A. Oudotte, J.L. Poulet, et al.

► To cite this version:

Catherine Hurtaud, Laurence Bernard, P. Trossat, Marine Gele, S. Meurisse, et al.. LIPOMECE: towards a better understanding of ruminant milk lipolysis through an integrative biology approach in milk and mammary gland. FAO-CIHEAM 2022 Joint seminar of networks on pasture and forage crops and on sheep and goat nutrition, Sep 2022, Catania, Italy. hal-03796220

HAL Id: hal-03796220

<https://hal.inrae.fr/hal-03796220v1>

Submitted on 4 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

LIPOMECC: towards a better understanding of ruminant milk lipolysis through an integrative biology approach in milk and mammary gland

C. Hurtaud¹, L. Bernard², P. Trossat³, M. Gelé⁴, S. Meurisse⁴, A. Barbat⁵, D. Boichard⁵, H. Larroque⁶, A. Oudotte³, J.L. Poulet⁴, A. Thierry⁷, M. Boutinaud¹, M. Delosière², A. Rau⁵, M. Bonnet², and C. Cebo⁵

¹PEGASE, INRAE, Institut Agro, 35590, Saint-Gilles, France

²Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, 63122 Saint-Genes-Champanelle, France

³Actalia-Cécalait, Rue de Versailles, 39800 Poligny, France

⁴Institut de l'Elevage, Maison Nationale des Eleveurs, 149 rue de Bercy, 75595, Paris cedex 12, France

⁵Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350, Jouy-en-Josas, France

⁶GenPhySE, Université de Toulouse, INRA, INPT, ENVT, Chemin de Borde Rouge, F-31326 Castanet-Tolosan Cedex, France

⁷UMR1253 Science et Technologie du Lait et de l'Œuf, INRA, Agrocampus Ouest, 65 rue de Saint Brieuc, 35000 Rennes, France

Take home Message LIPOMECC is the first large-scale integrative project to study lipolysis mechanisms in the milk and mammary gland in cow, goat and ewe.




Figure 1: A schematic overview of the LIPOMECC project

CE21-0010) and dairy stakeholders (APIS-GENE), is the first large-scale integrative project to study the lipolytic system in the milk and mammary gland in the three dairy species.

Material and methods An overview of the LIPOMECC project is depicted in **Figure 1**. A feed restriction study was conducted in the 3 dairy species (cow, goat and ewe). For each experiment, two levels of feeding were applied: “non-restricted” with animals fed 100% of the dry matter intake (DMI) ad libitum and “restricted” with animals fed at 65% of the DMI ad libitum. The impact of the milking frequency and milking intervals on lipolysis were also studied on cow. Lipolysis values were measured by the reference methods in each species, i.e. the copper-soap method for cow, and the BDI method for goat and ewe. Samples generated (milk, blood, mammary gland biopsies) are currently being used to characterize mechanisms of lipolysis in milk and mammary gland through an integrative approach in the three dairy species. Our approaches include animal science, milk biochemistry and biophysics, proteomics, lipidomics, genetics, transcriptomics, cell biology techniques, data modelling, and statistics. Interactions between genetics, physiology, farming practices and technological factors related to the milking process will be considered.

Results and discussion Milk yield decreased in the three species in response to the nutritional challenge. For cows, feed restriction caused as expected an increase in spontaneous lipolysis of milk (+0.20 Meq/100 g of fat, $P < 0.001$). Conversely, feed restriction induced a decrease in lipolysis in ewe milk (-0.27 Meq/100 g of fat, $P < 0.01$), whereas no significant effect was observed for goats under feed restriction. This strongly suggests that lipolytic systems and their adaptations to physiological challenges sharply differ between species. Omics studies on the samples collected during animal experiments are currently in progress to decipher molecular mechanisms occurring in milk and mammary gland with regard to lipolysis in the three species. In addition, an equation based on mid-infra-red (MIR) spectra has been developed to explore the genetic determinism of spontaneous lipolysis in dairy cows. A similar approach is currently in progress in dairy goats within the frame of the LIPOMECC project.

Conclusion Taken together, these results will improve understanding of the lipolytic systems in the three dairy species and highlight inter-species peculiarities with regard to the regulation of lipolysis in milk.

Acknowledgements LIPOMECC receives financial support from Agence Nationale de la Recherche and APIS-GENE, which brings together professionals from the French ruminant industry (<https://www.apis-gene.com/>).

References

Vanbergue E., Peyraud J. L., Ferlay A., Miranda G., Martin P., Hurtaud C. 2018. *Livestock Science* 217, 116-125.

Chilliard, Y., Toral P. G., Shingfield K. J., Rouel J., Leroux C., Bernard L. 2014. *Small Ruminant Research* 122, 31-37.