



HAL
open science

An equation based on MIR spectra to explore the genetic determinism of spontaneous lipolysis in dairy cows

Marine Gelé, Sofia Meurisse, Philippe Trossat, Aurore Oudotte, Didier Boichard, Laurence Bernard, Catherine Hurtaud, Anne Barbat, Christelle Cebo

► **To cite this version:**

Marine Gelé, Sofia Meurisse, Philippe Trossat, Aurore Oudotte, Didier Boichard, et al.. An equation based on MIR spectra to explore the genetic determinism of spontaneous lipolysis in dairy cows. 2022 ICAR/Interbull Conference, May 2022, Montréal, Canada. pp.132-136. hal-03796270

HAL Id: hal-03796270

<https://hal.inrae.fr/hal-03796270v1>

Submitted on 4 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An equation based on MIR spectra to explore the genetic determinism of spontaneous lipolysis in dairy cows

M. Gelé¹, S. Meurisse¹, P. Trossat², A. Oudotte², D. Boichard³, L. Bernard⁴, C. Hurtaud⁵, A. Barbat³ and C. Cebo³

¹*Institut de l'Élevage, Maison Nationale des Éleveurs, 149 rue de Bercy, 75595, Paris cedex 12, France (marine.gele@idele.fr)*

²*Actalia-Cécalait, Rue de Versailles, 39800 Poligny, France*

³*Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350 Jouy-en-Josas, France*

⁴*INRAE, Université Clermont Auvergne, VetAgroSup, UMR Herbivores, 63122, Saint-Genès-Champanelle, France*

⁵*PEGASE, INRAE, Institut Agro Rennes-Angers, 35590, Saint-Gilles, France*

Abstract (400 words maximum)

Mid infrared (MIR) spectra have been used since the late 2000s to phenotype new traits at large scale through prediction equations. It has been used in France since 2012 to quantify free fatty acids on herd milks in the framework of quality-based milk payment system. In addition to the loss of income, a high lipolysis level leads to a degradation of organoleptic (rancid taste) and technological (processing inability) properties of milk. Monitoring lipolysis is required to maintain the quality of milk and dairy products.

Lipolysis depends on many factors: genetics, farming practices, milking equipment, transformation process. Regarding genetics, Vanbergue highlighted in 2017 the existence of a genetic susceptibility to spontaneous lipolysis. The present study aims to investigate this issue further, by studying the genetic determinism of spontaneous lipolysis in a larger number of cows. This work is carried out within the framework of the LIPOMECA project which aims to better understand the molecular mechanisms controlling the degradation of milk fat.

Studying the genetic determinism of lipolysis requires the phenotyping of a large number of cows. As the equation initially developed on herd milk was not fully appropriate for individual milks (range, precision), a new prediction equation was estimated from the MIR spectra of individual milks. For this purpose, 432 milk samples were collected in 4 experimental farms in 2018 (approximately 40 cows per farm sampled 2 or 3 times a year) to maximize the variability of breeds (Holstein, Normande, Montbéliarde) and diets. A joint analysis of lipolysis according ISO/TS 22113 standard (BDI method) and by MIR spectrometry was carried out on each sample. Lipolysis measured by BDI method averaged 0.53 mmol/100 g fat (sd=0.41 mmol/100 g fat). The equation was developed by Partial Least Square regression after LOG transformation. Its coefficient of determination R^2 reached 0.72, with a residual standard deviation $S_{y,x}$ of 0.19 mmol/100 g of fat. The equation was then applied to obtain phenotypes on more than 300,000 MIR milk spectra from Holstein, Normande and Montbéliarde breeds. Genetic parameters were estimated using a repeatability animal model. Heritability and repeatability estimates were 0.21 and 0.40 in both Normande and Hostein breeds but higher (0.31 and 0.46) in Montbéliarde breed.

This work opens the opportunity to new uses of MIR spectra to improve the control of lipolysis in farm, by a closer management of the herd, or even by selection.

The LIPOMECA project was funded by APIS-GENE.

Keywords: lipolysis, dairy cattle, mid infrared spectroscopy, genetic