



HAL
open science

Les déterminants du climat. Cours Pour l'école Bordeaux-Sciences Agro (2e année). 2022

Denis Loustau

► **To cite this version:**

Denis Loustau. Les déterminants du climat. Cours Pour l'école Bordeaux-Sciences Agro (2e année). 2022. École d'ingénieur. France. 2022. hal-03799287

HAL Id: hal-03799287

<https://hal.inrae.fr/hal-03799287>

Submitted on 5 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Climate « change » ?

What is climate ?

The climate - Section 1

1. Climate: definitions and determinants
2. Quantitative aspects
3. The vegetation- climate equilibrium: the global ecosystem

The climate disturbance. Section 2

- Past and recent changes
- The state of the Earth system in 2022
- Scenarios for 2022-2120

Part 1. Climate: usual definitions

- **Climate**

- Climate is commonly defined as **the weather averaged over a “long” period.** The standard averaging period is **30 years** (WMO) that is allowing to define statistics: a **normal (mean value over 30 years)**, magnitudes of day-to-day or year-to-year variations, events frequency, min. max. etc .

- **Microclimate**

- Climate corresponding to the boundary layer coating an homogeneous surface (corn field, lake, forest plot...)

- **Climatology**

- Interdisciplinary scientific research discipline related to the climate and aiming at the characterisation of the climate, classification of climate types, their location on the earth ,their determinants and the analysis of their temporal variability .

- **Meteorology**

- Interdisciplinary scientific research discipline related to the atmosphere chemistry and physics which explains and forecasts weather events. Meteorology has application in many diverse fields such as the military, energy production, transport, agriculture and construction, health, ...

Part 1. The determinants of the temperature

Climate being summarised by the air temperature, what ingredients do we need for making a climate ?

Answer: first, energy !

Planck, Stefan Boltzmann laws:

$$j = \varepsilon \sigma T^4$$

j energy emitted or radiant emittance ($\text{J s}^{-1} \text{m}^{-2}$ or W m^{-2}),

ε emissivity

σ S-B constant,



T surface temperature (K)

Part 1. The determinants of the temperature: 1. solar radiation

The **sun and earth radiation emissions** are the almost unique, ultimate sources of energy

(Earth geothermic energy is ~0.03% of earth surface energy balance)


1. Earth rotation and obliquity causes the short term variations of earth insolation (hour-year).




Part 1. The determinants of the temperature: 1. solar radiation

Global distribution of incident solar radiation (SW ↓)

<http://climvis.org/anim/maps/global/nswrs.html>



Mean SW Radiation absorbed




Part 1. The determinants of the temperature: 1. solar radiation

2. Long term variations of incoming solar radiation are caused by changes in:

- Eccentricity (10 000 yr \pm 1%)
- Obliquity (41 000 yr, 22.5 to 24°)
- Axial precession, for high latitudes irradiance (26 000yr)

That were described first by Milankovitch




Credit: NASA/JPL-Caltech

<https://climate.nasa.gov/blog/2949/why-milankovitch-orbital-cycles-cant-explain-earths-current-warming/>


Part 1. The determinants of the temperature: 2. Earth radiation

Global distribution of LW radiation balance

<http://climvis.org/anim/maps/global/nswrs.html>



— SW radiation absorbed
— LW radiation emitted



SW + LW earth radiative balance


- Imbalance between the poles (-) and equator(+)
- Creates the air convective mass transportation from Equator to the poles

Supports the fast transportation in fluid envelopes of:

- Heat
- Water vapour ; liquid water
- Greenhouse gases: CO_2 , CH_4 , N_2O , O_3
- Pollutants: NO_x , NH_4 , SO_3

Part 1. The temperature determinants: 3. circulation of the earth fluid envelopes


ATMOSPHERE



Part 1. The climate determinants: 4. the « greenhouse » effect

Radiative exchanges between the earth surface and the atmosphere

1. Any body emits a radiation depending on its surface temperature that determines:
 - its spectral composition (Wien's law)
 - Its energy (Stefan-Bolzmann's law).
2. 70% of the solar radiation ($T=5600\text{ K}$) enters into the atmosphere, 30% is reflected.
3. The earth surface $T\sim 288\text{ K}$.
4. 40% of earth radiation is absorbed by the atmosphere and transformed in heat : this is the **greenhouse effect**.
30%



The climate determinants: 5. Surface feedbacks –radiative properties

Ice sheet extent variations affect surface albedo


The configuration of Northern Hemisphere ice sheets through the Quaternary

Christine L. Batchelor , Martin Margold, Mario Krapp, Della K. Murton, April S. Dalton, Philip L. Gibbard, Chris R. Stokes, Julian B. Murton & Andrea Manica


Nature Communications 10, Article number: 3713 (2019) | [Cite this article](#)

Fig. 2

From: The configuration of Northern Hemisphere ice sheets through the Quaternary



and sea level, so continental surface area.




The climate determinants: 5. Surface feedbacks –radiative properties



— Ok, Hornbeam

— Pines

— Holm Oak + Pines



✓ The albedo varies from 0.06 to 2.25 according to species.

The climate determinants: 5. Surface feedbacks – air humidity effects


Heat exchanges between surface and atmosphere: the latent heat case.

1st Feed back +: the air humidity increases the emissivity


ETR \rightarrow air humidity \uparrow \rightarrow LW radiation \uparrow \rightarrow T min \uparrow

2nd Feed back +: condensation is a net source of heat

ETR \rightarrow air humidity \uparrow \rightarrow dew point T \uparrow \rightarrow Latent heat for condensation \uparrow & Tmin \uparrow




Correlation between atmospheric emissivity and min. Temperature for 222 stations in N Americ.



Corrélation between dew point temperature and min Temperature for 222 met. stations .

The climate determinants: 5. Surface feedbacks – air humidity effects



Tmin = 27 °C

Tmin = 12 °C

25
g.m⁻³

11
g.m⁻³

18
g.m⁻³

18
g.m⁻³

Observations


Simulation with no
evapotranspiration

Part 1. Synthesis: the earth surface energy balance

Annual mean energy balance components over land


(Wild et al. 2015. Clim. Dynamics, DOI 10.1007/s00382-014-2430-z)

Solar radiation



albedo

Terrestrial radiation



Convective fluxes

H₂O Heat CO₂

Momentum

Bilan de la surface

$$\begin{aligned}
 +20 + 22 + 94 &= +136 \\
 -114 - 12 - 10 &= -136
 \end{aligned}$$





Bilan de l'atmosphère

$$\begin{aligned}
 +20 + 4 + 101 + 12 + 10 + 6 &= +153 \\
 -9 - 50 - 94 &= -153
 \end{aligned}$$

6
from ocean

Evaporation sensible heat

Part 1. The earth energy balance



The determinants of the climate: questions

Bilan de l'atmosphère

$$+20+4+101+12+10+6 = +153$$

$$-9-50-94 = -153$$

Bilan de la surface

$$+20+22+94 = +136$$


$$-114-12-10 = -136$$

What would happen in case the energy budget is not balanced ?

Part 2. Quantitative aspects : meteorological variables and their measurement

Temperature: degree Celsius, Kelvin.

Air water content : water vapor pressure (e in Pa), saturation pressure ,
saturation deficit of water vapor (δe).



→ The water vapor saturation deficit increases with the air temperature

→ δe is the driver of plant and animal transpiration :

→ when $e_{(surface)} - e_{(air)} = 0$, transpiration = 0

→ List the situation allowing transpiration to be >0 . Same for transpiration <0

NB. The air relative humidity ($HR = P_{air} / P_{sat}$) has no physical meaning. Should be avoided.

Part 2. The climate , quantitative aspects : meteorological variables and their measurement

Radiation ($J.T^{-1}.L^{-2}$):

quantum(Photons. m^{-2})

energy (Joule . s^{-1} . m^{-2} = $W . m^{-2}$).

Pressure

Pressure is force applied on a surface. Unit force per unit area, with SI unit = pascal

(Pa = $N.m^{-2}$ = $J.m^{-3}$ = $Kg.m^{-1}.s^{-2}$):

NB 1mBar = 1hPa =100 Pa.

Wind: a displacement of air caused by pressure differences. Wind flow can be zero, laminar (rare) or turbulent. Most often, only the horizontal component is given.


Unit ($L^3.L^{-2}.T^{-1}$) = $L.T^{-1}$ = $m.s^{-1}$

Precipitations: downward flow of solid or liquid water on to the surface. Type (snow, ice, rain, fog) intensity, interception, ($mass.T^{-1}.L^{-2}$)

Part 2. How often do climate variables change?

Power spectra of climate variables over the Fontainebleau forest, France, 2007-2012. (Moreaux et al. Tellus b, 2020).

- Radiation (SW) and temperature (Ta) variation patterns are forwarded to the air water vapour saturation deficit (VPD)
- Precipitation and soil water (REW), influenced by surface and local conditions shows distinctive patterns (weak high-frequency variations).



Which variables are relevant for describing the atmosphere ?

Physician answer : all of them !

Practical answer : the ones we can measure ! (Temperature, Precipitations,...)

Which statistics ?

Average values ? Min Max ? n^{th} order moments ?

Part 2. The climate , a quantitative assessment : temporal spectrum

The time characteristics of meteorological variables:

Temporal spectrum of meteorological variables matches the time characteristics of drivers:

PERIOD	DRIVER	EFFECT
10^4 yr	Milankovich cycles	glaciation cycles
10^{2-3} yr	Fossil GHG emissions	« climate change »
1 yr	Earth ellipsoid around Sun	Annual period
0.25 yr	Surface inclination angle	Season (mid latitude)
Week-day	Frontal wind regime	Anticyclone, depression
12 hours	Earth rotation	Night – Day sea / earth breeze
6 hours (tide)	Sun – Moon – Earth alignment	Tides
hour	Cloud passage duration	Sky fraction, direct/diffuse light
mn-sec-0.1 sec	Surface roughness length	Turbulence

Part 2. The climate , a quantitative assessment : temporal spectrum

Which is the relevant temporal scale for defining a climate?

→ pragmatic answer: 30 yrs (man's memory ?)

→ Mathematical answer : the shortest duration at which statistics (n^{th} order moments) are steady.

Part 3. The connection between vegetation and climate types:

1) phytogeography





Fig. 8: **TERRESTRIAL BIOGEOGRAPHIC REALMS AND BIOMES**



Part 3. Climate classification / types



Af	BWh	Csa	Cwa	Cfa	Dsa	Dwa	Dfa	ET
Am	BWk	Csb	Cwb	Cfb	Dsb	Dwb	Dfb	EF
Aw	BSh	Cwc	Cfc	Dsc	Dwc	Dfc		
BSk		Dsd		Dwd	Dfd			

DATA SOURCE : GHCN v2.0 station data
Temperature (N = 4,844) and
Precipitation (N = 12,396)

PERIOD OF RECORD : All available

MIN LENGTH : ≥30 for each month.


RESOLUTION : 0.1 degree lat/long

Contact : Murray C. Peel (mpeel@unimelb.edu.au) for further information



THE UNIVERSITY OF
MELBOURNE

Part 3. Climates: linking biomes to climate types




A **biome** is a biotic community defined by its dominant vegetation (« vegetation belts »).

The distribution of the vegetation in biomes can be deduced from the climate types. (NB y axis reversed).

Part 3. The connection between vegetation and climate :

2) biogeochemistry


Net Primary Productivity



Data taken from: BIS Simulation
(Rubsark, et al. 2000)
(Foley, et al. 1996)

Atlas of the Biosphere
Center for Sustainability and the Global Environment
University of Wisconsin - Madison

Part 3. The connection between vegetation and climate



La pression de sélection climatique donne lieu au phénomène de **convergence fonctionnelle ou convergence adaptative** qui est l'apparition des mêmes caractéristiques adaptatives chez les espèces non apparentées soumises à une pression identique.

Ex. : métabolisme CAM, Succulence, sclérophyllie, métabolisme C3 / C4, phénotypes de Raunkier etc...




Part 3. Vegetation and climate : a story of reciprocal interactions

Concise history of the GHG atmospheric concentration:

Link with land colonisation

- CO₂ atmospheric concentration →
- Conductive plant tissues (tracheids) →
- Wood (lignin), leaves →
- Trees (coniferous) →
- Soil →


Beerling D J , Berner R A
PNAS 2005;102:1302-1305
©2005 by National Academy of Sciences



Part 3. Vegetation and climate : reciprocal interactions


High frequency variations of Ecosystem physiology are controlled by meteo variables.

- The climate control (radiation) is tighter at high frequency
- Biological adjustments dampen the low frequency variability of ecosystem activity
- Respiration is more closely linked to temperature



Part 3. The global ecosystem concept

- Time constants = 1 to $>10^6$ yrs (from superfast to ultra slow)
- The connection between vegetation and atmosphere : a model of ecosystem



Interactions :
Biophysical

Biogeochemical

Hydrological

Biocoenosis

Biotope

Negative feedback

Climate: few thoughts


The concept of climate is based upon the **sensual perception by Man** of the weather. The climate classification is based on statistics calculated over a typical 30yr time series (the memory of a human being).

Values of all climate variables are varying actually at different temporal resolutions along a variance spectrum from 10^5 to 10^{-12} Hz

This variability is well understood. It was shaped for the last 800 000 years by the fluctuations of mostly cosmic and secondarily atmospheric drivers.

Supplementary slides.

The climate – vegetation correspondence: the Holridge triangle.



Climatology: The Koeppen classification

- 3 letters code based upon the temperature and precipitations
- Statistics calculated over 30-yr temporal series

1^{re} lettre : type de climat [\[modifier \]](#) [\[modifier le code \]](#)

Code	Type	Description
A	Climat tropical	<ul style="list-style-type: none">• Température moyenne de chaque mois de l'année > 18 °C• Pas de saison hivernale• Fortes précipitations annuelles (supérieure à l'évaporation annuelle)
B	Climat sec	<ul style="list-style-type: none">• Évaporation annuelle supérieure aux précipitations annuelles. Ce seuil est calculé de la manière suivante :<ul style="list-style-type: none">• Si moins de 30 % des précipitations tombent en été (avril à septembre dans l'hémisphère nord) : Précipitations annuelles moyennes (mm) < 20 × température annuelle moyenne (°C)• Si plus de 70 % des précipitations tombent en été : Précipitations annuelles moyennes (mm) < 20 × température annuelle moyenne + 280• Autrement : Précipitations annuelles moyennes (mm) < 20 × température annuelle moyenne + 140
C	Climat tempéré	<ul style="list-style-type: none">• Température moyenne du mois le plus froid comprise entre 0 °C et 18 °C• Température moyenne du mois le plus chaud > 10 °C• Les saisons été et hiver sont bien définies
D	Climat continental	<ul style="list-style-type: none">• Température moyenne du mois le plus froid < 0 °C• Température moyenne du mois le plus chaud > 10 °C• Les saisons été et hiver sont bien définies
E	Climat polaire	<ul style="list-style-type: none">• Température moyenne du mois le plus chaud < 10 °C• La saison d'été est très peu marquée

Climatology: The Koeppen classification

- 3 letters code based upon the temperature and precipitations
- Statistics calculated over 30-yr temporal series

2^e lettre : régime pluviométrique [\[modifier | modifier le code \]](#)

Code	Description	S'applique à
S	<ul style="list-style-type: none">• Climat de steppe• Précipitations annuelles comprises entre 50 et 100 % du seuil calculé	B
W	<ul style="list-style-type: none">• Climat désertique• Précipitations annuelles < 50 % du seuil	B
w	<ul style="list-style-type: none">• Saison sèche en hiver<ul style="list-style-type: none">• Pour A : climat de la savane, P du mois le hivernal le plus sec < 60 mm et < $[100 - (\text{précipitations annuelles moyennes})/25]$• pour C et D : P du mois hivernal le plus sec < 1/10 du mois le plus humide	A-C-D
s	<ul style="list-style-type: none">• Saison sèche en été<ul style="list-style-type: none">• Pour A : climat de la savane, P du mois le estival le plus sec < 60 mm et < $[100 - (\text{précipitations annuelles moyennes})/25]$³• Pour C et D : P du mois estival le plus sec < 40 mm et < 1/3 du mois hivernal le plus humide	A-C-D
f	<ul style="list-style-type: none">• Climat humide, précipitations tous les mois de l'année<ul style="list-style-type: none">• Pour A : climat de la forêt tropicale, P du mois le plus sec > 60 mm• Pour C et D : pas de saison sèche, ni « w » ni « s »	A-C-D
m	<ul style="list-style-type: none">• Climat de mousson :• P du mois le plus sec < 60 mm et > $[100 - (\text{précipitations annuelles moyennes})/25]$	A
T	<ul style="list-style-type: none">• Température moyenne du mois le plus chaud comprise entre 0 °C et 10 °C	E
F	<ul style="list-style-type: none">• Température moyenne du mois le plus chaud < 0 °C	E
M	<ul style="list-style-type: none">• Précipitations abondantes• Hiver doux (Température moyenne du mois le plus froid > -10 °C)	E

Climatology: The Koeppen classification

- 3 letters code based upon the temperature and precipitations
- Statistics calculated over 30-yr temporal series

3^e lettre : variations de température

Pour affiner les types B, C et D, une troisième lettre précise l'amplitude du cycle annuel des températures :

Code	Description	S'applique à
a : été chaud	<ul style="list-style-type: none">• Température moyenne du mois le plus chaud > 22 °C	C-D
b : été tempéré	<ul style="list-style-type: none">• Température moyenne du mois le plus chaud < 22 °C• Températures moyennes des 4 mois les plus chauds > 10 °C	C-D
c : été court et frais	<ul style="list-style-type: none">• Température moyenne du mois le plus chaud < 22 °C• Températures moyennes mensuelles > 10 °C pour moins de 4 mois• Température moyenne du mois le plus froid > -38 °C	C-D
d : hiver très froid	<ul style="list-style-type: none">• Température moyenne du mois le plus froid < -38 °C	D
h : sec et chaud	<ul style="list-style-type: none">• Température moyenne annuelle > 18 °C	B
k : sec et froid	<ul style="list-style-type: none">• Température moyenne annuelle < 18 °C	B

Climatology: The Koeppen classification

Köppen climate classification scheme symbols description table^[1]


1st	2nd	3rd
A (Tropical)	f (Rainforest)	
	m (Monsoon)	
	w (Savanna, Wet)	
	s (Savanna, Dry)	
B (Arid)	W (Desert)	
	S (Steppe)	
		h (Hot)
		k (Cold)
		n (With frequent fog) ^[10]
C (Temperate)	s (Dry summer)	
	w (Dry winter)	
	f (Without dry season)	
		a (Hot summer)
		b (Warm summer)
		c (Cold summer)
D (Cold (continental))	s (Dry summer)	
	w (Dry winter)	
	f (Without dry season)	
		a (Hot summer)
		b (Warm summer)
		c (Cold summer)
		d (Very cold winter)
E (Polar)	T (Tundra)	
	F (Eternal winter (ice cap))	

Climatology: The Koeppen classification

Table 1: Key to calculate the climate formula of Köppen and Geiger for the main climates and subsequent precipitation conditions, the first two letters of the classification. Note that for the polar climates (E) no precipitation differentiations are given, only temperature conditions are defined. This key implies that the polar climates (E) have to be determined first, followed by the arid climates (B) and subsequent differentiations into the equatorial climates (A) and the warm temperate and snow climates (C) and (D), respectively. The criteria are explained in the text.

Type	Description	Criterion
A	Equatorial climates	$T_{\min} \geq +18 \text{ }^\circ\text{C}$
Af	Equatorial rainforest, fully humid	$P_{\min} \geq 60 \text{ mm}$
Am	Equatorial monsoon	$P_{\text{ann}} \geq 25(100 - P_{\min})$
As	Equatorial savannah with dry summer	$P_{\min} < 60 \text{ mm in summer}$
Aw	Equatorial savannah with dry winter	$P_{\min} < 60 \text{ mm in winter}$
B	Arid climates	$P_{\text{ann}} < 10 P_{\text{th}}$
BS	Steppe climate	$P_{\text{ann}} > 5 P_{\text{th}}$
BW	Desert climate	$P_{\text{ann}} \leq 5 P_{\text{th}}$
C	Warm temperate climates	$-3 \text{ }^\circ\text{C} < T_{\min} < +18 \text{ }^\circ\text{C}$
Cs	Warm temperate climate with dry summer	$P_{\text{smin}} < P_{\text{wmin}}, P_{\text{wmax}} > 3 P_{\text{smin}}$ and $P_{\text{smin}} < 40 \text{ mm}$
Cw	Warm temperate climate with dry winter	$P_{\text{wmin}} < P_{\text{smin}}$ and $P_{\text{smax}} > 10 P_{\text{wmin}}$
Cf	Warm temperate climate, fully humid	neither Cs nor Cw
D	Snow climates	$T_{\min} \leq -3 \text{ }^\circ\text{C}$
Ds	Snow climate with dry summer	$P_{\text{smin}} < P_{\text{wmin}}, P_{\text{wmax}} > 3 P_{\text{smin}}$ and $P_{\text{smin}} < 40 \text{ mm}$
Dw	Snow climate with dry winter	$P_{\text{wmin}} < P_{\text{smin}}$ and $P_{\text{smax}} > 10 P_{\text{wmin}}$
Df	Snow climate, fully humid	neither Ds nor Dw
E	Polar climates	$T_{\text{max}} < +10 \text{ }^\circ\text{C}$
ET	Tundra climate	$0 \text{ }^\circ\text{C} \leq T_{\text{max}} < +10 \text{ }^\circ\text{C}$
EF	Frost climate	$T_{\text{max}} < 0 \text{ }^\circ\text{C}$

Climatology: the French case



K-G climate in metropolitan France

Cf: Temperate

Cs: Mediterranean



Dw : Subarctic

ET: Tundra

Climatology: examples

Maritime (Cf)

Mediterranean (Cs)



Wet temperate forests

clerophyllous forests

Average Annual Temperature (oC) =	9.8
Annual Temperature Range (oC) =	16
Total Annual Precipitation (mm) =	1048
Summer Precipitation (mm) =	277
Winter Precipitation (mm) =	771

13.75
9
475
54
421





Climatology: examples

Tropical (Af)

Subarctic

Polar

Desert (Bw)



Annual T (°C)	26.1	-10	-17	22.8
Range (°C)	1.4	63	23	21.2

Annual Prec. (mm.yr ⁻¹)	2879	213	7.8	44
Summer Prec. (mm)	1583	157	3.7	12
Winter Prec. (mm)	1294	56	4.1	32

Vegetation type

Tropical
Rainforest

Coniferous

none

none