



HAL
open science

Data from a proteomic comparative analysis highlight differential adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* to cow milk versus to soy milk environments

Gwénaél Jan, Florian Tarnaud, Fillipe Luiz Rosa Do Carmo, Nassima Illikoud, Fanny Canon, Julien Jardin, Fanny Guyomarc'H, Valérie Briard-Bion, Valérie Gagnaire

► To cite this version:

Gwénaél Jan, Florian Tarnaud, Fillipe Luiz Rosa Do Carmo, Nassima Illikoud, Fanny Canon, et al.. Data from a proteomic comparative analysis highlight differential adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* to cow milk versus to soy milk environments. *Data in Brief*, 2022, 45, pp.108653. 10.1016/j.dib.2022.108653 . hal-03822271

HAL Id: hal-03822271

<https://hal.inrae.fr/hal-03822271>

Submitted on 20 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Data Article

Data from a proteomic comparative analysis highlight differential adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* to cow milk versus to soy milk environments



Gwénaél Jan*, Florian Tarnaud, Fillipe Luiz Rosa do Carmo, Nassima Illikoud, Fanny Canon, Julien Jardin, Valérie Briard-Bion, Fanny Guyomarc'h, Valérie Gagnaire

STLO, INRAE, Institut Agro, Rennes F-35000, France

ARTICLE INFO

Article history:

Received 18 August 2022

Revised 6 September 2022

Accepted 28 September 2022

Available online 4 October 2022

Dataset link: [Data from a proteomic comparative analysis highlight differential adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* to cow milk versus to soy milk environments \(Original data\)](#)

Keywords:

Yogurt

Starters

Soy milk

Cow milk

Proteomic

Mass spectrometry

Probiotic

Stress

ABSTRACT

The article presents a proteomic dataset generated by a comparative analysis, using gel-free nanoLC-MS/MS, of the cellular proteome of *Lactobacillus delbrueckii* subsp. *bulgaricus*, a yogurt starter, when cultivated in soy milk versus in cow milk. The CIRM-BIA1592 strain was cultivated in the aqueous phase of soy milk, or of cow milk. Whole-cell proteins were extracted, trypsinolyzed and analyzed by nano LC-MS/MS, prior to identification and to classification by function using the X!Tandem pipeline software and the proteomic data from NCBI.nlm.nih.gov. Quantification of the proteins was moreover performed to evidence changes in their expression, depending on the culture medium. Data are available via ProteomeXchange with the identifier PXD033905 (<http://www.proteomexchange.org/>). This article is related to the research article entitled “The stressing life of *Lactobacillus delbrueckii* subsp. *bulgaricus* in soy milk”, by G.Jan et al. in Food Microbiology, 2022. This proteomic differential analysis indeed revealed major modulation of the stress proteome, with many stress proteins upregulated in the soy environment.

* Corresponding author.

E-mail address: Gwenael.jan@inrae.fr (G. Jan).

Specifications Table

Subject	Food Science: Food Microbiology
Specific subject area	food microbiology, dairy starter, yogurt starter, probiotic, stress adaptation
Type of data	Mass spectrometry data
How the data were acquired	Mass spectrometry. Whole cell protein extracts were subjected to trypsinolysis prior to nano LC-MS/MS analysis.
Data format	Raw, Analyzed, and Filtered
Description of data collection	<i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> , strain CIRM-BIA1592, was cultivated either in soy milk, or in cow milk, environment. Whole-cell protein extracts were extracted, trypsinolized, and analyzed by nanoLC-MS/MS in a gel-free strategy. Each peptide identified by tandem mass spectrometry was quantified using the free MassChroQ software before data treatment and statistical analysis under R software.
Data source location	The strain of <i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> was provided by the CIRM-BIA (https://collection-cirmbia.fr/), international center for microbial resources, located in the STLO research unit (https://www6.rennes.inrae.fr/stlo). The work was performed at STLO, a research unit of: Institution: INRAE & Institut Agro City: Rennes Country: France
Data accessibility	Data are available via ProteomeXchange with the identifier PXD033905. (http://www.proteomexchange.org/). Repository name: PRIDE (Proteomics Identifications Database) Data identification number: PXD033905 Direct URL to data: https://www.ebi.ac.uk/pride/archive/projects/PXD033905/
Related research article	Jan G, Tarnaud F, Rosa do Carmo FL, Illikoud N, Canon F, Jardin J, Briard-Bion V, Guyomar'h F, Gagnaire V. 2022. The stressing life of <i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> in soy milk. Food Microbiol 106:104042. https://doi.org/10.1016/j.fm.2022.104042

Value of the Data

- These data constitute an important step in understanding the molecular mechanisms responsible for the adaptation of yogurt starters to the soy environment.
- These data should be considered in the development of innovative fermented soy products.
- These data evidence that the yogurt starter *L. bulgaricus* is stressed when cultivated in soymilk, compared to milk.

1. Data Description

This article presents a dataset generated during a comparison of the cellular proteome of the yogurt starter *Lactobacillus delbrueckii* subsp. *bulgaricus* cultivated in (1) cow milk environment and (2) soy milk environment. The list of all the cellular proteins that were identified in this study is provided in a table entitled “protein_quantification_results.csv” available via ProteomeXchange with the identifier PXD033905. Project Name: Data from a proteomic comparative analysis highlight differential adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* to cow milk versus to soy milk environments. Project accession: PXD033905.

The abundance ratios of the differentially expressed proteins, displaying a significantly different abundance when bacteria were cultivated in soy versus in cow milk environment, classified in different COG categories, are published in the following article: [1]. The stressing life of *Lactobacillus delbrueckii* subsp. *bulgaricus* in soy milk. Food Microbiol 106:104042. <https://doi.org/10.1016/j.fm.2022.104042>.

For each biological and each technical replicate, a raw data file is presented in the deposited dataset as mentioned above.

This article presents data from the research article entitled “The stressing life of *Lactobacillus delbrueckii* subsp. *bulgaricus* in soy milk”. The cellular proteome, according to the growth medium, reveals major induction of stress proteins, in the soy environment.

2. Experimental Design, Materials and Methods

2.1. Strain and Pre-Cultures

De Man, Rogosa, Sharpe (MRS) medium [2] was used to grow liquid precultures of *Lactobacillus delbrueckii* subsp. *bulgaricus* CIRM-BIA1592 at 42 °C without agitation. This strain was provided by CIRM-BIA, which is an International Centre for Microbial Resources dedicated to bacteria of food interest called CIRM-BIA, (INRAE Rennes, France, <https://collection-cirmbia.fr/>). These precultures were used to inoculate cultures, either in MRS, or in ultrafiltrate media, as described below.

2.2. Culture in Cow Milk or Soy Milk Ultrafiltrate

Raw cow milk was skimmed and then ultrafiltered on the INRAE STLO dairy platform as described previously [3,4]. In that aim, we used a filtration pilot, containing a ceramic membrane. This membrane has a molecular weight cut-off point of 8 kDa. This ultrafiltration concentrates dairy proteins in the retentate and generates a permeate, MUF (milk ultrafiltrate), which constitutes the aqueous phase of cow milk. The overall composition of MUF was as follows: carbohydrate 5%; non-protein nitrogen 0.28%, minerals 0.75% and dry matter 6.14%. As an additional nitrogen source, 5 g.L⁻¹ food grade casein hydrolysate (Casein Peptone Plus, Organotechnie, La Courneuve, France) was added. After adjustment of pH to 7.0 using NaOH, the MUF medium was sterilized using 0.2 µm filters (Nalgene, Roskilde, Denmark) prior to storage at 4 °C.

Soy milk ultrafiltrate (SUF) is a by-product of industrial concentration of soy proteins during the manufacture of soymilk. It was provided by a local company, Sojasun Technologies Triballat Noyal (Noyal-sur-Vilaine, France) and prepared according to patent N° EP 1 983 844 B1 [5]. This SUF is commercially available under the designation BASOSOY (Triballat ingredients Triballat, Noyal-sur-Vilaine, France). The overall composition of SUF was as follows: carbohydrates 2.5%, protein 0.55%, non-protein nitrogen 0.20%, mineral 1%, dry matter 5%. As an additional nitrogen source, 5 g.L⁻¹ food grade soy hydrolysate (Bacto Soytone, BD Bioscience) was added. After adjustment of pH to 7.0 using NaOH, the SUF medium was sterilized by autoclaving (110 °C, 10 min) prior to sterile centrifugation (28,000 × g, 30 min) and filtration in order to remove insoluble compounds. It was then sterilized using 0.2 µm filters (Nalgene, Roskilde, Denmark) prior to storage at 4 °C.

In this work, 4 independent cultures of *Lactobacillus delbrueckii* subsp. *bulgaricus* CIRM-BIA1592 were prepared at 42 °C without agitation in MUF. In addition, 4 independent cultures of *Lactobacillus delbrueckii* subsp. *bulgaricus* CIRM-BIA1592 were prepared at 42 °C without agitation in SUF. Cultures were incubated until the end of growth and of acidification, i.e. during 16 h. These 8 cultures were then subjected to whole-cell label-free proteomic analysis as described below. For each culture (4 biological replicates), 3 mass spectrometry analyses (3 technical replicates) were performed.

2.3. Label-Free Proteomics

Label-free proteomics was performed as described in details in the corresponding research article [1] and adapted from a previous study [6]. Briefly, lactobacilli were grown either in milk or soy ultrafiltrate and then lysed in lysis solution (50 mM Tris-HCl [pH 7.5], 0.3% SDS, 200 mM dithiothreitol (DTT), 0.4 mM phenyl methyl sulfonyl fluoride, PMSF). Protein extracts were cleaned and quantified using the 2-D Clean-Up kit (GE Healthcare) and the 2-D Quant Kit, respectively. Whole-cell tryptic digestion was done using Sequencing Grade Modified Trypsin (Promega, Madison, USA) as described previously [7] prior to addition of spectrophotometric-grade trifluoroacetic acid (TFA) (Sigma-Aldrich, USA).

Nano-LC-MS/MS was done as previously described [7] using a nano RSLC Dionex U3000 system fitted to a Q-Exactive mass spectrometer (Thermo Scientific, San Jose, USA). Spectra of eluted peptides were recorded in full MS mode, selected between 250–2000 m/z, with a resolution of 70,000 at m/z 200.

Identification of proteins was as previously described [7], peptides being identified using the X!Tandem pipeline software (Langella et al., 2017), querying the proteome of *L. delbrueckii* subsp. *bulgaricus* (proteome UP000001259 downloaded from uniprot.org).

Protein quantification was as described [7], each peptide being quantified using the MassChroQ software prior to data treatment and statistical analysis under the R software (R 3.2.2, Project for statistical computing) using the specific R package called 'MassChroQR'. Variance analysis was performed on proteins with a minimum peak ratio of 1.5 for peak counting analysis. We considered significantly different proteins with an adjusted p -value < 0.05.

Proteins with a significant ($p < 0.05$, ANOVA) change in expression of ≥ 2 -fold (\log_2 ratio ≥ 1.5) were considered differentially expressed. We generated a volcano plot to illustrate differentially expressed proteins in *L. delbrueckii* subsp. *bulgaricus*. Functional annotation and Clusters of Orthologous Groups (COGs) were done with eggNOG-mapper v2 web tool (Huerta-Cepas et al., 2017, 2019).

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [8] partner repository with the dataset identifier PXD033905.

Ethics Statements

This manuscript adheres to Ethics in publishing standards (<https://www.elsevier.com/journals/data-in-brief/2352-3409/guide-for-authors>).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

Data from a proteomic comparative analysis highlight differential adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* to cow milk versus to soy milk environments (Original data) (PRIDE).

CRedit Author Statement

Gwénaél Jan: Writing – review & editing, Visualization, Methodology, Supervision; **Florian Tarnaud:** Writing – review & editing, Methodology; **Fillipe Luiz Rosa do Carmo:** Writing – review & editing, Data curation; **Nassima Illikoud:** Writing – review & editing, Data curation;

Fanny Canon: Writing – review & editing, Methodology; **Julien Jardin:** Writing – review & editing, Methodology; **Valérie Briard-Bion:** Writing – review & editing, Methodology; **Fanny Guyomarc’h:** Writing – review & editing, Methodology; **Valérie Gagnaire:** Writing – review & editing, Visualization, Methodology, Supervision.

Acknowledgments

N. Illikoud and F. Rosa do Carmo are recipients of post-doctoral fellowships from the French FUI (Fond Unique Interministériel) program, N°25, ProBioVeg. This study is part of the PhD project of F. Canon funded by the French National Research Institute for Agriculture, Food and Environment (INRAE) and the Brittany region. The authors offer their thanks to Victoria Chuat, from CIRM-BIA, for storing, checking and maintaining the bacterial strains.

References

- [1] G. Jan, F. Tarnaud, F.L. Rosa do Carmo, N. Illikoud, F. Canon, J. Jardin, V. Briard-Bion, F. Guyomarc’h, V. Gagnaire, The stressing life of *Lactobacillus delbrueckii* subsp. *bulgaricus* in soy milk, *Food Microbiol.* 106 (2022) 104042, doi:10.1016/j.fm.2022.104042.
- [2] J.D. De Man, M. Rogosa, M.E. Sharpe, A medium for the cultivation of lactobacilli, *J. Appl. Bacteriol.* 23 (1960) 130–135.
- [3] F.J. Cousin, S. Louesdon, M.B. Maillard, S. Parayre, H. Falentin, S.M. Deutsch, G. Boudry, G. Jan, The first dairy product exclusively fermented by *Propionibacterium freudenreichii*: a new vector to study probiotic potentialities in vivo, *Food Microbiol.* 32 (2012) 135–146.
- [4] M.C. Michalski, N. Leconte, V. Briard-Bion, J. Fauquant, J.L. Maubois, H. Goudedranche, Microfiltration of raw whole milk to select fractions with different fat globule size distributions: process optimization and analysis, *J. Dairy Sci.* 89 (2006) 3778–3790.
- [5] Efstathiou, T., Driss, F., 2010. Method for production of active extracts from soya beans and corresponding uses of the produced extracts. EP1983844B1. Available at: <https://patents.google.com/patent/EP1983844B1/en>. Accessed February 3, 2020.
- [6] F. Gaucher, S. Bonnassie, H. Rabah, P. Leverrier, S. Pottier, J. Jardin, V. Briard-Bion, P. Marchand, R. Jeantet, P. Blanc, G. Jan, Benefits and drawbacks of osmotic adjustment in *Propionibacterium freudenreichii*, *J. Proteom.* 204 (2019) 103400, doi:10.1016/j.jprot.2019.103400.
- [7] S. Huang, F. Gaucher, C. Cauty, J. Jardin, Y. Le Loir, R. Jeantet, X.D. Chen, G. Jan, Growth in hyper-concentrated sweet whey triggers multi stress tolerance and spray drying survival in *Lactobacillus casei* BL23: from the molecular basis to new perspectives for sustainable probiotic production, *Front. Microbiol.* 9 (2018) 1–12.
- [8] Y. Perez-Riverol, J. Bai, C. Bandla, D. García-Seisdedos, S. Hewapathirana, S. Kamatchinathan, D.J. Kundu, A. Prakash, A. Frericks-Zipper, M. Eisenacher, M. Walzer, S. Wang, A. Brazma, J.A. Vizcaino, The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences, *Nucl. Acids Res.* 50 (2022) D543–D552, doi:10.1093/nar/gkab1038.