



HAL
open science

A portable and unilateral NMR device to measure tree water content and locate conductive tissues

Shannan Blystone, Guilhem Pagès, Hervé H. Cochard, Pierre Conchon

► To cite this version:

Shannan Blystone, Guilhem Pagès, Hervé H. Cochard, Pierre Conchon. A portable and unilateral NMR device to measure tree water content and locate conductive tissues. Italian-French International Conference on Magnetic Resonance, Sep 2022, Milan, Italy. hal-03822537

HAL Id: hal-03822537

<https://hal.inrae.fr/hal-03822537>

Submitted on 20 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A PORTABLE AND UNILATERAL NMR DEVICE TO MEASURE TREE WATER CONTENT AND LOCATE CONDUCTIVE TISSUES

Blystone, Shannan^{††}, Pagès, Guilhem[‡], Cochard, Hervé[†], Conchon, Pierre[†]

[‡]INRAE, QuaPA, F-63122 St Genes Champanelle, France


[†]INRAE, PROBE research infrastructure, AgroResonance facility, F-63122 St Genes Champanelle, France

^{††}Université Clermont Auvergne, INRAE, PIAF, 63000 Clermont-Ferrand, France

We evaluated the NMR capacities of a portable, unilateral magnet having the potential to evaluate tree water dynamics *in-situ*: The Nuclear Magnetic Resonance Mobile Universal Surface Explorer (NMR-MOUSE). We tested the capacity of this device to measure tree water content and to differentiate the conductive tissues, i.e. the xylem and phloem fluxes, in the laboratory.




Materials and Methods

- 31 branches were cut, representing 6 tree species and 2 functional types.
- Branches were measured using the NMR-MOUSE immediately after sampling, and subsequently measured again, over time, to follow their dehydration dynamic.
- The water content of each branch was calculated by taking its fresh weight before each NMR measurement, and dry weight at the end of all experiments.
- Three models were constructed to test for linear correlations between water content and the integral of the raw NMR profiles.
- Branches were imaged using x-ray microtomography in order to locate anatomical structures and to correlate them with water peaks on normalized NMR profiles.





Results

There was a linear correlation between the integral of the raw NMR profiles obtained and the water content of the branches. This correlation was present regardless of tree species or functional type:



The normalized NMR profile of a fir branch (*A. alba*) presenting distinct peaks which corresponded to the xylem and phloem tissues, whose location was validated with x-ray microtomography imaging:



Conclusions and Perspectives

- We were able to observe a linear correlation between the NMR signal and the water content of branches, as well as locate conductive tissues.
- The NMR-MOUSE is a promising candidate for measuring plant water dynamics in the field. Future work will test the capacity of this device to measure tree water content *in-situ*, and to measure the speed of both the xylem and phloem fluxes.