



**HAL**  
open science

## **Integrating TGF-beta signalling pathways and epigenetic modulations for controlling sex determination in medaka**

Alexandra Depince, Florent Murat, Mateus Contar Adolfi, Violette Thermes,  
Yann Guiguen, Manfred Scharl, Amaury Herpin

### ► **To cite this version:**

Alexandra Depince, Florent Murat, Mateus Contar Adolfi, Violette Thermes, Yann Guiguen, et al.. Integrating TGF-beta signalling pathways and epigenetic modulations for controlling sex determination in medaka. 2. European Symposium on Sex Determination in Vertebrates, Oct 2022, Giens, France. hal-03833209

**HAL Id: hal-03833209**

**<https://hal.inrae.fr/hal-03833209>**

Submitted on 28 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Integrating TGF- $\beta$ signalling pathways and epigenetic modulations for controlling sex determination in medaka

Alexandra Depincé<sup>1</sup>, Florent Murat<sup>1</sup>, Mateus Contar Adolfi<sup>2</sup>, Violette Thermes<sup>1</sup>, Yann Guiguen<sup>1</sup>, Manfred Schartl<sup>2</sup> and Amaury Herpin<sup>1</sup>

[amaury.herpin@inra.fr](mailto:amaury.herpin@inra.fr)

<sup>1</sup> INRAE, UR1037 Fish Physiology and Genomics, Rennes F-35000, France.

<sup>2</sup> Department of Physiological Chemistry, University of Würzburg, D-97074 Würzburg, Germany.

In stark contrast to birds and mammals, which feature extreme conservative sex-determination mechanisms across species, fish display an amazing broad range of systems and master sex-determining (SD) genes. Thus, the molecular pathways of sex-determination and differentiation must have recurrently and independently adjusted and adapted during the course of evolution. Interestingly, although being clearly subordinates of the sex-regulatory network in mammals, TGF- $\beta$  family members (*Amh*, *Gsdf*, *Gdf6* or *Bmpr1bb*, *Amhr2*...) have nevertheless been the source of most of the SD genes in fish.

Facing this rolling wave of TGF- $\beta$  signaling in primary SD, and taking advantage of this “evolution in action”, our aim is to decipher the key role of those TGF- $\beta$  signaling pathways and their functional evolution for sex determination and maintenance of gonad sexual identity: -How is signaling specificity achieved among the different gonadal TGF- $\beta$  signal transducing factors? -What are the underlying molecular mechanisms determining the sex functions of these factors? and -How do they branch into the canonical downstream sex regulatory network?

Focusing on medaka *Gdf6* and *Amh* for approaching these conceptual questions, and using *in-vivo* and *in-vitro* approaches, combining gene editing together with transcriptomics and functional studies, we find that, in a context of sex determination:

- *Amh* and *Gdf6* likely act through different signaling pathways and are, in a physiological manner, distinctly integrated.
- *Amh* and *Gdf6* pathways do not necessarily branch together with the canonical sex-determining gene regulatory network for exerting their respective functions.
- Medaka *Gdf6* paralogs control variations in the methylation landscape, leading to epigenetic modifications strictly regulating sex determination.

Finally, evidence is provided for an unexpected link between (i) a growth factor, (ii) transcription factors and (iii) the modulation of the methylation landscape in the context of sex determination.