



**HAL**  
open science

## Assessing the state of marine biodiversity in the Northeast Atlantic

Abigail McQuatters-Gollop, Laurent Guérin, Nina Larissa Arroyo, Anais Aubert, Luis Felipe Artigas, Jacob Bedford, Emily Corcoran, Volker Dierschke, Sophie Elliott, Steve C V Geelhoed, et al.

► **To cite this version:**

Abigail McQuatters-Gollop, Laurent Guérin, Nina Larissa Arroyo, Anais Aubert, Luis Felipe Artigas, et al.. Assessing the state of marine biodiversity in the Northeast Atlantic. *Ecological Indicators*, 2022, 141, pp.109148. 10.1016/j.ecolind.2022.109148 . hal-03837035

**HAL Id: hal-03837035**

**<https://hal.inrae.fr/hal-03837035>**

Submitted on 4 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Assessing the state of marine biodiversity in the Northeast Atlantic

A. McQuatters-Gollop<sup>a,\*</sup>, L. Guérin<sup>b,c</sup>, N.L. Arroyo<sup>d</sup>, A. Aubert<sup>c,e</sup>, L.F. Artigas<sup>e</sup>, J. Bedford<sup>a,f</sup>, E. Corcoran<sup>g,h</sup>, V. Dierschke<sup>i</sup>, S.A.M. Elliott<sup>b,j</sup>, S.C.V. Geelhoed<sup>k</sup>, A. Gilles<sup>l</sup>, J.M. González-Irusta<sup>d</sup>, J. Haelters<sup>m</sup>, M. Johansen<sup>n</sup>, F. Le Loc'h<sup>o</sup>, C.P. Lynam<sup>p</sup>, N. Niquil<sup>q</sup>, B. Meakins<sup>r</sup>, I. Mitchell<sup>r</sup>, B. Padegimas<sup>g</sup>, R. Pesch<sup>s,t</sup>, I. Preciado<sup>d</sup>, I. Rombouts<sup>c,e</sup>, G. Safi<sup>q</sup>, P. Schmitt<sup>s</sup>, U. Schückel<sup>u</sup>, A. Serrano<sup>d</sup>, P. Stebbing<sup>v</sup>, A. De la Torre<sup>d</sup>, C. Vina-Herbon<sup>r</sup>

<sup>a</sup> Marine Conservation Research Group, School of Biological and Marine Science, University of Plymouth, Drake Circus, Plymouth PL4 8AA, United Kingdom

<sup>b</sup> Muséum National d'Histoire Naturelle, Service des Stations Marines. Station Marine de Dinard. 38 rue du Port Blanc, Dinard, Brittany 35800, France

<sup>c</sup> Patrimoine Naturel Joint Unit (OFB, MNHN, CNRS), Station Marine de Dinard, 38 Rue du Port Blanc, Dinard, Brittany 35800, France

<sup>d</sup> Instituto Español de Oceanografía (CNIEO-CSIC), Centro Oceanográfico de Santander, Promontorio San Martín, s/n, 39080 Santander, Spain

<sup>e</sup> Laboratoire d'Océanologie et de Géosciences (LOG), UMR 8187 CNRS, Université du Littoral Côte d'Opale, Université de Lille, 62930 Wimereux, France

<sup>f</sup> UN Environment Programme World Conservation Monitoring Centre, 219 Huntingdon Road, Cambridge CB3 0DL, United Kingdom

<sup>g</sup> OSPAR Secretariat, 63 Southampton Row, London WC1B 4DA, United Kingdom

<sup>h</sup> Independent Consultant, Sövdevägen 220-5, Blentarps 275 64, Sweden

<sup>i</sup> Gavia EcoResearch, Tönnhäuser Dorfstr. 20, 21423 Wüsen (Luhe), Germany

<sup>j</sup> DECOD (Ecosystem Dynamics and Sustainability), Institut Agro, Ifremer, INRAE, Rennes, France

<sup>k</sup> Wageningen Marine Research, NL-1781 AG Den Helder, The Netherlands

<sup>l</sup> Institute for Terrestrial and Aquatic Wildlife Research, University of Veterinary Medicine Hannover, Foundation, 25761 Bisum, Germany

<sup>m</sup> Royal Belgian Institute of Natural Sciences (RBINS), Operational Directorate Nature, B-8400 Ostend, Belgium

<sup>n</sup> Swedish Meteorological and hydrological institute, Sven Källfelts gata 31, 42671 Gothenburg, Sweden

<sup>o</sup> Institut de Recherche pour le Développement (IRD), University of Brest, CNRS, Ifremer, LEMAR, Plouzané, France

<sup>p</sup> Centre for Environment, Fisheries and Aquaculture Science, Lowestoft Laboratory, Lowestoft, Suffolk NR33 0HT, United Kingdom

<sup>q</sup> University of Caen Normandie, UMR 8067 BOREA MNHN, Sorbonne University, UCN, CNRS, IRD, UA. Esplanade de la Paix, CS 14032, 14032 Caen, France

<sup>r</sup> Joint Nature Conservation Committee, Monkstone House, City Road, Peterborough PE1 1JY, United Kingdom

<sup>s</sup> BioConsult GmbH & Co. KG, Auf der Muggenburg 30, 28217 Bremen, Germany

<sup>t</sup> Institute for Applied Photogrammetry and Geoinformatics, Jade University of Applied Sciences, Ofener Str. 16, 26121 Oldenburg, Germany

<sup>u</sup> Schleswig-Holstein Agency for Coastal Defence, National Park and Marine Conservation, National Park Authority, 25832 Tönning, Germany

<sup>v</sup> APEM Limited, International House, International Business Park, Southampton SO18 2RZ, United Kingdom

### ARTICLE INFO

#### Keywords:

Food webs  
Indicators  
Holistic assessment  
Ecosystem approach  
Marine Strategy Framework Directive  
Anthropogenic pressures

### ABSTRACT

The Northeast Atlantic, a highly productive maritime area, has been exposed to a wide range of direct human pressures, such as fishing, shipping, coastal development, pollution, and non-indigenous species (NIS) introductions, in addition to anthropogenically-driven global climate change. Nonetheless, this regional sea supports a high diversity of species and habitats, whose functioning provides a variety of ecosystem services, essential for human welfare. In 2017, OSPAR, the Northeast Atlantic Regional Seas Commission, delivered an assessment of marine biodiversity for the Northeast Atlantic. This assessment examined biodiversity indicators separately to identify changes in Northeast Atlantic biodiversity, but stopped short of determining the status of biodiversity for many species and habitats. Here, we expand on this work and for the first time, a semi-quantitative approach is applied to evaluate holistically the state of Northeast Atlantic marine biodiversity across marine food webs, from plankton to top predators, via fish, pelagic and benthic habitats, including xenobiodiversity (i.e. NIS). Our analysis reveals widespread degradation in marine ecosystems and biodiversity, particularly for marine birds and coastal bottlenose dolphins, as well as for benthic habitats and fish in some regions. The poor biodiversity status of these ecosystem components is likely the result of cumulative effects of human activities, such as habitat destruction or disturbance, overexploitation, eutrophication, the introduction of NIS, and climate change. Bright spots are also revealed, such as recent signs of recovery in some fish and marine bird communities and recovery in harbour and grey seal populations and the condition of coastal benthic

\* Corresponding author.

E-mail address: [Abigail.mcquatters-gollop@plymouth.ac.uk](mailto:Abigail.mcquatters-gollop@plymouth.ac.uk) (A. McQuatters-Gollop).

<https://doi.org/10.1016/j.ecolind.2022.109148>

Received 6 March 2022; Received in revised form 30 June 2022; Accepted 3 July 2022

Available online 12 July 2022

1470-160X/© 2022 Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

communities in some regions. The status of many indicators across all ecosystem components, but particularly for the novel pelagic habitats, food webs and NIS indicators, however, remains uncertain due to gaps in data, unclear pressure-state relationships, and the non-linear influence of some pressures on biodiversity indicators. Improving monitoring and data access and increasing understanding of pressure-state relationships, including those that are non-linear, is therefore a priority for enabling future assessments, as is consistent and stable resourcing for expert involvement.

## 1. Introduction

In 2019 the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Service (IPBES) declared a global biodiversity emergency recognising the degraded, and in many cases worsening, state of global biodiversity, including that of the marine environment (IPBES, 2019). Climate change, ocean acidification, and direct anthropogenic pressures such as overfishing, pollution, non-indigenous species (NIS), and intensive coastal development are affecting much of the global ocean, resulting in declines in, and even loss of, marine species and habitats (Halpern et al., 2008; IPBES, 2019; IPCC, 2019; Secretariat of the Convention on Biological Diversity, 2014). Such degradation affects ecosystem structure, processes, and functioning, consequently affecting essential marine ecosystem services including food provision, carbon and nutrient cycling, coastal protection, and recreation (IPBES, 2019). Global (e.g., Convention on Biological Diversity, Sustainable Development Goal 14) and regional (e.g., EU Marine Strategy Framework Directive, Northeast United States Shelf (NOAA, 2022), European Regional Seas Conventions (i.e. OSPAR's Northeast Atlantic Biodiversity Strategy (OSPAR, 2021)) policy mechanisms recognise this critical link between healthy marine biodiversity and human welfare, and have instated goals to assess, maintain, and improve marine biodiversity. A clear picture of the current state and past trends of marine biodiversity (including xeno-biodiversity (i.e. NIS) is therefore required as a baseline from which to both detect ongoing and future changes, design effective management measures and track progress towards policy targets. Furthermore, establishing the state of marine biodiversity at ecologically-relevant spatial and temporal scales may help to measure regional effects of global climate change effects and distinguish those from localised pressures.

The Northeast Atlantic has a long history of intensive human use (Roberts, 2007), which along with climate change, has caused large-scale degradation of marine ecosystems and biodiversity from historic baselines. During the last century Northeast Atlantic marine ecosystems have experienced changes in ecosystem functioning (Capuzzo et al., 2018), alterations to marine food webs (Piroddi et al., 2021), the loss and extinction of species (Lotze et al., 2017; Reynolds et al., 2009; Sguotti et al., 2016), colonisation by NIS (Castro et al., 2017; Tsiamis et al., 2019), and coastal and marine habitat degradation and loss (Airoldi and Beck, 2007; Bennema et al., 2020; Torbjørn et al., 2003). To halt the decline of marine biodiversity and improve its state, the European Union passed into force the Marine Strategy Framework Directive (MSFD) in 2008. The MSFD applies a holistic, ecosystem approach to managing the marine environment with key descriptors of marine biodiversity (e.g. ecosystem functioning, NIS, food webs, and sea floor integrity) explicitly named as some aspects of the marine environment which must reach the objective of Good Environmental Status (GES). The MSFD represents one of the world's most ambitious applications of a comprehensive ecosystem approach, looking across multiple biodiversity components (i.e. pelagic and benthic habitats, fish, marine birds, marine mammals, food webs, and NIS) based on a variety of indicators. As part of this process, 22 indicators have been developed to assess the state of marine biodiversity at the Northeast Atlantic regional scale (OSPAR, 2017m). These biodiversity indicators are assessed to evaluate progress towards targets representing the overall objective of GES for each ecosystem component. In 2017, OSPAR (the Northeast Atlantic regional seas commission) evaluated each indicator individually as part

of a Northeast Atlantic ecosystem assessment (OSPAR, 2017m). In many cases, the assessment was challenging due to the scale of assessment, methodological or data limitations, or due to the novel approach of assessing biodiversity across datasets, ecosystems, and multiple scientific experts from OSPAR Contracting Parties at this scale. It was often equally challenging to identify pressures causing changes in the marine environment, either because of scientific uncertainty or due to resource and time limitations. In most cases (e.g. pelagic habitats, food webs, NIS) no quantitative targets for GES were agreed at the OSPAR level at the time, so no overall assessment of the state of marine biodiversity was made during IA2017. In other words, in many cases the indicators were not formally assessed for Good Environmental Status, but biodiversity change was merely described (OSPAR, 2017m).

This paper aims, therefore, for the first time, to assess the state of marine biodiversity across multiple ecosystem components at the Northeast Atlantic regional scale, by using a suite of policy-relevant indicators. Biodiversity status is assessed semi-quantitatively by linking indicator change to pressures or evaluating against threshold values (where available) to reveal a broadscale assessment of Northeast Atlantic marine biodiversity state. Challenges to this process are then identified and recommendations for achieving a more complete and holistic assessment of regional marine biodiversity are given.

## 2. Methods

### 2.1. Spatial area

The Northeast Atlantic OSPAR maritime area is bordered by 12 countries, ten of which were European Union Member States at the time of this work. Countries bordering the Northeast Atlantic have a total population of approximately 329 million people. Consequently, Northeast Atlantic marine waters support a wide variety of human activities including fishing, shipping, oil and gas provision, and recreation (ICES, 2020).

Within OSPAR, the Northeast Atlantic maritime area can be divided into five regions (Arctic Waters, Greater North Sea, Celtic Seas, Bay of Biscay and Iberian Coast, and Wider Atlantic; Fig. 1). These regions comprise a wide range of environmental conditions and marine ecosystems, from deep ocean to shelf and coastal waters. Where ecologically relevant or where data were limited, specific indicator assessments have used further geographic divisions, referred to here as sub-regions (e.g., the English Channel, the Southern North Sea). Here we focus on the Celtic Seas, Greater North Sea, and Bay of Biscay and Iberian Coast regions, where most of the recently developed indicators were adopted and applied by OSPAR Contracting Parties. Marine birds were also assessed for Arctic Waters, and one fish and one food web indicator for the Wider Atlantic, in 2017, but are outside the focus of this paper.

### 2.2. Science-policy process

Through indicator assessments, the 2017 OSPAR Intermediate Assessment (IA2017) further developed our understanding of the marine environment of the Northeast Atlantic and its current ecological status. IA2017 also provided an opportunity for OSPAR Contracting Parties, most of which are also European Union Member States, to use the information and methodology in their reporting obligations under the MSFD.



**Fig. 1.** The OSPAR Maritime Area is divided into five regions for assessment purposes: Arctic Waters, Greater North Sea, Celtic Seas, Bay of Biscay and Iberian Coast, and Wider Atlantic. Source: OSPAR website (<https://oap.ospar.org/en/ospar-assess/>).

The IA2017 biodiversity assessments were carried out through OSPAR's Intersessional Correspondence Group on Biodiversity and Monitoring (ICG-COBAM). Due to the science-policy nature of OSPAR, IA2017 enabled indicators to be co-developed with policy-makers to meet policy needs. The process was organised according to seven ecosystem components (pelagic habitats, benthic habitats, fish, marine birds, marine mammals, food webs, and NIS), each of which was represented by an expert group consisting of scientific experts from OSPAR Contracting Party countries. The expert groups drove scientific indicator development and assessment, engaging in the wider OSPAR process to ensure policy alignment. Science-policy interaction primarily occurred at workshops where scientists and policy-makers worked together to review indicator advances and subsequently draft assessments. This collaborative process promoted scientific progress on indicator development with continual feedback and steer from policy-makers throughout the OSPAR area. For food webs and pelagic and benthic habitats, collaborative work was also progressed under the EU-funded "Applying an Ecosystem Approach to (sub) Regional Habitat Assessments: Addressing gaps in biodiversity indicator development for the OSPAR Region from data to ecosystem assessment" (EcAprHA) project, which enabled the acceleration of indicator development. This process produced indicators and assessments that are scientifically robust; can be communicated appropriately to scientists, policy makers, and the public; and are useful to policy-makers for supporting contributions to national MSFD assessments. The marine biodiversity portion of the IA2017 process mobilised considerable scientific effort, with approximately 200 scientists and 40 policy-makers involved (<https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/acknowledgements/>).

### 2.3. Biodiversity indicators: Background context and key characteristics

Biodiversity indicators were developed to represent key aspects of the seven OSPAR/MSFD ecosystem components (Tables 1–7). The biodiversity indicator development process started in 2010 with >140 theoretical, published, or national indicators identified as relevant for biodiversity assessment in the Northeast Atlantic (OSPAR, 2011). A combination of policy relevance and scientific robustness was used to create a shortlist of indicators, some of which had previously been used

in assessments and some of which were purely conceptual. The common methodological principles and approaches to determining GES (for OSPAR or MSFD), setting environmental targets and selecting indicators were published as an advice manual (OSPAR, 2012). Additional scrutiny was given to the indicator suite selected by ICES working groups (e.g. ICES, 2013; ICES, 2014b) and in the scientific literature where many new indicators and assessment methods were developed and published during this decade, notably influenced by the progressive implementation of the MSFD in European seas (Borja et al., 2019; Heiskanen et al., 2016; Uusitalo et al., 2016). Expert groups developed these indicators into operational indicators, through a process of testing, validation, and science and policy peer-review. For IA2017, each indicator was assessed individually, from relevant data made available by Contracting Parties, representing a significant milestone in describing the ecological status of Northeast Atlantic biodiversity across multiple ecosystem components. Each indicator was given a short code (PH = Pelagic Habitats, BH = Benthic Habitats, FC = Fish and Cephalopods (although cephalopod data were not used here), B = (Marine) Birds, M = Mammals, FW = Food Webs, NIS = Non-Indigenous Species) and identifying number (Tables 1–7). Some indicators were not operational, nor agreed by OSPAR Contracting Parties, at the time of IA2017, often as a result of lack of development or testing, and were designated as "candidate" indicators that might be used for future assessments (Elliott et al., 2017b; Padegimas et al., 2017). The entire process was performed in collaboration with policy-makers, to ensure policy-relevance, and with (inter) national expert networks to ensure scientific consensus via peer-review.

A variety of indicator types was developed to assess marine biodiversity status. The basic DPSIR (Driver-Pressure-State-Impact-Response) framework can be used to classify indicator types for the purpose of this work (Elliott et al., 2017a). In brief, Drivers of basic human needs lead to Pressures on the marine ecosystem. The Pressures cause State change on the natural system which then leads to Impacts on human welfare. These impacts then require management Responses. The majority of the biodiversity indicators here assess ecological state, although some assess the effect of pressure on state, combining data on extent and intensity of pressures with biological data (e.g. The common conceptual approach to assessing condition of benthic habitat-defining communities (BH2), Extent of physical damage to predominant special habitats (BH3), Harbour porpoise bycatch (M6); Tables 2, 5). Some indicators are comprised of multiple sub indicators, depending on the pressure or ecological information to be assessed (e.g. Changes in plankton diversity (PH3) and Condition of benthic habitat-defining communities (BH2) (Tables 1, 2)). Food web indicators are central to this analysis and represent an important step towards holistic assessment. For that reason, some food web indicators span multiple ecosystem components (e.g., Changes in phytoplankton and zooplankton communities (PH1/FW5) is both a pelagic and a food web indicator; Production of phytoplankton (FW2) encompasses primary productivity, an aspect of pelagic habitats; Size composition in fish communities (FW3) is related to fish community size structure, Change in mean trophic level of consumers or marine predators (FW4) includes invertebrate and fish data; Table 6). A brief synopsis of each indicator, including its underlying data, data analysis methods, time span, spatial coverage of data, link to detailed methodological guidelines under the OSPAR Coordinated Environmental Monitoring Programme (CEMP), and further references are provided in Tables 1–7.

### 2.4. Data

Data were gathered through OSPAR formal data calls, requests to ICES Working Groups and ICES data calls, national expert group contributions, and direct requests to data providers, many of whom were members of the OSPAR Expert Groups. Spatial and temporal data coverage varied by dataset and indicator due to differences in data availability; for this reason, and because ecological dynamics vary throughout the ecosystem, a common temporal baseline was not used

**Table 1**

The pelagic habitats ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region(s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Pelagic habitats (plankton)	PH1/FW5	Changes in phytoplankton and zooplankton communities	Zooplankton and phytoplankton taxa abundance time-series	2004–2014	Greater North Sea  Celtic Seas  Bay of Biscay & Iberian Coast	Fixed point stations in Greater North Sea and Celtic Sea. CPR Data aggregated by ecohydrodynamic zones (as per Van Leeuwen et al. (2015)) throughout Celtic Seas and Greater North Sea regions. Bay of Biscay treated as one region.	Plankton Index performed on plankton lifeform pair time-series	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/</a> (OSPAR, 2017t) CEMP: <a href="https://www.ospar.org/documents?v=39001">https://www.ospar.org/documents?v=39001</a> (OSPAR, 2018h)  Further references: McQuatters-Gollop et al. (2019a); Bedford et al. (2020a); Bedford et al. (2020b)
	PH2	Changes in phytoplankton biomass and zooplankton abundance	Proxies of total phytoplankton biomass and total copepod abundance time-series	1958–2013	Greater North Sea  Celtic Seas  Bay of Biscay & Iberian Coast	Fixed point stations in Greater North Sea. CPR Data aggregated by ecohydrodynamic zones (as per Van Leeuwen et al. (2015)) throughout Greater North Sea, and separately for the Bay of Biscay and the Celtic Sea.	Time-series anomalies	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/</a> (OSPAR, 2017u)  CEMP: <a href="https://www.ospar.org/documents/d=40972">https://www.ospar.org/documents/d=40972</a> (OSPAR, 2019a)
	PH3	Changes in plankton diversity	Phytoplankton taxa abundance time-series	1987–2014	Greater North Sea (coastal)  Celtic Seas (coastal)  Bay of Biscay & Iberian Coast (coastal)	Data from 4 stations near the French coast	Species richness (Menhnick Index) Species dominance (Hulbert Index) Local Contribution to Beta Diversity (LCBD)	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/pilot-assessment-changes-plankton/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/pilot-assessment-changes-plankton/</a> (OSPAR, 2017v)  CEMP: <a href="https://www.ospar.org/documents/d=40973">https://www.ospar.org/documents/d=40973</a> (OSPAR, 2019b)  Further references: Rombouts et al. (2019), Bedford et al. (2020c)

across indicators and ecosystem components. For example, broad-scale benthic habitat, cetacean abundance, and plankton abundance and biomass data were used throughout several regions, while some other datasets, such as benthic community condition, harbour porpoise bycatch data or changes in plankton diversity, were available only sub-regionally. For NIS, data were available on a national scale and then processed to meet the regional scale. Approximately 86 datasets were used to develop the indicators, but in some cases (e.g. pelagic habitats, benthic habitats) lack of formal data contribution by Contracting Parties restricted the inclusion of all possible datasets.

A wide spectrum of data types was used to construct indicators. The harmonisation of diverse datasets was often required (e.g. to construct pelagic and benthic habitats and food web indicators). Data and

metadata can be obtained from each indicator's IA2017 webpage (see Tables 1–7) and associated Coordinated Environmental Monitoring Programme (CEMP) guidelines, which contain official and detailed methodologies for all indicators contributing to IA2017 (see Tables 1–7; further information in CEMP at <https://www.ospar.org/work-areas/cross-cutting-issues/cemp>).

#### 2.4.1. Pelagic habitat data

Two types of plankton time-series data were used to construct pelagic habitat indicators: fixed-point sampling data from the UK, France, and Sweden, and underway data from the UK-based Continuous Plankton Recorder (CPR) survey. Differences in sampling and analysis methods exist between all time-series datasets. For example, fixed-point

**Table 2**

The benthic habitats ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region(s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Benthic habitats	BH2	The common conceptual approach to assessing condition of benthic habitat-defining communities	Benthic communities species relative abundances and pressure extent and intensity	N/A	Not assessed per se	Not assessed per se	Common conceptual approach, assessment scales, and parameters	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/common-conceptual-approach/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/common-conceptual-approach/</a> (OSPAR, 2017c)  CEMP: <a href="https://www.ospar.org/documents?v=39000">https://www.ospar.org/documents?v=39000</a> (OSPAR, 2018b)
	BH2-A	Assessment of coastal habitats in relation to nutrient and/or organic enrichment	Benthic macro fauna and macro algae and angiosperms communities' species and relative abundances	2003–2015	Greater North Sea (coastal) Celtic Seas (coastal) Bay of Biscay & Iberian Coast (coastal)	WFD stations (coastal) throughout regions	Multi-metric indices (notably reported for Water Framework Directive (WFD))	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/condition-benthic-habitat-communities-assessment-coastal-habitat/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/condition-benthic-habitat-communities-assessment-coastal-habitat/</a> (OSPAR, 2017d)  CEMP: <a href="https://www.ospar.org/documents?v=39000">https://www.ospar.org/documents?v=39000</a> (OSPAR, 2018b)
	BH2-B	Subtidal habitats of the southern North Sea	Benthic macro fauna communities' species relative abundances and abrasion pressure extent and intensity.	2009–2013	Southern part of the Greater North Sea	Data throughout this sub-region	Multi-metric indices	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/subtidal-habitats-southern-north-sea/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/subtidal-habitats-southern-north-sea/</a> (OSPAR, 2017e)  CEMP: <a href="https://www.ospar.org/documents?v=39000">https://www.ospar.org/documents?v=39000</a> (OSPAR, 2018b)  Further references: Van Loon et al. (2018)
	BH3	Extent of physical damage to predominant special habitats	Habitat maps and sensitivities matrices and abrasion pressure extent and intensity. Physical pressures currently limited to bottom-trawling fisheries surface and sub-surface abrasion.	2010–2015	Greater North Sea Celtic Seas Bay of Biscay & Iberian Coast (coastal)	Data throughout regions	Spatial analysis of habitat distribution and sensitivity, versus pressure extent and intensity	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/extent-physical-damage-predominant-and-special-habitats/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/extent-physical-damage-predominant-and-special-habitats/</a> (OSPAR, 2017f)  CEMP: <a href="https://www.ospar.org/documents?v=37641">https://www.ospar.org/documents?v=37641</a> (OSPAR, 2017s)

phytoplankton data were collected with a variety of methods, while fixed-point zooplankton data were collected using nets. The CPR survey is a phyto- and zooplankton monitoring programme which samples plankton using a device towed behind merchant navy ships on commercial routes spanning the Northeast Atlantic (Batten et al., 2003). As a result of these differences, pelagic data could not be simply aggregated across datasets. Instead, pelagic indicators were purposely developed to deal with this wide range of plankton data types, regardless of sampling and enumeration method or level of taxonomic identification (see Table 1). Indicator Changes in phytoplankton and zooplankton

communities (PH1/FW5) aggregates plankton taxa into “lifeforms” based on common functional traits. Changes in lifeforms can be interrogated temporally, spatially, or as ecologically-relevant lifeform pairs (McQuatters-Gollop et al., 2019a).

#### 2.4.2. Benthic habitat data

The data requirements for all OSPAR benthic indicators can be classified into three main categories: a) benthic species' community composition; b) benthic habitat mapping data; and c) spatial data of anthropogenic pressures. These data streams support the assessment of

**Table 3**

The fish ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region(s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Fish	FC1	Recovery in the population abundance of sensitive fish species	Time series of abundance-based catch rates for selected species assessed relative to historic quantiles of abundance (25th and 75th) aggregated into a single indicator and representative of change in sensitive species overall	1983–2016  Specific time coverage dependent on survey	Greater North Sea  Celtic Seas  Bay of Biscay & Iberian Coast	Data throughout the regions	Time series with probabilistic integration within surveys and within regions	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/recovery-sensitive-fish/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/recovery-sensitive-fish/</a> (OSPAR, 2017g)  CEMP: <a href="https://www.ospar.org/documents?v=38999">https://www.ospar.org/documents?v=38999</a> (OSPAR, 2018a)
	FC2	Proportion of large fish – Large Fish Index	Time series of biomass by size class of fish aggregated into a single indicator representative of change in the proportion of large individuals in the community	1983–2016  Specific time coverage dependent on survey	Greater North Sea  Celtic Seas	Data throughout the regions	Time series of Large Fish Index (LFI) by survey	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/proportion-large-fish-large-fish-index/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/proportion-large-fish-large-fish-index/</a> (OSPAR, 2017h)  CEMP: <a href="https://www.ospar.org/documents?v=38999">https://www.ospar.org/documents?v=38999</a> (OSPAR, 2018a)
	FC3	Mean Maximum Length of fish	Time series of biomass by species aggregated into a single indicator where species are weighted by their ultimate (maximum) body size. Indicator representative of species composition.	1983–2016  Specific time coverage dependent on survey	Greater North Sea  Celtic Seas  Bay of Biscay & Iberian Coast	Data throughout the regions	Time series of Mean Maximum Length (MML) by spatial strata and survey	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/mean-maximum-length/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/mean-maximum-length/</a> (OSPAR, 2017i)  CEMP: <a href="https://www.ospar.org/documents?v=38999">https://www.ospar.org/documents?v=38999</a> (OSPAR, 2018a)

each indicator and in combination provided for an integrated assessment (Table 2). Data on benthic macro-fauna and macro-flora species and communities were collected via standardised methods (See The common conceptual approach to assessing condition of benthic habitat-defining communities (BH2) CEMP, Table 2 (OSPAR, 2018b)). Benthic habitat mapping data provide information on seabed type, classified according to EUNIS (European Nature Information System) levels, and distribution. Benthic habitat maps were produced using EMODNet methods and data (EUNIS level 3), combined with local datasets, at different spatial resolutions, depending on methods used (e.g. ground-truthed sampling, acoustic mapping, and modelling) (Diaz et al., 2004; Ellwood, 2014; Kenny et al., 2003). The information on environmental variables (e.g. depth, exposure, sediment type, etc) was used to classify and model habitats and produce habitat maps (Brown et al., 2011). Information on the extent and distribution of benthic habitats is used alongside the associated sensitivity, in terms of resilience and resistance capacity of benthic species, biotopes and habitat types. Data for surface and sub-surface abrasion by bottom-trawling fisheries provided an indirect measure of pressure on the seabed by fishing. These data were obtained from ICES (2016b). The pressure layer on abrasion by bottom trawling was produced from bottom trawl data provided by OSPAR Contracting Parties, the assessment of related pressure by ICES, and finally an assessment of disturbance of habitats by the OSPAR benthic habitat expert group.

#### 2.4.3. Fish data

Quantitative spatio-temporal data across the OSPAR area were available from scientific trawl surveys coordinated by ICES and supporting the fisheries management process. These surveys collect data on the number and weight of each species of fish along with measurements of body length and mass. To maintain transparency and for quality assurance, a selection of surveys was used for the analyses. The data resulting from the surveys were made available through the routinely updated Database of Trawl Surveys (DATRAS). Relevant ICES Working groups (ICES, 2021a; ICES, 2021b) were consulted on the appropriate usage of these data and a single data product was subsequently created and, following discussion within ICG-COBAM, stored in an online repository to support the assessment process with comprehensive documentation available (Greenstreet and Moriarty, 2017; Moriarty et al., 2017; Moriarty et al., 2019). Cephalopods and other data collected through these surveys (e.g. benthic invertebrates) were excluded from the data product due to inconsistencies in species identification and recording practises between countries.

#### 2.4.4. Marine bird data

The assessment used data on marine bird species that, at some point in their annual life cycle, rely on coastal and/or offshore marine areas; the term ‘marine birds’ therefore includes seabirds and waterbirds. Taxonomic groups considered include Anseriformes (ducks, geese, swans), Procellariiformes (fulmars, petrels, shearwaters), Suliformes (gannets, cormorants), Gaviiformes (divers), Podicipediformes (grebes)

**Table 4**

The marine birds ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region (s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Marine birds	B1-A	Marine birds: non-breeding abundance	Marine birds: non-breeding abundance	1991–2014/15	Greater North Sea Celtic Seas	Data from individual migration and wintering sites; or aggregated data for large stretches of coastline.	Trends in non-breeding abundance	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/bird-abundance/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/bird-abundance/</a> (OSPAR, 2017a)  CEMP: <a href="https://www.ospar.org/documents?v=38978">https://www.ospar.org/documents?v=38978</a> (OSPAR, 2016a)
	B1-B	Marine birds: breeding abundance	Marine birds: breeding abundance	1991–2014/15	Greater North Sea Celtic Seas	Data from individual breeding colonies; or aggregated data for large stretches of coastline.	Trends in breeding abundance	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/bird-abundance/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/bird-abundance/</a> (OSPAR, 2017a)  CEMP: <a href="https://www.ospar.org/documents?v=38978">https://www.ospar.org/documents?v=38978</a> (OSPAR, 2016a)
	B3	Marine bird breeding success/failure	Annual colony failure rate of each species	1992–2015	Greater North Sea Celtic Seas	Data from individual breeding colonies	Number of years in which 'widespread colony failure' occurs (percentage of colonies failing to produce >0.1 chick per pair per year)	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/marine-bird-breeding-success-failure/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/marine-bird-breeding-success-failure/</a> (OSPAR, 2017b)  CEMP: <a href="https://www.ospar.org/documents?v=38979">https://www.ospar.org/documents?v=38979</a> (OSPAR, 2016b)

and Charadriiformes (shorebirds, skuas, gulls, terns, auks). Data were available from national monitoring schemes, which each country submitted to the OSPAR Seabird Database via the ICES Biodiversity Data Portal (<https://www.ices.dk/data/data-portals/Pages/Biodiversity.aspx>). Breeding abundance was assessed using trends in annual estimates of breeding bird abundance derived from counts of adult birds or pairs when they aggregate on land at coastal breeding colonies or sites, where they are easier to monitor than when being dispersed at sea over large areas for foraging. Some colonially breeding species were also assessed using data on breeding success – the average number of chicks fledged per breeding pair per year. Non-breeding bird abundance did not derive from at-sea surveys, but was restricted to counts of birds in intertidal areas or close to the shore and surveyed from land during migration or over the winter. During these times the birds in these habitats aggregate in large numbers and are much easier to count than during the breeding season, when they are dispersed over large areas of remote nesting habitat, such as the Arctic tundra. Many marine bird species assessed in the non-breeding season (mainly shorebirds, ducks, geese, swans, divers, and grebes) breed inland and in areas outside the OSPAR Maritime Area but are reliant on the Northeast Atlantic most of the year during migration and/or over the winter. Indicator Marine bird breeding success/failure (B3) was populated with seabird data for IA2017, although it is anticipated that waterbird data will be included in future assessments.

#### 2.4.5. Marine mammal data

Abundance and distribution data for both cetaceans and phocid seals were available from dedicated national monitoring programmes as well as from international large-scale surveys such as SCANS (Small Cetacean Abundance in the European Atlantic and North Sea; [Hammond et al., 2013](#)). These surveys were timed for key periods, i.e. for grey and

harbour seal during key life stages such as breeding (pupping) and moulting, whereas the surveys for cetaceans concentrated on the summer months. The sampling methods included line-transect distance sampling as well as mark-recapture analyses of photo-identification data, and for the seals, aerial photographic surveys, and land- and ship-based counts ([Brasseur et al., 2015](#); [Buckland et al., 2001, 2004](#); [Galatius et al., 2021](#)). ICES Working groups ([ICES, 2014a](#)) were consulted on relevant population-specific assessment units as well as threshold setting methods for population trends and bycatch. Data on bycatch rates of harbour porpoises were primarily obtained through observer schemes that operated according to European Union Council Regulation (EC) 812/2004, and were analysed by the ICES Working Group on Bycatch of Protected Species (WGBYC). [ICES advise \(2015a\)](#) was used for assessment of this pressure indicator.

#### 2.4.6. Food web data

Food web indicators differ in their data requirements according to the complexity of the indicator and the number of ecosystem components considered. Different types of phytoplankton primary production data were used to construct food web indicators. For Production of phytoplankton (FW2), for example, the estimation of primary production comes from different data collection and processing methods such as carbon or oxygen isotope techniques, fluorometric techniques and modelling methods. Since the time-series were collected from fixed stations and transects, a step-by-step approach was developed to integrate these datasets for the indicator calculation ([Kromkamp et al., 2017](#)). The data requirements for the Change in mean trophic level of consumers or marine predators (FW4) food web indicator can be classified into two main categories: a) biomass data per species for fish and invertebrates from surveys and landings and b) data on the trophic level of each species. Trophic level values were estimated from stomach



**Table 5**

The marine mammals ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region(s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Marine mammals	M3	Seal abundance and distribution (M3)	Harbour seal and Atlantic grey seal abundance time series (on haul-outs)	1992–2014	Greater North Sea Celtic Seas	Data from individual haul-out sites (both species) in the Greater North Sea and the United Kingdom part of the Celtic Sea	Trends in abundance, occupancy rate, distributional shift	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/seal-abundance-and-distribution/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/seal-abundance-and-distribution/</a> (OSPAR, 2017n)  CEMP: <a href="https://www.ospar.org/documents?v=38980">https://www.ospar.org/documents?v=38980</a> (OSPAR, 2018d)
	M4-A	Abundance and distribution of cetaceans	Pilot assessment on abundance and distribution of killer whales	1999–2016	Greater North Sea Celtic Seas Bay of Biscay & Iberian coast	no assessment of abundance due to lack of historical information; only pilot assessment of distribution	Pilot assessment of distribution	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/abundance-distribution-cetaceans/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/abundance-distribution-cetaceans/</a> (OSPAR, 2017o)
	M4A-1		Abundance and distribution of coastal bottlenose dolphins	1994–2016	Greater North Sea Celtic Seas Bay of Biscay & Iberian coast	pilot assessment on five populations, indicative assessment provided for one other population	Trends in abundance and distribution	CEMP: <a href="https://www.ospar.org/documents?v=39019">https://www.ospar.org/documents?v=39019</a> (OSPAR, 2018e)
	M4-B		Abundance and distribution of cetaceans other than killer whales and coastal bottlenose dolphins	1994–2016	Greater North Sea Celtic Seas Bay of Biscay & Iberian coast	Data throughout regions; trends only possible for species with more than two (regional) abundance estimates	Trends in abundance and distribution	
	M5	Grey seal pup production	Atlantic grey seal pup abundance at breeding sites	1992–2014	Greater North Sea Celtic Seas (parts)	Data from breeding sites in the Greater North Sea and the United Kingdom part of the Celtic Sea	Trends in grey seal pup production	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/grey-seal-pup/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/grey-seal-pup/</a> (OSPAR, 2017p)  CEMP: <a href="https://www.ospar.org/documents?v=38981">https://www.ospar.org/documents?v=38981</a> (OSPAR, 2018f)
	M6	Harbour porpoise bycatch	Numbers of harbour porpoise bycaught in commercial nets	2006–2014	Greater North Sea Celtic Seas (parts)	Kattegat & Belt Seas, North Sea, Celtic and Irish Seas	Bycatch estimates from annual fishing effort estimates and counts of bycaught harbour porpoises (ICES advice)	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/harbour-porpoise-bycatch/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/harbour-porpoise-bycatch/</a> (OSPAR, 2017q)  ICES advice: (ICES, 2015b, 2016a)

**Table 6**

The food webs ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region (s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Food webs	FW2	Production of phytoplankton	Phytoplankton primary production time-series data	1988–2014 (different time-series)	Greater North Sea Celtic Seas	Pilot assessment of nine sites	Assessment of phytoplankton production measured using different methods	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/phytoplankton-production/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/phytoplankton-production/</a> (OSPAR, 2017j)  CEMP: not currently available, as this was a pilot assessment
	FW3	Size composition in fish communities	Time series of geometric mean length of fish (size measurements weight by biomass by size class) representative of change in the size structure of fish	1983–2016 Specific time coverage dependent on survey	Greater North Sea Celtic Sea Bay of Biscay & Iberian Coast	Data throughout the regions	Typical Length index (TyL) by spatial strata and survey	Further reference: Kromkamp (2017) IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/size-fish-composition/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/size-fish-composition/</a> (OSPAR, 2017k)  CEMP: <a href="https://www.ospar.org/documents?v=38999">https://www.ospar.org/documents?v=38999</a> (OSPAR, 2018a)
	FW4	Change in mean trophic level of consumers or marine predators	Biomass survey data and fisheries landings data (benthic-demersal community)  Region-specific TL values for species concerned	1992–2015	Bay of Biscay & Iberian Coast	Data throughout regions of the Bay of Biscay and Iberian Coast	MTL indices including all consumers (MTL2) or the highest trophic levels from 2 different thresholds (MTL3.25 and MTL4)	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/mtl-bay-biscay/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/mtl-bay-biscay/</a> (OSPAR, 2017l)  CEMP: <a href="https://www.ospar.org/documents?v=39002">https://www.ospar.org/documents?v=39002</a> (OSPAR, 2018c)
	PH1/FW5	Changes in phytoplankton and zooplankton communities	Zooplankton and phytoplankton taxa abundance time-series	2004–2014	Greater North Sea Celtic Seas Bay of Biscay & Iberian Coast		Plankton Index performed on plankton lifeform time-series	A2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/</a> (OSPAR, 2017t)  CEMP: <a href="https://www.ospar.org/documents?v=39001">https://www.ospar.org/documents?v=39001</a> (OSPAR, 2018h)  Further references: McQuatters-Gollop et al. (2019a); Bedford et al. (2020b); Bedford et al. (2020a)

**Table 7**

The Non-Indigenous species ecosystem components and associated indicator characteristics used for the OSPAR 2017 intermediate assessment. The underlying data, method, time period, region(s)/sub-region(s) assessed, and spatial coverage are also identified. Lastly, links to its use in OSPAR IA2017, its methodology, published as part of the OSPAR Coordinated Environmental Monitoring Programme (CEMP) or ICES advice, and further publications using the indicator are included.

Ecosystem component	Indicator short code	Indicator name	Underlying data	Time period	(Sub)-region(s)	Spatial coverage of data	Data analysis method	IA2017, CEMP method guidelines, and additional references
Non-Indigenous species	NIS3	Trends in new records of non-indigenous species introduced by human activities	New records of non-indigenous species	2003–2014	Greater North Sea Celtic Sea Bay of Biscay & Iberian Coast	Data throughout regions	Cumulative species count T-test	IA2017: <a href="https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/pressures-human-activities/non-indigenous/">https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/pressures-human-activities/non-indigenous/</a> (OSPAR, 2017r)  CEMP: <a href="https://www.ospar.org/documents?v=38992">https://www.ospar.org/documents?v=38992</a> (OSPAR, 2018g)

**Table 8**

Biodiversity status categories and colors used for the interpretation, by expert judgement, of indicator biodiversity state.

Poor	Indicator value is below assessment threshold, <b>or</b> change in indicator represents a declining state, <b>or</b> indicator change is linked to increasing effect of anthropogenic pressures (including climate change), <b>or</b> indicator shows no change but state is considered unsatisfactory
Uncertain	No assessment threshold <b>and/or</b> unclear if change represents declining or improving state, <b>or</b> indicator shows no change but uncertain if state represented is satisfactory
Good	Indicator value is above assessment threshold, <b>or</b> indicator represents improving state, <b>or</b> indicator shows no change but state is satisfactory
Unassessed	Indicator was not assessed in a region due to lack of data, lack of expert resource, or lack of policy support.

content analyses, stable isotope analysis, models or collected from published literature (Table 6). Food web assessments drew upon the already available data for the fish and plankton indicators (Tables 1, 3, 6).

#### 2.4.7. NIS data

Most OSPAR parties provided data on NIS newly recorded within their waters between the 1st of January 2003 and the 31st of December 2014. A distinction was made between newly recorded NIS and new introductions to account for limitations in regional monitoring programmes (at time of data collection) and the frequent time delay between introduction and detection of NIS especially in aquatic systems. The period assessed was selected because it represents two six-year periods (2003–2008 and 2009–2014), from which Contracting Parties are most likely to have complete data sets, enabling comparison of new NIS records between these two six-year reporting periods. This was in order to test the indicator for further use within the six-year rolling programme scenario of the MSFD. Data included the date on which a NIS was first recorded within each of the regions and the location of the observation.

### 2.5. Creating a holistic overview of marine biodiversity in the Northeast Atlantic


Building on the work of the 2017 OSPAR Intermediate Assessment (IA2017) which, in many cases, merely described, rather than assessed, biodiversity change, here, for the first time, we extend the assessment of biodiversity through application of an interpretive assessment for each indicator to classify indicator change within the wider ecosystem context (Table 8). This categorical assessment used expert interpretation of indicator change (from the same experts who led the development of the 2017 OSPAR indicator assessments) with respect to assessment thresholds (where available), links to pressures, and knowledge of indicator state to categorise indicators as in poor, uncertain, or good biodiversity status (Table 8). Assigning indicators to these three

biodiversity status categories has been done for the purpose of this research and related scientific interpretation, and has currently no formal link to any policy regulation (e.g. OSPAR, MSFD) for GES assessments. However, it is expected that this exercise will aid in preparation and delivery of the next OSPAR biodiversity Quality Status Report assessment to be published in 2023, which will in turn contribute to MSFD national reporting due in 2024.

Some indicators were assessed against thresholds in IA2017 and so their biodiversity status was straightforward to determine (see summarised threshold information in Table A1). More information can be found in the IA2017 assessment links in Tables 1–7, but key information is summarised here. Indicator Assessment of coastal habitats in relation to nutrient and/or organic enrichment (BH2-A) was assessed against the Water Framework Directive's benthic quality elements (for multimetric indices and associate quality ranges) for each nation's coastal waterbodies. For Recovery in the population abundance of sensitive fish species (FC1), each sensitive species was assessed relative to a percentile of its annual catch rate from all available years of data, with the upper 25th percentile suggesting "recovery" (good status), and the lower 25th percentile indicating continued decline (poor status) following Greenstreet et al. (2012). Sensitive species were then grouped within surveys and overall change for the sensitive fish group was assessed for each survey through application of a binomial test with significance level of 0.05, where significant change depends on number of species monitored and a probability of each to reach the percentile threshold by chance based on a random walk model, following Greenstreet et al. (2012). Multiple survey-based assessments were grouped within OSPAR regions and an overall assessment made using an averaging integration approach. For the indicator Proportion of large fish – Large Fish Index (FC2), a temporal baseline (a recent historic period) where available was taken; this baseline differed by survey. Where a baseline was not available a trend was assessed, with a positive trend in abundance assessed as in good status.

For the marine bird abundance indicators, Marine birds: Non-breeding abundance (B1-A) and Marine birds: Breeding abundance (B1-B), thresholds were applied as follows. A species was in good status if 70% of baseline abundance was achieved (80% in species laying only 1 egg/year); good status of a species group was then determined if at least 75% of species assessed were in good status. For Marine bird breeding success/failure (B3) a partial threshold was used in IA2017, which was then expanded here. All species in the indicator apart from terns were assessed as in good status if widespread breeding failure (<0.1 chicks/pair in 5% of colonies or more) happened in fewer than four of the last six years. For terns, widespread breeding failure was defined as < 0.1 chicks/pair in the mean percentage of colonies failing per year, over the preceding 15 years. To categorise the status of Marine bird breeding success/failure (B3) here, good status of the indicator was then determined if at least 75% of species assessed were in good status; otherwise the status was categorised as poor.

Two marine mammal indicators were assessed against thresholds in



**Fig. 2.** The status of Northeast Atlantic marine biodiversity indicators varies between indicators, ecosystem components, and OSPAR regions. Indicator short codes are given – full indicator names are in Tables 1–7. Icons in red represent poor status, orange represent uncertain status, and green represent good status. Some indicators were not assessed in some regions (grey) due to lack of data, lack of expert resource, or lack of policy support. The Greater North Sea region had the most assessed indicators, and the Bay of Biscay and Iberian Coast the fewest. The Celtic Seas region had the most indicators categorised as in good status, but most of the assessed biodiversity indicators are uncertain or poor for any of the OSPAR regions.

IA2017. For both Seal abundance and distribution (M3) and Grey seal pup production (M5), good status was defined as no decline in seal abundance of > 1% per year in the previous six-year period (a decline of approximately 6% over six years), or no decline in seal abundance of > 25% since the fixed baseline in 1992 (or closest value).

Most indicators didn't have thresholds, including all pelagic habitats, food web, and NIS indicators as well as some benthic indicators (Subtidal habitats of the southern North Sea (BH2-B), Extent of physical damage to predominant special habitats (BH3)), one fish indicator (Mean Maximum Length of fish (FC3)), and some mammal (Abundance and distribution of cetaceans (M4A), Abundance and distribution of coastal bottlenose dolphins (M4A1), Abundance and distribution of cetaceans other than killer whales and coastal bottlenose dolphins (M4B), and Numbers of harbour porpoise bycaught in commercial nets (M6)) indicators. Biodiversity status for these indicators was characterised according to Table 8. If an indicator did not have an assessment threshold, expert judgement was used to categorise indicator status based on if indicator change represented a declining/improving state (poor/good status) or there was no change in the indicator but its state was considered unsatisfactory/satisfactory (poor/good status). If it was unclear if indicator change represented a declining or improving state, or if an indicator showed no change, but it was uncertain if the state represented was satisfactory, the indicator was categorised as in uncertain status. If an indicator demonstrated change linked to increasing effect of anthropogenic pressure (including climate change), the indicator was categorised as in poor status.

Indicators were not assessed in all regions for several reasons. For some indicator-region combinations adequate data were not available to make an assessment. This was the case in the Bay of Biscay and Iberian Coast for Harbour porpoise bycatch (M6) and Production of phytoplankton (FW2) and also, since seals don't occur in the Bay of Biscay and Iberian Coast, Seal abundance and distribution (M3) and Grey seal pup production (M5). Similarly, indicator Subtidal habitats of the southern North Sea (BH2-B) was too nascent to be sufficiently developed and tested in other regions. Other indicators did not have sufficient policy support from countries bordering a region for their development. This was true in the Celtic Seas for Changes in plankton diversity (PH3) and in the Bay of Biscay and Iberian Coast for Recovery in the population abundance of sensitive fish species (FC1) and Proportion of large fish –

Large Fish Index (FC2). The indicator Change in mean trophic level of consumers or marine predators (FW4) lacked both policy support and expert resource for indicator development and assessment in the Greater North Sea and the Celtic Sea, as did Subtidal habitats of the southern North Sea (BH2-B), and all three marine bird indicators (Marine birds: Non-breeding abundance (B1-A), Marine birds: Breeding abundance (B1-B), and Marine bird breeding success/failure (B3)) for the Bay of Biscay and Iberian Coast. The focus of IA2017 was on the Greater North Sea, Bay of Biscay and Iberian Coast, and Celtic Seas region, but future assessments should aim to expand further the indicator assessments to Arctic Waters and the Wider Atlantic as resources, data, and policy support allow.

### 3. Results

The assessment of marine biodiversity indicators has revealed widespread changes across all parts of the Northeast Atlantic marine ecosystem at various scales. Here, we build on the work done under IA2017, which examined each indicator separately, to now holistically examine changes in each ecosystem component, assessing biodiversity status and drawing conclusions, where possible, across the whole OSPAR area (Fig. 2). This is a high-level summary focusing on key changes detected in each indicator in each region; for further detail on individual indicator change, refer to the references in Tables 1–7.

#### 3.1. Pelagic habitats

Statistically significant changes were observed in pelagic habitats across all regions (references in Table 1). Since these indicators did not have thresholds and pressures causing indicator change were not identified during the assessment, the biodiversity status of Changes in phytoplankton and zooplankton communities (PH1/FW5), Changes in phytoplankton biomass and zooplankton abundance (PH2), and Changes in plankton diversity (PH3) therefore was classified as “uncertain” (Fig. 2, Table A1). Most ecologically-relevant pairs of plankton (referred to here as lifeform pairs; Changes in phytoplankton and zooplankton communities (PH1/FW5)) experienced significant changes throughout the Greater North Sea, Celtic Sea, and Bay of Biscay and Iberian Coast regions, indicating alterations to pelagic habitat structure

and functioning. Among the most pronounced was the relative change in the interannual mean abundance of holoplankton and meroplankton, particularly in the Greater North and Celtic Seas, suggesting changes in the coupling of the benthic and pelagic components of the ecosystem. Although multiple time-series datasets were analysed (see Table 1), changes in phytoplankton biomass (Changes in phytoplankton biomass and zooplankton abundance (PH2)) were remarkably consistent across the OSPAR maritime area with anomalies of phytoplankton biomass proxies rapidly increasing from the 1980s, a manifestation of the well-documented region-wide regime shift (Beaugrand, 2004). Similarly, zooplankton abundance largely decreased across the whole area, although it did increase in some regions post-2000s. For plankton diversity (Changes in plankton diversity (PH3)), an increase in species dominance (the relative abundance of certain species compared to the overall community) was observed in 2007–2008 for the Greater North Sea stations, suggesting a shift in community composition, but the presence of a definitive trend in species dominance patterns is unclear. In the Bay of Biscay and Iberian Coast, phytoplankton communities showed an increase in species dominance over the time period of assessment.

### 3.2. Benthic habitats

The 2017 assessment of coastal benthic habitats indicated that 89% of the coastal water bodies in the OSPAR Maritime Area, including most of the Bay of Biscay and Iberian Coast and Celtic Seas regions, had benthic habitats meeting thresholds for good status in relation to nutrient and/or organic enrichment (Assessment of coastal habitats in relation to nutrient and/or organic enrichment (BH2-A)) with regard to macroalgae and angiosperms, and 74% with regard to benthic invertebrates (Fig. 2, Table A1). However, there were wide regional variations and several data gaps existed, with localised problem areas for nutrient and/or organic enrichment, notably in the Greater North Sea, which was assessed as uncertain status due to lack of data or uncertainties in response of the indicator to nutrients. No thresholds were agreed for IA2017, but the extent of physical damage to the seafloor (Extent of physical damage to predominant special habitats (BH3)) showed that 86% of the assessed areas in the Greater North Sea and the Celtic Seas had physical disturbance due to bottom trawling, of which 58% were assessed as highly disturbed and thus in poor status (Fig. 2, Table 2). Three quarters (74%) of all assessed areas experienced consistent fishing pressure year on year, which is very likely to affect the ability of habitats to recover. The absence of baseline habitat maps for most of the deeper areas of the Bay of Biscay and Iberian Coast prevented assessment of most of this area, with assessment only possible for shallow and coastal regions. Benthic habitat community quality was sub-regionally assessed in terms of species richness in the southern North Sea (Subtidal habitats of the southern North Sea (BH2-B)) where community quality was found to be generally lower in coastal areas than further offshore, partly due to higher fishing pressure in coastal areas. The indicator was therefore assessed here as in uncertain status due to spatial gaps in the data used; no threshold was agreed in IA2017. Other anthropogenic activities potentially causing pressures on benthic habitats were not assessed in 2017.

### 3.3. Fish:

The decline in abundance of fish species sensitive to fishing effects (Recovery in the population abundance of sensitive fish species (FC1)) was found to have halted in the Greater North Sea (i.e. a statistically significant proportion of species was above the 75th percentile of each

species' annual catch rate), but recovery (above the upper 25th percentile) had not occurred, meaning the primary assessment threshold was not met and so the indicator was assessed as in poor status in the region. Recovery (i.e. meeting the threshold of a statistically significant proportion of species above the 25th percentile of each species' annual catch rate) was apparent overall in the Celtic Seas, resulting in good status in that region (Fig. 2, Table A1).

For the wider demersal fish community, there were improvements in parts of the OSPAR area. Recovery in the Large Fish Index (LFI; Proportion of large fish – Large Fish Index (FC2)) in the northern Celtic Seas (west of Scotland) suggests achievement of good status in this subdivision, but elsewhere in the Celtic Seas region the increases observed did not meet targets, resulting in poor status as the threshold was not met. Increases in LFI (Proportion of large fish – Large Fish Index (FC2)) in the Greater North Sea suggested that recovery could be achieved before the next assessment if fishing pressure does not increase, but the region is not yet at good status and hence categorised as poor. Despite these improvements for the region overall, additional analyses of the species composition of demersal fish communities (Mean Maximum Length of fish (FC3)) found that in the south-eastern and central-western North Sea, some areas along the continental shelf edge to the northeast and southwest of Ireland and parts of the Iberian coast were at the minimum observed state (lowest recorded (Mean Maximum Length) values) and thus, the community was dominated by small-bodied species that are generally more resilient to fishing than large-bodied species. This indicator did not have an assessment threshold in IA2017, and was therefore categorised as of uncertain status in all regions (Fig. 2, Table A1). Thus, the increase in the LFI for demersal fish in the Greater North Sea overall is dependent on only part of the region surveyed and, without a further recovery of large-bodied species elsewhere, may be limited.

Community indicators for pelagic fish were newly developed, with a weaker assessment basis, and assessment outcomes were largely of uncertain status. Nevertheless, an increase in the dominance of smaller species in the southern North Sea and west of Scotland merits further surveillance.

### 3.4. Marine birds

All three marine bird indicators were assessed against thresholds. Breeding populations of marine birds were determined to be in poor status in the Greater North Sea and Celtic Seas regions (Marine birds: breeding abundance (B1-B); Fig. 2, Table A1). In all three regions, and for more than a quarter of the marine bird species assessed, there has been a considerable (>20–30%) drop in abundance compared to the levels observed 25 years ago (thresholds depending on clutch size, see section 2.5). Declines in breeding abundance started mainly in the mid-2000s. Frequent and widespread breeding failure has also been observed for > 25% of species assessed in the Greater North Sea during the period assessed (2010 to 2015 inclusive), resulting in poor status. In the Celtic Seas, 75% of species did not show frequent and widespread breeding failure, indicating good status (Marine bird breeding success/failure (B3)). Breeding abundance declines (Marine birds: breeding abundance (B1-B)) and breeding failures (Marine bird breeding success/failure (B3)) were most apparent in those species feeding on small fish in the surface waters of the Greater North Sea and Celtic Seas, highlighting food availability as an important driver in the population dynamics of marine birds in the Northeast Atlantic (ICES, 2017).

Similarly, the non-breeding abundance (Marine birds: non-breeding abundance (B1-A)) of more than a quarter of the waterbird species that visit the Celtic Seas during migration and/or during winter has also declined by >20–30% between 1992 and 2014/15, indicating poor

biodiversity status. In contrast, non-breeding populations of marine birds visiting the Greater North Sea were classed as in good biodiversity status, with >75% of species being present in higher numbers compared to levels in the early 1990s.

### 3.5. Marine mammals

Marine mammals, as top predators, can provide suitable indicators to reflect the state of the marine ecosystem. Grey seal abundance (Seal abundance and distribution (M3)) and pup production (Grey seal pup production (M5)) have increased in the Greater North Sea and eastern Celtic Seas since 1992, while the number of grey seal breeding colonies has increased or remained unchanged since 2003. Harbour seal abundance (Seal abundance and distribution (M3)) has been stable or has increased in most places but is in decline in others, though some areas have insufficient data. Indicators Seal abundance and distribution (M3) and Grey seal pup production (M5) have therefore been classed here as achieving good status in the Greater North Sea and eastern Celtic Sea as assessment thresholds were met (Fig. 2, Table A1). Most of the coastal bottlenose dolphin populations (Abundance and distribution of coastal bottlenose dolphins (M4-A1)) remained stable when compared to populations in 1990, but the number of animals in those populations remains low while the population is in decline in Portugal; this suggests poor status across the Greater North Sea, Celtic Seas, and Bay of Biscay and Iberian Coast (Fig. 2, Table A1). There is no evidence for changes in abundance of white-beaked dolphin, minke whale and harbour porpoise since 1994 while there is insufficient evidence for other species (Abundance and distribution of cetaceans other than killer whales and coastal bottlenose dolphins (M4-B)), resulting in uncertain status for this indicator. The distribution of harbour porpoise and minke whale has shifted southward in the Greater North Sea. Insufficient data and high levels of uncertainty prevented a robust assessment of harbour porpoise bycatch (Harbour porpoise bycatch (M6)), resulting in uncertain status, but at least 4000 porpoises (from a population of > 490,000) are incidentally caught in fishing nets annually and bycatch is considered one of the main direct human pressures on this species.

### 3.6. Food webs

The structure of marine food webs was found to be altered across the OSPAR area, but it is unclear how the ecosystem components are affecting each other or responding to the multiple and cumulative effects of human activities and climate change in the area. The food web indicators had no assessment thresholds in IA2017.

The pilot assessment on phytoplankton productivity (Production of phytoplankton (FW2)) was inconclusive across the whole OSPAR area, but showed changes in ecosystem functioning in some discrete sites and in parts of the North Sea, where a decreasing trend in annual phytoplankton production was reported after 1988. This trend is in contrast to the phytoplankton component of indicator Changes in phytoplankton biomass and zooplankton abundance (PH2) and suggests a change in phytoplankton community structure, which potentially could affect upper trophic levels. Scientific uncertainty and data gaps have resulted in uncertain status addressed by this indicator (Fig. 2, Table A1). However, the pilot assessment illustrates the potential of this indicator, representing the main production source of organic matter in the marine environment, to provide valuable information to a broader and holistic ecosystem assessment.

Typical length (Size composition in fish communities (FW3)) was used to assess the size structure within fish communities but as no threshold was proposed the status is uncertain. For demersal fish the

indicator was generally increasing across all OSPAR regions since 2010, while no long-term trend was found for pelagic fish, suggesting overall that recovery is ongoing (Fig. 2, Table A1). However, this follows long-term decreases in the Greater North Sea between 1980s and 2000s, with particular declines in the southern and central North Sea. In addition, the lack of recovery in species composition (as evidenced by indicator Mean Maximum Length of fish (FC3)) in the North Sea suggests the ongoing recovery in the size structure of the fish community may be limited. Long-term decrease was also evidenced between 1990s and 2005 in the Celtic Seas, especially in the western Irish Sea, in the continental shelf edge north-west of Ireland, and to the west of Brittany. In contrast, long term increases overall for typical length of demersal fish were evident in the Bay of Biscay and Iberian Coast.

Change in average trophic level of consumers and marine predators (Change in mean trophic level of consumers or marine predators (FW4)), applied in the Bay of Biscay and Iberian Coast, showed no apparent change in overall food web structure over recent decades. However, there are some signs of improvement when only predators (Mean Trophic Level (MTL) 3.25 and MTL4) were considered. The status of the average trophic level in the Bay of Biscay and Iberian Coast therefore remains uncertain (Fig. 2, Table A1). These changes occurred alongside the underlying shifts observed in plankton communities assessed through plankton lifeforms (Changes in phytoplankton and zooplankton communities (PH1/FW5)).

### 3.7. NIS

The number of newly recorded non-indigenous species (Trends in new records of non-indigenous species introduced by human activities (NIS3)) in the Greater North Sea, Celtic Seas and the Bay of Biscay and Iberian Coast regions over the period 2003–2014 varied by year and region (references in Table 7); no assessment thresholds were applied to this indicator in IA2017. In the Greater North Sea there was a relatively constant linear increase in the number of newly recorded NIS over time, whereas the other two regions had particular years with comparatively high numbers of new NIS recorded (for the Celtic Seas, 2006 and 2012, and in the Bay of Biscay and Iberian Coast, 2004) which meant a similar linear increase did not occur. Within the Celtic Seas this resulted in a stepwise increase in newly recorded NIS over the period, however, within the Bay of Biscay and Iberian Coast the cumulative number of recorded new NIS plateaued after 2004 with much fewer new NIS recorded after this date. These results are highly likely to be an artefact of varying monitoring and reporting efforts between the regions over this period rather than discrete episodic events leading to high levels of introductions occurring in particular years (Fig. 2, Table A1), resulting in uncertain status. The information presented does indicate a consistent rate of new introductions occurring within the Greater North Sea and the Celtic Seas. While this does highlight the limitations of assessing new records of NIS and the dependence of the indicator on accurate monitoring data, it also illustrates the effectiveness of comparing six-year blocks of data as an indicator for MSFD purposes.

## 4. Discussion

This first science-policy collaborative, regional sea scale effort at assessing the status of Northeast Atlantic marine biodiversity has revealed variation in biodiversity status between the biodiversity indicators, the seven ecosystem components, and the OSPAR regions assessed here (Fig. 2). The semi-quantitative method applied here used expert interpretation of change in a total of 52 indicator-region combinations with respect to assessment thresholds, links to pressures, and/or

knowledge of indicator state to categorise indicators as in poor, uncertain, or good biodiversity status. Indicators were not assessed in all regions, due to lack of data, lack of expert resource to conduct an assessment, or lack of policy support (icons shown as grey in Fig. 2). The status of most assessed indicator-region combinations was either uncertain ( $n = 30$ ; 58% of indicator-region combinations) or poor ( $n = 13$ ; 25% of indicator-region combinations) (Fig. 2). In fact, very few examples of good status ( $n = 9$ ; 17% of indicator-region combinations) were identified, which is not in line with meeting MSFD objectives, or wider policy initiatives for sustainability such as the Aichi Biodiversity targets (Convention on Biological Diversity Secretariat, 2010) or United Nations Sustainable Development Goals (United Nations, 2015). Some bright spots categorised in good status were revealed, such as with marine mammals (i.e. seals) in the Greater North Sea and Celtic Seas, coastal benthic habitats in the Bay of Biscay and Iberian Coast and Celtic Seas (with respect to organic and nutrient enrichment), some fish communities in the Greater North Sea and Celtic Seas, and some birds in the Celtic Seas. Of the three regions, the Celtic Seas had the most ( $n = 5$ ) indicators in good status, including those of coastal benthic habitats, fish, birds, and marine mammals, with 5 indicators categorised as poor status. Similarly, the Greater North Sea had the most ( $n = 6$ ) indicators with poor status but also had 3 indicators with good status. The Bay of Biscay and Iberian Coast had the highest proportion of indicators categorised as in uncertain status (9 out of 12 assessed indicators). This could be because the Greater North Sea and Celtic Seas have relatively good data coverage when compared to the Bay of Biscay and Iberian Coast, which reduced uncertainty and increased the ability to determine status.

Most of the assessed biodiversity indicator results per region ( $n = 30$ ; 58% of indicator-region combinations) fell into the uncertain status category as data limitations, such as the quality and quantity of information, prevented robust status categorisation. Data gaps were mostly a result of insufficient geographical and temporal coverage for both pressure and biodiversity data. For example, some marine mammal indicators could not be assessed with confidence since the available pressure data, i.e. harbour porpoise bycatch, and the biological data, i.e. the abundance of cetaceans other than bottlenose dolphins, were not at a wide enough spatial scale. In terms of the temporal scale of analysis, the length of the time-series is an important factor for the interpretation of the indicator and hence, biodiversity assessment. In the case of plankton, the available datasets have different timespans, from 14 to 60 years, making it challenging to establish a common reference period across datasets to assess potential large-scale long-term changes. Paucity of long-term benthic datasets hinders the development of reference points to measure change over time. In the past few years, however, the implementation of the Water Framework Directive (2000) and MSFD (2008) and increased numbers of environmental impact studies and Marine Protected Areas have slightly increased the number of standardised surveys and datasets collected. These datasets, used to assess benthic biodiversity status, improve quantitative and in situ validated knowledge on species and biotopes, but only for a limited number of geographic areas and for specific environmental management scales and objectives. This also highlights the need to develop more coordinated benthic monitoring. The variety of methods and protocols (gears, metadata, taxonomic detail, quality assurance and sampling designs) used to acquire the different datasets also limits the spatial aggregation and comparability of biodiversity data at (sub)regional scales. Likewise, the data collected for NIS were patchy both temporally and spatially due to limited, uncoordinated, and varying efforts in monitoring and species categories considered. The absence of a clear baseline from which assessments of new arrivals could be made has been indicated as an issue

for NIS which has subsequently been addressed (Tsiamis et al., 2019). The temporal scale at which indicators are assessed must be therefore appropriate to the policy question being addressed. In future assessments, reporting more indicators over multiple temporal scales (as was done for Recovery in the population abundance of sensitive fish species (FC1), Mean Maximum Length of fish (FC3), and Size composition in fish communities (FW3)) could provide further information relevant to management and use all available data, while still retaining wider ecological context, including multi-decadal trends cause by climate change (Bedford et al., 2020c).

Even where data were available, the novel nature of this work was another contributing factor to the high number of indicators with uncertain status as a result of their relatively low state of methodological development and/or limited previous application in a management context. Some indicators, such as the Proportion of large fish—Large Fish Index (FC2) and Marine birds: non-breeding abundance (B1-A), have a history of assessment. While they were further developed for IA2017, the previous assessment experience, including an established network of experts and data flows, enabled the status category addressed by these indicators to be clearly determined. Most OSPAR common benthic, pelagic and food web indicators, were developed explicitly for the MSFD and were assessed for the first time for the OSPAR Intermediate Assessment 2017. As part of this assessment process, expert networks and data flows had to be established and appropriate indicators had to be selected and developed. An important yet often complex aspect of biodiversity indicators is a clear understanding of the nature and extent of potential pressures that could drive community changes. Modelling and statistical analyses provide a clear route forward to develop our understanding (Lynam et al., 2016) and such studies have been furthered for fish and food web indicators since OSPAR Intermediate Assessment 2017 (Piroddi et al., 2021; Queirós et al., 2018; Spence et al., 2021; Thompson et al., 2020; Thorpe et al., 2022). This understanding is essential for policy-makers to manage human activities effectively to reduce pressures and restore biodiversity.

Currently, scientific knowledge gaps exist around the mechanisms through which multiple pressures enact change in these new benthic, pelagic and food web indicators; this can be due to lack of data availability and/or relatively low scientific understanding of pressure-state relationships for some benthic, pelagic and food web components. Such relationships are often not linear (e.g. Elliott et al., 2018; Mackey and Currie, 2001; McQuatters-Gollop et al., 2007; Noël et al., 2009; Samhuri et al., 2010) and pressures can act in a cumulative manner making it difficult to identify the most important causes of biodiversity change. For example, pelagic habitat indicators may be influenced at local scales by nutrients while climate change is also acting at wide spatial and long temporal scales (Bedford et al., 2020b; McQuatters-Gollop et al., 2007). Similarly, the cumulative effects of multiple anthropogenic activities, most notably land-based contamination, and fishing, along with climate change, are key pressures on the benthic (Condition of benthic habitat-defining communities (BH2), Extent of physical damage to predominant special habitats (BH3)), food web (Size composition in fish communities (FW3), Change in mean trophic level of consumers or marine predators (FW4)) and fish (Recovery in the population abundance of sensitive fish species (FC1), Proportion of large fish – Large Fish Index (FC2)) indicators.

The effects of pressures on benthic habitats can be also measured by evaluating the sensitivity ranges of the species and biotopes within each habitat type and the spatial and temporal overlaps with those activities causing the pressures (de Juan and Demestre, 2012; de Juan et al., 2020; Dupaix et al., 2021; González-Irusta et al., 2018; Tillin and Tyler-Walters, 2014). Activities operating at large scale, such as fishing, will

have stronger effects on the sensitivity of benthic habitats and in particular their ability to recover from pressures, than those limited to a geographical area or with a small footprint (Halpern et al., 2008; Jennings et al., 2012; OSPAR, 2017m). The Production of phytoplankton (FW2) indicator was a pilot indicator in development at the time of IA2017, and relationships with pressures had not yet been evaluated. However, the influence of climate change and riverine inputs of nutrients to the North Sea have elsewhere been found to be important factors influencing primary productivity (e.g. Capuzzo et al., 2018).

Climate change was identified as a common key pressure on Northeast Atlantic marine biodiversity. However, in many cases (particularly for pelagic habitat, benthic habitat, and food web indicators) the links between climate change and indicator change were not quantified in IA2017. Climate change was not a key focus of IA2017 and was at the time considered an outside pressure, or prevailing condition, on the system (e.g. Elliott et al., 2015), with its role in driving change unexplored or unquantified for many of the indicators. Climate change was identified as a major pressure causing change in plankton indicators, but resource limitation and scientific knowledge gaps prevented the quantification of this pressure-state relationship as part of IA2017. Since IA2017, however, temperature change has been identified as a key pressure, notably causing changes in phytoplankton and zooplankton communities (Bedford et al., 2020b) and the proportion of large fish (Queirós et al., 2018), knowledge that will progress the scientific robustness of the future assessments.

In benthic habitats, fishing is clearly driving change as evaluated in the Extent of physical damage to predominant special habitats (BH3) indicator, but the ability to quantify the relationship between fishing and changes in species diversity of subtidal benthic communities (Subtidal habitats of the southern North Sea (BH2-B)) is highly dependent on data available from different habitat types under a variety of levels of exposure to pressures (Pitcher et al., 2022; van Loon et al., 2018). For marine mammals, fishing by-catch is one of the main human pressures on harbour porpoises (Harbour porpoise bycatch (M6)). However, there are high levels of uncertainty in estimates of harbour porpoise by-catch rates, making it difficult to confidently categorise ecological status.

Indicators for some ecosystem components are responding to both fishing and climate change. Change in bird indicators (Marine birds: Non-breeding abundance (B1-A), Marine birds: breeding abundance (B1-B), Marine bird breeding success/failure (B3)) is driven by both climate change and fishing, with climate change having an indirect effect on birds through food supply (e.g. forage fish, which are responding to warming), while fishing directly removes fish as a food source for birds from the ecosystem (Cook et al., 2014; Frederiksen et al., 2004). The indicators for Recovery in the population abundance of sensitive fish species (FC1, Greenstreet et al., 2012), Proportion of large fish – Large Fish Index (FC2, Greenstreet et al., 2010), and Mean Maximum Length of fish (FC3, Greenstreet and Rogers, 2006) were constructed to be responsive to the effects of fishing. Nevertheless, in the long term, each indicator will likely be affected by climate since the recovery of sensitive species (Blumel et al., 2022) and communities (Queirós et al., 2018) will depend on prevailing environmental conditions. Similarly, the Change in mean trophic level of consumers or marine predators (FW4) indicator appears to be mainly sensitive to fishing pressure (e.g., Arroyo et al., 2019; Preciado et al., 2019), although lack of reliable landings data and the inaccuracy of the trophic levels assigned to predators has challenged the quantification of this relationship. Even though results of IA2017 showed a promising apparent increase in the MTL trend based on survey data, indicating a recovery of the Bay of Biscay and Iberian Coast's benthic-demersal system, further analyses have revealed spatial differences in MTL values and demonstrated the

negative effect of fishing pressure on the indicator at local scale (Preciado et al., 2019). These results suggest that the apparent stability revealed by the indicator based on landed catch data may be masking other processes such as the expansion of demersal fisheries to deeper waters, and an over-exploitation of, especially, pelagic species (Arroyo et al., 2019; Arroyo et al., 2017). Overall, the stability of the indicator when analysed at the regional level may be indicating that the ecosystem is resilient (withstanding the pressure) but consistently overexploited. The effects of climate change and shifts in productivity interact with fishing, making the effects of various pressures difficult to disentangle or identify targets and thresholds (Arroyo et al., 2019).

Some other anthropogenic pressures were also found to be important for Northeast Atlantic marine ecosystems. An increase in the Trends in new records of non-indigenous species introduced by human activities (NIS3) indicator is likely linked to ballast water, hull fouling and aquaculture (Tidbury et al., 2016). However, this pressure-state relationship was not quantified in IA2017, due to uncertainties and gaps in the data, so the status of the indicator is uncertain (Fig. 2, Table A1). For the three pelagic habitats indicators (Changes in phytoplankton and zooplankton communities (PH1/FW5), Changes in phytoplankton biomass and zooplankton abundance (PH2), and Changes in plankton diversity (PH3)), it is likely that nutrients are influencing change in pelagic indicators in some localised areas (McQuatters-Gollop et al., 2007), but, again, this relationship needs to be quantified in reference to wide scale changes caused by warming (e.g. Bedford et al., 2020b). Nutrients also influence other ecosystem components such as benthic habitats. The Assessment of coastal habitats in relation to nutrient and/or organic enrichment (BH2-A) indicator clearly showed links to nutrient loading; but data are only available for some coastal habitats, assessed through various different multimetric indices, resulting in uncertain status. Chemical and noise pollution affect individual marine mammals, but the effects of these pressures on populations are not yet well understood (Erbe et al., 2018; Jepson et al., 2016; Murphy et al., 2015). Habitat loss is an increasing pressure for marine bird species displaced from areas where offshore wind farms are operating (Mendel et al., 2019) or ship traffic is disturbing (Mercker et al., 2021; Schwemmer et al., 2011).

Finally, gaps in data or knowledge were not the only challenges for determining if indicator status in an area was 'good' or 'not good'. The original approach for the biodiversity components of the MSFD was based on traditional environmental aspects, such as contaminant effects on biota, with known pressure-state relationships (e.g. Sahlmann et al., 2017). In such cases, indicators are monitored and if the state indicator exceeds a threshold value, it is considered 'not Good Environmental Status'. The IA2017 experience has revealed that, for some ecosystem component indicators, this process is not so easy, simple, or clear and with the idea of 'good' or 'not good' difficult to define with a single threshold value allowing categorisation of status. For pelagic habitats (Changes in phytoplankton and zooplankton communities (PH1/FW5)), for example, the maintenance of ecosystem functioning is more important than changes in the abundance of plankton lifeforms or individual taxa, but is difficult to define with a threshold value. There has been a long history of development of thresholds for commercial fisheries, where the concept is relatively straightforward (Schaefer, 1991), but even here multi-species approaches that take into account food web interactions are considered problematic (Thorpe and De Oliveira, 2019). For non-commercially fished fish species of concern, it is challenging to categorise their environmental status due to the lack of a baseline and/or reference conditions and, often, limited data. Furthermore, discerning the influence of environmental/natural drivers versus fishing on predator-prey interactions and, ultimately, ecosystem functioning, is an



issue that is far from being solved but great progress has been made since IA2017 (e.g. Link and Watson, 2019; Piroddi et al., 2021; Thompson et al., 2020). Although it is possible to measure seafloor disturbance using a combination of sensitivity ranges with levels of exposure to pressures (Halpern et al., 2008; OSPAR, 2017m; Tillin and Tyler-Walters, 2014), it is difficult with the data available to define the boundaries for benthic habitat recovery and therefore to determine 'good' status. This lack of boundary definition presents challenges for quantifying the magnitude of habitat damage, although modelling approaches could be used to estimate the levels of pressure effects or the proportion of habitat area to be protected (Jennings et al., 2012; Rondinini, 2011; Rovellini and Shaffer, 2020). There are also still methodological gaps around how to extrapolate the fine scale measure of the state-pressure relationship (Condition of benthic habitat-defining communities (BH2)) to wider biological and biogeographical scales (Extent of physical damage to predominant special habitats (BH3)). However, the use of condition benthic indicators (Condition of benthic habitat-defining communities (BH2)) to assess habitat condition and develop pressure-state curves (Elliott et al., 2018) is a key step in defining these boundaries as well as in establishing condition (also called quality) thresholds. When this is not possible, a precautionary approach is required, stopping the pressure by managing the related activities (e.g. bottom trawling) to avoid continued habitat damage while environmental status is evaluated. This would notably be possible and important in Marine Protected Areas, even if the benthic habitats are often not the initial and key objective of management (Greathead et al., 2020). Similarly, the presence of newly arrived NIS does not directly relate to poor or degrading environmental conditions as there can be considerable delay, sometimes of several decades, between the introduction of a NIS and related effects on the environment. In many cases especially in the marine environment the effect that NIS present is poorly understood further restricting the capability to directly link new records and environmental effect. In terms of management, the prevention of NIS introduction is crucial, as it is very difficult locally, and almost impossible at wider scale, to avoid the proliferation of invasive marine species which lead to high socio-economic impacts.

Even though knowledge gaps and challenges remain around categorising the status of biodiversity indicators, the work presented here demonstrates that science-policy collaboration is a successful approach to delivering robust science to support policy needs. IA2017 represents one of, if not the, world's most comprehensive applications of the ecosystem approach, which looked across multiple ecosystem components using a variety of indicator types at multiple spatio-temporal scales to better understand and manage marine biodiversity and ultimately realise the ambitions of the MSFD. This process enabled synergistic working between 188 nominated scientists, as well as policy-makers from 15 Contracting Parties, to develop policy-useful biodiversity indicators, evaluate indicator change, assess environmental status where possible, and communicate the work appropriately (e.g. McQuatters-Gollop et al., 2019b). Furthermore, the IA2017 process established data flows which increased the level of participation in the assessment from contracting parties. It was found that top-down data calls, issued by OSPAR through the heads of Contracting Party delegations, work better at mobilising data into the assessment process than bottom-up data calls among experts. Stabilising and formalising data flows has facilitated the whole assessment process, particularly for indicator datasets that rely on individual national or even finer scale monitoring programmes (e.g. datasets that are maintained by non-governmental research laboratories). The working relationships, information flows, cross-cutting issues, and lessons learned during the IA2017 process will underpin and facilitate future biodiversity

assessments, and will be important to consider for future policy mechanisms such as the development of the CBD Post-2020 Global Biodiversity Framework. The process, however, does have room for improvement. For example, extending scientific expert contribution by enlarging the expert groups to gain an even more robust scientific consensus, and enabling official citable authorship of experts by OSPAR could contribute positively to the next assessment cycles.

#### 4.1. Recommendations for future assessments

To increase the robustness of future assessments, biodiversity monitoring should be improved by filling the gaps in observations. Monitoring should be extended and coordinated so that the entire communities of plankton, benthos, fish, and NIS can be assessed in a more comprehensive way, for example, by including data on the smaller size fraction of the plankton, by monitoring both benthic biomass and diversity, including cephalopods, and by incorporating NIS monitoring in risk areas across all biodiversity components. Also, in further indicator development, other metrics and parameters of novel sampling and monitoring techniques and technologies (e.g. Pitois et al., 2021) and their potential contribution towards the production of biodiversity data products should be considered (Danovaro et al., 2016; Hardisty et al., 2019).

Secondly, conceptual understanding is still evolving around pressure-state relationships for many biodiversity indicators. There are particular gaps around how multiple pressures act on benthic, pelagic, and marine mammal communities, and how climate change affects indicators. These pressure-state relationships need to be quantified, and the effects of multiple pressures on indicators' sensitivity disentangled, so that policy can implement effective management measures. Similar to the spatially-nested approach taken for pelagic habitats indicators (Bedford et al., 2020b; McQuatters-Gollop et al., 2019a), which considers pressures at multiple spatial scales, a set of indicators across ecosystem components, and assessed at the regional sea scale, could help to spatially separate areas experiencing direct pressure effects, such as from fishing or nutrients, from long-term change such as climate change or natural variability.

Third, additional work needs to be done on alternative methods of systematically assessing indicators for GES where threshold values are not meaningful. More thinking is needed to accommodate the nuance of interpreting indicators that capture magnitude of change and recovery after halting pressures, or complex indicators that represent multiple aspects of ecosystem functioning. This also includes the further development and testing of the already existing candidate indicators. For example, the candidate food web indicator Ecological Network Analysis Indices (FW9), is necessarily complex as it represents a set of indices across multiple trophic levels, which form an overall picture of ecosystem functioning and organization (de Jonge and Schückel, 2021; Safi et al., 2019). The simple method we applied here, of assessing indicator status by linking indicator results to pressures in the absence of threshold values, can underpin this development.

Fourth, the conceptual modelling of indicator interactions is needed to enable integrated ecosystem assessment. Links between indicators need to be defined and, ultimately, quantified with data. The conceptual understanding of these interconnections is developing but the data or understanding to define and quantify these relationships is lacking. This is likely to be challenging due to multiple spatial and temporal scales, but some work on conceptualising these relationships has already begun (Elliott et al., 2017b; Elliott et al., 2018; Lockerbie et al., 2018). Understanding the interactions between indicators will help us provide even more robust evidence for policy. The Ecological Network Analysis

approach (ENA; candidate indicator Ecological Network Analysis Indices (FW9)) could help to understand such interconnections since the set of indices encompasses all components of the benthic and pelagic systems (Fath et al., 2019; Safi et al., 2019). An approach similar to Raoux et al. (2019) and Noguez et al. (2021) could bring together ENA indices (Ecological Network Analysis Indices (FW9)) and the mean trophic level indicator (Change in mean trophic level of consumers or marine predators (FW4)) to assess the status of food webs. The challenge would then be to make the conceptual complexity of the ENA approach understandable to policy makers and the general public (de Jonge and Schückel, 2021).

Finally, there is a clear and demonstrable need for this type of work and so it must be adequately resourced. In many instances indicators could not be assessed in IA2017 due to lack of expert resource to develop indicators and carry out assessments. For benthic and pelagic habitats and food webs, specific recommendations were developed and published as an action plan to guide through collective priorities for how to better progress towards a more ecosystem-based approach to holistic assessment, but this work needs to be consistently resourced (Padegimas et al., 2017). Much of the indicator development thus far has been funded ad hoc through external projects. This meant that expert group leads and indicator leads spent much effort and time responding to competitive national and European funding calls by writing proposals, many of which were not funded. A more efficient way to resource this policy work would be through consistent investment from Contracting Parties.

## 5. Conclusions

This work adds value to OSPAR IA2017, by determining, for the first time, the status of biodiversity, as addressed by indicators, across multiple ecosystem components and at a Northeast Atlantic scale, based on an understanding of pressure-state relationships. Some 'good' status bright spots were identified, along with some 'poor' status lowlights. The status of most biodiversity indicators across much of the Northeast Atlantic, however, was found to be 'uncertain', due to data and knowledge gaps, and challenges around the nuanced nature of categorising status for complex biodiversity indicators. The authors hope that this work will contribute to the upcoming formal OSPAR Quality Status Report and Contracting Party MSFD assessment processes by interpreting changes reported in IA2017 with new perspectives and insights that can underpin future formal assessments, and even more importantly achievement, of Good Environmental Status.

More work urgently needs to be done to fill data gaps; identify and quantify pressure-state relationships; understand the influence of climate change on indicators; test options for categorising environmental status where threshold values are unavailable or inappropriate; and conceptualise and quantify links between indicators and ecosystem components. These issues are not straightforward to resolve, particularly when working across multiple spatio-temporal scales. Crucially, success depends on the adequate resourcing of this work, which becomes ever more urgent as climate change and direct pressures on Northeast Atlantic marine ecosystems accelerate.

## CRedit authorship contribution statement

**A. McQuatters-Gollop:** Conceptualization, Methodology, Writing – original draft. **L. Guérin:** Conceptualization, Methodology, Writing – original draft. **N.L. Arroyo:** Investigation, Writing – review & editing. **A. Aubert:** Investigation, Writing – review & editing. **L.F. Artigas:** Investigation, Writing – review & editing. **J. Bedford:** Investigation, Writing –

review & editing. **E. Corcoran:** Investigation, Writing – review & editing. **V. Dierschke:** Investigation, Writing – review & editing. **S.A.M. Elliott:** Investigation, Writing – review & editing. **S.C.V. Geelhoed:** Investigation, Writing – review & editing. **A. Gilles:** Conceptualization, Methodology, Writing – original draft. **J.M. González-Irusta:** Investigation, Writing – review & editing. **J. Haelters:** Investigation, Writing – review & editing. **M. Johansen:** Investigation, Writing – review & editing. **F. Le Loc'h:** Investigation, Writing – review & editing. **C.P. Lynam:** Investigation, Writing – review & editing. **N. Niquil:** Investigation, Writing – review & editing. **B. Meakins:** Investigation, Writing – review & editing. **I. Mitchell:** Conceptualization, Methodology, Writing – original draft. **B. Padegimas:** Investigation, Writing – review & editing. **R. Pesch:** Investigation, Writing – review & editing. **I. Preciado:** Investigation, Writing – review & editing. **I. Rombouts:** Investigation, Writing – review & editing. **G. Safi:** Investigation, Writing – review & editing. **P. Schmitt:** Investigation, Writing – review & editing. **U. Schückel:** Conceptualization, Methodology, Writing – original draft. **A. Serrano:** Investigation, Writing – review & editing. **P. Stebbing:** Investigation, Writing – original draft. **A. De la Torre:** Investigation, Writing – review & editing. **C. Vina-Herbon:** Conceptualization, Methodology, Writing – original draft.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Data availability

Data are available from OSPAR ODIMS <https://odims.ospar.org/en/>






## Acknowledgements



The authors of this paper are grateful to the networks of scientific experts who have enabled and progressed this work, to the OSPAR ICG-COBAM and BDC members and convenors, to the OSPAR secretariat, and to sunny bar and restaurant terraces across Europe which facilitated development of these indicators by fuelling discussions with wine and cheese. Benthic, food web, and pelagic authors are grateful to funding received from European Maritime and Fisheries Fund for the "Applying an Ecosystem Approach to (sub) Regional Habitat Assessments (EcAprHA): Addressing gaps in biodiversity indicator development for the OSPAR Region from data to ecosystem assessment", grant number: 11.0661/2015/712630/SUB/ENVC.2 OSPAR. AM-G is supported by the UK Natural Environment Research Council (NE/R002738/1) Knowledge Exchange fellowship scheme. AA, LG, SE, NN, GS, and IR received funding from the French Ministry of Environment, Energy, and the Sea (MEEM), the French National Museum of Natural History (MNHN) and the French National Centre for Scientific Research (CNRS-INEE, CNRS-INSU). LFA was partly supported by a CNRS-INSU - MEEM convention n° 2101893310 for the MSFD implementation in France. AG and VD received funding from the German Federal Agency for Nature Conservation. CPL received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 869300.

## Appendix 1

**Table A1**

Building on IA2017, the biodiversity status of each indicator has been assessed according to the criteria described in Table 8. Red cells represent ‘poor’ status for an indicator in a region, while amber cells denote ‘uncertain’ status and green cells indicate ‘good’ status. Indicators were not assessed as part of IA2017 for cells in grey. Indicator icons are also used in Fig. 2.

Ecosystem component	Indicator	Greater North Sea	Celtic Seas	Bay of Biscay
Pelagic habitats	Changes in phytoplankton and zooplankton communities (PH1/FW5)	No threshold applied	No threshold applied	No threshold applied
	Changes in phytoplankton biomass and zooplankton abundance (PH2)	No threshold applied	No threshold applied	No threshold applied
	Changes in plankton diversity (PH3)	No threshold applied		No threshold applied
	Assessment of coastal habitats in relation to nutrient and/or organic enrichment (BH2-A)	Threshold applied (lack of representative data)	Threshold applied	Threshold applied
	Subtidal habitats of the southern North Sea (BH2-B)	No threshold applied		
	Extent of physical damage to predominant special habitats (BH3)	No threshold applied	No threshold applied	No threshold applied
	Recovery in the population abundance of sensitive fish species (FC1)	Threshold applied	Threshold applied	
	Proportion of large fish – Large Fish Index (FC2)	Threshold applied		
	Mean Maximum Length of fish (FC3)	No threshold applied	No threshold applied	No threshold applied
	Marine birds: Non-breeding abundance (B1-A)	Threshold applied	Threshold applied	
	Marine birds: Breeding abundance (B1-B)	Threshold applied	Threshold applied	
	Marine bird breeding success/failure (B3)	Threshold applied	Threshold applied	
	Seal abundance and distribution (M3)	Threshold applied	Threshold applied	
	Abundance and distribution of cetaceans (M4A): Killer whales	No threshold applied	No threshold applied	No threshold applied
	Abundance and distribution of cetaceans (M4A1): Coastal bottlenose dolphins	No threshold applied	No threshold applied	No threshold applied
	Abundance and distribution of cetaceans (M4B):	No threshold applied	No threshold applied	No threshold applied

	Cetaceans other than killer whales and bottlenose dolphins			
	Grey seal pup production (M5)	Threshold applied	Threshold applied	
	Harbour porpoise bycatch (M6)	No threshold applied	No threshold applied	
 Food webs	Production of phytoplankton (FW2)	No threshold applied	No threshold applied	
	Size composition in fish communities (FW3)	No threshold applied	No threshold applied	No threshold applied
	Change in mean trophic level of marine predators (FW4)			No threshold applied
 Non-indigenous species	Trends in new records of non-indigenous species introduced by human activities (NIS3)	No threshold applied	No threshold applied	No threshold applied

References

Airoidi, L., Beck, M.W., 2007. Loss, status and trends for coastal marine habitats of Europe. *Oceanogr. Mar. Biol.* 45, 345–405.

Arroyo, N.-L., Safi, G., Vouriot, P., López-López, L., Niquil, N., Le Loc'h, F., Hattab, T., Preciado, I., Coll, M., 2019. Towards coherent GES assessments at sub-regional level: signs of fisheries expansion processes in the Bay of Biscay using an OSPAR food web indicator, the mean trophic level. *ICES J. Mar. Sci.* 76 (6), 1543–1553.

Arroyo, N.L., Le Loc'h, F., Preciado, I., Niquil, N., Vouriot, P., Safi, G., 2017. Implementation of the mean trophic level indicator (MTL FW4) and assessment of its use at a sub-regional level (OSPAR region IV), as a contribution to the EU Co-financed EcAprRHA project (Applying an ecosystem approach to (sub) regional habitat assessments). Deliverable 3.1. OSPAR, London.

Batten, S.D., Clark, R., Flinkman, J., Hays, G., John, E., John, A.W.G., Jonas, T., Lindley, J.A., Stevens, D.P., Walne, A., 2003. CPR sampling: the technical background, materials and methods, consistency and comparability. *Prog. Oceanogr.* 58, 193–215. <https://doi.org/10.1016/j.pocan.2003.08.004>.

Beaugrand, G., 2004. The North Sea regime shift: Evidence, causes, mechanisms and consequences. *Prog. Oceanogr.* 60, 245–262. <https://doi.org/10.1016/j.pocan.2004.02.018>.

Bedford, J., Johns, D.G., McQuatters-Gollop, A., Ji, R., 2020a. Implications of taxon-level variation in climate change response for interpreting plankton lifeform biodiversity indicators. *ICES J. Mar. Sci.* 77 (7–8), 3006–3015.

Bedford, J., Ostle, C., Johns, D.G., Atkinson, A., Best, M., Bresnan, E., Machairopoulou, M., Graves, C.A., Devlin, M., Milligan, A., Pitois, S., Mellor, A., Tett, P., McQuatters-Gollop, A., 2020b. Lifeform indicators reveal large-scale shifts in plankton across the North-West European shelf. *Glob. Change Biol.* 26, 3482–3497. <https://doi.org/10.1111/gcb.15066>.

Bedford, J., Ostle, C., Johns, D.G., Budria, A., McQuatters-Gollop, A., 2020c. The influence of temporal scale selection on pelagic habitat biodiversity indicators. *Ecol. Ind.* 114, 106311. <https://doi.org/10.1016/j.ecolind.2020.106311>.

Bennema, F.P., Engelhard, G.H., Lindeboom, H., Norkko, J., 2020. *Ostrea edulis* beds in the central North Sea: delineation, ecology, and restoration. *ICES J. Mar. Sci.* 77 (7–8), 2694–2705.

Blumel, J.K., Fischer, S.H., Kulka, D.W., Lynam, C.P., Ellis, J.R., 2022. Decline in Atlantic wolffish *Anarhichas lupus* in the North Sea: Impacts of fishing pressure and climate change. *J. Fish Biol.* 100, 253–267. <https://doi.org/10.1111/jfb.14942>.

Borja, A., Garmendia, J.M., Menchaca, I., Uriarte, A., Sagarmínaga, Y., 2019. Yes, We Can! Large-Scale Integrative Assessment of European Regional Seas, Using Open Access Databases. *Front. Mar. Sci.* 6 <https://doi.org/10.3389/fmars.2019.00019>.

Brasseur, S.M.J.M., van Polanen Petel, T.D., Gerrodette, T., Meesters, E.H.W.G., Reijnders, P.J.H., Aarts, G., 2015. Rapid recovery of Dutch gray seal colonies fueled by immigration. *Mar. Mammal Sci.* 31, 405–426. <https://doi.org/10.1111/mms.12160>.

Brown, C.J., Smith, S.J., Lawton, P., Anderson, J.T., 2011. Benthic habitat mapping: A review of progress towards improved understanding of the spatial ecology of the seafloor using acoustic techniques. *Estuar. Coast. Shelf Sci.* 92, 502–520. <https://doi.org/10.1016/j.ecss.2011.02.007>.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L., 2001. *Introduction to Distance Sampling*. Oxford University Press, Oxford.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L., 2004. *Advanced distance sampling*. Oxford University Press, New York, USA.

Capuzzo, E., Lynam, C.P., Barry, J., Stephens, D., Forster, R.M., Greenwood, N., McQuatters-Gollop, A., Silva, T., Leeuwen, S.M.v., Engelhard, G.H., 2018. A decline in primary production in the North Sea over 25 years, associated with reductions in zooplankton abundance and fish stock recruitment. *Global Change Biol.* 24, e352–e364. <https://doi.org/10.1111/gcb.13916>.

Castro, M.C.T.d., Fileman, T.W., Hall-Spencer, J.M., 2017. Invasive species in the Northeastern and Southwestern Atlantic Ocean: A review. *Mar. Pollut. Bull.* 116 (1–2), 41–47.

Convention on Biological Diversity Secretariat, 2010. *The strategic plan for biodiversity 2011–2020 and the Aichi biodiversity targets*, Document UNEP/CBD/COP/DEC/X/2. Secretariat of the Convention on Biological Diversity, Nagoya, Japan.

Cook, A.S.C.P., Dadam, D., Mitchell, I., Ross-Smith, V.H., Robinson, R.A., 2014. Indicators of seabird reproductive performance demonstrate the impact of commercial fisheries on seabird populations in the North Sea. *Ecol. Ind.* 38, 1–11. <https://doi.org/10.1016/j.ecolind.2013.10.027>.

Danovaro, R., Carugati, L., Berzano, M., Cahill, A.E., Carvalho, S., Chenuil, A., Corinaldesi, C., Cristina, S., David, R., Dell'Anno, A., Dzhenbekova, N., Garcés, E., Gasol, J.M., Goela, P., Féral, J.-P., Ferrera, I., Forster, R.M., Kurekin, A.A., Rastelli, E., Marinova, V., Miller, P.L., Moncheva, S., Newton, A., Pearman, J.K., Pitois, S.G., René, A., Rodríguez-Espeleta, N., Saggiomo, V., Simis, S.G.H., Stefanova, K., Wilson, C., Lo Martire, M., Greco, S., Cochrane, S.K.J., Mangoni, O., Borja, A., 2016. Implementing and Innovating Marine Monitoring Approaches for Assessing Marine Environmental Status. *Front. Mar. Sci.* 3 <https://doi.org/10.3389/fmars.2016.00213>.

de Jonge, V.N., Schückel, U., 2021. A comprehensible short list of ecological network analysis indices to boost real ecosystem-based management and policy making. *Ocean Coast. Manag.* 208, 105582. <https://doi.org/10.1016/j.ocecoaman.2021.105582>.

de Juan, S., Demestre, M., 2012. A Trawl Disturbance Indicator to quantify large scale fishing impact on benthic ecosystems. *Ecol. Ind.* 18, 183–190. <https://doi.org/10.1016/j.ecolind.2011.11.020>.

de Juan, S., Hinz, H., Sartor, P., Vitale, S., Bentes, L., Bellido, J.M., Musumeci, C., Massi, D., Gancitano, V., Demestre, M., 2020. Vulnerability of Demersal Fish Assemblages to Trawling Activities: A Traits-Based Index. *Front. Mar. Sci.* 7 <https://doi.org/10.3389/fmars.2020.00044>.

Diaz, R.J., Solan, M., Valente, R.M., 2004. A review of approaches for classifying benthic habitats and evaluating habitat quality. *J. Environ. Manage.* 73, 165–181. <https://doi.org/10.1016/j.jenvman.2004.06.004>.

Dupaix, A., Mérillet, L., Kopp, D., Mouchet, M., Birchenough, S., 2021. Using biological traits to get insights into the benthic-demersal community sensitivity to trawling in the Celtic Sea. *ICES J. Mar. Sci.* 78 (3), 1063–1073.

Elliott, M., Borja, A., McQuatters-Gollop, A., Mazik, K., Birchenough, S., Andersen, J.H., Painting, S., Peck, M., 2015. *Force majeure*: will climate change affect our ability to attain Good Environmental Status for marine biodiversity? *Mar. Pollut. Bull.* 95, 7–27. <https://doi.org/10.1016/j.marpolbul.2015.03.015>.

Elliott, M., Burdon, D., Atkins, J.P., Borja, A., Cormier, R., de Jonge, V.N., Turner, R.K., 2017a. “And DPSIR begat DAPSI(W)R(M)” - A unifying framework for marine environmental management. *Mar. Pollut. Bull.* 118, 27–40. <https://doi.org/10.1016/j.marpolbul.2017.03.049>.

- Elliott, S.A.M., Arroyo, A.L., Safi, G., Ostle, C., Guérin, L., McQuatters-Gollop, A., Aubert, A., Artigas, F., Pesch, R., Schmitt, P., Vina-Herbon, C., Meakins, B., González-Irusta, J.M., Preciado, I., López-López, L., Punzón, A., de la Torre, A., Serrano, A., Haraldsson, M., Capuzzo, E., Clauquin, P., Kromkamp, J., Niquil, N., Judd, A., Padegimas, B., Corcoran, E., 2017b. Proposed approaches for indicator integration. *EcAprRHA deliverable WP4.1*, p. 26. ISBN: 978-1-911458-25-8.
- Elliott, S.A.M., Guérin, L., Pesch, R., Schmitt, P., Meakins, B., Vina-Herbon, C., González-Irusta, J.M., de la Torre, A., Serrano, A., 2018. Integrating benthic habitat indicators: Working towards an ecosystem approach. *Marine Policy* 90, 88–94. <https://doi.org/10.1016/j.marpol.2018.01.003>.
- Ellwood, H., 2014. Creating a EUNIS level 3 seabed habitat map integrating data originating from maps from fileds surveys and the EUSeaMap model. *JNCC Report, JNCC, Peterborough, UK*.
- Erbe, C., Dunlop, R., Dolman, S., 2018. Effects of Noise on Marine Mammals, in: Slabbekoorn, H., Dooling, R.J., Popper, A.N., Fay, R.R. (Eds.), *Effects of Anthropogenic Noise on Animals*. Springer New York, New York, NY, pp. 277–309. [https://doi.org/10.1007/978-1-4939-8574-6\\_10](https://doi.org/10.1007/978-1-4939-8574-6_10).
- Fath, B.D., Asmus, H., Asmus, R., Baird, D., Borrett, S.R., de Jonge, V.N., Ludovisi, A., Niquil, N., Scharler, U.M., Schückel, U., Wolff, M., 2019. Ecological network analysis metrics: The need for an entire ecosystem approach in management and policy. *Ocean Coast. Manag.* 174, 1–14. <https://doi.org/10.1016/j.ocecoaman.2019.03.007>.
- Frederiksen, M., Wanless, S., Harris, M.P., Rothery, P., Wilson, L.J., 2004. The role of industrial fisheries and oceanographic change in the decline of North Sea black-legged kittiwakes. *J. Appl. Ecol.* 41, 1129–1139. <https://doi.org/10.1111/j.0021-8901.2004.00966.x>.
- Galatus, A., Brackmann, J., Brasseur, S.M., Diederichs, B., Jeß, A., Klöpffer, S., Körber, P., Schop, J., Siebert, U., Teilmann, J., Thøstesen, B., Schmidt, B., 2021. EG-Marine Mammals harbour seal surveys in the Wadden Sea and Helgoland 2021, in: Secretariat, C.W.S. (Ed.). *Common Wadden Sea Secretariat, Wilhelmshaven, Germany*.
- González-Irusta, J.M., De la Torre, A., Punzón, A., Blanco, M., Serrano, A., Birchenough, S., 2018. Determining and mapping species sensitivity to trawling impacts: the Benthos Sensitivity Index to Trawling Operations (BESITO). *ICES J. Mar. Sci.* 75 (5), 1710–1721.
- Greathead, C., Magni, P., Vanaverbeke, J., Buhl-Mortensen, L., Janas, U., Blomqvist, M., Craeymeersch, J.A., Dannheim, J., Darr, A., Degraer, S., Desroy, N., Donnay, A., Griffiths, Y., Guala, I., Guerin, L., Hinchey, H., Labruno, C., Reiss, H., Van Hoey, G., Birchenough, S.N.R., 2020. A generic framework to assess the representation and protection of benthic ecosystems in European marine protected areas. *Aquat. Conserv. Mar. Freshwater Ecosyst.* 30, 1253–1275. <https://doi.org/10.1002/aqc.3401>.
- Greenstreet, S., Moriarty, M., 2017. Manual for Version 3 of the Groundfish Survey Monitoring and Assessment Data Product. *Scottish Mar. Freshw. Sci.* 8 <https://doi.org/10.7489/1986-1>.
- Greenstreet, S.P.R., Rogers, S.I., 2006. Indicators of the health of the North Sea fish community: identifying reference levels for an ecosystem approach to management. *ICES J. Mar. Sci.* 63, 573–593. <https://doi.org/10.1016/j.icesjms.2005.12.009>.
- Greenstreet, S.P.R., Rogers, S.I., Rice, J.C., Piet, G.J., Guirey, E.J., Fraser, H.M., Fryer, R. J., 2010. Development of the EcoQO for the North Sea fish community. *ICES J. Mar. Sci.* 68, 1–11. <https://doi.org/10.1093/icesjms/fsq156>.
- Greenstreet, S.P.R., Rossberg, A.G., Fox, C.J., Le Quesne, W.J.F., Blasdale, T., Boulcott, P., Mitchell, I., Millar, C., Moffat, C.F., 2012. Demersal fish biodiversity: species-level indicators and trends-based targets for the Marine Strategy Framework Directive. *ICES J. Mar. Sci.* 69, 1789–1801. <https://doi.org/10.1093/icesjms/fss148>.
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A Global Map of Human Impact on Marine Ecosystems. *Science* 319, 948–952. <https://doi.org/10.1126/science.1149345>.
- Hammond, P.S., Macleod, K., Berggren, P., Borchers, D.L., Burt, L., Cañadas, A., Desportes, G., Donovan, G.P., Gilles, A., Gillespie, D., Gordon, J., Hiby, L., Kuklik, I., Leaper, R., Lehnert, K., Leopold, M., Lovell, P., Oien, N., Paxton, C.G.M., Ridoux, V., Rogan, E., Samarra, F., Scheidat, M., Sequeira, M., Siebert, U., Skov, H., Swift, R., Tasker, M.L., Teilmann, J., Van Canneyt, O., Vázquez, J.A., 2013. Cetacean abundance and distribution in European Atlantic shelf waters to inform conservation and management. *Biol. Conserv.* 164, 107–122. <https://doi.org/10.1016/j.biocon.2013.04.010>.
- Hardisty, A.R., Michener, W.K., Agosti, D., Alonso García, E., Bastin, L., Belbin, L., Bowser, A., Buttigieg, P.L., Canhos, D.A.L., Egloff, W., De Giovanni, R., Figueira, R., Groom, Q., Guralnick, R.P., Hobern, D., Hugo, W., Koureas, D., Ji, L., Los, W., Manuel, J., Manset, D., Poelen, J., Saarenmaa, H., Schigel, D., Uhlir, P.F., Kissling, W.D., 2019. The Bari Manifesto: An interoperability framework for essential biodiversity variables. *Ecol. Inf.* 49, 22–31. <https://doi.org/10.1016/j.ecoinf.2018.11.003>.
- Heiskanen, A.-S., Berg, T., Uusitalo, L., Teixeira, H., Bruhn, A., Krause-Jensen, D., Lynam, C.P., Rossberg, A.G., Korpinen, S., Uyarra, M.C., Borja, A., 2016. Biodiversity in Marine Ecosystems—European Developments toward Robust Assessments. *Front. Mar. Sci.* 3 <https://doi.org/10.3389/fmars.2016.00184>.
- ICES, 2020. *ICES Ecosystem Overview: North Sea Ecoregion, ICES Advice 2020*. ICES, Copenhagen, Denmark <https://doi.org/10.17895/ices.advice.7632>.
- ICES, 2021a. International Bottom Trawl Survey Working Group (IBTSWG). *ICES Scientific Reports* 3 (69), 201. <https://doi.org/10.17895/ices.pub.8219>.
- ICES, 2021b. Working Group on Beam Trawl Surveys (WGBEAM). *ICES Scientific Reports* 3 (46), 89. <https://doi.org/10.17895/ices.pub.8114>.
- ICES, 2016b. OSPAR request for further development of fishing intensity and pressure mapping, in: ICES (Ed.). *ICES, Copenhagen, Denmark*, p. 28.
- ICES, 2013. Report of the Working Group on Biodiversity Science (WGBIODIV), 18–22 February 2013. *ICES CM 2013/SSGEF:02*, Copenhagen, Denmark.
- ICES, 2014b. Report of the Workshop to develop recommendations for potentially useful Food Web Indicators (WKFooWI), 31 March–3 April 2014. *ICES CM 2014\ACOM:48* Copenhagen, Denmark.
- ICES, 2014a. OSPAR request on implementation of MSFD for marine mammals, Report of the ICES Advisory Committee, 2014. *ICES Advice 2014, Book 1, Section 1.6.6.1*. ICES, Copenhagen, Denmark.
- ICES, 2015a. Bycatch of small cetaceans and other marine animals – Review of national reports under Council Regulation (EC) No. 812/2004 and other published documents. *ICES Advice 2015, Book 1, 1.6.1.1: Advice Northeast Atlantic and adjacent seas* Published 15 April 2015. ICES, Copenhagen, Denmark.
- ICES, 2015b. Bycatch of small cetaceans and other marine animals – Review of national reports under Council Regulation (EC) No. 812/2004 and other published documents. *ICES Advice 2015, Book 1, 1.6.1.1: Advice Northeast Atlantic and adjacent seas* Published 15 April 2015. ICES, Copenhagen, Denmark.
- ICES, 2016a. Bycatch of small cetaceans and other marine animals – Review of national reports under Council Regulation (EC) No. 812/2004 and other published documents. *ICES Advice 2016, Book 1, 1.6.1.1: Advice Northeast Atlantic and adjacent seas* Published 15 April 2016. ICES, Copenhagen, Denmark.
- ICES, 2017. Report of the Joint OSPAR/HELCOM/ICES Working Group on Seabirds (JWGBIRD), 6–10 November 2017, Riga, Latvia. *ICES CM 2017\ACOM:49*. 97 pp. ICES, Copenhagen, Denmark.
- IPBES, 2019. Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, in: E. S. Brondizio, J.S., S. Díaz, and H. T. Ngo (Ed.). *IPBES secretariat, Bonn, Germany*.
- IPCC, 2019. Summary for Policymakers, in: Pörtner, H.-O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., Weyer, N.M. (Eds.), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. IPCC, p. 36.
- Jennings, S., Lee, J., Hiddink, J.G., 2012. Assessing fishery footprints and the trade-offs between landings value, habitat sensitivity, and fishing impacts to inform marine spatial planning and an ecosystem approach. *ICES J. Mar. Sci.* 69, 1053–1063. <https://doi.org/10.1093/icesjms/fss050>.
- Jepson, P.D., Deaville, R., Barber, J.L., Aguilar, A., Borrell, A., Murphy, S., Barry, J., Brownlow, A., Barnett, J., Berrow, S., Cunningham, A.A., Davison, N.J., ten Doeschate, M., Esteban, R., Ferreira, M., Foote, A.D., Genov, T., Giménez, J., Loveridge, J., Llavona, A., Martin, V., Maxwell, D.L., Papachlimitzou, A., Penrose, R., Perkins, M.W., Smith, B., de Stephanis, R., Tregenza, N., Verborgh, P., Fernandez, A., Law, R.J., 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.* 6, 18573. <https://doi.org/10.1038/srep18573>.
- Kenny, A.J., Cato, I., Desprez, M., Fader, G., Schüttenhelm, R.T.E., Side, J., 2003. An overview of seabed-mapping technologies in the context of marine habitat classification. *ICES J. Mar. Sci.* 60, 411–418. [https://doi.org/10.1016/s1054-3139\(03\)00006-7](https://doi.org/10.1016/s1054-3139(03)00006-7).
- Kromkamp, J., Capuzzo, E., Philippart, C.J.M., 2017. Measuring Phytoplankton Primary Production: Review of Existing Methodologies and Suggestions for a Common Approach: EcAprRHA Deliverable WP 3.2. OSPAR.
- Link, J.S., Watson, R.A., 2019. Global ecosystem overfishing: Clear delineation within real limits to production. *Sci. Adv.* 5, eaav0474. <https://doi.org/10.1126/sciadv.aav0474>.
- Lockerbie, E.M., Lynam, C.P., Shannon, L.J., Jarre, A., Link, J., 2018. Applying a decision tree framework in support of an ecosystem approach to fisheries: IndiSeas indicators in the North Sea. *ICES J. Mar. Sci.* 75 (3), 1009–1020.
- Lotze, H.K., Flemming, J.M., Magera, A.M., 2017. Critical factors for the recovery of marine mammals. *Conserv. Biol.* 31, 1301–1311. <https://doi.org/10.1111/cobi.12957>.
- Lynam, C.P., Uusitalo, L., Patrício, J., Piroddi, C., Queirós, A.M., Teixeira, H., Rossberg, A.G., Sagarmingaya, Y., Hyder, K., Niquil, N., Möllmann, C., Wilson, C., Chust, G., Galparsoro, I., Forster, R., Veríssimo, H., Tedesco, L., Revilla, M., Neville, S., 2016. Uses of Innovative Modeling Tools within the Implementation of the Marine Strategy Framework Directive. *Front. Mar. Sci.* 3 <https://doi.org/10.3389/fmars.2016.00182>.
- Mackey, R.L., Currie, D.J., 2001. The Diversity-Disturbance Relationship: Is It Generally Strong and Peaked? *Ecology* 82, 3479–3492. <https://doi.org/10.2307/2680166>.
- McQuatters-Gollop, A., Raitos, D.E., Edwards, M., Pradhan, Y., Mee, L.D., Lavender, S.J., Attrill, M.J., 2007. A long-term chlorophyll dataset reveals regime shift in North Sea phytoplankton biomass unconnected to nutrient levels. *Limnol. Oceanogr.* 52, 635–648. <https://doi.org/10.4319/lo.2007.52.2.0635>.
- McQuatters-Gollop, A., Atkinson, A., Aubert, A., Bedford, J., Best, M., Bresnan, E., Cook, K., Devlin, M., Gowen, R., Johns, D.G., Machairoupolou, M., McKinney, A., Mellor, A., Ostle, C., Scherer, C., Tett, P., 2019a. Plankton lifeforms as a biodiversity indicator for regional-scale assessment of pelagic habitats for policy. *Ecol. Ind.* 101, 913–925.
- McQuatters-Gollop, A., Mitchell, I., Vina-Herbon, C., Bedford, J., Addison, P.F.E., Lynam, C.P., Geetha, P.N., Vermeulan, E.A., Smit, K., Bayley, D.T.I., Morris-Webb, E., Niner, H.J., Otto, S.A., 2019b. From Science to Evidence – How Biodiversity Indicators Can Be Used for Effective Marine Conservation Policy and Management. *Front. Mar. Sci.* 6 <https://doi.org/10.3389/fmars.2019.00109>.
- Mendel, B., Schwemmer, P., Peschko, V., Müller, S., Schwemmer, H., Mercker, M., Garthe, S., 2019. Operational offshore wind farms and associated ship traffic cause

- profound changes in distribution patterns of Loons (*Gavia spp.*). *J. Environ. Manage.* 231, 429–438. <https://doi.org/10.1016/j.jenvman.2018.10.053>.
- Mercker, M., Dierschke, V., Camphuysen, K., Kreutle, A., Markones, N., Vanermen, N., Garthe, S., 2021. An indicator for assessing the status of marine-bird habitats affected by multiple human activities: A novel statistical approach. *Ecol. Ind.* 130, 108036. <https://doi.org/10.1016/j.ecolind.2021.108036>.
- Moriarty, M., Greenstreet, S.P.R., Rasmussen, J., de Boois, I., 2019. Derivation of Groundfish Survey Monitoring and Assessment Data Products for the Northeast Atlantic Area. *Scottish Mar. Freshw. Sci. Rep.* 8 <https://doi.org/10.7489/1984-1>.
- Moriarty, M., Greenstreet, S.P.R., Rasmussen, J., 2017. Assessing the State of Demersal Fish to Address Formal Ecosystem Based Management Needs: Making Fisheries Independent Trawl Survey Data 'Fit for Purpose'. *Front. Mar. Sci.* 6 <https://doi.org/10.3389/fmars.2019.00162>.
- Murphy, S., Barber, J.L., Learmonth, J.A., Read, F.L., Deaville, R., Perkins, M.W., Brownlow, A., Davison, N., Penrose, R., Pierce, G.J., Law, R.J., Jepson, P.D., 2015. Reproductive Failure in UK Harbour Porpoises *Phocoena phocoena*: Legacy of Pollutant Exposure? *PLOS ONE* 10, e0131085. <https://doi.org/10.1371/journal.pone.0131085>.
- NOAA, 2022. State of the Ecosystem 2022: Mid-Atlantic p. 48. <https://doi.org/10.25923/jd1w-dc26>.
- Noël, L.-M.-L.-J., Griffin, J.N., Moschella, P.S., Jenkins, S.R., Thompson, R.C., Hawkins, S.J., 2009. Changes in Diversity and Ecosystem Functioning During Succession. In: Wahl, M. (Ed.), *Marine Hard Bottom Communities: Patterns, Dynamics, Diversity, and Change*. Springer, Berlin Heidelberg, Berlin, Heidelberg, pp. 213–223. <https://doi.org/10.1007/b76710.15>.
- Nogues, Q., Raoux, A., Araignous, E., Chaalali, A., Hattab, T., Leroy, B., Ben Rais Lasram, F., David, V., Le Loc'h, F., Dauvin, J.-C., Niquil, N., 2021. Cumulative effects of marine renewable energy and climate change on ecosystem properties: Sensitivity of ecological network analysis. *Ecol. Ind.* 121, 107128.
- OSPAR, 2011. Report of the OSPAR/MSFD workshop on approaches to determining GES for biodiversity. ISBN 978-1-907390-94-4, Publication Number: 553/2011. 55 p. <https://www.ospar.org/meetings/archive/ospar-msfd-workshop-on-approaches-to-determining-ges-for-biodiversity>.
- OSPAR, 2012. OSPAR MSFD Advice Manual and Background Document on Biodiversity. Approaches to determining good environmental status, setting of environmental targets and selecting indicators for Marine Strategy Framework Directive descriptors 1, 2, 4 and 6. OSPAR Agreement 2012-3. 141 pp. Available at: <https://www.ospar.org/documents?d=7287>.
- OSPAR, 2017m. Intermediate Assessment 2017. Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017>.
- OSPAR, 2018a. OSPAR CEMP Guideline. Combined guideline for the common indicators FC1, FC2. In: FC3 and FW3 for fish and food webs, p. 5 pp.
- OSPAR, 2018b. OSPAR CEMP Guideline. Common indicator BH2: Condition of Benthic Habitat Communities. Available at: the Common Conceptual Approach 60, pp <https://www.ospar.org/documents?v=39000>.
- OSPAR, 2018c. Available at: OSPAR CEMP Guideline. Common indicator FW4: Change in average trophic level of marine predators. 19, pp <https://www.ospar.org/documents?v=39002>.
- Ospar Ospar OSPAR Northeast Atlantic Environment Strategy 2021 OSPAR Commission, London.
- OSPAR, 2016a. OSPAR CEMP Guideline. Common indicator B1: Marine Bird Abundance. OSPAR Agreement 2016-09. 26 pp. Available at: <https://www.ospar.org/documents?v=38978>.
- OSPAR, 2016b. OSPAR CEMP Guideline. Common indicator B3: Marine bird breeding success/failure. OSPAR Agreement 2016-10. 24 pp. Available at: <https://www.ospar.org/documents?v=38979>.
- OSPAR, 2017a. B1: Marine Bird Abundance, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/bird-abundance/>.
- OSPAR, 2017b. B3: Marine Bird Breeding Success / Failure, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-birds/marine-bird-breeding-success-failure/>.
- OSPAR, 2017c. BH2: Condition of Benthic Habitat Communities: the Common Conceptual Approach, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/common-conceptual-approach/>.
- OSPAR, 2017d. BH2a: Condition of Benthic Habitat Communities: Assessment of Coastal Habitats in relation to Nutrient and/or Organic Enrichment, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/condition-benthic-habitat-communities-assessment-coastal-habita/>.
- OSPAR, 2017e. BH2b: Condition of Benthic Habitat Communities: Subtidal Habitats of the Southern North Sea, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/condition-of-benthic-habitat-defining-communities/subtidal-habitats-southern-north-sea/>.
- OSPAR, 2017f. BH3: Extent of Physical Damage to Predominant and Special Habitats, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/extent-physical-damage-predominant-and-special-habitats/>.
- OSPAR, 2017g. FC1: Recovery in the Population Abundance of Sensitive Fish Species, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/recovery-sensitive-fish/>.
- OSPAR, 2017h. FC2: Proportion of Large Fish (Large Fish Index), in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/proportion-large-fish-large-fish-index/>.
- OSPAR, 2017i. FC3: Pilot Assessment of Mean Maximum Length of Fish, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/mean-maximum-length/>.
- OSPAR, 2017j. FW2: Pilot Assessment of Production of Phytoplankton, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/phytoplankton-production/>.
- OSPAR, 2017k. FW3: Size Composition in Fish Communities, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/size-fish-composition/>.
- OSPAR, 2017l. FW4: Change in Average Trophic Level of Marine Predators in the Bay of Biscay, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/fish-and-food-webs/mtl-bay-biscay/>.
- OSPAR, 2017n. M3: Seal Abundance and Distribution, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/seal-abundance-and-distribution/>.
- OSPAR, 2017o. M4: Abundance and Distribution of Cetaceans, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/abundance-distribution-cetaceans/>.
- OSPAR, 2017p. M5: Grey Seal Pup Production, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/grey-seal-pup/>.
- OSPAR, 2017q. M6: Harbour Porpoise Bycatch, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/marine-mammals/harbour-porpoise-bycatch/>.
- OSPAR, 2017r. NIS3: Trends in New Records of Non-Indigenous Species Introduced by Human Activities, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/pressures-human-activities/non-indigenous/>.
- OSPAR, 2017s. OSPAR CEMP Guideline. Common indicator: BH3 Extent of Physical damage to predominant and special habitats. 61 pp. Available at: <https://www.ospar.org/documents?v=37641>.
- OSPAR, 2017t. PH1/FW5: Changes in phytoplankton and zooplankton communities, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/>.
- OSPAR, 2017u. PH2: Changes in phytoplankton biomass and zooplankton abundance, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/>.
- OSPAR, 2017v. PH3: Pilot assessment of changes in plankton diversity, in: OSPAR (Ed.), OSPAR Intermediate Assessment OSPAR 2017 London UK Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/pilot-assessment-changes-plankton/>.
- OSPAR, 2018d. OSPAR CEMP Guideline. Common indicator M3: Seal abundance and distribution. 22 pp. Available at: <https://www.ospar.org/documents?v=38980>.
- OSPAR, 2018e. OSPAR CEMP Guideline. Common indicator M4: Abundance and distribution at the relevant spatio-temporal scale of cetacean species regularly present. 9 pp. Available at: <https://www.ospar.org/documents?v=39019>.
- OSPAR, 2018f. OSPAR CEMP Guideline. Common indicator M5: Grey seal pup production. 17 pp. Available at: <https://www.ospar.org/documents?v=38981>.
- OSPAR, 2018g. OSPAR CEMP Guideline. Common indicator NIS3: Changes to non-indigenous species communities. OSPAR Agreement 2018-04. 8 pp. Available at: <https://www.ospar.org/documents?v=38992>.
- OSPAR, 2018h. OSPAR CEMP Guideline. Common indicator PH1/FW5: Plankton lifetimes. OSPAR Agreement 2018-07. 17 pp. Available at: <https://www.ospar.org/documents?v=39001>.
- OSPAR, 2019a. OSPAR CEMP Guideline. Common indicator PH2: Changes in Phytoplankton Biomass and Zooplankton Abundance. OSPAR Agreement 2019-06. 6 pp. Available at: <https://www.ospar.org/documents?d=40972>.
- OSPAR, 2019b. OSPAR CEMP Guideline. Common indicator PH3: Changes in Plankton Diversity. OSPAR Agreement 2019-07. 7 pp. Available at: <https://www.ospar.org/documents?d=40973>.
- Padegimas, B., Artigas, F., Arroyo, N.L., Aubert, A., Budria, A., Capuzzo, E., Corcoran, E., Elliott, S.A.M., González-Irusta, J.M., Guérin, L., Judd, A., Kromkamp, J., McQuatters-Gollop, A., Meakins, B., Niquil, N., Ostle, C., Pesch, R., Preciado, I., Safi, G., Schmitt, P., Serrano, A., Thorpe, R., Torriente, A., Vina-Herbon, C., 2017. Action plan for the further implementation of habitat and food web indicators and progressing integrated assessments in OSPAR (sub)regions. EcAprHA deliverable WP5.6, p. 20. ISBN: 978-1-911458-30-2.

- Piroddi, C., Akoglu, E., Andonegi, E., Bentley, J.W., Celić, I., Coll, M., Dimarchopoulou, D., Friedland, R., de Mutser, K., Girardin, R., Garcia-Gorriz, E., Grizzetti, B., Hervann, P.-Y., Heymans, J.J., Müller-Karulis, B., Libralato, S., Lynam, C.P., Macias, D., Miladinova, S., Moullec, F., Paliolax, A., Parn, O., Serpetti, N., Solidoro, C., Steenbeek, J., Stips, A., Tomczak, M.T., Travers-Trolet, M., Tsikliras, A.C., 2021. Effects of Nutrient Management Scenarios on Marine Food Webs: A Pan-European Assessment in Support of the Marine Strategy Framework Directive. *Front. Mar. Sci.* 8 <https://doi.org/10.3389/fmars.2021.596797>.
- Pitcher, C.R., Hiddink, J.G., Jennings, S., Collie, J., Parma, A.M., Amoroso, R., Mazor, T., Sciberras, M., McConnaughey, R.A., Rijnsdorp, A.D., Kaiser, M.J., Suuronen, P., Hilborn, R., 2022. Trawl impacts on the relative status of biotic communities of seabed sedimentary habitats in 24 regions worldwide. *Proceedings of the National Academy of Sciences* 119, e2109449119. <https://doi.org/10.1073/pnas.2109449119>.
- Pitois, S.G., Graves, C.A., Close, H., Lynam, C., Scott, J., Tilbury, J., van der Kooij, J., Culverhouse, P., 2021. A first approach to build and test the Copepod Mean Size and Total Abundance (CMSTA) ecological indicator using in-situ size measurements from the Plankton Imager (PI). *Ecol. Ind.* 123, 107307. <https://doi.org/10.1016/j.ecolind.2020.107307>.
- Preciado, I., Arroyo, N.L., González-Irusta, J.M., López-López, L., Punzón, A., Muñoz, I., Serrano, A., 2019. Small-scale spatial variations of trawling impact on food web structure. *Ecol. Ind.* 98, 442–452. <https://doi.org/10.1016/j.ecolind.2018.11.024>.
- Queirós, A.M., Fernandes, J., Genevier, L., Lynam, C.P., 2018. Climate change alters fish community size-structure, requiring adaptive policy targets. *Fish Fish.* 19, 613–621. <https://doi.org/10.1111/faf.12278>.
- Raoux, A., Lassalle, G., Pezy, J.-P., Tecchio, S., Safi, G., Ernande, B., Mazé, C., Loc'h, F.L., Lequesne, J., Girardin, V., Dauvin, J.-C., Niquil, N., 2019. Measuring sensitivity of two OSPAR indicators for a coastal food web model under offshore wind farm construction. *Ecol. Ind.* 96, 728–738. <https://doi.org/10.1016/j.ecolind.2018.07.014>.
- Reynolds III, J.E., Marsh, H., Ragen, T.J., 2009. Marine mammal conservation. *Endangered Species Res.* 7, 23–28. <https://doi.org/10.3354/esr00179>.
- Roberts, C., 2007. *The Unnatural History of the Sea: The past and future of humanity and fishing*. Gaia, London.
- Rombouts, I., Simon, N., Aubert, A., Cariou, T., Feunteun, E., Guérin, L., Hoebeke, M., McQuatters-Gollop, A., Rigaut-Jalabert, F., Artigas, L.F., 2019. Changes in marine phytoplankton diversity: Assessment under the Marine Strategy Framework Directive. *Ecol. Ind.* 102, 265–277. <https://doi.org/10.1016/j.ecolind.2019.02.009>.
- Rondinini, C., 2011. A review of methodologies that could be used to formulate ecologically meaningful targets for marine habitat coverage within the UK MPA network. *JNCC Report No. 438*.
- Rovellini, A., Shaffer, M.R., 2020. Quantitative targets for marine protection: a review of the scientific basis and applications. Prepared for the Department of Conservation, Wellington, New Zealand. DOC Project 4792.
- Safi, G., Giebels, D., Arroyo, N.L., Heymans, J.J., Preciado, I., Raoux, A., Schückel, U., Tecchio, S., de Jonge, V.N., Niquil, N., 2019. Vitamine ENA: A framework for the development of ecosystem-based indicators for decision makers. *Ocean Coast. Manag.* 174, 116–130. <https://doi.org/10.1016/j.ocecoaman.2019.03.005>.
- Sahlmann, A., Wolf, R., Holth, T.F., Titelman, J., Hylland, K., 2017. Baseline and oxidative DNA damage in marine invertebrates. *J. Toxicol. Environ. Health Part A* 80, 807–819. <https://doi.org/10.1080/15287394.2017.1352179>.
- Samhoury, J.F., Levin, P.S., Ainsworth, C.H., Bograd, S.J., 2010. Identifying Thresholds for Ecosystem-Based Management. *PLoS ONE* 5 (1), e8907.
- Schaefer, M.B., 1991. Some aspects of the dynamics of populations important to the management of the commercial marine fisheries. *Bull. Math. Biol.* 53, 253–279. [https://doi.org/10.1016/S0092-8240\(05\)80049-7](https://doi.org/10.1016/S0092-8240(05)80049-7).
- Schwemmer, P., Mendel, B., Sonntag, N., Dierschke, V., Garthe, S., 2011. Effects of ship traffic on seabirds in offshore waters: implications for marine conservation and spatial planning. *Ecol. Appl.* 21, 1851–1860. <https://doi.org/10.1890/10.0615.1>
- Secretariat of the Convention on Biological Diversity, 2014. An Updated Synthesis of the Impacts of Ocean Acidification on Marine Biodiversity. Technical Series No. 75., in: S. Hennige, J.M.R.P.W. (Ed.) *Convention on Biological Diversity*, Montreal, p. 99p.
- Sguotti, C., Lynam, C.P., Garcia-Carreras, B., Ellis, J.R., Engelhard, G.H., 2016. Distribution of skates and sharks in the North Sea: 112 years of change. *Glob. Change Biol.* 22, 2729–2743. <https://doi.org/10.1111/gcb.13316>.
- Spence, M.A., Griffiths, C.A., Waggit, J.J., Bannister, H.J., Thorpe, R.B., Rossberg, A.G., Lynam, C.P., 2021. Sustainable fishing can lead to improvements in marine ecosystem status: an ensemble-model forecast of the North Sea ecosystem. *Mar. Ecol. Prog. Ser.* 680, 207–221.
- Thompson, M.S.A., Pontalier, H., Spence, M.A., Pinnegar, J.K., Greenstreet, S.P.R., Moriarty, M., Hélaouët, P., Lynam, C.P., O'Connor, N., 2020. A feeding guild indicator to assess environmental change impacts on marine ecosystem structure and functioning. *J. Appl. Ecol.* 57 (9), 1769–1781.
- Thorpe, R.B., De Oliveira, J.A.A., 2019. Comparing conceptual frameworks for a fish community MSY (FCMSY) using management strategy evaluation—an example from the North Sea. *ICES J. Mar. Sci.* 76, 813–823. <https://doi.org/10.1093/icesjms/fsz015>.
- Thorpe, R.B., Arroyo, N.L., Safi, G., Niquil, N., Preciado, I., Heath, M., Pace, M.C., Lynam, C.P., 2022. The Response of North Sea Ecosystem Functional Groups to Warming and Changes in Fishing. *Front. Mar. Sci.* 9 <https://doi.org/10.3389/fmars.2022.841909>.
- Tidbury, H.J., Taylor, N.G.H., Copp, G.H., Garnacho, E., Stebbing, P.D., 2016. Predicting and mapping the risk of introduction of marine non-indigenous species into Great Britain and Ireland. *Biol. Invasions* 18, 3277–3292. <https://doi.org/10.1007/s10530-016-1219-x>.
- H. Tillin H. Tyler-Walters Assessing the sensitivity of subtidal sedimentary habitats to pressures associated with marine activities - Phase 1 Report, JNCC Report 512 2014 <http://jncc.defra.gov.uk/page-6790>.
- Torbjørn, J., Kari, N., Frode, O., 2003. The North-east Atlantic Ocean - huge, deep and heavily exploited. *European Environment Agency*, p. 21.
- Tsiamis, K., Paliolax, A., Stefanova, K., Gladan, Ž.N., Skejić, S., Despalatović, M., Cvitković, I., Dragičević, B., Dulčić, J., Vidjak, O., Bojanić, N., Žuljević, A., Aplikioti, M., Argyrou, M., Josephides, M., Michailidis, N., Jakobsen, H.H., Staehr, P. A., Ojaveer, H., Lehtiniemi, M., Massé, C., Zenetos, A., Castriota, L., Livi, S., Mazziotti, C., Schembri, P.J., Evans, J., Bartolo, A.G., Kabuta, S.H., Smolders, S., Knežtering, E., Gittenberger, A., Gruszka, P., Kraśniewski, W., Bartilotti, C., Tuaty-Guerra, M., Canning-Clode, J., Costa, A.C., Parente, M.I., Botelho, A.Z., Micael, J., Miodonski, J.V., Carreira, G.P., Lopes, V., Chainho, P., Barberá, C., Naddafi, R., Florin, A.-B., Barry, P., Stebbing, P.D., Cardoso, A.C., 2019. Non-indigenous species refined national baseline inventories: A synthesis in the context of the European Union's Marine Strategy Framework Directive. *Mar. Pollut. Bull.* 145, 429–435. <https://doi.org/10.1016/j.marpolbul.2019.06.012>.
- United Nations, 2015. *Sustainable Development Goals: 17 Goals to transform our world*. United Nations.
- Uusitalo, L., Blanchet, H., Andersen, J.H., Beauchard, O., Berg, T., Bianchelli, S., Cantafaro, A., Carstensen, J., Carugati, L., Cochrane, S., Danovaro, R., Heiskanen, A.-S., Karvinen, V., Moncheva, S., Murray, C., Neto, J.M., Nygård, H., Pantazi, M., Papadopoulou, N., Simbora, N., Srébalienė, G., Uyarra, M.C., Borja, A., 2016. Indicator-Based Assessment of Marine Biological Diversity-Lessons from 10 Case Studies across the European Seas. *Front. Mar. Sci.* 3 <https://doi.org/10.3389/fmars.2016.00159>.
- S. van Leeuwen P. Tett D. Mills Molen, J.v.d., Stratified and nonstratified areas in the North Sea: Long-term variability and biological and policy implications *Journal of Geophysical Research Oceans* 120 2015 4670 4686 <https://doi.org/doi:10.1002/2014JC010485>.
- van Loon, W.M.G.M., Walvoort, D.J.J., van Hoey, G., Vina-Herbon, C., Blandon, A., Pesch, R., Schmitt, P., Scholle, J., Heyer, K., Lavaley, M., Phillips, G., Duineveld, G. C.A., Blomqvist, M., 2018. A regional benthic fauna assessment method for the Southern North Sea using Margalef diversity and reference value modelling. *Ecol. Ind.* 89, 667–679. <https://doi.org/10.1016/j.ecolind.2017.09.029>.