



HAL
open science

Redefining crop breeding strategy for effective use of nitrogen in cropping systems

Ignacio Ciampitti, Jean-Francois Briat, Francois Gastal, Gilles Lemaire

► To cite this version:

Ignacio Ciampitti, Jean-Francois Briat, Francois Gastal, Gilles Lemaire. Redefining crop breeding strategy for effective use of nitrogen in cropping systems. *Communications Biology*, 2022, 5 (1), pp.1-4. 10.1038/s42003-022-03782-2 . hal-03837220

HAL Id: hal-03837220

<https://hal.inrae.fr/hal-03837220>

Submitted on 7 Jun 2023




HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Redefining crop breeding strategy for effective use of nitrogen in cropping systems

Ignacio A. Ciampitti ^{1✉}, Jean-Francois Briat ², Francois Gastal³ & Gilles Lemaire ^{4✉}

More than a half-century after the initial conceptualization of nitrogen use efficiency, crop improvement progress for direct nitrogen gains remains an elusive target. A more relevant conceptual framework bridging soil and plant processes is needed for crop improvement and environmental stewardship.

Nitrogen (N) is a critical element to guarantee global food security and to reduce the environmental footprint of agriculture¹. Maintaining a balance between N applied and crop N harvested is critical to minimize the consequences of global N losses². The traditional metric, N use efficiency (NUE), refers to the responsiveness of crops to N fertilization^{3,4}. However, a myriad of indices for estimating NUE have been proposed^{5,6}, complicating the quantification of true N gains and comparison across cropping systems^{7,8}. Recently, a perspective summary of NUE indices stressed the need to rethink this index relative to research goals while targeting global productivity and sustainability⁹.

Traditional NUE: a metric for evaluating N fertilization efficiency

Crop NUE is traditionally defined as the ratio of the supplement of grain yield (ΔY) or aboveground biomass (ΔW) to the supplement of N fertilizer application (ΔNf)⁴. NUE can be separated into N uptake efficiency (NupE) and N conversion efficiency (NCE):

$$NUE = \Delta Y / \Delta Nf. \quad (1)$$

N uptake efficiency, NupE as the N taken up by the crop (ΔNup) per unit of ΔNf :

$$NupE = \Delta Nup / \Delta Nf. \quad (2)$$

N conversion efficiency, NCE as the ΔW per unit of ΔNup :

$$NCE = \Delta W / \Delta Nup. \quad (3)$$

Harvest Index, HI is the increase in ΔY to increase in ΔW :

$$HI = \Delta Y / \Delta W. \quad (4)$$

$$\text{Therefore, } NUE = NupE \times NCE \times HI. \quad (5)$$

The major drawback of these equations is that N uptake is co-regulated by both soil N availability and potential plant growth rate¹⁰, which determine root N absorption capacity. Thus, the dissection of NUE into NupE and NCE does not allow a relevant analysis of mutual roles of soil and/or plant processes. Consequently, NUE cannot be interpreted as a general property of a given genotype to efficiently use all its N resources.

¹Department of Agronomy, Kansas State University, Manhattan, KS, USA. ²Institute for Plant Sciences of Montpellier, Montpellier, France. ³INRA, UE FERLUS, Les Verrines CS80006, 86600 Lusignan, France. ⁴INRAE, 86600 Lusignan, France. ✉email: ciampitti@ksu.edu; gilles.lemaire.inra@gmail.com

Toward a re-definition of NUE for crop breeding objectives

It is imperative to re-define NUE for crop breeding objectives, maximizing grain yield (Y_{\max}), mainly driven by application of N fertilizers (Nf) and the contribution by soil N supply (Ns), while minimizing environmental N losses¹¹ (Fig. 1).

Therefore, if the goal is to assess the ability of a crop to produce yield or biomass per unit of N, then Eq. (1) should be reformulated as:

$$\text{NUE} = \Delta Y(\text{or } W) / \Delta(\text{Nf} + \text{Ns}). \quad (6)$$

However, Ns is also affected by Nf (as demonstrated by ¹⁵N studies¹²) and controlled by the plant itself. Therefore, attempts to compute Ns + Nf as an external resource are not relevant, as Ns and Nf are intertwined.

Figure 1 illustrates two strategies for increasing crop performance in N use:

- (1) Breeding for high Y_{\max} (or W_{\max}) at high N supply, leading to an increased slope dY/dNf , but with increased environmental risks linked to higher Nf applications.
- (2) Breeding for high crop N uptake capacity at low N supply, reducing the crop dependency on Nf and minimizing environmental risks

The first strategy is clearly to maximize NCE, while the second strategy is oriented toward maximizing NupE. However, the asymptotic response of Y (or W) to Nf would create a trade-off between NCE and NupE. This feature illustrates the impossibility to separately analyze NupE and NCE as relevant crop traits for NUE.

Holistic N approach considering the soil-plant system

The soil-plant system is an integrated, auto-regulated system within which plants interact with soil to determine the N available for root absorption¹³ (Na). Thus, as indicated in Fig. 2a, Na is the relevant bridge variable connecting soil to plant sub-systems but depending upon: (1) soil characteristics (organic matter and microbiome interactions), (2) plant root traits (density and architecture¹⁴) allowing forage of NH_4^+ and NO_3^- through active root development¹⁵, and (3) C substrate deposition and exudation within the N mineralization-organization turnover^{16,17}.

As illustrated in Fig. 2b, N uptake rate is regulated by Na within the rhizosphere but is also feedback-regulated by plant growth capacity¹⁰. This co-regulation of N uptake leads to an allometry of this process with W ¹⁸, with the critical $\text{Nup} = aW^b$ curve defined as the minimum N uptake required for maximum W. This critical curve has been demonstrated as stable across genotype \times environment (G \times E) scenarios in maize (*Zea mays* L.) and tall fescue (*Festuca arundinacea* Schreb.) crops^{18,19}, but divergent for C3 versus C4 species²⁰. When N becomes limiting, the intensity of crop N deficiency can be quantified by the distance of any data point (Nup-W) to the critical curve, i.e., the Nitrogen Nutrition Index (NNI)²¹. Determination of NNI, as proposed in recent literature²², is a prerequisite for deciphering NUE and its efficiency terms.

Metrics for effective use of N in field crops

The traditional NUE (Eq. (1)) index is a proper metric of the crop responsiveness to Nf. However, if the objective is to evaluate agro-ecological crop performance and provide relevant plant traits for breeding programs, the determination of crop NNI will be necessary to clearly separate plant traits related to:

- (1) increasing plant N demand as a consequence of plant crop W capacity.
- (2) increasing plant capacity to satisfy its own N demand via its effect on Na.

The determination of NNI facilitates this distinction and disentangles the confusing estimation of both NCE and NupE terms in NUE data^{22–24}.

New breeding strategies for improving acquisition of resources

Most NUE-focused plant breeding programs for different crop species utilize non-limiting, or at least large, Nf applications. As established for maize¹¹, breeding progress achieved through selectively increasing W_{\max} or Y_{\max} can have a small effect on Y_0 (Fig. 1) but necessitate higher N application rates to achieve Y_{\max} , potentially increasing the risk for N losses. Concurrently, crop sensitivity to N deficiency is enhanced when Ns is low.

This concept evolution occurs simultaneously with the development of new strategies to improve resource acquisition capacity of plant roots. A number of genes have been identified both in




Fig. 1 Schematic representation of crop yield (Y) response to N fertilizer application rate (Nf). The response curve $Y = f(Nf)$ is asymptotic and dY/dNf declines as Nf increases. Thus, Y increases from a value of Y_0 (yield without N fertilization) until a maximum yield (Y_{\max}) is reached for an optimum value of $Nf = Nf_{\text{opt}}$. The environmental risk for N losses is directly linked to Nf application and increases rapidly as Nf approaches and exceeds Nf_{opt} . Linear extrapolation of dY/dNf allows the estimation of an “apparent” contribution by soil N supply (Ns).




Fig. 2 Schematic representation of soil-plant system. **a** N_a , the N available for plant remains a virtual entity due to the fractal geometry of the rhizosphere. **b** Allometry between plant N uptake (N_{up}) and crop mass (W), $N_{up} = aW^b$, reflecting (i) the feedback control of the rate of root N absorption by both N availability (N_a) and potential plant growth rate in time (dW/dt) and (ii) the diminishing of the crop N demand as plant mass increase $dN_{up}/dW = abW^{b-1}$ leading to crop N dilution process. This allometry implies that both N uptake efficiency ($N_{up}E$) and N conversion efficiency (NCE) increases as crop W increases. NNI nitrogen nutrition index, $ActN_{up}$ (Actual N uptake)/ $CritN_{up}$ (Critical N uptake), N_f N fertilizer.

model and crop plants which facilitate enhanced nutrient acquisition²⁵. Meta-analysis and allele combination analysis indicated that root system depth and root spreading angle are valuable candidate traits for improving grain yield by pyramiding favorable alleles²⁶. Genome-wide association studies and quantitative trait loci are powerful tools for understanding genetic variation of root architecture and delivering markers to assist new breeding strategies to facilitate genetic improvement of water/nutrient efficiencies.

A forward-facing outlook

In summary, the root availability of N in the rhizosphere is clearly the bridge variable to connect soil and plant processes. In addition, NNI is an a posteriori proxy of the ability of plant roots to capture and utilize available rhizospheric N. Thus, NNI determination is fundamental to untangle true gains of NUE rather than trivial pseudo efficiency improvements solely based on yield. The integration of NNI in crop breeding programs can be more rapidly ingested (high-throughput phenotyping^{23,27}) with the utilization of new technologies (e.g., sensors, satellite) under varying $G \times E \times M$ conditions^{28,29}.

Although the concept presented here is mainly focused on N, the foundation and framework could be extended to other nutrients to improve our understanding in a more holistic approach of nutrient use efficiency at a broader soil-plant system scale. Lastly, crop improvement for effective use of nutrients will require the integration of key scientists (i.e., agronomists, crop physiologists, breeders)³⁰ to realize the potential benefits of direct selection and for rapid progress³¹ to overcome this complex challenge and address food security goals in a more sustainable way.

Code availability

The code used for this analysis is available from the corresponding author on request.

Received: 28 January 2022; Accepted: 1 August 2022;

Published online: 16 August 2022

References

- Zhang, X. et al. Managing nitrogen for sustainable development. *Nature* **528**, 51–59 (2015).
- Erisman, J. et al. How a century of ammonia synthesis changed the world. *Nat. Geosci.* **1**, 636–639 (2008).
- Steenbjerg, F. & Jakobsen, S. T. Plant nutrition and yield curves. *Soil Sci.* **95**, 69–90 (1963).
- Moll, R. H., Kamprath, E. J. & Jackson, W. A. Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agron. J.* **74**, 562–564 (1982).
- Cassman, K. G., Dobermann, A. & Walters, D. T. Agroecosystems, nitrogen-use efficiency, and nitrogen management. *Ambio*. **31**, 132–140 (2002).
- Dobermann, A. Nutrient use efficiency. Measurement and management. *Proceedings of the International Fertilizer Industry Association (IFA) Workshop on Fertilizer Best Management Practices*, 7–9 March, 1–28 (Brussels, 2007).
- Fixen, P. et al. Nutrient/fertilizer use efficiency: measurement, current situation and trends. In *Managing Water and Fertilizer for Sustainable Agricultural Intensification* (eds Drechsel, P. et al.) 8–38 (IFA, 2015).
- European Union Nitrogen Expert Panel. *Nitrogen Use Efficiency (NUE)—An Indicator for the Utilization of Nitrogen in Agriculture and Food Systems Prepared by the EU Nitrogen Expert Panel* (Wageningen University, 2015).
- Congreves, K. A. et al. Nitrogen use efficiency definitions of today and tomorrow. *Front. Plant Sci.* **12**, 637108 (2021).
- Briat, J. F. et al. Reappraisal of the central role of soil nutrient availability in nutrient management in light of recent advances in plant nutrition at crop and molecular levels. *Eur. J. Agron.* **116**, 126069 (2020).
- Ciampitti, I. A. & Lemaire, G. From use efficiency to effective use of nitrogen: a dilemma for maize breeding improvement. *Sci. Total Env.* **20**, 154125 (2022).
- Quan, Z., Zhang, X., Fang, Y. & Davidson, E. A. Different quantification approaches for nitrogen use efficiency lead to divergent estimates with varying advantages. *Nat. Food* **2**, 241–245 (2021).
- Araus, V. et al. A balancing act: how plants integrate nitrogen and water signals. *J. Exp. Bot.* **15**, 4442–4451 (2020).
- Lynch, J. P. Root phenotypes for improved nutrient capture: an underexploited opportunity for global agriculture. *N. Phytologist* **223**, 548–564 (2019).
- Duque, L. O. & Villordon, A. Root branching and nutrient efficiency: status and way forward in root and tuber crops. *Front. Plant Sci.* **10**, 237 (2019).
- Canarini, A. et al. Root exudation of primary metabolites: mechanisms and their roles in plant responses to environmental stimuli. *Front. Plant Sci.* **10**, 157 (2019).
- Elrys, A. S. et al. Global gross nitrification rates are dominantly driven by soil carbon-to-nitrogen stoichiometry and total nitrogen. *Glob. Change Biol.* **27**, 5950–5962 (2021).
- Ciampitti, I. A. et al. Does the critical N dilution curve for maize crop vary across genotype x environment x management scenarios?—A Bayesian analysis. *Eur. J. Agron.* **123**, 126202 (2021).

19. Fernandez, J. et al. Revisiting the critical nitrogen dilution curve for tall fescue: a quantitative synthesis. *Eur. J. Agron.* **131**, 126380 (2021).
20. Greenwood, D. J. et al. Decline in percentage N of C3 and C4 crops with increasing plant mass. *Ann. Bot.* **66**, 425–436 (1990).
21. Lemaire, G. et al. Diagnosis tool for plant and crop N status in vegetative stage: theory and practice for crop N management. *Eur. J. Agron.* **28**, 614–624 (2008).
22. Lemaire, G. & Ciampitti, I. A. Crop mass and N status as prerequisite covariables for unraveling nitrogen use efficiency across genotype-by-environment-by-management scenarios: a review. *Plants* **9**, 1309 (2020).
23. Sadras, V. O. & Lemaire, G. Quantifying crop nitrogen status for comparisons of agronomic practices and genotypes. *Field Crops Res.* **164**, 54–64 (2014).
24. Gastal, F. et al. *Quantifying Crop Responses to Nitrogen and Avenues to Improve Nitrogen-use Efficiency* ISBN:978-0-12-417104-6, 159–206 (Academic Press, 2014).
25. Jia, Z. & von Wieren, N. Signaling pathways underlying nitrogen-dependent changes in root system architecture: from model to crop species. *J. Exp. Bot.* **71**, 4393–4404 (2020).
26. Torti, S. et al. Transient reprogramming of crop plants for agronomic performance. *Nat. Plants* **7**, 159–171 (2021).
27. Ratjen, A. M. et al. Key variables for simulating leaf area and N status: biomass based relations versus phenology driven approaches. *Eur. J. Agron.* **100**, 110–117 (2018).
28. Peng, J. et al. Random forest regression results in accurate assessment of potato nitrogen status based on multispectral data from different platforms and the critical concentration approach. *Field Crops Res.* **268**, 10815 (2021).
29. Chen, P. A comparison of two approaches for estimating the wheat nitrogen nutrition index using remote sensing. *Remote Sens.* **7**, 4527–4548 (2015).
30. Hirel, B. et al. Improving nitrogen use efficiency in crops for sustainable agriculture. *Sustainability* **3**, 1452–1485 (2011).
31. Lammerts van Bueren, E. T. & Struik, P. C. Diverse concepts of breeding for nitrogen use efficiency. A review. *Agron. Sustain. Dev.* **37**, 50 (2017).

Acknowledgements

Contribution no. 23-008-J from the Kansas Agricultural Experiment Station.

Author contributions

I.A.C. and G.L. wrote the original draft of the manuscript. J.-F.B. and F.G. were involved in the interpretation of results. All authors contributed to writing—review and editing of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Ignacio A. Ciampitti or Gilles Lemaire.

Peer review information *Communications Biology* thanks Mina Devkota and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Primary Handling Editors: Caitlin Karniski and Christina Karlsson Rosenthal.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022