



**HAL**  
open science

## Efficacy of a targeted selective treatment in dairy herds affected by clinical dictyocaulosis.

T. Lurier, Thomas Hilaire, Philippe Camuset, G. Bourgoin, M.A. Arcangioli

### ► To cite this version:

T. Lurier, Thomas Hilaire, Philippe Camuset, G. Bourgoin, M.A. Arcangioli. Efficacy of a targeted selective treatment in dairy herds affected by clinical dictyocaulosis.. 31st WORLD BUIATRICS CONGRESS, National Association of Spanish Specialists in Bovine Medicine (ANEMBE); World Association for Buiatrics (WAB), Sep 2022, Madrid, Spain. hal-03838494

**HAL Id: hal-03838494**

**<https://hal.inrae.fr/hal-03838494>**

Submitted on 3 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives | 4.0  
International License

# EFFICACY OF A TARGETED SELECTIVE TREATMENT IN DAIRY HERDS AFFECTED BY CLINICAL DICTYOCAULOSIS

Thibaut LURIER<sup>1,2</sup>, Thomas HILAIRE<sup>1</sup>, Claire HENON<sup>1</sup>, Philippe CAMUSET<sup>3</sup>, Gilles  
BOURGOIN<sup>1,4</sup>, Marie-Anne ARCANGIOLI<sup>1</sup>

1-VetAgro Sup, 2-UMR EpiA, 3- Caux Seine veterinary practice 4-UMR 5558



# Bovine dictyocaulosis


- Dictyocaulosis is a worldwide parasitic disease reported in most temperate country
- Prevalence  $\approx$  80% of herds with enzootic cough in pasture during summer (France, *Lurier et al. 2018*)

- Economic importance


Clinical outbreaks cost = **159-167€/cows** (*Holzhauser et al., 2011*)

⇒ Decreased milk production (30/50% of cost), mortality (30/50% of cost), extra AI + veterinary expense (12-15%)

At EU level : **€ 139 million** [€ 86–225 million] in **cattle** (*Charlier et al. 2020*)



Country with reported cases of dictyocaulosis



Relative economic impact of dictyocaulosis among other parasitic disease according to the climate in Europe

## Diagnostic in adult cattle (*Lurier et al. 2018*)

- Routine method : **Baermann / Mc Kenna sedimentation**

**Sp = 100%, Se = 7.4%**

- **Broncho-alveolar Lavage (BAL)**

- Presence of eggs or larvae :


**Sp = 100%, Se = 24.7 %**

- **Eosinophils proportion > 4.77% in BALF cytology**

**Sp = 85.2% & Se = 85.2%**

- **Serology** not available in France, Seuil = 0.389 ODR


**Sp = 85.7% & Se=76.2%**



*D. Viviparus L3 in feces*



*D. Viviparus adult in BALF*



*BALF cytology with eosinophils*

# Treatment of dictyocaulosis

- Most anthelmintic are effective against *D.viviparus*
  - Very few treatment failure reported : eprinomectine (Rigaud et al. 2019), Albendazol (Coles et al. 2010), Abamectine and moxidectine (Molento et al. 2006)
- Routinely
  - Blanket treatment applied to all cows without confirmatory diagnostic
    - Potential **unnecessary treatment**
    - Potential selection of **anthelmintic resistance** in *D. viviparus* or in other gastrointestinal nematodes
- No **targeted selective treatment protocols** assessed for Dictyocaulosis control in cattle



# Pilot study design 1/2

## Objectives

- Assess the efficacy of a targeted selected treatment of dairy cattle for :
  - Resolving clinical manifestation of dictyocaulosis in the herd
  - Limit the incidence of the disease in non-treated animal

## Methods

- Inclusion of herds with a veterinary dictyocaulosis suspicion in two French regions
- **Confirmatory diagnosis** by bronchoalveolar lavage and Mc Kenna sedimentation
- **Targeted selective treatment of 50%** of the herd with eprinomectine injectable
- **Longitudinal follow up** until 3 month post treatment

## Pilot study design 2/2

D0

Diagnostic

Random selection of  
6 cows (3  
**primiparous** and 3  
**multiparous**)

BAL,  
Individual Mc Kenna  
Pooled Mc Kenna

D3-D8

Treatment

Treatment with  
Eprecis ND  
(0,2mg/kg)

**Primiparous**  
**Postpartum cows**  
**Clinically affected**  
**animals**

D30

Immediate  
follow up

Herd clinical signs according to the farmer

BAL,  
Individual Mc Kenna on the 6 followed  
cows

Pooled Mc Kenna on 10 primiparous cows

D90

Long-term  
follow-up

# Rational for treatment

## Eprecis ND CEVA (eprinomectine)

- Posology 0.2 mg/kg
- **Injectable** eprinomectine (suitable for selective treatment)
- 2 weeks protection against reinfestation by *D.viviparus*

## Treated animals

- Clinically affected (cough)
- **Primiparous** (naive cows => most at risk to amplify the presence *D.viviparus*)
- Post partum cows (until 15 days after calving)



## Results : Initial visit

- Only 6 herd enrolled
- Diagnostic positive in 4/6 (66%) herds
  - 1 herd excluded (non-compliance with treatment protocol)
- Animal level prevalence
  - **Primiparous** 6/9 LBA  
2/9 Mc Kenna
  - **Multiparous** 1/9 LBA  
1/9 Mc Kenna
- Only 1/3 pooled Mc Kenna positive

	Herd	Cows	Eosinophil %	Mc Kenna	Pooled Mc kenna
Primiparous	A	5824	27,75	0	0
		5834	20,75	0	
		5813	4,75	0	
	B	1782	15	1	1
		1808	7,5	0	
		1777	7,5	1	
	C	763	2,5	0	0
		4385	3,5	0	
		769	21,5	0	
	D	8969	0,1	0	0
		2708	0	0	
		4365	0,5	0	
	E	1301	0	0	0
		1465	0	0	
		4627	1,2	0	
Multiparous	A	7064	4,2	0	1
		3493	1	0	
		3489	16	1	
	B	1581	0	0	0
		1565	0	0	
		1642	2	0	
	C	3864	2	0	0
		8525	3	0	
		827	1,4	0	
	D	8670	0,2	0	0
		9912	0	0	
		6277	0	0	
	E	6679	0	0	0
		M3	1,2	0	

## Results : Follow-up


### • Primiparous

- All 9/9 primiparous (treated) were negative to LBA and Mc Kenna at D30 and D90

### • Multiparous

- 2 excluded cows
  - **1 treated** (cough) replaced by another cows at D30
  - 1 culled at D90
- 2/8 primiparous (not treated) were positive to LBA or Mc Kenna at D30 and D90

⇒ **Incidence rate : 4.4%/month in non treated multiparous  
2%/month for all susceptible cows**



# Clinical follow up : Tachypnea & lung auscultation modification

- **Abnormal clinical examination at D0**
  - 3/9 primiparous
  - 1/8 multiparous
- **During follow-up**
  - Persistence of the **abnormal clinical signs in 2/4** primiparous and **1/1 multiparous**
  - No newly clinically affected cows
- **According to the farmer**
  - **Resolution of the clinical signs in 2/3** herds
  - Persistence of **cough in 1 herd** without needs for supplementary treatments

# Discussion

- **Interest of the targeted selected treatment**
  - No re-infestation in primiparous cows
  - No new clinical manifestation after treatment
  - Low incidence (4.4%/month) of *D.viviparus* infestation in non-treated cows
- **Few herds corresponding to the inclusion criteria**
  - Dry weather limiting the contamination of the pasture
    - ⇒ Could have limited the number of lungworm outbreaks
    - ⇒ Could also have favored the good response to the targeted selected treatment
- **Only a proof of concept without control groups**
  - Need to be reproduced with a larger number of herds
  - The targeted selected treatment strategy needs to be compared with a blanket treatment strategy in a randomized control trial

## Conclusion and take-home message

- **Importance of the initial diagnosis**
  - 2/6 herds presenting enzootic cough at pasture during summer without dictyocaulosis
  - Confirmation of the higher sensitivity of the bronchoalveolar lavage compared to the Mc Kenna sedimentation in adult cows
- **Potential interest of a targeted selected treatment in dairy cows by treating**
  - Clinically affected animals
  - Primiparous and newly introduced animals
  - Early post-partum cows

# Acknowledgment

- This study was funded by CEVA France who provides the Eprecis ND and some financial supports
- We also acknowledge
  - The farmers who accept to participate to this study
  - Nathalie Menudier, Damien Remmy (CEVA) for their implication in this project



## Thank you for your attention



**RÉPUBLIQUE  
FRANÇAISE**

*Liberté  
Égalité  
Fraternité*



VetAgro Sup

31 **WBC**

WORLD BUiatrics CONGRESS

SEPTEMBER 4<sup>TH</sup> TO 8<sup>TH</sup>

MADRID 2022

[www.wbc-madrid2022.com](http://www.wbc-madrid2022.com)

