



HAL
open science

Determinants of survival and dispersal along the range expansion of a biological invasion

Eric Edeline, Agnès Starck, Yoann Bennevault, Jean-Marc Paillisson, Eric Petit

► **To cite this version:**

Eric Edeline, Agnès Starck, Yoann Bennevault, Jean-Marc Paillisson, Eric Petit. Determinants of survival and dispersal along the range expansion of a biological invasion. 2024. hal-03839947

HAL Id: hal-03839947

<https://hal.inrae.fr/hal-03839947v1>

Preprint submitted on 15 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Determinants of survival and dispersal along the range expansion of a biological invasion

Eric Edeline^{1†}, Agnès Starck², Yoann Bennevault², Jean-Marc Paillisson^{3*}, Eric J. Petit^{1*}

*: Joint last authors

1: DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro Rennes-Angers, Ifremer, Rennes, France.

2: U3E, Aquatic Ecology and Ecotoxicology Experimental facility, INRAE, Rennes, France

3: University of Rennes, CNRS, ECOBIO [(Ecosystèmes, biodiversité, évolution)] – UMR 6553, Rennes, France.

†: corresponding author: Eric.Edeline@inrae.fr

Abstract

20 Projecting future response of biological systems to global change requires a mechanistic
understanding of how climate and ecology jointly drive species demography and range dynamics.
22 Such knowledge is particularly crucial when it comes to invasive species, which expansion may
have far-reaching consequences for recipient ecosystems. Here, we use mark-recapture in replicated
24 outdoor mesocosms to examine how survival and dispersal, two key drivers of population and range
dynamics, respond to climate and ecology in the invasive red swamp crayfish (*Procambarus*
26 *clarkii*) along an invasion gradient. We show that crayfish survival probability increased with (i)
increasing body size at high (but not low) crayfish density and (ii) with warmer temperatures, and
28 decreased (i) with increasing body condition and (ii) under higher crayfish density. Overland
dispersal probability by crayfish increased with increasing (i) body-size, (ii) body condition and (iii)
30 temperatures. In contrast, crayfish from range-edge and range-core habitats had similar survival and
overland dispersal probabilities, suggesting no evolution of the crayfish expansion potential along
32 the invasion gradient. Our results highlight that species population dynamics and range shifts in a
changing world are driven by joint contributions from both climate and ecology. In *P. clarkii*, global
34 warming will simultaneously promote both a demographic increase and a geographic range
expansion, especially in populations dominated by large-bodied individuals.

36

Key words : Biological invasions, Body size, Condition-dependent dispersal, Invasion
38 management, Mesocosms, Multistate mark-recapture model, Population density, *Procambarus*
clarkii, Survival-dispersal trade off, Temperature.

40

42

INTRODUCTION

44 How species geographic distributions will change in a near future is a central question in ecology,
evolution, conservation and the management of biological invasions. To date, our ability to answer
46 this question often rests on correlative species distribution models, which provide a poor
mechanistic understanding and, arguably, have a low power to predict how organisms will respond
48 to global change. A recent approach to addressing these questions consists in explicitly modelling
the ecological and evolutionary processes that control geographic distributions in so-called process-
50 explicit models (Briscoe et al., 2019; Townsend Peterson et al., 2015; Travis & Dytham, 2012).
These models draw heavily on population-dynamic models, and on how demographic rates, i.e.,
52 survival, reproduction and dispersal, respond to environmental change (Fig. 1).




Fig. 1. Conceptual framework for process-explicit models of species range dynamics.

Global change may select upon individual state variables such as sex, body size, body condition or genotype (arrow 1a), or alter the environment (arrow 1b) which includes climate (e.g., temperature) and ecological variables (e.g., competitor density). Individual state (arrow 2) and environment variables (arrow 3) both influence demographic rates, which ultimately drive the dynamics of both populations and geographic ranges (arrow 4).

In turn, population and range dynamics entail environmental alterations (arrow 5), which influence individual states through both phenotypic plasticity and selection experienced by individuals (arrow 6). Bold arrows show the causality links that are specifically investigated in the present study.

66

68

70 The complexity of mechanistically predicting how demographic rates respond to environmental
change may be usefully tackled by separating the changes that act directly onto demographic rates,
72 or indirectly through a modification of individual state variables (Fig. 1). Global change may select
upon individual state variables that, in turn, influence demographic rates (arrow 1a-2 sequence in

74 Fig. 1, Table 1). For instance, a number of human activities, such as harvesting, are selective on sex
and/or body size (Biro & Post, 2008; Edeline, 2016; Kuparinen & Festa-Bianchet, 2017), which are
76 state variables that typically have strong effects on survival, reproduction or dispersal
(Charlesworth, 1994; Clobert et al., 2009; Clobert, Baguette, et al., 2012; Roff, 2002). At the same
78 time, global change disrupts a host of abiotic (e.g., climate) and ecological (i.e., density-dependent)
factors which have long been known to strongly influence demographic rates, including survival
80 and dispersal either through ecological processes (arrow 1b-3 sequence in Fig. 1, Begon et al.,
2005; Clobert, Baguette, et al., 2012; Kendall et al., 1999; Krebs, 2014; Stenseth et al., 2002), or
82 through an alteration of individual state variables (arrow 1b-4-2 sequence Fig. 1 Le Galliard et al.,
2012; Matthysen, 2012).

84

An additional complexity layer comes from demography and range dynamics responding not only
86 respond to direct and indirect effects of anthropogenic selection, climate and ecology, but also to
feedbacks between these different components: as both the population size and geographic range
88 change, so do the environmental conditions and selective pressures and/or plastic effects
experienced by individuals (e.g., competition, predation, habitat quality, arrow 5-6 sequence in Fig.
90 1). In particular, emigration from the core habitat, or from the introduction point in the case of
biological invasions, selects for increased dispersal propensity through spatial sorting, and is further
92 possibly a competition-avoidance response where poorer competitors disperse from high-density
core habitats to low-density marginal habitats (Burton et al., 2010; Chuang & Peterson, 2016;
94 Messenger & Olden, 2019; Phillips, 2016; Phillips et al., 2010). Hence, high dispersers and poor
competitors are likely to be overrepresented among immigrants at range edges. Accordingly, results
96 from spread experiments support the prediction that increased dispersal evolves at range edges
(Fronhofer & Altermatt, 2015; Ochocki & Miller, 2017; J. L. Williams et al., 2016), but evidence
98 for reduced competitive ability is mixed (Duckworth, 2008; Hudina et al., 2015).

100 Unfortunately, however, the development of process-explicit models of geographic ranges that
would integrate both direct and indirect effects of global change and feedbacks is impeded by
102 serious logistical and knowledge limitations. The data needed to estimate demographic rates, as
well as their state- and environment-dependency, are hard and costly to acquire in the wild, and are
104 thus typically lacking (Briscoe et al., 2019; Gaston, 2009). Even when present, such data may be
poor if detection or recapture rates are low, or if population size is small, resulting in inaccurate
106 parameter estimation. A further obstacle to implementing process-explicit models of species ranges
is that the eco-evolutionary feedback loops that may operate during range expansion (Fig. 1, arrow
108 5; see below) remain a frontier in biology and are poorly understood empirically (Angert et al.,
2020; Clobert, Massot, et al., 2012; Miller et al., 2020; Travis & Dytham, 2012), which makes it
110 uncertain which mechanisms should be incorporated into models.

112 In the present study, we propose to circumvent these logistical and knowledge limitations using, in
outdoor mesocosms, mark-recapture of wild-caught range-core and range-edge animals.
114 Specifically, we quantified the effects of key individual state variables (sex, body size, body
condition), climate (temperature), ecology (population density) and geographic range (core vs. edge
116 of an invaded area) on both survival and overland dispersal propensity in red swamp crayfish
(*Procambarus clarkii*). This species, well known for dispersing through overland movements (Cruz
118 & Rebelo, 2007; Huner & Barr, 1991; Ramalho & Anastácio, 2015), is considered as one of the
most invasive aquatic species worldwide (Oficialdegui et al., 2019; Savini et al., 2010), and poses
120 very serious management problems due to the profound reorganizations they cause in recipient
ecosystems (Lodge et al., 2012; Souty-Grosset et al., 2016; Twardochleb et al., 2013). From a
122 literature review, we formed detailed predictions for the state, temperature-, density- and range-
dependency of *P. clarkii* survival and dispersal (Table 1). Our experimental approach allowed us to
124 fully test all of these predictions.

126 **Table 1. Predicted and experimentally-observed in the present study effects of sex, body size, body condition, geographic range (core vs. edge), as well as**
 128 **population density and temperature on both survival and dispersal in the red swamp crayfish (*Procambarus clarkii*).** Detailed statistics for observed effects
 are provided in Table 2.

Factor	Relationship to Survival				Relationship to Dispersal			
	Effect sign		Explanation of predicted sign	References	Effect sign		Explanation of predicted sign	References
	Predicted	Observed			Predicted	Observed		
Sex male	-	0	Males invest less than females in maintenance and future reproduction due to higher hazards	(Kirkwood, 2001; G. C. Williams, 1957)	+ or -	0	Sign depends on relative strengths of opposing selective forces. Male-biased in <i>Pacifastacus leniusculus</i>	(Gros et al., 2008; Hudina et al., 2012; Wutz & Geist, 2013)
Body size	+	0	Larger-bodied individuals dominate competition	(Figler et al. 1999; Edeline and Loeuille 2021 and references therein)	+	+	Sign depends on relative strengths of opposing selective forces. In crayfish, overland and in-stream dispersal is often favoured by a large body size.	(Claussen et al., 2000; Clobert et al., 2009; Kisdi et al., 2012; Moorhouse & MacDonald, 2011; Ramalho & Anastácio, 2015; Thomas et al., 2018; Wutz & Geist, 2013)
Body condition	+	-	Higher physical quality increases survival	Multiple evidence, see e.g. Blums et al. (2005)	+ or -	+	Sign depends on relative strengths of opposing selective forces.	(Bonte & Peña, 2009; Clobert et al., 2009; Kisdi et al., 2012)
High population density	-	-	Increased competition decreases average	(Burton et al., 2010; Phillips,	+	0	Competition avoidance favours	(Clobert et al., 2009; Galib et

			survival	2016; Phillips et al., 2010) see also Appendix 8 for crayfish-specific studies			increased dispersal	al., 2022)
Temperature	-	+	Higher metabolic rate elevates mortality, but sign should depend on temperature relative to thermal optimum	(Angilletta Jr., 2009; Brown et al., 2004; Knies & Kingsolver, 2010; Westhoff & Rosenberger, 2016)	+ or -	+	Sign depends on thermal preference	(Angilletta Jr., 2009; Claussen et al., 2000; Le Galliard et al., 2012; Ramalho & Anastácio, 2015)
Range edge	-	0	A survival-dispersal trade off makes dispersers less able to survive	(Angert et al., 2020; Burton et al., 2010; Messenger & Olden, 2019; Miller et al., 2020; Phillips, 2016; Phillips et al., 2010)	+	0	Spatial filtering leads to more dispersive individuals at range edges	(Angert et al., 2020; Burton et al., 2010; Miller et al., 2020; Phillips, 2016; Phillips et al., 2010)
High density-by-size interaction	+	+	Benefits from a large body size increase under elevated competition	(Burton et al., 2010; Edeline & Loeuille, 2021; Phillips, 2016; Phillips et al., 2010)	-	0	Competition at high density favours dispersal in small, competitively-dominated individuals	(Clobert et al., 2009)
Range edge-by-size interaction	-	0	Lower competitive ability in Range-edge crayfish make body size less important to survival	(Burton et al., 2010; Edeline & Loeuille, 2021; Messenger & Olden, 2019; Phillips, 2016; Phillips et al.,	-	0	Lower competitive ability in Range-edge crayfish makes dispersal less size-dependent	(Clobert et al., 2009; Phillips et al., 2010)

Range edge-by-density interaction	-	0	Negative effect of high density on survival stronger in less competitive Range-edge crayfish	2010) (Burton et al., 2010; Edeline & Loeuille, 2021; Messenger & Olden, 2019; Phillips, 2016; Phillips et al., 2010)	+	0	Lower competitive ability in Range-edge crayfish makes them more sensitive to the positive effect of density	(Clobert et al., 2009; Phillips et al., 2010)
Range edge-by-density-by-size interaction	-	0	Weaker size-dependency of survival in Range-edge crayfish is more pronounced under high density	(Burton et al., 2010; Edeline & Loeuille, 2021; Phillips, 2016; Phillips et al., 2010)	-	0	Weaker size-dependency of dispersal in Range-edge crayfish is more pronounced under high density	(Clobert et al., 2009; Phillips et al., 2010)

132

MATERIALS AND METHODS

134 **Crayfish populations**

P. clarkii originates from south-central USA (Louisiana and Texas) and was introduced to Europe
136 through both USA-to-Kenya-to-Europe and USA-to-Spain routes (Oficialdegui et al., 2019). The
species is highly fecund, tolerant to a large range of environmental conditions (including drought),
138 generally spreads rapidly once introduced either through secondary human-mediated introductions
(Oficialdegui et al., 2020) or through active in-stream and overland dispersal (Cruz & Rebelo, 2007;
140 Tréguier et al., 2018), and often has major impacts on the invaded ecosystems (Souty-Grosset et al.,
2016; Twardochleb et al., 2013).

142

In this study, we used crayfish sampled from natural populations in the Brière area, north-western
144 France (47°23'N, 02°12'W) where crayfish belonging to the USA-to-Kenya-to-Europe introduction
route were introduced during the early 1980's (Bélouard et al., 2019; Oficialdegui et al., 2019).
146 Crayfish escaped from a farm and invaded the nearby marsh (90 km²), where they now form a large
and genetically-homogeneous population (Bélouard et al., 2019). From this range core, dispersers
148 used streams, ditches and, secondarily, overland dispersal to colonize ponds embedded in the
surrounding hedgerow landscape, where they founded multiple new populations that may be
150 considered as range-edge populations (Bélouard et al., 2019; Tréguier et al., 2018).

152 We sampled crayfish from three different range-core locations in the marsh, hereafter collectively
referred to as “Range-core crayfish”, and from three range-edge pond populations genetically
154 differentiated from the marsh source at neutral markers (CB7, I6 and I10 ponds as named in
Bélouard et al. 2019), and hereafter collectively referred to as “Range-edge crayfish”.

156

In addition to crayfish from the Brière, we sampled crayfish from a third locality, a pond located in
158 Rennes city (48°06'N, 1°42'W, about 80 km away from the Brière), hereafter referred to as an
“Outgroup”. These Rennes crayfish have a genetic signature typical of the native range, suggestive
160 of an introduction route that differs from what has been described for the marsh of Brière
(Appendix 1). We used these Outgroup crayfish in order to standardize the density treatments in the
162 experiment (see below), and also because we were able to capture Range-edge crayfish only in
limited numbers due to small population sizes (Bélouard et al., 2019).

164

Crayfish sampling, marking and maintenance

166 Crayfish in the Brière were sampled using non-baited traps on May 21st, 22nd and 24th 2019. Upon
capture, crayfish were determined as male or female from secondary sexual characters (first
168 pleopod pair transformed to gonopods in males), measured for body size (cephalothorax length to
the nearest mm, from the tip of the rostrum to the end of the cephalothorax), weighted to the nearest
170 10⁻¹g, and individually marked using 8 mm passive integrated transponders (Biomark HPT8 FDX
PIT tags) inserted in the abdominal musculature of the third abdominal segment in the ventral
172 surface. Crayfish were then transported to the PEARL, the INRAE experimental facility for aquatic
ecology and ecotoxicology in Rennes (https://www6.rennes.inrae.fr/u3e_eng/), and stocked in 8 m²
174 mesocosms. The same procedure was applied to Rennes crayfish, that were PIT-tagged on April
24th. In stocking mesocosms, crayfish were provided with an excess shelters made from PVC tubes,
176 and were fed with fish food in excess (Le Gouessant, Carpe Extrudée Coul 4).

178 Experimental mesocosms

Experimental mesocosms were 28, three-m² circular tanks covered with nets to prevent avian
180 predation, and equipped with six artificial refuge traps (ART, Green et al. 2018). ARTs consisted in
five PVC tubes (250 mm length) glued together and closed by a grid at one end (Appendix 2A),
182 using either 26.5 or 32.5 mm tubes (inner diameter) so as to fit with the needs of both small- and

184 large-bodied crayfish. ARTs sank on the bottom of the mesocosms and provided crayfish with
shelters in excess (i.e., one tube provided one shelter, 30 tubes per mesocosm).

186 Each experimental mesocosm was further equipped with an exit trap requiring for trapping that
crayfish actively walk out from water, i.e., that crayfish start overland dispersal. Specifically, exit
188 traps were made from a 50L bucket attached to the inner wall of the mesocosm and hanging 30 cm
underwater. Trapping required crayfish to climb a ramp made from a plastic net and extending from
190 the bottom of the mesocosm up to the rim of the bucket (Appendix 2B). Hence, exit traps captured
crayfish that clearly expressed active, out-of-water movements which may be interpreted as the
192 onset of overland dispersal. Crayfish falling into the trap could not escape back to the mesocosm.
The bottom of the bucket was pierced with holes to allow water circulation, and PVC tubes
194 provided trapped crayfish with shelters.

196 Throughout the experiment, crayfish were mildly fed with 418 ± 11 mg (mean \pm SD) of fish food
per experimental mesocosm per week.

198

Experimental treatments and recapture

200 On June 3rd 2019, we started the experiment by transferring crayfish into experimental mesocosms.
We introduced randomly-chosen Range-core and Range-edge crayfish at a constant density of 7
202 crayfish per mesocosm, and varied density using either 2 or 12 Outgroup crayfish per mesocosm,
yielding two density treatments (9 or 19 crayfish per mesocosm) and two range treatments (Range-
204 core with Outgroup or Range-edge with Outgroup) in a factorial design (7 replicate mesocosms per
treatment, 392 crayfish in total).

206

Experimental densities (3.0 vs. 6.3 individuals m^{-2}) were relatively high compared to naturally-
208 observed densities in several crayfish species, which range from $9.3 \cdot 10^{-3}$ to 20.0 individuals m^{-2} ($n =$

29, mean 2.5, median 1.0, SD 4.7, Bubb et al., 2004; Coignet et al., 2012; Correia & Bandeira,
212 2004; Galib et al., 2022; Guan, 2000; Lamontagne & Rasmussen, 1993; Pilotto et al., 2008; Wutz &
Geist, 2013), and in line with experimentally-manipulated densities in enclosures or mesocosms,
214 which range from 0.8 to 14.0 individuals m⁻² (mean 4.3, median 3.6, SD 3.3, Lodge et al. 1994;
Angeler et al. 2003; Rodríguez et al. 2003; Gherardi and Lazzara 2006; Gherardi and Acquistapace
216 2007; Correia and Anastácio 2008; Matsuzaki et al. 2009; Jackson et al. 2014; Rodríguez-Pérez et
al. 2016; Závorka et al. 2020).

218

At 1-7 days intervals (mean \pm SD = 2.4 \pm 0.8 days) starting from June 5th, both ARTs and exit traps
220 were cleared, and recaptured crayfish were recorded for PIT tag number and returned to their
mesocosm. Two crayfish lost their tags during the course of the experiment. These were identified
222 based on their sex, size and previous capture histories, and re-tagged. On November 5th, mesocosms
were drained and all surviving crayfish were recaptured and recorded for PIT tag number. In total,
224 there were 63 recapture occasions.

226 **Temperature**

Hourly air temperature data for the entire duration of the experiment was obtained from a 5km-
228 distant Meteo France weather station (Station météorologique de Rennes-St Jacques, 48°07'N,
1°74'W). Additionally, during 90 % of the duration of the experiment (from June 17th to November
230 5th) we recorded hourly water temperature in six of 28 experimental mesocosms using HOBO Tidbit
v2 temperature loggers. A cross-correlation analysis shows that water temperature in the mesocosms
232 was highly correlated with air temperature at the weather station with a 5-hour lag (Appendix 3).
For our analyses, we computed mean hourly air temperature during the transition interval between
234 two successive capture occasions.

236 **Multistate capture-recapture model**

We estimated the effects of individual and environmental factors on crayfish survival and overland dispersal probabilities using a multistate mark-recapture model (Lebreton et al., 2009), separating among three states: A for crayfish present in the mesocosm, B for crayfish present in the exit trap (i.e., overland dispersal), and Dead. We assumed that crayfish survived in state A between capture occasions t and $t+1$, and dispersed to B just prior to $t+1$. Hence, only survival in state A is considered. This assumption was justified by the fact that we did not find any dead crayfish in an exit trap. The state process can be represented by a transition matrix with departure states (time t) in rows and arrival states (time $t+1$) in columns:

$$\begin{array}{rcc}
 & \begin{array}{c} \text{State A} \\ \text{State B} \\ \text{Dead} \end{array} & \begin{array}{c} \text{State A} \\ \text{State B} \\ \text{Dead} \end{array} \\
 \begin{array}{c} \text{State A} \\ \text{State B} \\ \text{Dead} \end{array} & \begin{pmatrix} \phi_A(1-\psi_{A \rightarrow B}) & \phi_A \psi_{A \rightarrow B} & 1-\phi_A \\ \phi_A \psi_{B \rightarrow A} & \phi_A(1-\psi_{B \rightarrow A}) & 1-\phi_A \\ 0 & 0 & 1 \end{pmatrix} & (1),
 \end{array}$$

with ϕ_A = survival probability, $\psi_{A \rightarrow B}$ = transition probability from state A to state B (i.e., dispersal from mesocosm to exit trap), and $\psi_{B \rightarrow A}$ = transition probability from state B to state A.

Although all crayfish captured in an exit trap (state B) were in fact effectively returned to their experimental mesocosm (state A), i.e., although the experimental design made state B effectively absent in the rows of the state transition matrix (1), we chose to use this model parametrization so as to separate dispersal performed by previously-dispersing individuals. That is, $\psi_{B \rightarrow A}$ represented probability of settlement in the mesocosm for a crayfish having dispersed to the exit trap at the previous occasion, hereafter referred to as “post-dispersal settlement” probability. Non-symmetric patterns of $\psi_{A \rightarrow B}$ and $\psi_{B \rightarrow A}$ would be suggestive of a cost of dispersal. Note that we also ran a model in which transition probabilities were the same for the “state A” and “state B”

lines in the transition matrix (i.e., with $\psi_{B \rightarrow A}$ absent), with no change on posterior parameter
 260 estimates for ϕ_A and $\psi_{A \rightarrow B}$.

262 The observation process is conditional on underlying states, and is described via detection
 probabilities in each state p_A and p_B . In our experiment, ARTs provided imperfect detection
 264 (crayfish present in a mesocosm but not in ARTs when cleared), such that $0 < p_A < 1$, but crayfish
 present in exit traps were seen for sure, such that $p_B = 1$. These observations can be summarized
 266 by a row-stochastic observation matrix (i.e., probabilities in a row sum up to one):

	<i>Seen state A</i>	<i>Seen state B</i>	<i>Not seen</i>	
268 <i>State A</i>	p_A	0	$1 - p_A$	(2).
<i>State B</i>	0	1	0	
<i>Dead</i>	0	0	1	

270 Our multistate model included a “full-detection” formulation with one independent p_A
 parameter estimated at each occasion t for each individual i . The model further included three
 272 independent Bernoulli GLMMs for the effects of individual- and environment-level factors on
 probabilities ϕ_A , $\psi_{A \rightarrow B}$, and $\psi_{B \rightarrow A}$ (generically symbolized by Ω):

274

$$\begin{aligned}
 \text{logit}(\Omega_{i,t}) = & \alpha_0 + \beta_{\text{Sex}[i]} + \alpha_1 \cdot \text{BS}_i + \alpha_2 \cdot \text{Cond}_i + \gamma_{\text{Density}[i]} + \alpha_3 \cdot \text{Temp}_t + \delta_{\text{Range}[i]} + \epsilon_{\text{Density}[i]} \cdot \text{BS}_i \\
 & + \zeta_{\text{Range}[i]} \cdot \text{BS}_i + \eta_{\text{Range}[i], \text{Density}[i]} + \theta_{\text{Range}[i], \text{Density}[i]} \cdot \text{BS}_i + \alpha_4 \cdot \text{Time}_t + \iota_{\text{Meso}[i]} + \kappa_i \quad (3), \\
 & \iota_{\text{Meso}[i]} \sim N(0, \sigma_\iota) \\
 & \kappa_i \sim N(0, \sigma_\kappa)
 \end{aligned}$$

276 where subscripts i and t stand for crayfish individuals and capture occasions, respectively, α_0 is
 278 model intercept (see below), *Sex* is a two-level factor (Male or Female), *Range* has three levels
 (Range-core, Range-edge, Outgroup), *Density* has two levels (9 or 19 crayfish per mesocosm), *BS* is
 280 body size (cephalothorax length, mm), *Cond* is individual body condition measured as the residual

from the $\ln(\text{Mass}_i) / \ln(\text{BS}_i)$ linear regression, Temp is the mean of hourly air temperature during the
282 transition interval between two capture occasions, Time is duration of the transition interval in days,
and Meso is mesocosm identity (numbered from 1 to 28). The ι and κ random effects
284 accounted for mesocosm and individual (overdispersion) effects. Note that ϕ_A was not
overdispersed and, hence, its GLMM did not include a κ parameter. Eq. 3 was fitted using an
286 “effect” parametrization as is the default in R, such that the intercept α_0 corresponds to mean
response in Range-core females at low density. To evaluate the main effects of covariates, we also
288 fitted a version of Eq. 3 with no interaction term.

290 To test for nonlinearities in the effects of temperature on crayfish survival and dispersal, we also
fitted a version of the model in which Eq. 3 included a *Density-by-Temp*, a *BS-by-Temp* and a
292 *Density-by-BS-by-Temp* interactions. None of these interactions were statistically significant, and
we thus discarded this version of the model from our subsequent analyses.

294

We fitted the multistate model using a state-space formulation (Gimenez et al., 2012; Kéry &
296 Schaub, 2012) and Markov chain Monte Carlo (MCMC) in JAGS (Plummer, 2003) through the
jagsUI package (Kellner, 2019) for R 4.2.1 (R Core Team, 2022). All numeric covariates were
298 centred to zero mean and 0.5 standard deviation as recommended by Gelman (2008) for logistic
regressions. We used vague priors on all parameter distributions. Specifically, we chose Cauchy
300 robust priors with scale 2.5 for regression parameters (Gelman et al., 2008), uniform distributions
on the [0,1] interval for probabilities, and uniform distributions on the [0,5] interval for variance
302 parameters. Additionally, to test posterior sensitivity to the priors, we also ran the model using as
priors uniform distributions on the [-10,10] interval for regression parameters in Eq. 3.

304

We ran three parallel MCMC chains for 10000 iterations (burnin of 2000 iterations) thinned at a 4-
306 iteration interval. We assessed parameter convergence using the Gelman-Rubin statistic which
compares the within- to the between-variability of chains started at different and dispersed initial
308 values (Gelman & Rubin, 1992). We tested for the significance of effects in Eq. (3) through
computing MCMC p-values as twice the proportion of the posterior which sign was opposite to the
310 sign of the posterior mode. We further evaluated the identifiability of model parameters by
computing the overlap between prior and posterior distributions using the BEST package in R
312 (Kruschke & Meredith, 2021). A high prior-posterior overlap (PPO) indicates that information from
the data is weak, such that the posterior parameter distribution is strongly dependent on the prior
314 distribution. A PPO greater than 35 % is considered as indicative of a weak parameter identifiability
(Gimenez et al., 2009).

316

318

RESULTS

320 Phenotypes

Crayfish had a balanced sex ratio (51 % females). Cephalothorax length ranged from 24.9 to 66.2
322 mm (mean \pm SD = 44.7 ± 6.8 mm), and was similar among Range-core (42.7 ± 7.1 mm) and Range-
edge crayfish (42.2 ± 6.4 mm), but was significantly larger in Outgroup crayfish (46.9 ± 6.2 mm,
324 Appendix 4; estimate = 4.22, ddl = 392, res. ddl = 389, t-value = 5.01, p-value < 0.001). Outgroup
crayfish also had a significantly higher body condition than Brière crayfish (estimate = 0.10, ddl =
326 392, res. ddl = 389, t-value = 7.31, p-value < 0.001).

328 Temperature

Mean air temperature during a recapture interval ranged from 9.3 to 26.9°C (Fig. 2). Maximum air
330 temperature was 39.4°C. There was a high positive correlation between mean and maximum
temperature (cor = 0.93) and between mean and variance temperature (cor = 0.68, data not shown).

332




Fig. 2. Daily air temperature during the experiment. Red, solid line: median daily temperature. Red
334 ribbon: 95 % confidence intervals.

336 **Survival**

Posteriors for regression parameters in Eq. 3 were insensitive to the choice of Cauchy or uniform
338 priors (compare Table 2 with Appendix 5). Prior-posterior overlaps (PPOs) for survival parameters
were below 35 %, except for η [Range edge, HD] (PPO = 36 %), indicating no major identifiability
340 problem (Appendix 6A). In particular, none of the posteriors associated with significant effects
were weakly identifiable.

342

Crayfish survival probability during a time interval ranged from 0.9 to 1 (Figs. 3a-d). These high
344 probabilities were due to the short duration of a time interval (2.4 days on average). Over the entire
duration of experiment (153 days), 51.5 % of crayfish survived.

346

Larger-bodied crayfish had increased survival probability under high (but not low) density
348 conditions, a trend mostly present in Range-core crayfish (Figs. 3b vs. 3a, α [1] vs. ϵ [HD] parameters

in Table 2). Specifically, average survival probability during a 2.4-day time interval was $9.57 \cdot 10^{-1}$ in
 350 the smallest crayfish (24.9 mm thorax length) and $9.98 \cdot 10^{-1}$ in the largest crayfish (66.2 mm,
 probabilities predicted for Range-core female crayfish under high density, all other quantitative
 352 covariates fixed to their average value).




Fig. 3. Predicted effects of body size-by-density interaction, body condition and temperature on
 354 **crayfish survival probability Φ_A , dispersal probability $\Psi_{A \rightarrow B}$ and post-dispersal settlement**
probability $\Psi_{B \rightarrow A}$. Lines represent median prediction, ribbons indicate 95 % credibility intervals. Green,
 356 solid lines: Range-core crayfish. Red, dashed lines: Range-edge crayfish. Blue, dotted lines: Outgroup
 crayfish. LD and HD stand for low and high crayfish density, respectively. All probabilities are for a
 358 transition interval of 2.4 ± 0.8 days (mean \pm SD). Predictions were made from parameters estimates of Eq. 3
 were further made by setting the Density factor to HD. Credible intervals are not displayed for clarity
 360 reasons.

362
 Higher crayfish density (Figs. 3b vs. 3a, γ [HD] parameter in Table 2) and higher individual body
 364 condition (Fig. 3c, α [2] parameter in Table 2) both decreased crayfish survival probability. Note that

the negative effect of density on crayfish survival was weaker and not significant when evaluated as
366 a main effect in a model that did not include any interaction (Appendix 7).

368 Warmer temperatures had a positive effect on crayfish survival probability (Fig. 3d, $\alpha[3]$ parameter
in Table 2). When all other quantitative covariates were fixed to their average value, raising mean
370 temperature from its minimum (9.3°C) to its maximum (26.9°C) increased mean survival
probability during a 2.4-day time interval from $9.81 \cdot 10^{-1}$ to $9.95 \cdot 10^{-1}$ (predictions computed for a
372 Range-core female crayfish at low density).

374 There was no significant effect of the geographic range on crayfish survival probability (δ [Range
edge] parameter in Table 2), nor there was any significant range-by-body size (ζ [Range edge]
376 parameter in Table 2), range-by-density (η [Range edge, HD] parameter in Table 2, a weakly
identifiable parameter) or range-by-body size-by-density interaction (θ [Range edge, HD] parameter
378 in Table 2). Note that we also tested for range-by-temperature and range-by-temperature-by-density
interactions and did not find any significant interaction (results not shown).

380

382 **Table 2. Parameter estimates from the multistate mark-recapture model.** Φ_A : survival probability, $\psi_{A \rightarrow B}$:
 dispersal probability, $\psi_{B \rightarrow A}$: post-dispersal settlement probability. The model is described in Eq. 3 and was
 384 fitted using an “effect” parametrization as the default in R. MCMC p-values were computed as twice the
 proportion of the posterior which sign was opposite to the sign of the posterior mode. P-values significant at
 386 the 5 % risk are bold faced. MCMC p-values are not relevant for variance parameters (random effects) that
 are constrained to be non-zero. HD: High density, LD: Low density. Parameter estimates for a model
 388 including only main effects in Eq. 3 (i.e., no interaction) are provided in Appendix 7. NB: The survival
 model for Φ_A was not overdispersed and did not include an “individual” random effect. Quantitative
 390 covariates were standardized to zero mean and 0.5 standard deviation.

Transition probability	Link	Covariate	Parameter	mean	sd	Rhat	MCMC p-value
Φ_A	logit	Intercept	$\alpha[0]$	4.50	0.25	1.00	0.000
		Male	$\alpha[\text{Sex Male}]$	-0.04	0.15	1.00	0.809
		Outgroup	$\beta[\text{Outgroup}]$	0.25	0.38	1.00	0.534
		Range edge	$\beta[\text{Range edge}]$	-0.02	0.33	1.00	0.952
		High density	$\gamma[\text{HD}]$	-0.62	0.31	1.01	0.046
		Outgroup-by-HD interaction	$\delta[\text{Outgroup, HD}]$	0.58	0.43	1.01	0.185
		Range edge-by-HD interaction	$\delta[\text{Range edge, HD}]$	0.57	0.45	1.01	0.222
		Body size in Range core	$\alpha[1]$	-0.08	0.35	1.00	0.806
		Outgroup on Body-size effect	$\epsilon[\text{Outgroup}]$	-0.55	0.68	1.01	0.412
		Range edge on Body-size effect	$\epsilon[\text{Range edge}]$	0.44	0.57	1.00	0.450
		HD on Body-size effect in range core	$\zeta[\text{HD}]$	1.02	0.49	1.00	0.033
		Outgroup-by-HD interaction on Body-size effect	$\eta[\text{Outgroup, HD}]$	-1.38	0.77	1.01	0.069
		Range edge-by-HD interaction on Body-size effect	$\eta[\text{Range edge, HD}]$	-0.90	0.75	1.00	0.225
		Body condition	$\alpha[2]$	-0.51	0.17	1.00	0.004
		Temperature	$\alpha[3]$	0.54	0.18	1.00	0.002
Time interval	$\alpha[4]$	-0.43	0.17	1.01	0.033		
Mesocosm	$\sigma[\theta]$	0.25	0.14	1.05			
$\psi_{A \rightarrow B}$	logit	Intercept	$\alpha[0]$	-3.44	0.20	1.00	0.000
		Male	$\alpha[\text{Sex Male}]$	-0.10	0.14	1.00	0.507
		Outgroup	$\beta[\text{Outgroup}]$	-0.61	0.33	1.00	0.056
		Range edge	$\beta[\text{Range edge}]$	-0.10	0.27	1.00	0.726
		High density	$\gamma[\text{HD}]$	0.08	0.28	1.01	0.776
		Outgroup-by-HD interaction	$\delta[\text{Outgroup, HD}]$	0.06	0.40	1.01	0.894
		Range edge-by-HD interaction	$\delta[\text{Range edge, HD}]$	-0.15	0.40	1.01	0.698
		Body size in Range core	$\alpha[1]$	1.58	0.34	1.00	0.000
		Outgroup on Body-size effect	$\epsilon[\text{Outgroup}]$	-0.04	0.60	1.01	0.952
		Range edge on Body-size effect	$\epsilon[\text{Range edge}]$	0.06	0.58	1.00	0.931

		HD on Body-size effect in range core	ζ [HD]	-0.15	0.52	1.00	0.739
		Outgroup-by-HD interaction on Body-size effect	η [Outgroup, HD]	-0.23	0.73	1.00	0.767
		Range edge-by-HD interaction on Body-size effect	η [Range edge, HD]	-0.40	0.80	1.00	0.626
		Body condition	α [2]	0.58	0.16	1.00	0.001
		Temperature	α [3]	0.92	0.08	1.00	0.000
		Time interval	α [4]	-0.23	0.11	1.00	0.034
		Mesocosm	σ [θ]	0.12	0.10	1.02	
		Individual	σ [ι]	0.84	0.07	1.00	
		Intercept	α [0]	1.79	0.57	1.01	0.000
		Male	α [Sex Male]	-0.11	0.40	1.01	0.847
		Outgroup	β [Outgroup]	1.81	0.97	1.02	0.038
		Range edge	β [Range edge]	0.64	0.73	1.00	0.369
		High density	γ [HD]	0.55	0.70	1.01	0.352
		Outgroup-by-HD interaction	δ [Outgroup, HD]	-0.30	1.00	1.02	0.730
		Range edge-by-HD interaction	δ [Range edge, HD]	0.51	0.98	1.00	0.611
		Body size in Range core	α [1]	-1.68	0.67	1.00	0.012
		Outgroup on Body-size effect	ϵ [Outgroup]	-0.59	1.18	1.01	0.595
	$\psi_{B \rightarrow A}$	logit Range edge on Body-size effect	ϵ [Range edge]	-0.40	1.25	1.00	0.781
		HD on Body-size effect in range core	ζ [HD]	0.23	1.02	1.01	0.851
		Outgroup-by-HD interaction on Body-size effect	η [Outgroup, HD]	-0.41	1.39	1.00	0.869
		Range edge-by-HD interaction on Body-size effect	η [Range edge, HD]	1.33	1.80	1.00	0.440
		Body condition	α [2]	-1.18	0.46	1.00	0.006
		Temperature	α [3]	-1.17	0.23	1.00	0.000
		Time interval	α [4]	-0.12	0.18	1.00	0.492
		Mesocosm	σ [θ]	0.71	0.36	1.00	
		Individual	σ [ι]	1.39	0.25	1.00	

392 Dispersal

394 PPOs were below 35 % for all dispersal parameters except for η [Outgroup, HD] (PPO = 37 %) and
for η [Range edge, HD] (PPO = 40 %), indicating that the identifiability of our model parameters
was generally satisfying (Appendix 6B). The probability for an individual crayfish to disperse to an
396 exit trap during a time interval ranged from 0 to 0.3 (Figs. 3 e-h), while post-dispersal settlement

probability ranged from 0.3 to 1 (Figs. 3 i-l). Hence, crayfish were mainly sedentary during our
398 experiment.

400 Dispersal probability was almost 0 in crayfish smaller than 45 mm in cephalothorax length, but
increased up to 0.1-0.3 in the largest-bodied individuals, a significant body-size effect (Figs. 3e and
402 3f, $\alpha[1]$ parameter in Table 2). Specifically, mean dispersal probability increased from $3.07 \cdot 10^{-3}$ in a
small-bodied (25 mm cephalothorax length) crayfish to $2.86 \cdot 10^{-1}$ in a large-bodied crayfish (66 mm,
404 predictions computed for a Range-core female at low density, all other covariates were fixed to their
average value).

406
Increasing body condition, as well as warmer temperatures, also had significantly positive effects
408 on crayfish dispersal probability (Figs. 3g and 3h, respectively, $\alpha[2]$ and $\alpha[3]$ parameters in Table
2). When all other quantitative covariates were fixed to their average value, raising mean
410 temperature from its minimum (9.3°C) to its maximum (26.9°C) increased mean dispersal
probability during a 2.4-day time interval from $1.17 \cdot 10^{-2}$ to $9.46 \cdot 10^{-2}$ (predictions computed for a
412 Range-core female at low density)

414 Neither crayfish density nor geographic range did influence crayfish dispersal probability, either
directly or in interaction with body size (Figs 3e and 3f, Table 2). Interestingly, however, Outgroup
416 crayfish, which had an overall higher survival probability when “Range” was included only as a
main effect in the non-interaction model, were also those having the lowest dispersal probability
418 (Appendix 7).

420 The identifiability of model parameters for post-dispersal settlement probability was low, since 9
over 16 parameters had prior-posterior overlaps exceeding 35 % (Appendix 6C). However, the
422 specific response of post-dispersal settlement to major covariates (body size, body condition and

temperature, as displayed in Fig. 3) was identifiable ($\alpha[1]$, $\alpha[2]$ and $\alpha[3]$ parameters, PPOs ranging
424 from 13 to 27 %, Appendix 6C), and seemed generally symmetric to the response of dispersal
probability (Figs. 3i-l vs 3e-h, Table 2), suggesting that dispersing at time t did not alter dispersal
426 probability at time $t+1$ with respect to covariates.

428 DISCUSSION

430 Our study shows in the red swamp crayfish that survival and dispersal are under the control of both
individual- and environment-level factors (Fig. 1). Below, we first discuss these results separately
432 for each set of factors, and then we discuss the benefits of our findings for range-dynamics
modelling and applied issues with invasive *P. clarkii*.

434

Individual state variables

436 As predicted (Table 1), a larger body size increased crayfish survival probability, but this effect was
significant only under a high density and in Range-core crayfish. This result highlights density-
438 dependent selection for a larger body size in *P. clarkii*, as previously reported in other taxa
(collembolans, fish and lizards), and presumably resulting from a dominance of large-bodied
440 individuals in interference competition and cannibalism (see Edeline and Loeuille 2021 and
references therein). Accordingly, larger-bodied crayfish dominate in competition for shelters (Figler
442 et al., 1999; Rabeni, 1985; Ranta & Lindström, 1993), and crayfish with larger chela have a higher
probability of winning agonistic encounters (Graham & Angilletta, 2020).

444

A larger body size further had a strong positive effect on crayfish dispersal probability in our
446 experiment, a finding that is in line with a previous report of overland movements being
predominantly performed by sexually mature (i.e., adult) individuals in *P. clarkii* (Ramalho &
448 Anastácio, 2015). In *P. leniusculus*, both overland and in-stream dispersal also increase with

450 increasing body size (Bubb et al., 2004; Moorhouse & MacDonald, 2011; Thomas et al., 2018;
Wutz & Geist, 2013), suggesting that a larger body size elevates a general propensity for
movements, and might represent a major component of dispersal syndromes in crayfish.

452

This strong and taxonomically-conserved positive effect of body size on crayfish movements is
454 probably a response to multiple convergent selective pressures, in link with physiological and
ecological constraints. The mass-specific energetic cost of transport (in $\text{J kg}^{-1} \text{m}^{-1}$) of walking or
456 running animals decreases with body size (Peters, 1983), and a larger body size further confers a
higher body volume-to-surface ratio and, consequently, increases resistance to desiccation during
458 overland movements (Antoľ et al., 2021; Claussen et al., 2000). Additionally, walking speed also
increases with body size (Claussen et al., 2000), and larger-bodied dispersers thus have increased
460 probability to find a water body before desiccation. Finally, larger-bodied individuals dominate in
interference competition for food and contests for shelters (see above), and thus probably also enjoy
462 reduced costs of settlement during the colonization of novel habitats. Hence, larger-bodied crayfish
thus probably enjoy a strongly decreased cost-to-benefit ratio of dispersal.

464

In accordance with the general expectation that increased energy stores decrease the survival cost of
466 long-distance movements (Table 1), we found that body condition was positively linked to dispersal
probability in crayfish. However, in contrast with our predictions we found that a higher body
468 condition was associated with a lower survival probability. In shrimps, a higher nutritional status is
associated with increased moulting probability (Lemos & Weissman, 2021; Sharawy et al., 2019).
470 Maybe, in crayfish also a higher body condition was possibly indicative of a higher energetic status
and of a higher probability of moulting, which entails increased mortality in crayfish (Taugbøl &
472 Skurdal, 1992). In particular, newly-moulted crayfish have a soft shell and are strongly exposed to
cannibalism (E.E. pers. obs.).

474

Also contrary to our expectations (Table 1), survival and dispersal were both sex-independent. This
476 result suggests that, in *P. clarkii*, hazards and costs of dispersal are similar among males and
females. In *P. leniusculus*, dispersal was reported to be sometimes sex-independent (Bubb et al.,
478 2004; Galib et al., 2022), and sometimes male-biased (Wutz & Geist, 2013), which might suggest
that, in crayfish in general, the sex-dependency of dispersal is context-dependent and versatile.

480

Our predictions were also refuted regarding the effects of spatial sorting on competitive ability and
482 dispersal propensity. Range-core and Range-edge crayfish did not differ in either survival or
dispersal probabilities, nor they did in their response to crayfish density variation. Such an absence
484 of any effect of the geographic range contrasts with theoretical expectations (Table 1), and is maybe
surprising given the large genetic divergence of Range-edge crayfish at neutral markers (Bélouard
486 et al., 2019), but agrees with previous results showing no divergence among Range-core and Range-
edge crayfish populations in the Brière for the fluctuating asymmetry of morphological traits
488 (Bélouard et al., 2019). This apparent paradox may result from both genetic and selective processes.

490 Without selection, divergence in quantitative traits should equal the divergence at neutral marker
loci (Leinonen et al., 2008, 2013). However, this equality does not hold for complex traits such as
492 dispersal syndromes (Mackay et al., 2009; Saastamoinen et al., 2018), because allelic interactions
within (dominance) or between (epistasis) loci reduce the divergence at quantitative traits compared
494 to the neutral expectation (Goudet & Martin, 2007; Whitlock, 2008). Additionally, density-
dependent selection in edge habitats may have rapidly erased the effects of spatial sorting to make
496 Range-edge crayfish similarly dispersive and competitive as their Range-core counterparts. Strong
density-dependent selection is particularly likely to occur in the “boom” phase of large
498 demographic increase that sometimes immediately follows a colonization event, due to colonists
arriving in an empty ecological niche with lots of resources available (Strayer et al., 2017). Which

500 of these different mechanisms best explains phenotypic similarity among Range-core and Range-
edge crayfish in the Brière requires further investigation.

502

Environmental variables

504 As expected (Table 1), crayfish density in mesocosms had a marginally-significant negative effect
on the survival of Range-core crayfish. However, this effect was weaker than expected since it was
506 not significant when evaluated simultaneously across all crayfish origins or sex (main-effects
model, γ [High density] parameter, Appendix 7). This weakness probably reflects limited density
508 variation (from 3.0 to 6.3 individuals m^{-2}) in presence of an excess shelters (30 shelters for a
maximum of 19 crayfish introduced per mesocosm). Accordingly, an average relationship between
510 survival probability and stocking density for different experiment durations, as reconstructed from
literature data (Appendix 8), predicts largely overlapping survival probabilities at these two
512 densities and for an average time interval of 153 days corresponding to the duration of our
experiment (0.64-0.85 at 3.0 individuals m^{-2} and 0.58-0.80 at 6.3 individuals m^{-2}). Hence, we
514 conclude that the density variation we applied in our experiment, although located in the upper
range of naturally-observed densities, entailed relatively mild changes in competition and, hence,
516 resulted in weak effects on survival and no effect on dispersal.

518 Finally, temperature was the only environmental factor that had consistent effects on crayfish
demographic rates during our experiment: both survival and dispersal probabilities increased with
520 increasing temperature. All biological rates are expectedly maximized at intermediate temperatures
(Table 1), which correspond to a specific thermal optimum that may vary among species (Angilletta
522 Jr., 2009; Knies & Kingsolver, 2010). Crayfish are no exception and reach maximal rates of somatic
growth (Westhoff & Rosenberger, 2016) and dispersal (Claussen et al., 2000; Ramalho & Anastácio,
524 2015) at intermediate temperatures. However, when biological rates are measured over a narrow
temperature range, their hump-shaped relationship to temperature may not be observed, and authors

526 report either a positive or negative relationship. Accordingly, warmer temperatures are reported to
either increase or decrease rates of crayfish survival rate (Harlioğlu, 2009; Verhoef & Austin, 1999;
528 Webster et al., 2004). In *P. clarkii*, somatic growth rate, which is a putative surrogate for survival
rate, is maximal in the 22-30°C temperature range (Westhoff & Rosenberger, 2016), while overland
530 dispersal seems maximal at temperatures ranging from 16 to 24°C (Ramalho & Anastácio, 2015).
These temperatures will more often be met in the future, as hot days will get hotter and more
532 frequent (IPCC, 2021).

534 In Europe for instance, climate projections suggest that, by 2081-2100 and compared to the 1995-
2014 period, temperatures will increase by 1.25 to 2.65°C under the SSP1-2.6 scenario (low GHG
536 emissions), and by 2.45 to 4.85°C under the SSP5-8.5 scenario (high GHG emissions, IPCC, 2021,
page 14). Assuming an annual average reference temperature equal to 14.3°C, the mean summer
538 (May-September) temperature in Rennes city¹, these two warming scenarios are predicted by our
model to result in a 1.5 to 5.2 % increase in *P. clarkii* monthly survival probability, and in a 14 to 44
540 % decrease in the number of days needed for *P. clarkii* to reach an overland dispersal probability
equal to 1 (Appendix 9). Hence, potentially, climate warming will strongly expand the crayfish
542 geographic range.

544 *Implications for the projection and management of geographic ranges*

Correlative species distribution models based on climate envelopes also predict a geographic range
546 expansion of *P. clarkii* under projected climate change scenarios (Capinha et al., 2013; Liu et al.,
2011; Zhang et al., 2020). Our results are consistent with this prediction and, in addition, provide a
548 mechanistic, demography-based understanding of the underlying processes, that may serve as a
starting point to building process-explicit models of range dynamics for *P. clarkii* (Fig. 4).

550

1 <https://fr.wikipedia.org/wiki/Rennes>

In this context of an expanding geographic range of *P. clarkii*, improvement of containment strategies appears as a crucial management target. Our results suggest that crayfish containment could be based on manipulating crayfish body size. Specifically, mechanically culling large-bodied crayfish through, e.g., selective trapping, might promote an immediate crayfish containment through reducing both the average survival and dispersal probabilities in the population (Fig. 4). Culling-based containment might become even more efficient over the long term because, after several generations, selective removal of large-bodied individuals induces an evolutionary change towards smaller body sizes and earlier maturation, which would not only cause crayfish to disperse less, but are also traits generally associated with lower population growth rates (Evangelista et al., 2020, 2021; Heino et al., 2015; Hutchings, 2005). Such harvest-induced evolutionary changes are often considered as slowly reversible (Law, 2000), which is desirable from a long-term management perspective.




Fig. 4. Adaptation of the conceptual framework in Fig. 1 to *P. clarkii* range dynamics. Increased survival and dispersal are assumed to both increase the geographic range. Bold arrows indicate causality links that were investigated in the present study. The arrow pointing from population density to the positive effect of body size on survival indicates that this positive effect was present only under high crayfish density.

However, whether or not long-term culling of large-bodied individuals represents an efficient containment strategy remains to be effectively tested in the field. On the short term, removal of large-bodied individuals relaxes competition and favours a plastic increase in rates of somatic growth and reproduction of survivors, which led some authors to conclude that culling is inefficient at controlling crayfish populations (Gherardi et al., 2011). Additionally, locally-lower population densities may possibly trigger increased immigration from surrounding, non-culled areas where

densities are higher (Moorhouse & MacDonald, 2011). Hence, selective culling of large-bodied
580 crayfish might be better envisioned as a large-scale management strategy which, after an initially
strong culling effort, may provide good results if milder removal may be sustained over temporal
582 and spatial scales encompassing multiple crayfish generations across the whole (meta)population
range.

584

Conclusions

586 Our study demonstrates the power of individual mark-recapture in mesocosms to disentangle
climate, ecological and evolutionary drivers of demographic rates. Our results show that survival
588 and overland dispersal in *P. clarkii* are under the control of both climatic and ecological factors that
act at both the individual- and population levels, but suggest no rapid trait evolution at range edges.
590 Future studies should extend the approach to other demographic rates, i.e., somatic-growth and
maturation rates, as well as to measuring their dependency to a gradient of environmental
592 conditions. Complementary research is also needed to include other means of dispersal, because *P.*
clarkii actively disperses not only overland or through waterways (Cruz & Rebelo, 2007; Tréguier
594 et al., 2018), but also passively, as evidenced by the role played by humans or animals in their
current geographic distribution (Acevedo-Limón et al., 2020; Anastácio et al., 2014; Capinha et al.,
596 2013). Finally, future comprehensive mark-recapture experimental data could be coupled to more
readily-available count data from natural populations in integrated population models (Briscoe et
598 al., 2019; Pagel & Schurr, 2012; Schaub & Kéry, 2021). This new class of models will provide more
robust estimates of demographic rates under a wider range of ecological and climatic scenarios and,
600 we hope, will represent high-class tools for the management of established invasive crayfish
populations under global change.

602

Acknowledgements: We are grateful to Flavie Perrier and Owen Nino for help with collecting
604 crayfish in the marsh of Brière, and to Mattéo Le Goues for help with the genetic analysis of

crayfish from Rennes. Permission for crayfish capture and transport were provided by the
606 Préfecture de la Loire-Atlantique (license number 2019/SEE-53), and authorization for crayfish
retention was granted to U3E by the Direction Départementale de la Cohésion Sociale et de la
608 Protection des Populations d'Ille et Vilaine. EE received support from Rennes Métropole (AIS
18C0356). Through the use of the U3E INRAE facility, this work received support under the
610 program 'Investissements d'Avenir' launched by the French government and implemented by ANR
with the reference ANR-11-INBS-0001 AnaEE France.

612

Author contribution:

614 E.E., J-M.P. and E.P. designed the experiment, Y.B. and A.S. collected the data, E.E. analysed the
data and led manuscript writing. All authors participated in manuscript editing and revision.

616

Data availability statement:

618 Data and codes will be made available online upon acceptance of this manuscript.

References

Acevedo-Limón, L., Oficialdegui, F. J., Sánchez, M. I., & Clavero, M. (2020). Historical, human,
and environmental drivers of genetic diversity in the red swamp crayfish (*Procambarus*
clarkii) invading the Iberian Peninsula. *Freshwater Biology*, 65(8), 1460–1474.

<https://doi.org/10.1111/fwb.13513>

Anastácio, P. M., Ferreira, M. P., Banha, F., Capinha, C., & Rabaça, J. E. (2014). Waterbird-
mediated passive dispersal is a viable process for crayfish (*Procambarus clarkii*). *Aquatic*
Ecology, 48(1), 1–10. <https://doi.org/10.1007/s10452-013-9461-0>

Angeler, D. G., Sanchez-Carrillo, S., Alvarez-Cobelas, M., Cirujano, S., & Medina, L. (2003).
Exotic crayfish activity and its effects on water quality: Preliminary implications for the

alternative stable equilibria in Mediterranean wetlands. *Journal of Mediterranean Ecology*, 4, 13–22.

Angert, A. L., Bontrager, M. G., & Ågren, J. (2020). What do we really know about adaptation at range edges? *Annual Review of Ecology, Evolution, and Systematics*, 51(1), 341–361.

<https://doi.org/10.1146/annurev-ecolsys-012120-091002>

Angilletta Jr., M. J. (2009). *Thermal adaptation: A theoretical and empirical synthesis*. Oxford University Press.

<http://www.oxfordscholarship.com/view/10.1093/acprof:oso/9780198570875.001.1/acprof-9780198570875>

Antoń, A., Berg, M. P., & Verberk, W. C. (2021). Effects of body size and lung type on desiccation resistance, hypoxia tolerance and thermal preference in two terrestrial isopods species.

Journal of Insect Physiology, 132, 104247. <https://doi.org/10.1016/j.jinsphys.2021.104247>

Begon, M., Townsend, C. R., & Harper, J. L. (2005). *Ecology: From individuals to ecosystems* (4th ed.). Blackwell. <https://www.wiley.com/en-us/Ecology>

[https://www.wiley.com/en-us/Ecology](https://www.wiley.com/en-us/Ecology+From+Individuals+to+Ecosystems%2C+4th+Edition-p-9781405111171)

Bélouard, N., Paillisson, J.-M., Oger, A., Besnard, A.-L., & Petit, E. J. (2019). Genetic drift during the spread phase of a biological invasion. *Molecular Ecology*, 28(19), 4375–4387.

<https://doi.org/10.1111/mec.15238>

Biro, P. A., & Post, J. R. (2008). Rapid depletion of genotypes with fast growth and bold personality traits from harvested fish populations. *Proceedings of the National Academy of Sciences of the United States of America*, 105(8), 2919–2922. <https://doi.org/10.1073/pnas.0708159105>

<https://doi.org/10.1073/pnas.0708159105>

Blums, P., Nichols, J. D., Hines, J. E., Lindberg, M. S., & Mednis, A. (2005). Individual quality, survival variation and patterns of phenotypic selection on body condition and timing of nesting in birds. *Oecologia*, 143(3), 365–376. <https://doi.org/10.1007/s00442-004-1794-x>

<https://doi.org/10.1007/s00442-004-1794-x>

Bonte, D., & Peña, E. D. L. (2009). Evolution of body condition-dependent dispersal in metapopulations. *Journal of Evolutionary Biology*, 22(6), 1242–1251.

<https://doi.org/10.1111/j.1420-9101.2009.01737.x>

Briscoe, N. J., Elith, J., Salguero-Gómez, R., Lahoz-Monfort, J. J., Camac, J. S., Giljohann, K. M., Holden, M. H., Hradsky, B. A., Kearney, M. R., McMahon, S. M., Phillips, B. L., Regan, T.

- J., Rhodes, J. R., Vesk, P. A., Wintle, B. A., Yen, J. D. L., & Guillera-Aroita, G. (2019). Forecasting species range dynamics with process-explicit models: Matching methods to applications. *Ecology Letters*, 22(11), 1940–1956. <https://doi.org/10.1111/ele.13348>
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology*, 85(7), 1771–1789. <https://doi.org/10.1890/03-9000>
- Bubb, D. H., Thom, T. J., & Lucas, M. C. (2004). Movement and dispersal of the invasive signal crayfish *Pacifastacus leniusculus* in upland rivers. *Freshwater Biology*, 49(3), 357–368. <https://doi.org/10.1111/j.1365-2426.2003.01178.x>
- Burton, O. J., Phillips, B. L., & Travis, J. M. J. (2010). Trade-offs and the evolution of life-histories during range expansion. *Ecology Letters*, 13(10), 1210–1220. <https://doi.org/10.1111/j.1461-0248.2010.01505.x>
- Capinha, C., Brotons, L., & Anastácio, P. (2013). Geographical variability in propagule pressure and climatic suitability explain the European distribution of two highly invasive crayfish. *Journal of Biogeography*, 40(3), 548–558. <https://doi.org/10.1111/jbi.12025>
- Charlesworth, B. (1994). *Evolution in age-structured populations* (2nd ed.). Cambridge University Press.
- Chuang, A., & Peterson, C. R. (2016). Expanding population edges: Theories, traits, and trade-offs. *Global Change Biology*, 22(2), 494–512. <https://doi.org/10.1111/gcb.13107>
- Claussen, D. L., Hopper, R. A., & Sanker, A. M. (2000). The effects of temperature, body size, and hydration state on the terrestrial locomotion of the crayfish *Orconectes rusticus*. *Journal of Crustacean Biology*, 20(2), 218–223. <https://doi.org/10.1163/20021975-99990033>
- Clobert, J., Baguette, M., Benton, T. G., & Bullock, J. M. (Eds.). (2012). *Dispersal ecology and evolution*. Oxford University Press. <http://www.oxfordscholarship.com/view/10.1093/acprof:oso/9780199608898.001.0001/acprof-9780199608898>
- Clobert, J., Le Galliard, J.-F., Cote, J., Meylan, S., & Massot, M. (2009). Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. *Ecology Letters*, 12(3), 197–209. <https://doi.org/10.1111/j.1461-0248.2008.01267.x>

- Clobert, J., Massot, M., & Galliard, J.-F. L. (2012). Multi-determinism in natal dispersal: The common lizard as a model system. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal Ecology and Evolution*. Oxford University Press.
<https://doi.org/10.1093/acprof:oso/9780199608898.003.0003>
- Coignet, A., Pinet, F., & Souty-Grosset, C. (2012). Estimating population size of the red swamp crayfish (*Procambarus clarkii*) in fish-ponds (Brenne, Central France). *Knowledge and Management of Aquatic Ecosystems*, 406. <https://doi.org/10.1051/kmae/2012019>
- Correia, A. M., & Anastácio, P. M. (2008). Shifts in aquatic macroinvertebrate biodiversity associated with the presence and size of an alien crayfish. *Ecological Research*, 23(4), 729–734. <https://doi.org/10.1007/s11284-007-0433-5>
- Correia, A. M., & Bandeira, N. (2004). Seasonal availability of *Procambarus clarkii* in the Tejo River basin, Portugal. *Freshwater Crayfish*, 14(1), 190–196.
- Cruz, M. J., & Rebelo, R. (2007). Colonization of freshwater habitats by an introduced crayfish, *Procambarus clarkii*, in Southwest Iberian Peninsula. *Hydrobiologia*, 575(1), 191–201. <https://doi.org/10.1007/s10750-006-0376-9>
- Duckworth, R. A. (2008). Adaptive dispersal strategies and the dynamics of a range expansion. *The American Naturalist*, 172(S1), S4–S17. <https://doi.org/10.1086/588289>
- Edeline, E. (2016). Life history evolution, human impacts on. In R. Kliman (Ed.), *The encyclopedia of evolutionary biology* (1st ed., pp. 335–342). Academic Press.
- Edeline, E., & Loeuille, N. (2021). Size-dependent eco-evolutionary feedbacks in harvested systems. *Oikos*, 130(10), 1636–1649. <https://doi.org/10.1111/oik.08592>
- Evangelista, C., Dupeu, J., Sandkjenn, J., Pauli, B. D., Herland, A., Meriguet, J., Vøllestad, L. A., & Edeline, E. (2021). Ecological ramifications of adaptation to size-selective mortality. *Royal Society Open Science*, 8(10), 210842. <https://doi.org/10.1098/rsos.210842>
- Evangelista, C., Vøllestad, L. A., Diaz Pauli, B., & Edeline, E. (2020). Density-dependent consequences of size-selective induced life-history changes to population fitness in medaka (*Oryzias latipes*). *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 1741–1748. <https://doi.org/10.1139/cjfas-2019-0406>

- Figler, M. H., Cheverton, H. M., & Blank, G. S. (1999). Shelter competition in juvenile red swamp crayfish (*Procambarus clarkii*): The influences of sex differences, relative size, and prior residence. *Aquaculture*, 178(1), 63–75. [https://doi.org/10.1016/S0044-8486\(99\)00114-3](https://doi.org/10.1016/S0044-8486(99)00114-3)
- Fronhofer, E. A., & Altermatt, F. (2015). Eco-evolutionary feedbacks during experimental range expansions. *Nature Communications*, 6. <http://dx.doi.org/10.1038/ncomms7844>
- Galib, S. M., Sun, J., Twiss, S. D., & Lucas, M. C. (2022). Personality, density and habitat drive the dispersal of invasive crayfish. *Scientific Reports*, 12(1), 1114. <https://doi.org/10.1038/s41598-021-04228-1>
- Gaston, K. J. (2009). Geographic range limits: Achieving synthesis. *Proceedings of the Royal Society B: Biological Sciences*, 276(1661), 1395–1406. <https://doi.org/10.1098/rspb.2008.1480>
- Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics in Medicine*, 27(15), 2865–2873. <https://doi.org/10.1002/sim.3107>
- Gelman, A., Jakulin, A., Pittau, M. G., & Su, Y.-S. (2008). A weakly informative default prior distribution for logistic and other regression models. *Annals of Applied Statistics*, 2(4), 1360–1383.
- Gelman, A., & Rubin, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 4, 457–472. <https://doi.org/10.1214/ss/1177011136>
- Gherardi, F., & Acquistapace, P. (2007). Invasive crayfish in Europe: The impact of *Procambarus clarkii* on the littoral community of a Mediterranean lake. *Freshwater Biology*, 52(7), 1249–1259. <https://doi.org/10.1111/j.1365-2427.2007.01760.x>
- Gherardi, F., Aquiloni, L., Diéguez-Urbeondo, J., & Tricarico, E. (2011). Managing invasive crayfish: Is there a hope? *Aquatic Sciences*, 73(2), 185–200. <https://doi.org/10.1007/s00027-011-0181-z>
- Gherardi, F., & Lazzara, L. (2006). Effects of the density of an invasive crayfish (*Procambarus clarkii*) on pelagic and surface microalgae in a Mediterranean wetland. *Archiv Für Hydrobiologie*, 165(3), 401–414. <https://doi.org/10.1127/0003-9136/2006/0165-0401>

- Gimenez, O., Lebreton, J.-D., Gaillard, J.-M., Choquet, R., & Pradel, R. (2012). Estimating demographic parameters using hidden process dynamic models. *Theoretical Population Biology*, 82(4), 307–316. <https://doi.org/10.1016/j.tpb.2012.02.001>
- Gimenez, O., Morgan, B. J. T., & Brooks, S. P. (2009). Weak identifiability in models for mark-recapture-recovery data. In D. L. Thomson, E. G. Cooch, & M. J. Conroy (Eds.), *Modeling Demographic Processes In Marked Populations* (pp. 1055–1067). Springer US. https://doi.org/10.1007/978-0-387-78151-8_48
- Goudet, J., & Martin, G. (2007). Under neutrality, $QST \leq FST$ when there is dominance in an island model. *Genetics*, 176(2), 1371–1374. <https://doi.org/10.1534/genetics.106.067173>
- Graham, Z. A., & Angilletta, M. J. (2020). Claw size predicts dominance within and between invasive species of crayfish. *Animal Behaviour*, 166, 153–161. <https://doi.org/10.1016/j.anbehav.2020.06.021>
- Green, N., Bentley, M., Stebbing, P., Andreou, D., & Britton, R. (2018). Trapping for invasive crayfish: Comparisons of efficacy and selectivity of baited traps versus novel artificial refuge traps. *Knowledge and Management of Aquatic Ecosystems*, 419(15). <https://doi.org/10.1051/kmae/2018007>
- Gros, A., Hovestadt, T., & Poethke, H. J. (2008). Evolution of sex-biased dispersal: The role of sex-specific dispersal costs, demographic stochasticity, and inbreeding. *Ecological Modelling*, 219(1), 226–233. <https://doi.org/10.1016/j.ecolmodel.2008.08.014>
- Guan, R.-Z. (2000). Abundance and production of the introduced signal crayfish in a British lowland river. *Aquaculture International*, 8(1), 59–76. <https://doi.org/10.1023/A:1009272916339>
- Harhoğlu, M. (2009). A comparison of the growth and survival of two freshwater crayfish species, *Astacus leptodactylus* Eschscholtz and *Pacifastacus leniusculus* (Dana), under different temperature and density regimes. *Aquaculture International*, 17(1), 31–43. <https://doi.org/10.1007/s10499-008-9177-7>
- Heino, M., Díaz Pauli, B., & Dieckmann, U. (2015). Fisheries-induced evolution. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 461–480. <https://doi.org/10.1146/annurev-ecolsys-112414-054339>

- Hudina, S., Hock, K., Žganec, K., & Lucić, A. (2012). Changes in population characteristics and structure of the signal crayfish at the edge of its invasive range in a European river. *International Journal of Limnology*, 48(1), 3–11. <https://doi.org/10.1051/limn/2011051>
- Hudina, S., Žganec, K., & Hock, K. (2015). Differences in aggressive behaviour along the expanding range of an invasive crayfish: An important component of invasion dynamics. *Biological Invasions*, 17(11), 3101–3112. <https://doi.org/10.1007/s10530-015-0936-x>
- Huner, J. V., & Barr, J. E. (1991). *Red swamp crayfish: Biology and exploitation* (Fisheries & Aquaculture, p. 136) [Technical Report]. National Oceanic and Atmospheric Administration.
- Hutchings, J. A. (2005). Life history consequences of overexploitation to population recovery in Northwest Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences*, 62(4), 824–832. <https://doi.org/10.1139/f05-081>
- IPCC. (2021). Summary for policymakers. In V. Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, Y. Chen, L. Goldfarb, M. I. Gomis, J. B. R. Matthews, S. Berger, M. Huang, O. Yelekçi, R. Yu, B. Zhou, E. Lonnoy, T. K. Maycock, T. Waterfield, K. Leitzell, & N. Caud (Eds.), *Climate Change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the Intergovernmental Panel on Climate Change* (1st ed., p. 40). Cambridge University Press.
- Jackson, M. C., Jones, T., Milligan, M., Sheath, D., Taylor, J., Ellis, A., England, J., & Grey, J. (2014). Niche differentiation among invasive crayfish and their impacts on ecosystem structure and functioning. *Freshwater Biology*, 59(6), 1123–1135. <https://doi.org/10.1111/fwb.12333>
- Kellner, K. (2019). *JagsUI: a wrapper around “rjags” to streamline “JAGS” analyses*. <https://CRAN.R-project.org/package=jagsUI>
- Kendall, B. E., Briggs, C. J., Murdoch, W. W., Turchin, P., Ellner, S. P., McCauley, E., Nisbet, R. M., & Wood, S. N. (1999). Why do populations cycle? A synthesis of statistical and mechanistic modeling approaches. *Ecology*, 80(6), 1789–1805. [https://doi.org/10.1890/0012-9658\(1999\)080\[1789:WDPCAS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1789:WDPCAS]2.0.CO;2)
- Kéry, M., & Schaub, M. (2012). *Bayesian population analysis using WinBUGS* (1st ed.). Academic Press.

- Kirkwood, T. B. L. (2001). Sex and ageing. *Experimental Gerontology*, 36(3), 413–418.
[https://doi.org/10.1016/S0531-5565\(00\)00255-2](https://doi.org/10.1016/S0531-5565(00)00255-2)
- Kisdi, E., Utz, M., & Gyllenberg, M. (2012). Evolution of condition-dependent dispersal. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal Ecology and Evolution* (1st ed.). Oxford University Press.
<https://doi.org/10.1093/acprof:oso/9780199608898.003.0011>
- Knies, J. L., & Kingsolver, J. G. (2010). Erroneous Arrhenius: Modified Arrhenius model best explains the temperature dependence of ectotherm fitness. *The American Naturalist*, 176(2), 227–233. <https://doi.org/10.1086/653662>
- Krebs, C. J. (2014). *Ecology: The experimental analysis of distribution and abundance* (6th ed.). Pearson.
- Kruschke, J. K., & Meredith, M. (2021). *BEST: Bayesian Estimation Supersedes the t-Test* (0.5.4). <https://CRAN.R-project.org/package=BEST>
- Kuparinen, A., & Festa-Bianchet, M. (2017). Harvest-induced evolution: Insights from aquatic and terrestrial systems. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1712), 20160036. <https://doi.org/10.1098/rstb.2016.0036>
- Lamontagne, S., & Rasmussen, J. B. (1993). Estimating crayfish density in lakes using quadrats: Maximizing precision and efficiency. *Canadian Journal of Fisheries and Aquatic Sciences*, 50(3), 623–626. <https://doi.org/10.1139/f93-071>
- Law, R. (2000). Fishing, selection, and phenotypic evolution. *ICES Journal of Marine Science: Journal Du Conseil*, 57(3), 659–668. <https://doi.org/10.1006/jmsc.2000.0731>
- Le Galliard, J.-F., Massot, M., & Clobert, J. (2012). Dispersal and range dynamics in changing climates: A review. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal Ecology and Evolution* (pp. 317–335).
<https://doi.org/10.1093/acprof:oso/9780199608898.003.0025>
- Lebreton, J.-D., Nichols, J. D., Barker, R. J., Pradel, R., & Spendelov, J. A. (2009). Modeling individual animal histories with multistate capture–recapture models. *Advances in Ecological Research*, 41, 87–173.

- Leinonen, T., McCairns, R. J. S., O'Hara, R. B., & Merila, J. (2013). QST-FST comparisons: Evolutionary and ecological insights from genomic heterogeneity. *Nature Reviews Genetics*, *14*(3), 179–190. <https://doi.org/10.1038/nrg3395>
- Leinonen, T., O'Hara, R. B., Cano, J. M., & Merilä, J. (2008). Comparative studies of quantitative trait and neutral marker divergence: A meta-analysis. *Journal of Evolutionary Biology*, *21*(1), 1–17. <https://doi.org/10.1111/j.1420-9101.2007.01445.x>
- Lemos, D., & Weissman, D. (2021). Moulting in the grow-out of farmed shrimp: A review. *Reviews in Aquaculture*, *13*(1), 5–17. <https://doi.org/10.1111/raq.12461>
- Liu, X., Guo, Z., Ke, Z., Wang, S., & Li, Y. (2011). Increasing potential risk of a global aquatic invader in Europe in contrast to other continents under future climate change. *PloS One*, *6*(3), e18429. <https://doi.org/10.1371/journal.pone.0018429>
- Lodge, D. M., Deines, A., Gherardi, F., Yeo, D. C. J., Arcella, T., Baldrige, A. K., Barnes, M. A., Chadderton, W. L., Feder, J. L., Gantz, C. A., Howard, G. W., Jerde, C. L., Peters, B. W., Peters, J. A., Sargent, L. W., Turner, C. R., Wittmann, M. E., & Zeng, Y. (2012). Global introductions of crayfishes: Evaluating the impact of species invasions on ecosystem services. *Annual Review of Ecology, Evolution, and Systematics*, *43*(1), 449–472. <https://doi.org/10.1146/annurev-ecolsys-111511-103919>
- Lodge, D. M., Kershner, M. W., Aloï, J. E., & Covich, A. P. (1994). Effects of an omnivorous crayfish (*Orconectes rusticus*) on a freshwater littoral food web. *Ecology*, *75*(5), 1265–1281. <https://doi.org/10.2307/1937452>
- Mackay, T. F. C., Stone, E. A., & Ayroles, J. F. (2009). The genetics of quantitative traits: Challenges and prospects. *Nature Reviews Genetics*, *10*(8), 565–577. <https://doi.org/10.1038/nrg2612>
- Matsuzaki, S. S., Usio, N., Takamura, N., & Washitani, I. (2009). Contrasting impacts of invasive engineers on freshwater ecosystems: An experiment and meta-analysis. *Oecologia*, *158*(4), 673–686. <https://doi.org/10.1007/s00442-008-1180-1>
- Matthysen, E. (2012). Multicausality of dispersal: A review. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal Ecology and Evolution* (pp. 3–18). Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780199608898.003.0001>

- Messenger, M. L., & Olden, J. D. (2019). Phenotypic variability of rusty crayfish (*Faxonius rusticus*) at the leading edge of its riverine invasion. *Freshwater Biology*, 64(6), 1196–1209.
<https://doi.org/10.1111/fwb.13295>
- Miller, T. E. X., Angert, A. L., Brown, C. D., Lee-Yaw, J. A., Lewis, M., Lutscher, F., Marculis, N. G., Melbourne, B. A., Shaw, A. K., Szűcs, M., Tabares, O., Usui, T., Weiss-Lehman, C., & Williams, J. L. (2020). Eco-evolutionary dynamics of range expansion. *Ecology*, 101(10), e03139. <https://doi.org/10.1002/ecy.3139>
- Moorhouse, T. P., & MacDonald, D. W. (2011). Immigration rates of signal crayfish (*Pacifastacus leniusculus*) in response to manual control measures. *Freshwater Biology*, 56(5), 993–1001.
<https://doi.org/10.1111/j.1365-2427.2010.02528.x>
- Ochocki, B. M., & Miller, T. E. X. (2017). Rapid evolution of dispersal ability makes biological invasions faster and more variable. *Nature Communications*, 8(1), 14315.
<https://doi.org/10.1038/ncomms14315>
- Oficialdegui, F. J., Clavero, M., Sánchez, M. I., Green, A. J., Boyero, L., Michot, T. C., Klose, K., Kawai, T., & Lejeusne, C. (2019). Unravelling the global invasion routes of a worldwide invader, the red swamp crayfish (*Procambarus clarkii*). *Freshwater Biology*, 64(8), 1382–1400. <https://doi.org/10.1111/fwb.13312>
- Oficialdegui, F. J., Sánchez, M. I., Lejeusne, C., Pacini, N., & Clavero, M. (2020). Brought more than twice: The complex introduction history of the red swamp crayfish into Europe. *Knowledge and Management of Aquatic Ecosystems*, 421.
<https://doi.org/10.1051/kmae/2019044>
- Pagel, J., & Schurr, F. M. (2012). Forecasting species ranges by statistical estimation of ecological niches and spatial population dynamics. *Global Ecology and Biogeography*, 21(2), 293–304.
<https://doi.org/10.1111/j.1466-8238.2011.00663.x>
- Peters, R. H. (1983). *The ecological implications of body size*. Cambridge University Press; Cambridge Core. <https://www.cambridge.org/core/books/ecological-implications-of-body-size/4D86337571D7F26E76F885B2548FCBFB>

- Phillips, B. L. (2016). Behaviour on invasion fronts, and the behaviour of invasion fronts. In J. L. Weis & D. Sol (Eds.), *Biological Invasions and Animal Behaviour* (pp. 82–95). Cambridge University Press; Cambridge Core. <https://doi.org/10.1017/CBO9781139939492.007>
- Phillips, B. L., Brown, G. P., & Shine, R. (2010). Life-history evolution in range-shifting populations. *Ecology*, *91*(6), 1617–1627. <https://doi.org/10.1890/09-0910.1>
- Pilotto, F., Free, G., Crosa, G., Sena, F., Ghiani, M., & Cardoso, A. C. (2008). The invasive crayfish *Orconectes limosus* in Lake Varese: Estimating abundance and population size structure in the context of habitat and methodological constraints. *Journal of Crustacean Biology*, *28*(4), 633–640. <https://doi.org/10.1651/07-2967.1>
- Plummer, M. (2003). *JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling*. Proceedings of the 3rd international workshop on distributed statistical computing, Vienna, Austria.
- R Core Team. (2022). *R: a language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rabeni, C. F. (1985). Resource partitioning by stream-dwelling crayfish: The influence of body size. *The American Midland Naturalist*, *113*(1), 20–29. JSTOR. <https://doi.org/10.2307/2425344>
- Ramalho, R. O., & Anastácio, P. M. (2015). Factors inducing overland movement of invasive crayfish (*Procambarus clarkii*) in a ricefield habitat. *Hydrobiologia*, *746*(1), 135–146. <https://doi.org/10.1007/s10750-014-2052-9>
- Ranta, E., & Lindström, K. (1993). Body size and shelter possession in mature signal crayfish, *Pacifastacus leniusculus*. *Annales Zoologici Fennici*, *30*(2), 125–132. JSTOR.
- Rodríguez, C. F., Bécares, E., & Fernández-Aláez, M. (2003). Shift from clear to turbid phase in Lake Chozas (NW Spain) due to the introduction of American red swamp crayfish (*Procambarus clarkii*). *Hydrobiologia*, *506*(1), 421–426. <https://doi.org/10.1023/B:HYDR.0000008626.07042.87>
- Rodríguez-Pérez, H., Hilaire, S., & Mesléard, F. (2016). Temporary pond ecosystem functioning shifts mediated by the exotic red swamp crayfish (*Procambarus clarkii*): A mesocosm study. *Hydrobiologia*, *767*(1), 333–345. <https://doi.org/10.1007/s10750-015-2523-7>

- Roff, D. A. (2002). *The evolution of life histories: Theory and analysis* (1st ed.). Chapman & Hall.
- Saastamoinen, M., Bocedi, G., Cote, J., Legrand, D., Guillaume, F., Wheat, C. W., Fronhofer, E. A., Garcia, C., Henry, R., Husby, A., Baguette, M., Bonte, D., Coulon, A., Kokko, H., Matthysen, E., Niitepõld, K., Nonaka, E., Stevens, V. M., Travis, J. M. J., ... del Mar Delgado, M. (2018). Genetics of dispersal. *Biological Reviews*, 93(1), 574–599. <https://doi.org/10.1111/brv.12356>
- Savini, D., Occhipinti–Ambrogi, A., Marchini, A., Tricarico, E., Gherardi, F., Olenin, S., & Gollasch, S. (2010). The top 27 animal alien species introduced into Europe for aquaculture and related activities. *Journal of Applied Ichthyology*, 26, 1–7. <https://doi.org/10.1111/j.1439-0426.2010.01503.x>
- Schaub, M., & Kéry, M. (2021). *Integrated population models: Theory and ecological applications with R and JAGS* (1st ed.). Academic Press.
- Sharawy, Z. Z., Hufnagl, M., & Temming, A. (2019). A condition index based on dry weight as a tool to estimate in-situ moult increments of decapod shrimp: Investigating the effects of sex, year and measuring methods in brown shrimp (*Crangon crangon*). *Journal of Sea Research*, 152, 101762. <https://doi.org/10.1016/j.seares.2019.05.004>
- Souty-Grosset, C., Anastácio, P. M., Aquiloni, L., Banha, F., Choquer, J., Chucholl, C., & Tricarico, E. (2016). The red swamp crayfish *Procambarus clarkii* in Europe: Impacts on aquatic ecosystems and human well-being. *Limnologica*, 58, 78–93. <https://doi.org/10.1016/j.limno.2016.03.003>
- Stenseth, N. Chr., Mysterud, A., Ottersen, G., Hurrell, J. W., Chan, K. S., & Lima, M. (2002). Ecological effects of climate fluctuations. *Science*, 297(5585), 1292–1296. <https://doi.org/10.1126/science.1071281>
- Strayer, D. L., D’Antonio, C. M., Essl, F., Fowler, M. S., Geist, J., Hilt, S., Jarić, I., Jöhnk, K., Jones, C. G., Lambin, X., Latzka, A. W., Pergl, J., Pyšek, P., Robertson, P., von Schmalensee, M., Stefansson, R. A., Wright, J., & Jeschke, J. M. (2017). Boom-bust dynamics in biological invasions: Towards an improved application of the concept. *Ecology Letters*, 20(10), 1337–1350. <https://doi.org/10.1111/ele.12822>

- Taugbøl, T., & Skurdal, J. (1992). Growth, mortality and moulting rate of noble crayfish, *Astacus astacus* L., juveniles in aquaculture experiments. *Aquaculture Research*, 23(4), 411–420. <https://doi.org/10.1111/j.1365-2109.1992.tb00785.x>
- Thomas, J. R., Fisher, J., Cable, J., & Griffiths, S. W. (2018). Terrestrial dispersal of invasive signal crayfish during vulnerable life stages. *Behavioural Processes*, 157, 204–207. <https://doi.org/10.1016/j.beproc.2018.09.014>
- Townsend Peterson, A., Papeş, M., & Soberón, J. (2015). Mechanistic and correlative models of ecological niches. *European Journal of Ecology*, 1(2), 28–38. <https://doi.org/10.1515/eje-2015-0014>
- Travis, J. M. J., & Dytham, C. (2012). Dispersal and climate change: A review of theory. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal Ecology and Evolution* (1st ed., pp. 337–348). Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780199608898.003.0026>
- Tréguier, A., Roussel, J.-M., Bélouard, N., & Paillisson, J.-M. (2018). Is it a hindrance for an invasive aquatic species to spread across scattered habitat patches? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28(3), 610–618. <https://doi.org/10.1002/aqc.2887>
- Twardochleb, L. A., Olden, J. D., & Larson, E. R. (2013). A global meta-analysis of the ecological impacts of nonnative crayfish. *Freshwater Science*, 32(4), 1367–1382. <https://doi.org/10.1899/12-203.1>
- Verhoef, G. D., & Austin, C. M. (1999). Combined effects of temperature and density on the growth and survival of juveniles of the Australian freshwater crayfish, *Cherax destructor* Clark,; Part 1. *Aquaculture*, 170(1), 37–47. [https://doi.org/10.1016/S0044-8486\(98\)00394-9](https://doi.org/10.1016/S0044-8486(98)00394-9)
- Webster, C. D., Thompson, K. R., Muzinic, L. A., Yancey, D. H., Dasgupta, S., Xiong, Y. L., Rouse, D. B., & Manomaitis, L. (2004). A preliminary assessment of growth, survival, yield, and economic return of Australian red claw crayfish, *Cherax quadricarinatus*, stocked at three densities in earthen ponds in a cool, temperate climate. *Journal of Applied Aquaculture*, 15(3–4), 37–50. https://doi.org/10.1300/J028v15n03_03

- Westhoff, J. T., & Rosenberger, A. E. (2016). A global review of freshwater crayfish temperature tolerance, preference, and optimal growth. *Reviews in Fish Biology and Fisheries*, 26(3), 329–349. <https://doi.org/10.1007/s11160-016-9430-5>
- Whitlock, M. C. (2008). Evolutionary inference from QST. *Molecular Ecology*, 17(8), 1885–1896. <https://doi.org/10.1111/j.1365-294X.2008.03712.x>
- Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of senescence. *Evolution*, 11(4), 398–411. <https://doi.org/10.2307/2406060>
- Williams, J. L., Kendall, B. E., & Levine, J. M. (2016). Rapid evolution accelerates plant population spread in fragmented experimental landscapes. *Science*, 353(6298), 482. <https://doi.org/10.1126/science.aaf6268>
- Wutz, S., & Geist, J. (2013). Sex- and size-specific migration patterns and habitat preferences of invasive signal crayfish (*Pacifastacus leniusculus* Dana). *Limnologica*, 43(2), 59–66. <https://doi.org/10.1016/j.limno.2012.02.002>
- Závorka, L., Lassus, R., Britton, J. R., & Cucherousset, J. (2020). Phenotypic responses of invasive species to removals affect ecosystem functioning and restoration. *Global Change Biology*, 26(10), 5693–5704. <https://doi.org/10.1111/gcb.15271>
- Zhang, Z., Capinha, C., Usio, N., Weterings, R., Liu, X., Li, Y., Landeira, J. M., Zhou, Q., & Yokota, M. (2020). Impacts of climate change on the global potential distribution of two notorious invasive crayfishes. *Freshwater Biology*, 65(3), 353–365. <https://doi.org/10.1111/fwb.13429>

APPENDIX 1. Genetics of the Rennes (“Outgroup”) crayfish population.

Aim — The invasion history of the pond from which our Outgroup individuals originate was unknown. We tested the hypothesis that this invasion followed the same introduction route as red swamp crayfish (*Procambarus clarkii*) from Brière, namely, the USA-Kenya introduction route (Oficialdegui et al., 2019). Besides looking for the origin of this specific pond, we also tested new ways for obtaining DNA material from crayfish. In that aim, we collected three kinds of samples on all individuals: we sampled with sterile cotton swabs (i) the buccal region and (ii) haemolymph leaking out of the abdomen while PIT-tagging. We also collected (iii) muscle from the abdomen when individuals were sacrificed as a classical way to obtain genomic DNA.

Methods — 30 individuals were collected and stocked in an outdoor mesocosm at the PEARL (https://www6.rennes.inrae.fr/u3e_eng/). Swabs were collected on all individuals at two capture occasions to check whether results were reproducible. At the first capture event, individuals were marked with Biomark HTP8 PIT-tags and leaking haemolymph was collected with a cotton swab. At the second capture event three days later, crayfish were jabbed again with a PIT-tag syringe and a second haemolymph swab was collected, then individuals were sacrificed and their muscle was sampled.

DNA was extracted from all (n=150) samples with the NucleoSpin Tissue kit (Macherey-Nagel), following manufacturer’s instructions. This protocol was slightly modified for swabs, for which we doubled buffer volumes: we added 360µl of the initial (T1) lysis buffer and 25µl of proteinase K to the swab, and then used 400µl of buffer B3 and 420µl of ethanol in the following steps. All extracts were quantified using a DS-11 spectrophotometer (DeNovix).

Because at least one of our sampling protocol targeted extra-corporal DNA (buccal swabs), which is potentially degraded, we chose to amplify and sequence a short mitochondrial DNA fragment (Broquet et al., 2007). Primers were designed from the alignment of published CO1 sequences: *P. clarkii* haplotypes from Oficialdegui et al. (2019) (NCBI accession numbers MK026671 to MK026718) and six CO1 sequences taken from other crayfish species (species names and NCBI accession numbers: *Procambarus virginalis*, LR884234.1; *Procambarus fallax*, LC228303.1; *Procambarus riojae*, KX238226.1; *Procambarus hoffmanni*, KX238193.1; *Procambarus geminus*, JX514457.1; *Pacifastacus leniusculus*, MK439898.1). The resulting primers, PC-FCO1202 (5'-GGGGTATAGTTGAGAGAGGAGT-3') and PC-RCO1202 (5'-

GGTATTCGATCCATGGTTATCCC -3'), amplify a 202bp CO1 fragment in *P. clarkii*. PCR amplifications were carried out in a total volume of 50 µl containing 1µM of each primer, 1mM MgCL₂, 1X GoTaq G2 (Promega) PCR Mix, 1.25 units of GoTaq G2 (Promega) Taq Polymerase, 2µl DNA, the whole being completed with sterilized water. DNA samples obtained from muscles were diluted 10 times to reach DNA concentrations that are comparable to DNA concentrations obtained from swabs (see below). Thermocycling used a touchdown approach: one 2-min step at 95°C, 30 cycles of 30 s at 95°C, 30 s at 65°C to 59°C, 90 s at 72°C and one final 5-min step at 72°C before incubation at 15°C. The touchdown protocol implies that the first cycle had a hybridization temperature at 65°C, the following six cycles with hybridization temperatures decreasing down to 59°C by steps of 1°C, and the remaining 24 cycles at 59°C. Amplification success was evaluated by electrophoresis and BET visualization of 5µl of amplification products in 1.75% agarose gels run 30 min at 100V in TAE buffer.

Twelve PCR products issued in balanced numbers (n=4) from muscles, buccal and haemolymph swabs from different individuals were then sent to GenoScreen (<https://www.genoscreen.fr/>) for Sanger sequencing. Sequencing was performed from both ends.

Results — Genomic DNA was detectable in all samples. As expected, it was more concentrated (but this concentration was also more variable) in samples obtained from muscles (mean +/- standard error of the mean: 181,1+/-15,1 ng/µl) as compared to buccal and haemolymph swabs (14.5+/-1.4 ng/µl and 37.1+/-3 ng/µl, respectively). These results also show that there is on average more than double DNA when swabbing haemolymph than when swabbing the buccal region. All DNA samples obtained from muscle and haemolymph yielded a visible band at the expected size (202 bp), resulting in a 100% +/- 0% amplification success of this particular locus for these samples. By contrast, amplification with the DNA samples obtained from buccal swabs was less successful (61%, binomial confidence interval: [47%;73%]).

As explained above, we sequenced twelve samples from the successful amplifications. These samples were chosen to represent 12 different individuals and the three types of sampling (muscle, haemolymph and buccal swabs), with four samples each.

The twelve samples yielded high quality sequences. Oficialdegui et al. (2019) found only one haplotype among the 20 individuals they sequenced for a 701 bp fragment while we obtained three different haplotypes from 12 individuals over a 202 bp fragment (data not shown). The first striking

result is thus that our pond in Rennes (Outgroup crayfish in the main text) harbours much more genetic diversity than the entire marsh of Brière. The three haplotypes we observed were not in equal frequencies. A first haplotype, obtained from one individual only, cannot be distinguished from Hap_04 and Hap_11 from Oficialdegui et al. (2019). Hap_11 is the only haplotype found in Brière. The two other haplotypes, found in 4 and 7 individuals, belong to the group defined by Hap_09 in Oficialdegui et al. (2019). This haplotype is found in high frequency in Louisiana but also in low frequency in Spain and France.


Conclusion — (1) Genomic DNA can reliably be obtained from haemolymph captured on sterile cotton swabs when PIT-tagging crayfish. (2) Crayfish from the pond situated in Rennes display more genetic variability than Brière, all sequences but one tracing an origin that is different from what has been described for the Brière region.

References

- Broquet, T., Ménard, N., & Petit, E. (2007). Noninvasive population genetics: A review of sample source, diet, fragment length and microsatellite motif effects on amplification success and genotyping error rates. *Conservation Genetics*, 8(1), 249–260. <https://doi.org/10.1007/s10592-006-9146-5>
- Oficialdegui, F. J., Clavero, M., Sánchez, M. I., Green, A. J., Boyero, L., Michot, T. C., Klose, K., Kawai, T., & Lejeusne, C. (2019). Unravelling the global invasion routes of a worldwide invader, the red swamp crayfish (*Procambarus clarkii*). *Freshwater Biology*, 64(8), 1382–1400. <https://doi.org/10.1111/fwb.13312>


APPENDIX 2. Trapping devices used in the mesocosms.

A: artificial refuge trap (ART) with one crayfish inside. B: exit trap.




APPENDIX 3. Link between air and water temperature.

Cross-correlation between hourly air temperature at a 5 km-distant weather station and hourly water temperature in six of the 28 the experimental mesocosms. The horizontal blue, dotted lines show the values beyond which the autocorrelation is significantly different from zero at a 5 % risk. Water temperature was highly correlated with air temperature and maximal at a 5-hour lag.



APPENDIX 4. Crayfish body-size distributions in the three groups.

Range core crayfish are from the Brière marsh, Range edge crayfish are from ponds in the Brière area, and Outgroup crayfish are from a pond in Rennes city.



APPENDIX 5. Parameter estimates from the multistate mark-recapture model using uniform priors on the [-10,10] interval for regression parameters in Eq. 3. For more details, see Table 2 legend in the main text.


Transition probability	Link	Covariate	Parameter	mean	sd	Rhat	MCMC P-val
Φ_A	logit	Intercept	$\alpha[0]$	4.57	0.26	1.00	0.000
		Male	$\alpha[\text{Sex Male}]$	-0.06	0.16	1.00	0.701
		Outgroup	$\beta[\text{Outgroup}]$	0.19	0.42	1.00	0.658
		Range edge	$\beta[\text{Range edge}]$	-0.07	0.34	1.00	0.841
		High density	$\gamma[\text{HD}]$	-0.69	0.32	1.00	0.032
		Outgroup-by-HD interaction	$\delta[\text{Outgroup, HD}]$	0.66	0.47	1.00	0.165
		Range edge-by-HD interaction	$\delta[\text{Range edge, HD}]$	0.64	0.47	1.00	0.164
		Body size in Range core	$\alpha[1]$	-0.12	0.39	1.01	0.760
		Outgroup on Body-size effect	$\epsilon[\text{Outgroup}]$	-0.50	0.74	1.02	0.503
		Range edge on Body-size effect	$\epsilon[\text{Range edge}]$	0.56	0.64	1.00	0.370
		HD on Body-size effect in range core	$\zeta[\text{HD}]$	1.17	0.56	1.01	0.034
		Outgroup-by-HD interaction on Body-size effect	$\eta[\text{Outgroup, HD}]$	-1.56	0.87	1.02	0.072
		Range edge-by-HD interaction on Body-size effect	$\eta[\text{Range edge, HD}]$	-1.14	0.86	1.00	0.176
		Body condition	$\alpha[2]$	-0.50	0.17	1.00	0.003
		Temperature	$\alpha[3]$	0.55	0.18	1.00	0.002
		Time interval	$\alpha[4]$	-0.43	0.19	1.01	0.048
		Mesocosm	$\sigma[\theta]$	0.26	0.13	1.007	
		Ψ_{A-B}	logit	Intercept	$\alpha[0]$	-3.43	0.21
Male	$\alpha[\text{Sex Male}]$			-0.09	0.15	1.00	0.520
Outgroup	$\beta[\text{Outgroup}]$			-0.68	0.36	1.00	0.054
Range edge	$\beta[\text{Range edge}]$			-0.13	0.28	1.00	0.639
High density	$\gamma[\text{HD}]$			0.05	0.29	1.00	0.881
Outgroup-by-HD interaction	$\delta[\text{Outgroup, HD}]$			0.13	0.43	1.00	0.769
Range edge-by-HD interaction	$\delta[\text{Range edge, HD}]$			-0.11	0.40	1.00	0.802

		Body size in Range core	$\alpha[1]$	1.65	0.38	1.05	0.000
		Outgroup on Body-size effect	$\epsilon[\text{Outgroup}]$	-0.08	0.69	1.09	0.932
		Range edge on Body-size effect	$\epsilon[\text{Range edge}]$	0.02	0.67	1.01	0.982
		HD on Body-size effect in range core	$\zeta[\text{HD}]$	-0.20	0.62	1.04	0.741
		Outgroup-by-HD interaction on Body-size effect	$\eta[\text{Outgroup, HD}]$	-0.22	0.88	1.08	0.788
		Range edge-by-HD interaction on Body-size effect	$\eta[\text{Range edge, HD}]$	-0.42	0.95	1.02	0.660
		Body condition	$\alpha[2]$	0.59	0.16	1.00	0.000
		Temperature	$\alpha[3]$	0.92	0.08	1.00	0.000
		Time interval	$\alpha[4]$	-0.23	0.11	1.00	0.038
		Mesocosm	$\sigma[\theta]$	0.12	0.09	1.01	
		Individual	$\sigma[\iota]$	0.84	0.07	1.00	
		Intercept	$\alpha[0]$	1.795	0.666	1.015	0.000
		Male	$\alpha[\text{Sex Male}]$	-0.092	0.425	1.007	0.847
		Outgroup	$\beta[\text{Outgroup}]$	3.030	1.367	1.007	0.038
		Range edge	$\beta[\text{Range edge}]$	0.850	0.937	1.015	0.369
		High density	$\gamma[\text{HD}]$	0.734	0.876	1.024	0.352
		Outgroup-by-HD interaction	$\delta[\text{Outgroup, HD}]$	-1.599	1.464	1.014	0.730
		Range edge-by-HD interaction	$\delta[\text{Range edge, HD}]$	0.302	1.342	1.022	0.611
		Body size in Range core	$\alpha[1]$	-1.777	0.840	1.028	0.012
		Outgroup on Body-size effect	$\epsilon[\text{Outgroup}]$	-1.910	1.913	1.006	0.595
$\psi_{B \rightarrow A}$	logit	Range edge on Body-size effect	$\epsilon[\text{Range edge}]$	-1.205	1.770	1.019	0.781
		HD on Body-size effect in range core	$\zeta[\text{HD}]$	0.013	1.426	1.039	0.851
		Outgroup-by-HD interaction on Body-size effect	$\eta[\text{Outgroup, HD}]$	1.204	2.306	1.014	0.869
		Range edge-by-HD interaction on Body-size effect	$\eta[\text{Range edge, HD}]$	3.172	2.651	1.022	0.440
		Body condition	$\alpha[2]$	-1.415	0.495	1.005	0.006
		Temperature	$\alpha[3]$	-1.222	0.233	1.005	0.000
		Time interval	$\alpha[4]$	-0.131	0.177	1.000	0.492
		Mesocosm	$\sigma[\theta]$	0.802	0.411	1.017	
		Individual	$\sigma[\iota]$	1.441	0.277	1.043	

APPENDIX 6. Densities of prior (yellow bars) and posterior (blue bars) distributions for all main parameters in Eq. (3). The green colour shows overlap areas. HD = high-density treatment.


A. Survival model parameters.

Survival probability Φ_A




B. Dispersal model parameters.

Dispersal probability $\Psi_{A \rightarrow B}$



C. Post-dispersal settlement probability.

Post-dispersal settlement probability $\Psi_{B \rightarrow A}$




APPENDIX 7. Parameter estimates for a model including only main effects on crayfish probabilities of survival Φ_A , dispersal $\psi_{A \rightarrow B}$ and post-dispersal settlement $\psi_{B \rightarrow A}$. For more details, see Table 2 legend.

Transition	Link	Covariate	Parameter	mean	sd	Rhat	MCMC p-value
Φ_A	logit	Intercept	$\alpha[0]$	4.291	0.199	1.001	0.000
		Male	$\beta[\text{Sex Male}]$	0.020	0.151	1.001	0.886
		Body size	$\alpha[1]$	-0.217	0.150	1.000	0.147
		Body condition	$\alpha[2]$	-0.452	0.172	1.000	0.008
		High density	$\gamma[\text{HD}]$	-0.322	0.198	1.000	0.103
		Temperature	$\alpha[3]$	0.564	0.182	1.004	0.001
		Outgroup	$\delta[\text{Outgroup}]$	0.484	0.191	1.001	0.012
		Range edge	$\delta[\text{Range edge}]$	0.191	0.217	1.001	0.373
		Time interval	$\alpha[4]$	-0.407	0.176	1.008	0.056
		Mesocosm	$\sigma[\iota]$	0.242	0.128	1.002	
$\psi_{A \rightarrow B}$	logit	Intercept	$\alpha[0]$	-3.404	0.185	1.000	0.000
		Male	$\beta[\text{Sex Male}]$	-0.114	0.139	1.002	0.402
		Body size	$\alpha[1]$	1.324	0.146	1.000	0.000
		Body condition	$\alpha[2]$	0.554	0.152	1.002	0.000
		High density	$\gamma[\text{HD}]$	0.033	0.168	1.002	0.834
		Temperature	$\alpha[3]$	0.918	0.082	1.000	0.000
		Outgroup	$\delta[\text{Outgroup}]$	-0.564	0.192	1.002	0.002
		Range edge	$\delta[\text{Range edge}]$	-0.135	0.201	1.001	0.493

		Time interval	$\alpha[4]$	-0.231	0.112	1.002	0.037
		Mesocosm	$\sigma[\iota]$	0.128	0.089	1.023	
		Individual	$\sigma[\kappa]$	0.824	0.068	1.002	
<hr/>							
		Intercept	$\alpha[0]$	1.809	0.528	1.011	0.000
		Male	$\beta[\text{Sex Male}]$	-0.119	0.368	1.003	0.734
		Body size	$\alpha[1]$	-1.917	0.378	1.002	0.000
		Body condition	$\alpha[2]$	-1.066	0.413	1.004	0.010
		High density	$\gamma[\text{HD}]$	0.576	0.473	1.005	0.216
$\psi_{B \rightarrow A}$	logit	Temperature	$\alpha[3]$	-1.151	0.223	1.001	0.000
		Outgroup	$\delta[\text{Outgroup}]$	1.167	0.473	1.001	0.015
		Range edge	$\delta[\text{Range edge}]$	0.654	0.541	1.000	0.219
		Time interval	$\alpha[4]$	-0.125	0.181	1.000	0.492
		Mesocosm	$\sigma[\iota]$	0.701	0.349	1.008	
		Individual	$\sigma[\kappa]$	1.308	0.240	1.019	
<hr/>							

APPENDIX 8. Effects of stocking density and experiment duration on crayfish survival rate in aquaculture studies.

Density varied from 1 to 1200 individuals m^{-2} , and duration from 3 to 306 days. $N = 436$ observations from 29 studies. Species included: *Astacus leptodactylus*: $n = 107$, *Austropotamobius pallipes*: $n = 32$, *Cherax destructor*: $n = 55$, *Cherax quadricarinatus*: $n = 110$, *Pacifastacus leniusculus*: $n = 45$, *Pontastacus leptodactylus*: $n = 3$, *Procambarus clarkii*: $n = 84$. Relationship modelled from a beta GLMM (logit link) with $\log(\text{density})$ and duration and their interaction as fixed effects, and with the study as random intercept in the `glmmTMB` library of R (Brooks et al., 2017). The effect of crayfish species increased model AIC and was thus discarded. Reference list cited below.



- Barki, A., & Karplus, I. (2000). Crowding female red claw crayfish, *Cherax quadricarinatus*, under small-tanks hatchery conditions: What is the limit? *Aquaculture*, *181*(3), 235–240.
[https://doi.org/10.1016/S0044-8486\(99\)00235-5](https://doi.org/10.1016/S0044-8486(99)00235-5)
- Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). GlmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, *9*(2), 378–400. <https://doi.org/10.32614/RJ-2017-066>
- Farhadi, A., & Jensen, M. A. (2016). Effects of photoperiod and stocking density on survival, growth and physiological responses of narrow clawed crayfish (*Astacus leptodactylus*). *Aquaculture Research*, *47*(8), 2518–2527. <https://doi.org/10.1111/are.12700>
- Figiel, C. R., & Miller, G. L. (1995). The frequency of chela autotomy and its influence on the growth and survival of the crayfish *Procambarus clarkii* (Girard, 1852) (Decapoda, Cambaridae). *Crustaceana*, *68*(4), 472–483.
- García-Ulloa, G. M., Pérez-Moreno, M. R., Rodríguez-González, D., Gallo-García, M. C., Ponce-Palafox, J. T., Rodríguez-González, H., & Góngora-Gómez, A. M. (2012). Stocking density for nursery production of redclaw crayfish, *Cherax quadricarinatus*, in a recirculating system. *Journal of Applied Aquaculture*, *24*(1), 8–15.
<https://doi.org/10.1080/10454438.2012.650599>
- Geddes, M. C., & Smallridge, M. (1993). Survival, growth and yield of the Australian freshwater crayfish *Cherax destructor* in extensive aquaculture ponds. *Aquaculture*, *114*(1), 51–70.
[https://doi.org/10.1016/0044-8486\(93\)90250-3](https://doi.org/10.1016/0044-8486(93)90250-3)
- González, R., Celada, J. D., González, A., García, V., Carral, J. M., & Sáez-Royuela, M. (2010). Stocking density for the intensive rearing of juvenile crayfish, *Pacifastacus leniusculus* (Astacidae), using *Artemia* nauplii to supplement a dry diet from the onset of exogenous feeding. *Aquaculture International*, *18*(3), 371–378. <https://doi.org/10.1007/s10499-009-9250-x>
- Goyert, J. C., & Avault Jr., J. W. (1978). Effects of stocking density and substrate on growth and survival of crayfish (*procambarus clarkii*) grown in a recirculating culture system. *Journal*

of the *World Aquaculture Society*, 9(1–4), 731–735. <https://doi.org/10.1111/j.1749-7345.1978.tb00287.x>

- Harlioğlu, M. (2009). A comparison of the growth and survival of two freshwater crayfish species, *Astacus leptodactylus* Eschscholtz and *Pacifastacus leniusculus* (Dana), under different temperature and density regimes. *Aquaculture International*, 17(1), 31–43. <https://doi.org/10.1007/s10499-008-9177-7>
- Jones, C. M. (1995). Production of juvenile redclaw crayfish, *Cherax quadricarinatus* (von Martens) (Decapoda, Parastacidae) III. Managed pond production trials. *Aquaculture*, 138(1), 247–255. [https://doi.org/10.1016/0044-8486\(95\)00067-4](https://doi.org/10.1016/0044-8486(95)00067-4)
- Jones, C. M., & Ruscoe, I. M. (2000). Assessment of stocking size and density in the production of redclaw crayfish, *Cherax quadricarinatus* (von Martens) (Decapoda: Parastacidae), cultured under earthen pond conditions. *Aquaculture*, 189(1), 63–71. [https://doi.org/10.1016/S0044-8486\(00\)00359-8](https://doi.org/10.1016/S0044-8486(00)00359-8)
- Karplus, I., Harpaz, S., Hulata, G., Segev, R., & Barki, A. (2001). Culture of the Australian red-claw crayfish (*Cherax quadricarinatus*) in Israel—IV. Crayfish incorporation into intensive tilapia production units. *Israeli Journal of Aquaculture - Bamidgeh*, 53(1), 23–33.
- Lutz, Ch. G., & Wolters, W. R. (1986). The effect of five stocking densities on growth and yield of red swamp crayfish *Procambarus clarkii*. *Journal of the World Aquaculture Society*, 17(1–4), 33–36. <https://doi.org/10.1111/j.1749-7345.1986.tb00551.x>
- Mazlum, Y. (2007). Stocking density affects the growth, survival, and cheliped injuries of third instars of narrow-clawed crayfish, *Astacus leptodactylus* Eschscholtz, 1823 Juveniles. *Crustaceana*, 80(7), 803–815. JSTOR.
- Mazlum, Y., Yazici, M., & Gurlek, O. (2020). Evaluation of stocking densities and feed types on growth and survival of narrow-clawed crayfish *Pontastacus leptodactylus* (escholtz, 1823) reared under laboratory conditions. *Fresenius Environmental Bulletin*, 29(9A), 8283–8293.
- Mills, B. J., & McCloud, P. I. (1983). Effects of stocking and feeding rates on experimental pond production of the crayfish *Cherax destructor* clark (Decapoda: Parastacidae). *Aquaculture*, 34(1), 51–72. [https://doi.org/10.1016/0044-8486\(83\)90291-0](https://doi.org/10.1016/0044-8486(83)90291-0)

- Naranjo-Páramo, J., Hernandez-Llamas, A., & Villarreal, H. (2004). Effect of stocking density on growth, survival and yield of juvenile redclaw crayfish *Cherax quadricarinatus* (Decapoda: Parastacidae) in gravel-lined commercial nursery ponds. *Aquaculture*, 242(1), 197–206. <https://doi.org/10.1016/j.aquaculture.2004.05.017>
- Nightingale, J., Stebbing, P., Taylor, N., McCabe, G., & Jones, G. (2018). Determining an effective density regime for rearing juvenile *Austropotamobius pallipes* in a small-scale closed system hatchery. *Aquaculture Research*, 49(9), 3055–3062. <https://doi.org/10.1111/are.13766>
- Pinto, G. F., & Rouse, D. B. (1996). Growth and survival of the Australian red claw crayfish *Cherax quadricarinatus* at three densities in earthen ponds. *Journal of the World Aquaculture Society*, 27(2), 187–193. <https://doi.org/10.1111/j.1749-7345.1996.tb00268.x>
- Ramalho, R. O., Correia, A. M., & Anastácio, P. M. (2008). Effects of density on growth and survival of juvenile red swamp crayfish, *Procambarus clarkii* (Girard), reared under laboratory conditions. *Aquaculture Research*, 39(6), 577–586. <https://doi.org/10.1111/j.1365-2109.2008.01907.x>
- Rodgers, L. J., Saoud, P. I., & Rouse, D. B. (2006). The effects of monosex culture and stocking density on survival, growth and yield of redclaw crayfish (*Cherax quadricarinatus*) in earthen ponds. *Aquaculture*, 259(1), 164–168. <https://doi.org/10.1016/j.aquaculture.2005.11.056>
- Romaire, R. P., & Villágran, E. R. (2010). Evaluation of stocking density and feeding regime on production of red swamp crawfish in outdoor mesocosms. *Journal of the World Aquaculture Society*, 41(3), 298–307. <https://doi.org/10.1111/j.1749-7345.2010.00372.x>
- Sáez-Royuela, M., Carral, J. M., Celada, J. D., & Pérez, J. R. (2001). Effects of shelter type and food supply frequency on survival and growth of stage-2 juvenile white-clawed crayfish (*Austropotamobius pallipes* Lereboullet) under laboratory conditions. *Aquaculture International*, 9(6), 489–497. <https://doi.org/10.1023/A:1020509627870>
- Savolainen, R., Ruohonen, K., & Railo, E. (2004). Effect of stocking density on growth, survival and cheliped injuries of stage 2 juvenile signal crayfish *Pasifastacus leniusculus* Dana. *Aquaculture*, 231(1), 237–248. <https://doi.org/10.1016/j.aquaculture.2003.09.045>

- Shun, C., Yong-yi, J., Mei-li, C., Jian-bo, Z., Shi-li, L., & Zhi-min, G. (2020). Culture model of *Cherax quadricarinatus*: Temporary shelter in shed and pond culture. *Aquaculture*, 526, 735359. <https://doi.org/10.1016/j.aquaculture.2020.735359>
- Ulikowski, D., Krzywosz, T., & Śmietana, P. (2006). A comparison of survival and growth in juvenile *Astacus leptodactylus* (Esch.) and *Pacifastacus leniusculus* (Dana) under controlled conditions. *Knowledge and Management of Aquatic Ecosystems*, 380–381, 1245–1253. <https://doi.org/10.1051/kmae:2006023>
- Verhoef, G. D., & Austin, C. M. (1999). Combined effects of temperature and density on the growth and survival of juveniles of the Australian freshwater crayfish, *Cherax destructor* Clark,: Part 1. *Aquaculture*, 170(1), 37–47. [https://doi.org/10.1016/S0044-8486\(98\)00394-9](https://doi.org/10.1016/S0044-8486(98)00394-9)
- Webster, C. D., Thompson, K. R., Muzinic, L. A., Yancey, D. H., Dasgupta, S., Xiong, Y. L., Rouse, D. B., & Manomaitis, L. (2004). A preliminary assessment of growth, survival, yield, and economic return of Australian red claw crayfish, *Cherax quadricarinatus*, stocked at three densities in earthen ponds in a cool, temperate climate. *Journal of Applied Aquaculture*, 15(3–4), 37–50. https://doi.org/10.1300/J028v15n03_03
- Yu, J., Xiong, M., Ye, S., Li, W., Xiong, F., Liu, J., & Zhang, T. (2020). Effects of stocking density and artificial macrophyte shelter on survival, growth and molting of juvenile red swamp crayfish (*Procambarus clarkii*) under experimental conditions. *Aquaculture*, 521, 735001. <https://doi.org/10.1016/j.aquaculture.2020.735001>

APPENDIX 9. Predicted crayfish survival and overland dispersal probabilities by 2071-2100 under the SSP1-2.6 and SSP5-8.5 scenarios of IPCC.

Predictions were formed from parameter estimates for the multistate mark-recapture model (Table 2), for an average-sized Range-core female at low crayfish density. Annual mean reference temperature was 10.7°C (“Historical” situation), which is average summer (May-September) temperature in Rennes city France for the 1981-2010 period (<https://fr.wikipedia.org/wiki/Rennes>).

	Historical	SSP1-2.6 scenario		SSP5-8.5 scenario	
		Low	High	Low	High
Monthly survival probability	0.84	0.86	0.87	0.87	0.89
Time to dispersal probability = 1 (days)	111	95	81	83	62