

Spatio-Temporal Domains of Wildfire-Prone Teleconnection Patterns in the Western Mediterranean Basin

Marcos Rodrigues, Michela Mariani, Ana Russo, Michele Salis, Luiz Felipe Galizia, Adrian Cardil

▶ **To cite this version:**

Marcos Rodrigues, Michela Mariani, Ana Russo, Michele Salis, Luiz Felipe Galizia, et al.. Spatio-Temporal Domains of Wildfire-Prone Teleconnection Patterns in the Western Mediterranean Basin. Geophysical Research Letters, 2021, 48 (19), pp.1-14. 10.1029/2021gl094238 . hal-03847569

HAL Id: hal-03847569
https://hal.inrae.fr/hal-03847569

Submitted on 10 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Check for updates

Geophysical Research Letters

RESEARCH LETTER
10.1029/2021GL094238

Special Section:
Fire in the Earth System

Spatio-Temporal Domains of Wildfire-Prone Teleconnection Patterns in the Western Mediterranean Basin

Marcos Rodrigues[1,2,3], Michela Mariani[4], Ana Russo[5], Michele Salis[6], Luiz Felipe Galizia[7], and Adrian Cardil[2,8,9]

[1]Department of Agricultural and Forest Engineering, University of Lleida, Lleida, Spain, [2]Joint Research Unit CTFC – AGROTECNIO, Solsona, Spain, [3]Department of Geography and Land Management, GEOFOREST Group, University of Zaragoza, Zaragoza, Spain, [4]School of Geography, University of Nottingham, Nottingham, UK, [5]Instituto Dom Luiz (IDL), Faculdade de Ciências da Universidade de Lisboa, Lisboa, Portugal, [6]National Research Council, Institute of BioEconomy (CNR-IBE), Sassari, Italy, [7]INRAE, RECOVER, Aix-Marseille University, Aix-en-Provence, France, [8]Department of Crop and Forest Sciences, University of Lleida, Lleida, Spain, [9]Technosylva Inc., La Jolla, CA, USA

Abstract This work explores the main climate teleconnections influencing the Western Mediterranean Basin to outline homogeneous fire-prone weather domains combining cross-correlation time series and cluster analysis. We found a zonal effect of the Scandinavian pattern over the entire region with an interesting alternation of phases from positive during winter-spring (increased rainfall leading to fuel accumulation) to negative (dry conditions) modes during summer controlling burned area and fire size. The North Atlantic Oscillation (NAO) dominates the number of fires over the Iberian Peninsula (IP) while the Western Mediterranean Oscillation pattern modulates fire activity over the Mediterranean coast in the IP (linked to westerly winds), Southern France, Corsica and Sardinia (rainfall regulation). These distinctive influence traits resulted in three different domains splitting the IP into a Mediterranean rim along the coast (from southern Spain to southwestern France) and an inland and western region (Portugal plus western Spain); and a third in southeastern France, Corsica and Sardinia.

Plain Language Summary We synthetized the climate influence on wildfire activity over the Western Mediterranean Basin into three distinctive configurations. We applied computer-assisted statistical analysis on historical fire records from national/regional agencies and climate-related data [the so-called climate teleconnection indexes; e.g., "El Niño" or the North Atlantic Oscillation (NAO)]. Burned area size over the entire basin tends to be larger under the confluence of rainy winter-spring and sudden heat waves and dry spells during summer, which is governed by the Scandinavian pattern. In the Mediterranean coast of Spain fires occur more frequently under sustained calm and warm conditions while they grow larger when winds blow from the western side of the Iberian Peninsula. These conditions depend on the NAO and Western Mediterranean Oscillation (WeMOi) pattern, respectively. Fire activity in Southern France, Corsica and Sardinia seems to be linked to the lack of rainfall, which is regulated by the atmospheric conditions over the Mediterranean sea, mostly controlled by the WeMOi pattern.

Key Points:

- We found three distinctive but homogenous domains of influence of climate teleconnections patterns
- The Scandinavian pattern exerts a zonal influence over the western Mediterranean basin. North Atlantic Oscillation controls the Iberian Peninsula and Western Mediterranean Oscillation the Mediterranean coast
- Positive moisture winter-spring anomalies coupled to short-term dry and warm conditions boost most of the wildfire activity

Correspondence to:
M. Rodrigues,
marcos.rodrigues@udl.cat

Citation:
Rodrigues, M., Mariani, M., Russo, A., Salis, M., Galizia, L. F., & Cardil, A. (2021). Spatio-temporal domains of wildfire-prone teleconnection patterns in the Western Mediterranean Basin. *Geophysical Research Letters*, *48*, e2021GL094238. https://doi.org/10.1029/2021GL094238

Received 13 MAY 2021
Accepted 20 SEP 2021

© 2021. The Authors.
This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

1. Introduction

Wildfires are a key ecosystem process that modulates vegetation distribution and evolution (Bond et al., 2005) and impacts the global carbon cycle (Jones et al., 2019), whilst having substantial economic and social impacts (Moritz et al., 2014). During the last two decades, most of the extreme wildfires reported as being economically or socially catastrophic were concentrated in suburban areas intermixed with flammable forests, particularly in the western United States (Radeloff et al., 2018), southeastern Australia (Bowman et al., 2017) and the Mediterranean Basin (Modugno et al., 2016). The latter was specifically identified as a disaster-prone landscape with projections suggesting an increase of extreme fire weather of 50%–100% by the end of the current century (Bowman et al., 2017). Currently, most of the total burned area in Europe occurs in this region during the summer, with an average of about 4,500 km^2/yr, an area that is expected to increase (Ganteaume et al., 2013; Turco et al., 2017).

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Mediterranean Europe occupies a climatic transitional area under the alternate influence of sub-tropical and temperate climates (Lionello, 2012). In this area, ecosystems and human populations are affected by frequent weather-driven natural hazards, such as droughts (Hoerling et al., 2012; Russo et al., 2017), heat waves (Sánchez-Benítez et al., 2018; Sousa et al., 2019) and wildfires (Ruffault et al., 2020). The frequency and severity of these extreme weather-driven events are likely to increase under climate change (Dupuy et al., 2020; Lionello and Scarascia, 2020). Additionally, the compound occurrence of drought and heatwave events has recently increased (Vogel et al., 2021), a trend that will persist according to future forecasts (AghaKouchak et al., 2020; Zscheischler et al., 2018). Additionally, the summer fire activity in Mediterranean-type ecosystems is thought to be sensitive to antecedent winter rainfall pulses that could lead to an accumulation of fuels, while concurrent droughts and/or hot-dry winds are the short-term driver of wildfires (Moritz et al., 2012). Predicting future fire activity does not only hinge on understanding the short-term fire weather patterns, but also the longer-term of climate variability (Rodrigues et al., 2020; Vieira et al., 2020), particularly on biomass accumulation mediated by regional-scale drivers (Moritz et al., 2012).

Previous studies suggest that coincident drought conditions and high temperatures promote large wildfires across southern Europe (Camiá et al., 2009; Bedia et al., 2013; Gouveia et al., 2016; Urbieta et al., 2015). Nevertheless, the role of antecedent large-scale climate conditions remains a debated topic across much of the Mediterannean Europe. Previous analyses of the linkages between fire activity and meteorological variables in southern France (Ruffault et al., 2016), Greece (Gouveia et al., 2016; Koutsias et al., 2013), and the Iberian Peninsula (Turco et al., 2013; Vieira et al., 2020) reveal significant correlations with both same-summer and lagged climate variables.

Climate teleconnections (CTs) have a synchronous influence on weather at a regional scale (sub-continental) while playing a key role in modulating temperature and precipitation patterns at both interannual and decadal scales (e.g., Ascoli et al., 2017; Harris and Lucas, 2019). These patterns can subsequently dictate plant growth (fuel amount) and its dryness, and hence modulate the occurrence and spread of wildfires (Cai et al., 2014; Mariani et al., 2018). CTs patterns are expected to become more extreme in the future (Cai et al., 2014; Power et al., 2013), potentially exacerbating their effects on wildfires (Mariani et al., 2018). Thus, the links between CTs and fire activity have been investigated in various regions (Cardil et al., 2021; Kitzberger et al., 2007; Mariani et al., 2018; Rodrigues et al., 2021). However, the role of large-scale CTs on wildfire occurrence and spread has not yet been fully explored in the western Mediterranean region (Royé et al., 2020). Within this area, various CTs influence weather, including the North Atlantic Oscillation (NAO), the East Atlantic (EA), the Atlantic Multidecadal Oscillation, the El Niño Southern Oscillation (ENSO), the Mediterranean Oscillation (MOI), the Pacific Decadal Oscillation (PDO), the Scandinavian (SCAND) pattern and the Western Mediterranean oscillation (WeMOi). In a recent study carried out in the Iberian Peninsula, fire danger patterns were found to be significantly correlated to the WeMOi, MOI, NAO, ENSO, and SCAND indexes (Rodrigues et al., 2021).

Investigating mechanisms behind climate-fire dynamics, including lagged relationships between climatic conditions and wildfire activity (i.e., fuel built-up fostered by above average antecedent moist conditions), is critical for understanding the future of Mediterranean ecosystems. Despite the clear importance of wildfires in shaping vegetation and impacting human lives under the projected climate change (Bowman et al., 2017; Moritz et al., 2012), the climatic drivers of fire activity through time remain poorly understood in many regions on Earth, including the European Mediterranean Basin (Turco et al., 2017). In this study, we provide a new assessment of the climatic drivers of wildfire occurrence and spread within the western Mediterranean region over the period 1980–2015. We expand the spatial and temporal scales of work carried out so far in the Iberian Peninsula (IP) on the links between wildfires and CTs, while focusing on actual fire features (namely, number of ignitions, burned area and fire size) rather than fire-weather indexes (Rodrigues et al., 2021). The main goal of the present work is assessing spatio-temporal climate domains reflecting the interaction of CTs and their potential influence in wildfire activity across western Mediterranean Europe. We aim to address the following research questions: (a) Which are the leading CTs over wildfire occurrence and spread across western Mediterranean Europe? (b) Can we identify transboundary domains of CT influence?, and (c) Is biomass limitation generating a lagged relationship between CTs and wildfires? Answering these questions will provide a better understanding of the regional climate-fire linkages, which would allow a better preparation for potential future wildfires or prompt timely fuel load reduction programmes.

2. Materials and Methods

We calculated the Pearson's R correlation coefficient between monthly time series of fire features (number of fires, burned area, and fire size) and CTs in each grid cell. Thus we retrieved a set of Pearson correlation coefficients for each combination of fire feature, CT and lag (0, 3, 6, and 9 months) for each grid cell. The spatial pattern of correlations was summarized in maps depicting the direction (either positive or negative) and the significance ($p < 0.05$, $R = \pm 0.34$) of the association.

To outline potential climatic domains resulting from the interaction of CT at multiple time scales we submitted the correlation coefficients to a cluster analysis. Identifying such domains will enable us to better understand the climatic forcing of wildfire activity. Our approach is heavily focused on identifying transnational climatic regions under the premise that homogenous patterns of both wildfire activity and climate factors do exist. We adopted a hierarchical cluster approach using *Euclidean Distance* as dissimilarity measure and *ward.D2* as agglomeration criterion, which aims to create groups such that variance is minimized within them. Clusters were trained for each fire feature separately and the optimal number of clusters (i.e., climate domains) was determined from the set of criteria available in the *nbClust* package (Charrad et al., 2014). Climate domains were then characterized according to the observed distribution of CT correlation, indicating the most influential patterns in terms of significance and direction of the relationship.

3. Data

3.1. Study Region and Fire Data

The region under study extends over the Western Mediterranean Basin, including Portugal, Spain, Southern France, Corsica and Sardinia (Italy), which are among the most fire-affected region of Europe. Fire data were obtained from the national/regional wildfire databases of Portugal (ICNF, http://www2.icnf.pt/portal/florestas/dfci/inc/estat-sgif), Spain (EGIF; https://www.miteco.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/incendios-forestales.aspx), France (Prométhée; https://www.promethee.com/incendies) and Sardinia (CFVA; http://www.sardegnageoportale.it/webgis2/sardegnamappe/?map=aree_tutelate). We retrieved all fire records in the period 1980–2015, retaining only those larger than 100 ha to prevent inhomogeneities related to fire detection and compilation. For each fire record we extracted the date and place of ignition and the final size. Fire events were aggregated into a regular grid of 0.5° resolution to prevent undesired effects due to partial/inaccurate location information. For each cell, we calculated the burned area, the total number of fires and the 95th percentile of fire size on a monthly basis. Fire features were aggregated as the sum of the number of fires and burned areas, and the 95th percentile of fire size, in the period May–October. Finally, we applied an unweighted moving window procedure using a bandwidth of 500 km. This procedure helps smoothing the spatial distribution of fire activity to facilitate the identification of spatial patterns of association with large-scale CTs (Andela et al., 2017; Koutsias et al., 2016). Because of the unweighted moving window, a given pixel does not represent its actual location but rather the 500-km region surrounding it, increasing the average number of fire events per pixel to sufficient amount (from 0.07 to 18.16) for the investigation of extremes.

3.2. CT Patterns

We investigated seven CT patterns, known to be among the most influential factors modulating wildfires in the Western Mediterranean Basin.

1. The NAO computed as the Gibraltar- Reykjavik normalized Sea Level Pressure (Jones et al., 1997); https://crudata.uea.ac.uk/cru/data/nao/nao.dat.
2. The EA teleconnection pattern index, which is structurally similar to the NAO, and consists of a north-south dipole of anomaly centers spanning the North Atlantic from east to west (Barnston and Livezey, 1987); http://www.cpc.ncep.noaa.gov/data/teledoc/ea.shtml.
3. The SCAND, which consists of a primary circulation center over Scandinavia, with weaker centers of opposite sign over western Europe and eastern Russia/western Mongolia, and known there as Eurasia-1 (Barnston and Livezey, 1987); https://www.cpc.ncep.noaa.gov/data/teledoc/scand.shtml.

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

4. The ENSO, which was calculated from the HadISST1. It is the area averaged sea surface temperature (SST) from 5 S–5 N and 170–120 W (Rayner et al., 2003); https://psl.noaa.gov/gcos_wgsp/Timeseries/Nino34/.
5. The PDO, which was derived as the leading PC of monthly SST anomalies in the North Pacific Ocean, poleward of 20N (Mantua et al., 1997); https://psl.noaa.gov/data/climateindices/list/.
6. The MOI, defined by Conte et al. (1989) as the normalized pressure difference between Algiers and Cairo; https://crudata.uea.ac.uk/cru/data/moi/moi1.output.dat.- The WeMOi, which is an index measuring the difference between the standardized atmospheric pressure recorded at Padua in northern Italy, and San Fernando, Cádiz in Southwestern Spain (Martin-Vide and Lopez-Bustins, 2006); http://www.ub.edu/gc/documents/Web_WeMOi-2020.txt.

CTs were averaged into the corresponding 6-month time frame. In addition to synchronous correlation (i.e., lag = 0 May–October), we investigated asynchronous effects in the CTs signal by exploring a lag of 3 (February–July), 6 (November–May) and 9 (September–February) months before the summertime.

4. Results

4.1. Influence of CTs on Fire Features

The spatial patterns of correlation between CTs and the selected fire features (burned area, number of fires, and fire size) depicted varied spatial arrangements. It should also be highlighted that the statistical significance was non-stationary over space and differed among CTs (Figures 1–3). The CT indices depicting the strongest positive or negative correlations were SCAND, WeMOI and NAO.

The SCAND index showed significant negative synchronous correlations (lag 0) with burned area almost in the entire study area. Surprisingly, we identified a sharp transition across the different time lags from positive association at lag 6 (antecedent winter), though statistically significant only in midland Spain. The NAO index was positively correlated above 40° latitude in the IP and Southern France at lag 0. NAO also denoted a significant lagged signal (6 and 9 months before summer) in Southern France, Corsica and Sardinia. The WeMOI index is correlated to increased burned area in Sardinia, Corsica, and the Mediterranean coast of Spain and France from lag 3 to 9 (Figure 1). The correlations with MOI and PDO were lower than the aforementioned indices and only significant in specific enclaves (i.e., MOI in the south of Sardinia at lag 9 and PDO in Spain at lag 9).

In terms of correlation results with the number of fires (Figure 2), NAO was clearly the dominant pattern, being positively correlated in the IP and Southern France at time lag 0 and Southern France and Sardinia at lag 9. The effect of the SCAND index differed, in spatial terms, from the burned area analysis. This index was only significant (negatively) in Corsica, the north of Sardinia, Southern France and NE Spain at lag 0. It was found to be positively correlated in Sardinia, Southern France and the Spanish Mediterranean coast at lag 3. The contribution of the persistent positive WeMOI phase (and to a lesser extent PDO) was significantly evident in the Mediterranean at all lags. The MOI was also significant in Sardinia, Corsica and some coastal Mediterranean areas of France at lag 6 and 9.

The significance of all indices explaining fire size was lower compared to burned area and fire ignitions (Figure 3). Similarly to the burned area, the SCAND index significantly explained the size of fires, transitioning as well from positive association in lag 6 to antiphase correlation at lag 0 in most of the study area. NAO shows significant correlations with fire size in Sardinia, Southern France and Corsica at lag 3 (antecedent spring) and the WeMOI index showed a positive correlation along the Spanish Mediterranean coast, though only significant at lag 3.

The other CT modes (EA and ENSO) did not show significant effects anywhere in the study region.

4.2. Spatio-Temporal Domains of CTs

We identified three clusters based on most influential CT indices (SCAND, WeMOI and NAO) that consistently outlined coincident domains among the three fire features considered in this work (Figure 4). These homogeneous zones reflect the climatic gradient from Atlantic (Zone 1) to Mediterranean conditions

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

10.1029/2021GL094238




Figure 1. Spatial distribution of Pearson's *R* correlation coefficient between climate teleconnection and burned area (fires > 100 ha), 1980–2015. Columns indicate lagged cross-correlation intervals of months before summer season.

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

10.1029/2021GL094238




Figure 2. Spatial distribution of Pearson's *R* correlation coefficient between climate teleconnection and number of fires (fires > 100 ha), 1980–2015. Columns indicate lagged cross-correlation intervals of months before summer season.

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

10.1029/2021GL094238




Figure 3. Spatial distribution of Pearson's *R* correlation coefficient between climate teleconnection and 95th percentile of fire size (fires > 100 ha), 1980–2015. Columns indicate lagged cross-correlation intervals of months before summer season.

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




Figure 4. Climate teleconnection domains. Maps show the spatial distribution of climate teleconnection domains for burned area (left), number of fires (middle) and fire size (right). Violin plots display the correlation strength (Pearsons' R) between North Atlantic Oscillation, Scandinavian and Western Mediterranean Oscillation per feature and lag. The solid black line indicates the 0 correlation threshold. Inner and outer dashed lines indicate the significance threshold $p < 0.05$.

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

(Zones 2 and 3). Despite the geographical concordance of the patterns, we observed clear differences in the inner contribution of CTs in terms of association and significance of CTs and temporal scales of influence, which significantly varied across fire features.

Zone 1 gathered the influence of SCAND and NAO patterns on all fire features This CT domain extends over the western IP, excluding the Mediterranean coast. Positive NAO at lag 0 and a sharp transition from positive to negative SCAND (also observed in Zones 2 and 3) are the distinctive traits of the region. Zone 2 seems to act as a transition area from Atlantic to Mediterranean conditions. It covers the Spanish Mediterranean coast, extending over the western portion of southern France. The SCAND pattern is still exerting a strong influence, showing negative and synchronous associations. Zone 2 better relates to positive association with the WeMOI pattern, which boosts both burned area and fire size but showed weaker influence in fire ignitions, still strongly linked to synchronic NAO.

The third domain (Zone 3), was clearly led by the WeMOI pattern, exerting a strong positive association with burned area and number of fire ignitions. The influence of WeMOI extends from lag 9 to 0 in the case of fire ignitions, shortening in the case of burned area (3–9). Again, the SCAND index depicted a synchronous negative influence in the entire study region and a sharp transition. NAO significantly explained fire features at different time lags: burned area (lag 0), number of fires (lag 3), fire size (lag 9).

5. Discussion and Conclusions

Several studies already prompt the importance of anticipating the pernicious effects of future shifts in fire regimes, with a wide consensus about the growing impacts of climate warming cascading into increased extreme wildfire events (Bedia et al., 2013, 2014, 2015; Ruffault et al., 2020; Turco et al., 2018). In this study, we provide insights into the dominant CT patterns modulating wildfire activity in the Western Mediterranean basin.

In line with Royé et al. (2020), we identified three CT domains spanning west-to-east and being mainly associated with the SCAND, the NAO and the WeMOi climate modes. From a spatial standpoint, the mediterranean coast CT domain (zone 2) largely matches fire regime zones delineated in the region (Rodrigues et al., 2020; Rodrigues, Costafreda-Aumedes, et al., 2019; Rodrigues, Jiménez-Ruano, & de la Riva, 2019; Rodrigues, González-Hidalgo, et al., 2019; Trigo et al., 2016; Vieira et al., 2020). According to Calheiros et al. (2021), this region is likely to persist over time and the frequency of days experiencing extreme fire-weather conditions will increase; an analysis that should consider the aforementioned CT modes influencing this domain in future analysis. Our domain delimitation assigns the rest of the IP into a single cluster (Zone 1), despite the known variety of driving forces and different fire regimes (Nunes et al., 2016; Rodrigues et al., 2018). However, from a climate perspective, large fires in the region (>100 ha) seem to respond to similar spatial (Rodrigues et al., 2020) and temporal (Silva et al., 2019) patterns. The domains observed in France span from zone 3 in the East, to Zone 2 in the west. They match to a certain degree the "pyroclimate" delimitation by Curt et al. (2014). However, the coarse spatial resolution of our fire data (50×50 km compared to the 2×2 km resolution of the French database) precluded us from detecting the fine-grained mosaic of local regions.

These CTs differently interacted across space and time (time lags between the CT and fire season), thus leading to different driving weather patterns described below. The negative relationship with the SCAND observed 0–3 months before summer involves reduced precipitation, suggesting drought spells influencing dead fuel moisture content as the main climate driver of wildfires (Ruffault et al., 2018). One of the most striking results was the observed phase transition of the SCAND (from positive to negative, i.e., wet to dry conditions) related to increased fire activity. This sharp shift suggests fuel build-up during +SCAND (increased rainfall) coupled to dry spells (−SCAND) fostering the accumulation of low dead fuel moisture content in the months leading up to summer. This indicates a certain limiting effect of dead biomass/fuel availability (herbaceous fuel load) in the ultimate dimensions of the burning (Gouveia et al., 2016; Littell et al., 2018) while supporting the notion that wildfires in the western Mediterranean are usually events fostered by short-term fire-prone conditions modulated by the moisture status and abundance of herbaceous dead fuels. This is in line also with the results of Russo et al. (2017) for the IP, which highlight the relationship between wildfires and drought is better explained by the influence of spring precipitation on

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

the central sector, and by the influence of temperature and precipitation during summer on most of the Portuguese provinces. Thus we should not overlook antecedent positive water balance anomalies (+SCAND), which may play an equally important role as dry spells or heat waves do (Pausas & Fernández-Muñoz, 2012; Pausas & Paula, 2012).

The NAO is linked to a higher number of fires in the entire region. Positive NAO is linked to anticyclonic conditions boosting temperature and limiting precipitation potentially leading to extreme drought episodes (García-Herrera et al., 2007; Vicente-serrano &Cuadrat, 2007). Eventually, this situation may evolve into thermal lows and thunderstorms fostering lightning strikes and fires (Pineda & Rigo, 2017). Likewise, NAO may connect also with sub saharan intrusions due to Atlantic blocking (Sousa et al., 2018, 2019), boosting temperatures in the Mediterranean area. In fact, the largest fires in the Mediterranean side of the IP seem to be associated with southeastern advections conducive to extreme heat waves (Rodrigues, Costafreda-Aumedes, et al., 2019; Rodrigues, Jiménez-Ruano, & de la Riva, 2019; Rodrigues, González-Hidalgo, et al., 2019; Rodrigues et al., 2020). Likewise, fires are known to be concomitant with thermal anomalies in the northeastern end of the IP (Cardil et al., 2015; Duane and Brotons, 2018) or Sardinia and Corsica (Ager et al., 2014; Salis et al., 2021). Though weaker than NAO's, the WeMOI also exerts moderate influence in the Mediterranean rim of the IP. During the +WeMOI the prevailing winds on the IP are typically from the West and Northwest. These winds, by the time they reach the Mediterranean sea, have crossed the peninsular continental areas reaching the leeward side of the coastal mountain systems, thus becoming warm and dry westerly winds (Rodrigues, Costafreda-Aumedes, et al., 2019; Rodrigues, Jiménez-Ruano, & de la Riva, 2019; Rodrigues, González-Hidalgo, et al., 2019) or cool but equally dry northwesterly winds (Duane and Brotons, 2018; Ruffault et al., 2017).

Nonetheless, while fires in the IP seem to be modulated by NAO, fire activity in southern France, Corsica and Sardinia (Zone 3) is better associated with the WeMOI pattern. For instance, in the period 1998–2016, Sardinia presented the most relevant wildfire spread conditions in days with southern and southwestern winds. Slightly more than 50% of the area burned by very large wildfires (>200 ha) during these wind regimes (Salis et al., 2021). Moreover, the advection of hot and dry air masses from the south through the inner parts and the north end of the island resulted in an evident increasing gradient in wildfire size and risk from southern to inner and northern Sardinia. On the other hand, strong mistral winds from the west and north-west (dominant during the positive phase of the WeMOI) promoted an increase in wildfire size and area burned in eastern and southern zones of Sardinia (Salis et al., 2021).

By identifying the key CT patterns boosting wildfire activity and its spatial-temporal domains of influence we can leverage operational climate services to provide seasonal forecasts that can be used to complement early warning systems (Lledó et al., 2020). While early warning systems based on short-term forecasted weather conditions are useful to anticipate hazardous fire behavior and danger in operational environments, CTs modulate weather patterns at monthly scales with lagged effects as shown in this manuscript, facilitating the identification of adverse fire windows during the fire season. Furthermore, it is possible to produce long-term (up to 3 months) predictions of CT patterns (Świerczyńska-Chlaściak & Niedzielski, 2020; Wang et al., 2017) which can allow for some predictability and therefore contributing to the improvement of alert systems. Finally, the ultimate effects of CTs on key weather parameters influencing fire behavior such as fuel loading, dead and live fuel moisture content or wind fields should be further analyzed.

Data Availability Statement

Data used in the study are available at Zenodo via https://10.5281/zenodo.5138095 (https://zenodo.org/record/5138095) with Creative Commons Attribution 4.0 International license. Fire data were obtained from the following national/regional wildfire databases: Portugal (ICNF, http://www2.icnf.pt/portal/florestas/dfci/inc/estat-sgif). Spain (EGIF; https://www.miteco.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/incendios-forestales.aspx). France (Prométhée; https://www.promethee.com/incendies). Sardinia (CFVA; http://www.sardegnageoportale.it/webgis2/sardegnamappe/?map=aree_tutelate). Data for climate teleconnection patterns were provided by the Climatic Research Unit from the University of East Anglia, the Climatology Group from the Universitat de Barcelona, and the US's Climate Prediction Center. Individual datasets and calculations can be accessed as follows: The NAO computed as

the Gibraltar- Reykjavik normalized Sea Level Pressure (Jones et al., 1997); https://crudata.uea.ac.uk/cru/data/nao/nao.dat. The EA teleconnection pattern index, which is structurally similar to the NAO, and consists of a north-south dipole of anomaly centers spanning the North Atlantic from east to west (Barnston and Livezey, 1987). http://www.cpc.ncep.noaa.gov/data/teledoc/ea.shtml. The SCAND, which consists of a primary circulation center over Scandinavia, with weaker centers of opposite sign over western Europe and eastern Russia/ western Mongolia, and known there as Eurasia-1 (Barnston & Livezey, 1987). https://www.cpc.ncep.noaa.gov/data/teledoc/scand.shtml. The ENSO, which was calculated from the HadISST1. It is the area averaged sea surface temperature (SST) from 5S-5N and 170-120W (Rayner et al., 2003). https://psl.noaa.gov/gcos_wgsp/Timeseries/Nino34/. The PDO, which was derived as the leading PC of monthly SST anomalies in the North Pacific Ocean, poleward of 20N (Mantua et al., 1997). https://psl.noaa.gov/data/climateindices/list/. The MOI, defined by Conte, Giuffrida and Tedesco (1989) as the normalized pressure difference between Algiers and Cairo; https://crudata.uea.ac.uk/cru/data/moi/moi1.output.dat. The WeMOi, which is an index measuring the difference between the standardized atmospheric pressure recorded at Padua in northern Italy, and San Fernando, Cádiz in Southwestern Spain (Martin-Vide and Lopez-Bustins, 2006); http://www.ub.edu/gc/documents/Web_WeMOi-2020.txt. All analyses, maps and plots were conducted using the R's framework for stastistical computing, Version 4.1.0, available via https://cran.r-project.org/ (R Core Team, 2021).

Acknowledgments
This work was partially supported by national funds through FCT (Fundação para a Ciência e a Tecnologia, Portugal) through projects IMPECAF (PTDC/CTA–CLI/28902/2017) and UIDB/50019/2020 - Instituto Dom Luiz.

References

Ager, A. A., Preisler, H. K., Arca, B., Spano, D., & Salis, M. (2014). Wildfire risk estimation in the Mediterranean area. *Environmetrics*, *25*(6), 384–396. https://doi.org/10.1002/env.2269

AghaKouchak, A., Chiang, F., Huning, L. S., Love, C. A., Mallakpour, I., Mazdiyasni, O., et al. (2020). Climate extremes and compound hazards in a warming world. *Annual Review of Earth and Planetary Sciences*, *48*(1), 519–548. https://doi.org/10.1146/annurev-earth-071719-055228

Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., et al. (2017). A human-driven decline in global burned area. *Science*, *356*(6345), 1356–1362. https://doi.org/10.1126/science.aal4108

Ascoli, D., Vacchiano, G., Turco, M., Conedera, M., Drobyshev, I., Maringer, J., et al. (2017). Inter-annual and decadal changes in teleconnections drive continental-scale synchronization of tree reproduction. *Nature Communications*, *8*(1), 2205. https://doi.org/10.1038/s41467-017-02348-9

Barnston, A. G., & Livezey, R. E. (1987). Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Monthly Weather Review*, *115*(6), 1083–1126. https://doi.org/10.1175/1520-0493(1987)115<1083:csapol>2.0.co;2

Bedia, J., Herrera, S., Camia, A., Moreno, J. M., & Gutiérrez, J. M. (2014). Forest fire danger projections in the Mediterranean using ENSEMBLES regional climate change scenarios. *Climatic Change*, *122*(1–2), 185–199. https://doi.org/10.1007/s10584-013-1005-z

Bedia, J., Herrera, S., Gutiérrez, J. M., Benali, A., Brands, S., Mota, B., & Moreno, J. M. (2015). Global patterns in the sensitivity of burned area to fire-weather: Implications for climate change. *Agricultural and Forest Meteorology*, *214*(215), 369–379. https://doi.org/10.1016/J.AGRFORMET.2015.09.002

Bedia, J., Herrera, S., Martín, D. S., Koutsias, N., & Gutiérrez, J. M. (2013). Robust projections of fire weather index in the Mediterranean using statistical downscaling. *Climatic Change*, *120*, 229–247. https://doi.org/10.1007/s10584-013-0787-3

Bond, W. J., Woodward, F. I., & Midgley, G. F. (2005). The global distribution of ecosystems in a world without fire. *New Phytologist*, *165*(2), 525–538. https://doi.org/10.1111/j.1469-8137.2004.01252.x

Bowman, D. M. J. S., Kolden, C. A., Abatzoglou, J. T., Johnston, F. H., van der Werf, G. R., & Flannigan, M. (2020). Vegetation fires in the Anthropocene. *Nature Reviews Earth & Environment*, *1*(10), 500–515. https://doi.org/10.1038/s43017-020-0085-3

Bowman, D. M. J. S., Williamson, G. J., Abatzoglou, J. T., Kolden, C. A., Cochrane, M. A., & Smith, A. M. S. (2017). Human exposure and sensitivity to globally extreme wildfire events. *Nature Ecology & Evolution*, *1*(3), 0058. https://doi.org/10.1038/s41559-016-0058

Brönnimann, S. (2007). Impact of El Niño–Southern Oscillation in European climate. *Reviews of Geophysics*, *45*, 1–28. https://doi.org/10.1029/2006RG000199.1.INTRODUCTION

Brunner, L., Hegerl, G. C., & Steiner, A. K. (2017). Connecting atmospheric blocking to European temperature extremes in spring. *Journal of Climate*, *30*(2), 585–594. https://doi.org/10.1175/JCLI-D-16-0518.1

Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., et al. (2014). Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature Climate Change*, *4*(2), 111–116. https://doi.org/10.1038/nclimate2100

Calheiros, T., Pereira, M. G., & Nunes, J. P. (2021). Assessing impacts of future climate change on extreme fire weather and pyro-regions in Iberian Peninsula. *The Science of the Total Environment*, *754*, 142233. https://doi.org/10.1016/j.scitotenv.2020.142233

Camiá, A., & Amatulli, G. (2009). Weather factors and fire danger in the Mediterranean. In E. Chuvieco (Ed.), *Earth Observation of Wildland fires in mediterranean ecosystems* (pp. 71–82). VerlagSpringer-Verlag. https://doi.org/10.1007/978-3-642-01754-4_6

Cardil, A., Eastaugh, C. S., & Molina, D. M. (2015). Extreme temperature conditions and wildland fires in Spain. *Theoretical and Applied Climatology*, *122*(1–2), 219–228. https://doi.org/10.1007/s00704-014-1295-8

Cardil, A., Rodrigues, M., Ramirez, J., de-Miguel, S., Silva, C. A., Mariani, M., & Ascoli, D. (2021). Coupled effects of climate teleconnections on drought, Santa Ana winds and wildfires in southern California. *The Science of the Total Environment*, *765*, 142788. https://doi.org/10.1016/j.scitotenv.2020.142788

Charrad, M., Ghazzali, N., Boiteau, V., & Niknafs, A. (2014). NbClust: An R package for determining the relevant number of clusters in a data set. *Journal of Statistical Software*, *61*(6), 1–36. https://doi.org/10.18637/jss.v061.i06

Conte, M., Giuffrida, A., & Tedesco, S. (1989). *The mediterranean oscillation. Impact on precipitation and hydrology in Italy climate water*. Publications of the Academy of Finland.

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Curt, T., Fréjaville, T., & Bouillon, C. (2014). *Characterizing pyroregions in south-eastern France. Characterizing pyroregions in south-eastern France* (pp. 1093–1101). Imprensa da Universidade de Coimbra. https://doi.org/10.14195/978-989-26-0884-6_119

Duane, A., & Brotons, L. (2018). Synoptic weather conditions and changing fire regimes in a Mediterranean environment. *Agricultural and Forest Meteorology*, *253–254*, 190–202. https://doi.org/10.1016/j.agrformet.2018.02.014

Dupuy, J. L., Fargeon, H., Martin-StPaul, N., Pimont, F., Ruffault, J., Guijarro, M., et al. (2020). Climate change impact on future wildfire danger and activity in southern Europe: A review. *Annals of Forest Science*, *77*(2). https://doi.org/10.1007/s13595-020-00933-5

Fréjaville, T., & Curt, T. (2015). Spatiotemporal patterns of changes in fire regime and climate: Defining the pyroclimates of south-eastern France (Mediterranean Basin). *Climatic Change*, *129*(1–2), 239–251. https://doi.org/10.1007/s10584-015-1332-3

Gallego, M. C., García, J. A., & Vaquero, J. M. (2005). The NAO signal in daily rainfall series over the Iberian Peninsula. *Climate Research*, *29*, 103–109. https://doi.org/10.3354/cr029103

Ganteaume, A., Camia, A., Jappiot, M., San-Miguel-Ayanz, J., Long-Fournel, M., & Lampin, C. (2013). A review of the main driving factors of forest fire ignition over Europe. *Environmental Management*, *51*(3), 651–662. https://doi.org/10.1007/s00267-012-9961-z

García-Herrera, R., Hernández, E., Barriopedro, D., Paredes, D., Trigo, R. M., Trigo, I. F., & Mendes, M. A. (2007). The outstanding 2004/05 drought in the Iberian Peninsula: Associated atmospheric circulation. *Journal of Hydrometeorology*, *8*(3), 483–498. https://doi.org/10.1175/JHM578.1

Gouveia, C. M., Bistinas, I., Liberato, M. L. R., Bastos, A., Koutsias, N., & Trigo, R. (2016). The outstanding synergy between drought, heatwaves and fuel on the 2007 Southern Greece exceptional fire season. *Agricultural and Forest Meteorology*, *218–219*, 135–145. https://doi.org/10.1016/j.agrformet.2015.11.023

Harris, S., & Lucas, C. (2019). Understanding the variability of Australian fire weather between 1973 and 2017. *PLoS One*, *14*(9), 1–33. https://doi.org/10.1371/journal.pone.0222328

Hoerling, M., Eischeid, J., Perlwitz, J., Quan, X., Zhang, T., & Pegion, P. (2012). On the increased frequency of Mediterranean drought. *Journal of Climate*, *25*(6), 2146–2161. https://doi.org/10.1175/JCLI-D-11-00296.1

Jones, M. W., Santín, C., van der Werf, G. R., & Doerr, S. H. (2019). Global fire emissions buffered by the production of pyrogenic carbon. *Nature Geoscience*, *12*(9), 742–747. https://doi.org/10.1038/s41561-019-0403-x

Jones, P. D., Jonsson, T., & Wheeler, D. (1997). Extension to the North Atlantic oscillation using early instrumental pressure observations from Gibraltar and south-west Iceland. *International Journal of Climatology*, *17*(13), 1433–1450. https://doi.org/10.1002/(sici)1097-0088(19971115)17:13<1433::aid-joc203>3.0.co;2-p

Kitzberger, T., Brown, P. M., Heyerdahl, E. K., Swetnam, T. W., & Veblen, T. T. (2007). Contingent Pacific–Atlantic Ocean influence on multicentury wildfire synchrony over western North America. *Proceedings of the National Academy of Sciences*, *104*(2), 543–548. https://doi.org/10.1073/pnas.0606078104

Koutsias, N., Allgöwer, B., Kalabokidis, K., Mallinis, G., Balatsos, P., & Goldammer, J. (2016). Fire occurrence zoning from local to global scale in the European Mediterranean basin: Implications for multi-scale fire management and policy. *Forest*, *9*, 195–204. https://doi.org/10.3832/ifor1513-008

Koutsias, N., Xanthopoulos, G., Founda, D., Xystrakis, F., Nioti, F., Pleniou, M., et al. (2013). On the relationships between forest fires and weather conditions in Greece from long-term national observations (1894–2010). *International Journal of Wildland Fire*, *22*(4), 493. https://doi.org/10.1071/WF12003

Lionello, P. (2012). *The climate of the mediterranean Region : From the past to the future*. Elsevier Science.

Lionello, P., & Scarascia, L. (2020). The relation of climate extremes with global warming in the Mediterranean region and its north versus south contrast. *Regional Environmental Change*, *20*, 31. https://doi.org/10.1007/s10113-020-01610-z

Littell, J. S., McKenzie, D., Wan, H. Y., & Cushman, S. A. (2018). Climate change and future wildfire in the western United States: An ecological approach to nonstationarity. *Earth's Future*, *6*(8), 1097–1111. https://doi.org/10.1029/2018EF000878

Lledó, L., Cionni, I., Torralba, V., Bretonnière, P.-A., & Samsó, M. (2020). Seasonal prediction of Euro-Atlantic teleconnections from multiple systems. *Environmental Research Letters*, *15*(7), 074009. https://doi.org/10.1088/1748-9326/ab87d2

Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., & Francis, R. C. (1997). A pacific interdecadal climate oscillation with impacts on salmon production*. *Bulletin of the American Meteorological Society*, *78*(6), 1069–1079. https://doi.org/10.1175/1520-0477(1997)078<1069:apicow>2.0.co;2

Mariani, M., Veblen, T. T., Williamson, G. J., Williamson, G., Fletcher, M. S., & Bowman, D. M. J. S. (2018). Climate change amplifications of climate-fire teleconnections in the southern climate change amplifications of climate-fire teleconnections in the Southern Hemisphere. *Geophysical Research Letters*, *45*, 5071–5081. https://doi.org/10.1029/2018GL078294

Martin-Vide, J., & Lopez-Bustins, J.-A. (2006). The western Mediterranean oscillation and rainfall in the Iberian Peninsula. *International Journal of Climatology*, *26*(11), 1455–1475. https://doi.org/10.1002/joc.1388

Modugno, S., Balzter, H., Cole, B., & Borrelli, P. (2016). Mapping regional patterns of large forest fires in Wildland–Urban Interface areas in Europe. *Journal of Environmental Management*, *172*(Supplement C), 112–126. https://doi.org/10.1016/j.jenvman.2016.02.013

Moritz, M. A., Batllori, E., Bradstock, R. A., Gill, A. M., Handmer, J., Hessburg, P. F., et al. (2014). Learning to coexist with wildfire. *Nature*, *515*, 58–66. https://doi.org/10.1038/nature13946

Moritz, M. A., Parisien, M.-A., Batllori, E., Krawchuk, M. A., Van Dorn, J., Ganz, D. J., & Hayhoe, K. (2012). Climate change and disruptions to global fire activity. *Ecosphere*, *3*(6), art49. https://doi.org/10.1890/ES11-00345.1

Nunes, A. N., Lourenço, L., & Meira, A. C. C. (2016). Exploring spatial patterns and drivers of forest fires in Portugal (1980–2014). *The Science of the Total Environment*, *573*(Supplement C), 1190–1202. https://doi.org/10.1016/j.scitotenv.2016.03.121

OrtizBevia, M. J., RuizdeElvira, A., Alvarez-Garcia, F. J., & Tasambay-Salazar, M. (2016). The multidecadal component of the Mediterranean summer variability. *Climate Dynamics*, *47*(9–10), 3373–3386. https://doi.org/10.1007/s00382-016-3341-y

Pausas, J. G., & Paula, S. (2012). Fuel shapes the firez-climate relationship: Evidence from Mediterranean ecosystems. *Global Ecology and Biogeography*, *21*(11), 1074–1082. https://doi.org/10.1111/j.1466-8238.2012.00769.x

Pausas, J. G. G., & Fernández-Muñoz, S. (2012). Fire regime changes in the Western Mediterranean Basin: From fuel-limited to drought-driven fire regime. *Climatic Change*, *110*(1–2), 215–226. https://doi.org/10.1007/s10584-011-0060-6

Pereira, M. G., Malamud, B. D., Trigo, R. M., & Alves, P. I. (2011). The history and characteristics of the 1980–2005 Portuguese rural fire database. *Natural Hazards and Earth System Sciences*, *11*(12), 3343–3358. https://doi.org/10.5194/nhess-11-3343-2011

Pineda, N., & Rigo, T. (2017). The rainfall factor in lightning-ignited wildfires in Catalonia. *Agricultural and Forest Meteorology*, *239*, 249–263. https://doi.org/10.1016/j.agrformet.2017.03.016

Power, S., Delage, F., Chung, C., Kociuba, G., & Keay, K. (2013). Robust twenty-first-century projections of El Niño and related precipitation variability. *Nature*, *502*(7472), 541–545. https://doi.org/10.1038/nature12580

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

R Core Team (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Radeloff, V. C., Helmers, D. P., Kramer, H. A., Mockrin, M. H., Alexandre, P. M., Bar-Massada, A., et al. (2018). Rapid growth of the US wildland-urban interface raises wildfire risk. *Proceedings of the National Academy of Sciences*, *115*(13), 3314–3319. https://doi.org/10.1073/pnas.1718850115

Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., et al. (2003). Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research*, *108*(D14). https://doi.org/10.1029/2002JD002670

Rodrigues, M., Costafreda-Aumedes, S., Comas, C., & Vega-García, C. (2019). Spatial stratification of wildfire drivers towards enhanced definition of large-fire regime zoning and fire seasons. *The Science of the Total Environment*, *689*, 634–644. https://doi.org/10.1016/j.scitotenv.2019.06.467

Rodrigues, M., González-Hidalgo, J. C., Peña-Angulo, D., & Jiménez-Ruano, A. (2019). Identifying wildfire-prone atmospheric circulation weather types on mainland Spain. *Agricultural and Forest Meteorology*, *264*, 92–103. https://doi.org/10.1016/j.agrformet.2018.10.005

Rodrigues, M., Jiménez-Ruano, A., & de la Riva, J. (2019). Fire regime dynamics in mainland Spain. Part 1: Drivers of change. *The Science of the Total Environment*, *721*, 135841. https://doi.org/10.1016/j.scitotenv.2019.135841

Rodrigues, M., Jiménez-Ruano, A., Peña-Angulo, D., & de la Riva, J. (2018). A comprehensive spatial-temporal analysis of driving factors of human-caused wildfires in Spain using Geographically Weighted Logistic Regression. *Journal of Environmental Management*, *225*, 177–192. https://doi.org/10.1016/j.jenvman.2018.07.098

Rodrigues, M., Peña-Angulo, D., Russo, A., Zúñiga-Antón, M., & Cardil, A. (2021). Do climate teleconnections modulate wildfire-prone conditions over the Iberian Peninsula? *Environmental Research Letters*, *16*. Retrieved from http://iopscience.iop.org/article/10.1088/1748-9326/abe25d

Rodrigues, M., Trigo, R. M., Vega-García, C., & Cardil, A. (2020). Identifying large fire weather typologies in the Iberian Peninsula. *Agricultural and Forest Meteorology*, *280*, 107789. https://doi.org/10.1016/j.agrformet.2019.107789

Royé, D., Tedim, F., Martin-Vide, J., Salis, M., Vendrell, J., Lovreglio, R., et al. (2020). Wildfire burnt area patterns and trends in Western Mediterranean Europe via the application of a concentration index. *Land Degradation & Development*, *31*(3), 311–324. https://doi.org/10.1002/ldr.3450

Ruffault, J., Curt, T., Martin-StPaul, N. K., Moron, V., & Trigo, R. M. (2018). Extreme wildfire events are linked to global-change-type droughts in the northern Mediterranean. *Natural Hazards and Earth System Sciences*, *18*(3), 847–856. https://doi.org/10.5194/nhess-18-847-2018

Ruffault, J., Curt, T., Moron, V., Trigo, R. M., Mouillot, F., Koutsias, N., et al. (2020). Increased likelihood of heat-induced large wildfires in the Mediterranean Basin. *Scientific Reports*, *10*(1), 13790. https://doi.org/10.1038/s41598-020-70069-z

Ruffault, J., Moron, V., Trigo, R. M., & Curt, T. (2016). Objective identification of multiple large fire climatologies: An application to a Mediterranean ecosystem. *Environmental Research Letters*, *11*(7), 075006. https://doi.org/10.1088/1748-9326/11/7/075006

Ruffault, J., Moron, V., Trigo, R. M., & Curt, T. (2017). Daily synoptic conditions associated with large fire occurrence in Mediterranean France: Evidence for a wind-driven fire regime. *International Journal of Climatology*, *37*(1), 524–533. https://doi.org/10.1002/joc.4680

Russo, A., Gouveia, C. M., Páscoa, P., DaCamara, C. C., Sousa, P. M., & Trigo, R. M. (2017). Assessing the role of drought events on wildfires in the Iberian Peninsula. *Agricultural and Forest Meteorology*, *237–238*, 50–59. https://doi.org/10.1016/j.agrformet.2017.01.021

Salis, M., Arca, B., Del Giudice, L., Palaiologou, P., Alcasena-Urdiroz, F., Ager, A., et al. (2021). Application of simulation modeling for wildfire exposure and transmission assessment in Sardinia, Italy. *International Journal of Disaster Risk Reduction*, *58*, 102189. https://doi.org/10.1016/j.ijdrr.2021.102189

Sánchez-Benítez, A., García-Herrera, R., Barriopedro, D., Sousa, P. M., & Trigo, R. M. (2018). June 2017: The earliest European summer mega-heatwave of reanalysis period. *Geophysical Research Letters*, *45*(4), 1955–1962. https://doi.org/10.1002/2018GL077253

Silva, J. M. N., Moreno, M. V., Le Page, Y., Oom, D., Bistinas, I., & Pereira, J. M. C. (2019). Spatiotemporal trends of area burnt in the Iberian Peninsula, 1975–2013. *Regional Environmental Change*, *19*(2), 515–527. https://doi.org/10.1007/s10113-018-1415-6

Sousa, P. M., Barriopedro, D., Ramos, A. M., García-Herrera, R., Espírito-Santo, F., & Trigo, R. M. (2019). Saharan air intrusions as a relevant mechanism for Iberian heatwaves: The record breaking events of August 2018 and June 2019. *Weather and Climate Extremes*, *26*, 100224. https://doi.org/10.1016/j.wace.2019.100224

Sousa, P. M., Trigo, R. M., Aizpurua, P., Nieto, R., Gimeno, L., & Garcia-Herrera, R. (2011). Trends and extremes of drought indices throughout the 20th century in the Mediterranean. *Natural Hazards and Earth System Sciences*, *11*(1), 33–51. https://doi.org/10.5194/nhess-11-33-2011

Sousa, P. M., Trigo, R. M., Barriopedro, D., Soares, P. M. M., & Santos, J. A. (2018). European temperature responses to blocking and ridge regional patterns. *Climate Dynamics*, *50*(1), 457–477. https://doi.org/10.1007/s00382-017-3620-2

Sousa, P. M., Trigo, R. M., Pereira, M. G., Bedia, J., & Gutiérrez, J. M. (2015). Different approaches to model future burnt area in the Iberian Peninsula. *Agricultural and Forest Meteorology*, *202*, 11–25. https://doi.org/10.1016/j.agrformet.2014.11.018

Świerczyńska-Chlaściak, M., & Niedzielski, T. (2020). Forecasting the North Atlantic Oscillation index using altimetric sea level anomalies. *Acta Geodaetica et Geophysica*, *55*(4), 531–553. https://doi.org/10.1007/s40328-020-00313-5

Trigo, R. M., Sousa, P. M., Pereira, M. G., Rasilla, D., & Gouveia, C. M. (2016). Modelling wildfire activity in Iberia with different atmospheric circulation weather types. *International Journal of Climatology*, *36*(7), 2761–2778. https://doi.org/10.1002/joc.3749

Turco, M., Llasat, M. C., von Hardenberg, J., & Provenzale, A. (2013). Impact of climate variability on summer fires in a Mediterranean environment (northeastern Iberian Peninsula). *Climatic Change*, *116*(3–4), 665–678. https://doi.org/10.1007/s10584-012-0505-6

Turco, M., Rosa-Cánovas, J. J., Bedia, J., Jerez, S., Montávez, J. P., Llasat, M. C., & Provenzale, A. (2018). Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. *Nature Communications*, *9*(1), 3821. https://doi.org/10.1038/s41467-018-06358-z

Turco, M., von Hardenberg, J., AghaKouchak, A., Llasat, M. C., Provenzale, A., & Trigo, R. M. (2017). On the key role of droughts in the dynamics of summer fires in Mediterranean Europe. *Scientific Reports*, *7*(1), 81. https://doi.org/10.1038/s41598-017-00116-9

Urbieta, I. R., Zavala, G., Bedia, J., Gutiérrez, J. M., Miguel-Ayanz, J. S., Camia, A., et al. (2015). Fire activity as a function of fire–weather seasonal severity and antecedent climate across spatial scales in southern Europe and Pacific western USA. *Environmental Research Letters*, *10*(11), 114013. https://doi.org/10.1088/1748-9326/10/11/114013

Vicente-serrano, S. M., & Cuadrat, J. M. (2007). North Atlantic oscillation control of droughts in north-east Spain: Evaluation since 1600 a.d. *Climatic Change*, *85*, 357–379. https://doi.org/10.1007/s10584-007-9285-9

Vieira, I., Russo, A., & Trigo, R. M. (2020). Identifying local-scale weather forcing conditions favorable to generating Iberia's largest fires. *Forests*, *11*(5), 1–14. https://doi.org/10.3390/f11050547

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Vogel, J., Paton, E., Aich, V., & Bronstert, A. (2021). Increasing compound warm spells and droughts in the Mediterranean Basin. *Weather and Climate Extremes*, *32*, 100312. https://doi.org/10.1016/j.wace.2021.100312

Wang, L., Ting, M., & Kushner, P. J. (2017). A robust empirical seasonal prediction of winter NAO and surface climate. *Scientific Reports*, *7*(1), 279. https://doi.org/10.1038/s41598-017-00353-y

Zscheischler, J., Westra, S., van den Hurk, B. J. J. M., Seneviratne, S. I., Ward, P. J., Pitman, A., et al. (2018). Future climate risk from compound events. *Nature Climate Change*, *8*(6), 469–477. https://doi.org/10.1038/s41558-018-0156-3

19448007, 2021, 19, Downloaded from https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL094238 by Cochrane France, Wiley Online Library on [10/11/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License