



HAL
open science

More diversity within livestock farming systems: an improvement of performances at all scales

Bertrand Dumont, Joël Aubin, Stéphane Ingrand, Guillaume Martin, Vincent Niderkorn, Laurence Puillet, Davi Savietto, Lucille Steinmetz, Marielle Thomas

► To cite this version:

Bertrand Dumont, Joël Aubin, Stéphane Ingrand, Guillaume Martin, Vincent Niderkorn, et al.. More diversity within livestock farming systems: an improvement of performances at all scales. Standing Committee on Agricultural Research (SCAR) - Sustainable Animal Production (SAP) Workshop, Oct 2022, Bonn, Germany. hal-03850970

HAL Id: hal-03850970

<https://hal.inrae.fr/hal-03850970v1>

Submitted on 14 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

More diversity within livestock farming systems: an improvement of performances at all scales

SCAR
CWG SAP
workshop
18/19
October
2022

INRAE

B. Dumont, J. Aubin, S. Ingrand, G. Martin, V. Niderkorn,
L. Puillet, D. Savietto, L. Steinmetz, M. Thomas



The 'Phase' division at INRAE (animal physiology & farming systems)

~1500 people

800 INRAE



460 partners

220 docs + post-docs

Scientific domains:

reproduction, nutrition, development, ethology, neurosciences, zootechnie, bio-informatic, stat. & modelisation

4 thematic fields






Diversity: which criteria? which organization? which goals?

Different components of diversity in LFS

❖ Resources

❖ Animals

❖ Systems



_01

DIVERSITY OF RESOURCES

Complementarity between plants in grasslands



Not really a scoop...

“It has been experimentally proved, if a plot of ground be sown with one species of grass, and a similar plot be sown with several distinct genera of grasses, a greater number of plants and a greater weight of dry herbage can be raised ...”

- Yield = 30% > average of the same species alone
- In 60% of sites = yield > to the best monoculture

Higher yields

Darwin (1872), *On the Origin of Species by Means of Natural Selection*, ch IV.



Complementarity between plants in grasslands (mechanisms)

Gains due to legumes / symbiotic fixation N_{atm}

- N = improve growth of grass
- Presence of grass = improve N fixation by legumes

(Nyfeler et al. 2011)

*Reduction of the
use of fertilizers*

Functional diversity reduces propagation of weeds

(Suter et al. 2016)





*Reduction of the use of
herbicides*

Improvement of resilience

- Portfolio effect *(Figge, 2004; Volaire et al., 2014)*
- Functional redundancy *(Biggs et al., 2012)*

Zootechnical added-values: both ecological & economic

Better performances when animals graze mixed grasslands

-  *Roca-Fernández et al., 2016*  *Grace et al., 2019*
- Greater motivation to eat (*Ginane et al., 2002; Niderkorn et al., 2017*)
- Anthelmintics properties of tannin plants : *Hoste et al. 2006; Collas et al. 2018* 

Reduction of concentrate

Reduction of veterinary products

Reduction of N leaching ($61 \Rightarrow 34$ kg N/ha; *Romera et al., 2017*)

Reduction of pollutions

Reduction in CH₄ emission with some species (*in vitro; Macheboeuf et al., 2017*)

Increase of income / ha ($2946 \Rightarrow 3062$ \$/ha; *Romera et al., 2017*)



_02

DIVERSITY OF ANIMALS


Rabbit breeding

Univ. Valence, epizootic rabbit enteropathy

Exp.1 : 1 single line (*Martinez-Vallespin et al., 2011*)

Exp.2 : 3 simultaneous lines (*Savietto et al., 2012*)


Exp.3 : Introduction of a resistant line among these \neq lines (*Garcia-Quiros et al., 2014*)



Towards herd immunity

Aquaculture

Favour multi-species systems to take advantage of the complementarity of the trophic niche of each one



Yield increase = 40%
(*Rahman et al., 2006; Wahab et al., 2011*)

... to which are added facilitation phenomena via the resuspension of nutrients, which favors plankton (*Milstein et al., 2006*)

_03

DIVERSITY OF SYSTEMS

Diversity of species in mixed livestock systems

2 territoires

40 enquêtes

PhD thesis Louise Forteau



11 élevages



10 élevages



10 élevages



10 élevages



Sward management



Parasitology



Massif Central

Horses for leisure + cattle (priority to generate income)

Normandy

Horses for sport, more productive swards

Management of parasitism in mixed systems




(Forteau et al., 2019)

- The hypothesis of dilution is validated: *Marley et al. 2006; Mahieu 2013*
- Only 35% of mixed farmers surveyed were aware of this property



Income mixed dairy (D) + beef cattle farms



- **Maximum income** for scenario 50D
- **Lowest variability** / market fluctuations for scenario 25D

(Diakité et al., 2019)

_04

**MY VIEW: DIVERSITY IS AN
ADDED VALUE WHEN WE HAVE
THE MEANS TO MANAGE IT**

To be able to track and to assess the status of the system

le: to mobilize ecological processes, we need to know them

Proxies, biomarkers, high throughput phenotyping, sensors

→ the dramatic expansion of digital sciences in farming systems

→ monitoring devices


To be able to act and to adapt practices in real time

Decision-making tools, rules, thresholds, adaptive management

→ artificial intelligence and modelling

→ robotics

➤ TEF Project (Testing and Experimental Facilities)



Conclusion

- ❖ **Diversity** has to be considered **at ≠ organisation levels**
- ❖ **Diversity is vital for geneticists**: no diversity = no possible selection
- ❖ **The clone fashion is over**
- ❖ Not only for extensive systems, **but also for intensive indoor systems** (rabbits, dairy goats, poultry)
- ❖ 3 steps are necessary:
 - Identification of the relevant parameters
 - Assessment of the relevant level of diversity for each parameter, according to the others
 - Definition of the management means & rules to drive the system throughout time

THANK YOU!

stephane.ingrand@inrae.fr

