



**HAL**  
open science

# Effect of starch content and rumen-protected amino acid supplementation on milk performance and persistency in dairy cows

Jean-Charles Anger, Christelle Loncke, R. Bidaux, P. Piantoni, K. Dieho,  
Sophie Lemosquet

## ► To cite this version:

Jean-Charles Anger, Christelle Loncke, R. Bidaux, P. Piantoni, K. Dieho, et al.. Effect of starch content and rumen-protected amino acid supplementation on milk performance and persistency in dairy cows. 7. EAAP International Symposium on Energy and Protein Metabolism and Nutrition (Isep), Sep 2022, Granada, Spain. pp.268-269, 10.1016/j.anscip.2022.07.030 . hal-03877812

**HAL Id: hal-03877812**

**<https://hal.inrae.fr/hal-03877812>**

Submitted on 6 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Effect of starch content and rumen-protected amino acid supplementation on milk performance and persistency in dairy cows**

J.C. Anger<sup>a,c</sup>, C. Loncke<sup>b</sup>, R. Bidaux<sup>c</sup>, P. Piantoni<sup>d</sup>, K. Dieho<sup>e</sup>, S. Lemosquet<sup>a</sup>

<sup>a</sup>PEGASE, INRAE, Institut Agro, Saint Gilles, France

<sup>b</sup>Université Paris-Saclay, INRAE, AgroParisTech, UMR Modélisation Systémique Appliquée aux Ruminants, 75005 Paris, France

<sup>c</sup>Provimi France, Cargill Animal Nutrition and Health, Crevin, France

<sup>d</sup>Cargill Animal Nutrition, Elk River, USA

<sup>e</sup>Cargill Animal Nutrition, Velddriël, Netherlands

**Key words** Dairy cow; Starch; Amino acid; Persistency; Efficiency

### **Introduction**

Reducing metabolizable protein (MP) supply increased protein efficiency but reduced milk yield (MY) and milk protein yield (MPY) in dairy cows. However, increasing dietary starch content or balancing Lys, Met and His could mitigate performance loss (Cantalapiedra-Hijaret al., 2014, Giallongo et al., 2017). The aim of the experiment was to evaluate the effects of diets varying in starch content and amino acid (AA: Lys, Met, His) supply on MY, MPY, and glucose and AA concentrations in a 18-wk study in dairy cows.

### **Material and methods**

Forty Holstein dairy cows were used in a randomized complete block design experiment with a 2-wk covariate (22 – 35 DIM) and a 18-wk treatment (55 – 180 DIM) period. A factorial arrangement of treatments was used with two levels of starch (high and low starch: HS and LS) and two levels of AA supplies (through rumen-protected AA; RPAA; AA<sup>-</sup> or AA<sup>+</sup>). Cows were blocked by parity, calving date and MY. Diets were formulated to be iso-nitrogenous and iso-energetic. The LS diet was based on maize and grass silages, whereas HS was based on maize silage and alfalfa hay. The LS and HS diets contained 37% of concentrate (wheat, barley, sugar beet pulp, molasses vs. corn grain, respectively), and experimentally provided (in g/kg DM): 182 vs. 265 of starch, 384 vs. 380 of NDF, 138 vs. 142 of CP, 90.9 vs. 94.4 of MP (i.e. PDI for Protein Digestible in the Intestine; INRA, 2018) and 6.53 vs. 6.19 MJ/kg DM of NEL, respectively. The AA<sup>-</sup> vs. AA<sup>+</sup> diets were formulated to provide [in % of MP]: 6.5 vs. 7.2% of Lys, 2.0 vs. 2.4% of Met, and 2.2 vs. 2.4% of His, respectively. Persistency was estimated as the slope of the declining phase of lactation. Plasma were sampled once a month and milk twice a week. Data were analyzed by ANCOVA using the MIXED procedure of SAS with week as the repeated measurement, cows as the random effect and orthogonal contrasts. Milk protein efficiency was calculated as proposed by INRA (2018).

### **Results and Discussion**

The HS diets increased MY (+ 3.0 kg/day; Table 1) and MPY (+ 100 g/day) despite decreased NEL intake (151 vs. 158 MJ/day;  $P < 0.01$ ) compared with LS. Supplementation of RPAA increased MPY (+ 36 g/day) and tended to increase DMI (+ 0.7 kg/day), consistent with Giallongo et al. (2017). Overall, MP efficiency was not affected by treatment (71.0%) because HS and AA<sup>+</sup> increased MP supply compared with LS and AA<sup>-</sup> (+ 97 g/day and + 83 g/day, respectively; both  $P < 0.05$ ). Milk persistency was higher with HS vs. LS (+ 0.28 kg/wk) or RPAA supplementation (+ 0.11 kg/wk). Plasma glucose concentration did not differ among treatments (3.9 mM). Plasma Lys and Met concentrations were higher in AA<sup>+</sup> vs. AA<sup>-</sup>. Histidine metabolism was modified by starch (carnosine, 1-methylhistidine) and RPAA (1-methylhistidine) or by starch × AA (tendency of interaction), and plasma His and 3-methylhistidine were lower in treatment LSAA<sup>-</sup> vs. other treatments. A higher mobilization of muscular His, indicated by higher carnosine and 3-methylhistidine concentrations, could at least partially explain the lack of effect of RPAA supplementation on plasma His concentration in the HS diet compared with LS.

### **Conclusion and implications**

In conclusion, dietary starch content and RPAA supplementation increased both MPY but with different mechanisms: dietary starch content through MY and RPAA supplementation through DMI. We hypothesized that the changes in His metabolism could contribute to the greater persistency observed both when increasing starch content or RPAA supplies.

### Financial support

Cargill Animal Nutrition & Health France financially supports J.C. Anger Ph.D. scholarship and trials.

### References

G. Cantalapiedra-Hijar, J.L. Peyraud, S. Lemosquet, E. Molina-Alcaide, H. Boudra, P. Noziere, I. Ortigues-Marty. Dietary carbohydrate composition modifies the milk N efficiency in late lactation cows fed low crude protein diets *Animal*, 8 (2) (2014), pp. 275-285

F. Giallongo, M.T. Harper, H.J. Oh, C. Parys, I. Shinzato, A.N. Hristov. Histidine deficiency has a negative effect on lactational performance of dairy cows *Journal of Dairy Science*, 100 (4) (2017), pp. 2784-2800

INRA. feeding system for ruminants, Wageningen Academic Publishers, Wageningen, Netherlands (2018), p. 640

Table 1. Effect of two levels of dietary starch or Lys, Met, His supplementation on cow performances and plasma metabolites

	LSAA <sup>-1</sup>	LSAA <sup>+1</sup>	HSAA <sup>-1</sup>	HSAA <sup>+1</sup>	Pooled SE	P value <sup>4</sup>		
						Starch	AA	Starch × AA
Dry matter intake, kg /d	24.1	24.4	23.9	25.0	0.37	NS	†	NS
Milk								
Yield, kg/d	32.3	32.8	35.3	35.8	0.54	***	NS	NS
Persistency, kg/week	-0.50	-0.31	-0.14	-0.11	0.031	***	**	NS
Protein yield, g/d	986	1023	1087	1122	18.4	***	*	NS
Protein, g/kg	30.8	31.5	31.0	31.6	0.42	NS	NS	NS
Plasma concentration, μM								
Lys	61.4	71.2	65.5	74.0	3.68	NS	**	NS
Met	20.7	29.9	22.1	29.2	1.33	NS	***	NS
His	23.9	34.0	42.7	40.7	3.24	***	NS	†
Carnosine	14.7	14.2	19.6	19.2	1.02	***	NS	NS
1-methylhistidine <sup>2</sup>	5.56	5.23	6.37	5.90	0.222	**	†	NS
3-methylhistidine <sup>3</sup>	4.22	4.50	5.39	4.50	0.251	*	NS	*

<sup>1</sup>LS: Low starch; HS: High starch; AA- vs AA+: without and with RPAA (Lys, Met, His) supplementation, CAS number: <sup>2</sup>332-80-9; <sup>3</sup>368-16-1

<sup>4</sup>P value: † : < 0.1, \* : < 0.5, \*\* : < 0.01, \*\*\* : < 0.001