



**HAL**  
open science

# Complexity of the tripartite interaction in the rhizosphere: case of soil-borne pathogenic fungi with the holoparasitic plant *Phelipanche ramosa* and its host plant

Dinïa Cartry, Stéphanie Gibot-Leclerc, Véronique Edel-Hermann, Christian Steinberg

## ► To cite this version:

Dinïa Cartry, Stéphanie Gibot-Leclerc, Véronique Edel-Hermann, Christian Steinberg. Complexity of the tripartite interaction in the rhizosphere: case of soil-borne pathogenic fungi with the holoparasitic plant *Phelipanche ramosa* and its host plant. 13èmes Rencontres de Phytopathologie - Mycologie, Sep 2022, Aussois, France. . hal-03887279

**HAL Id: hal-03887279**

**<https://hal.inrae.fr/hal-03887279>**

Submitted on 30 May 2024


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Complexity of the tripartite interaction in the rhizosphere: case of soil-borne pathogenic fungi with the holoparasitic plant *Phelipanche ramosa* and its host plant.

D. Cartry <sup>(1)</sup>, S. Gibot-Leclerc <sup>(2)</sup>, V. Edel-Hermann, C. Steinberg <sup>(1)</sup>

<sup>(1)</sup> INRAE, UMR Agroécologie, 17 rue Sully, 21000 Dijon, France; <sup>(2)</sup> Institut Agro, 21000 Dijon, France




Branched broomrape (*Phelipanche ramosa* (L.) Pomel) is an achlorophyllous root parasitic plant with a wide host range. Its complex management is leading to the abandonment of tobacco or oilseed rape cultivation in the most affected regions in France. Among broomrape regulation factors [1], soil fungi could be a relevant biocontrol lever.




**Hypotheses:** i) symptomatic broomrapes are indicators of a pathogenic fungus ii) the later will negatively impact broomrape's growth at young phenological stages but not rapeseed and iii) the biological candidate colonizes both plants but with a pathogenic action in broomrape, not rapeseed

**Objectives:** i) to identify soil fungi as potential biological agents ii) to evaluate their pathogenicity against broomrape and iii) to understand the tripartite interaction between broomrape, rapeseed, and pathogenic fungi.

A fungal collection was made from **symptomatic** broomrape. Among the **374** collected isolates (A), nearly 80% belonged to **19 *Fusarium*** species (B).




87 isolates representative of this diversity showed **functional diversity by inhibiting seed germination** of the parasite - The 20 best-performing isolates showed differences in germination inhibition of *P. ramosa* at the intraspecific level.



These isolates also show differences in their **necrotic activity** against the stems of the parasitic plant in *in vitro* conditions.

***Fusarium venenatum*** isolates showed dual competence, i.e. germination inhibition and necrotic activity, and were non-pathogenic to tobacco.



Cytological observations confirm that the fungi colonize the broomrape tissues (mycelium colored in red on photo A) and uses the starch reserves of the plant to ensure its colonization (see arrows, photo B, C).

**Conclusions:** The results highlight different and complex biotic interactions possible between broomrape, host plant and pathogenic fungi, more specifically *Fusarium* sp. in the rhizosphere [3]. A **metabarcoding** approach will enlighten the colonization mode of the fungus to understand its pathogenic virulence against broomrape. Understanding the *Fusarium* pathogenicity in this **tripartite interaction is essential to implement an integrated control solution.**