



HAL
open science

Pan-European multi-crop model ensemble simulations of wheat and grain maize under climate change scenarios

Heidi Webber, Diane Kathleene Cooke, Frank Ewert, Jørgen Eivind Olesen, Stefan Fronzek, Alex C Ruane, Pierre Martre, Brian Collins, Marco Bindi, Roberto Ferrise, et al.

► To cite this version:

Heidi Webber, Diane Kathleene Cooke, Frank Ewert, Jørgen Eivind Olesen, Stefan Fronzek, et al.. Pan-European multi-crop model ensemble simulations of wheat and grain maize under climate change scenarios. Open Data Journal for Agricultural Research, 2020, 6, pp.21-27. <10.18174/odjar.v6i0.16326>. <hal-03895315>

HAL Id: hal-03895315

<https://hal.inrae.fr/hal-03895315v1>

Submitted on 13 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Pan-European multi-crop model ensemble simulations of wheat and grain maize under climate change scenarios

Heidi Webber^{1*}, Diane Cooke¹, Frank Ewert^{1,2}, Jørgen E. Olesen³, Stefan Fronzek⁴, Alex C. Ruane⁵, Pierre Martre⁶, Brian Collins⁷, Marco Bindi⁸, Roberto Ferrise⁸, Nándor Fodor¹⁰, Clara Gabaldón-Leal¹¹, Thomas Gaiser², Mohamed Jabloun¹², Kurt-Christian Kersebaum^{1,17}, Jon I. Lizaso¹³, Ignacio J. Lorite¹¹, Loic Manceau⁶, Marco Moriondo⁹, Claas Nendel^{1,16}, Alfredo Rodríguez^{13,14}, Margarita Ruiz-Ramos¹³, Mikhail A. Semenov¹⁵, Tommaso Stella¹, Pierre Stratonovitch¹⁵ and Giacomo Trombi⁸

¹ Leibniz Centre for Agricultural Landscape Research (ZALF), Eberswalder Straße 84, 15374 Müncheberg, Germany

² Institute of Crop Science and Resource Conservation (INRES), University of Bonn, Bonn, Germany

³ Department of Agroecology, Aarhus University, Tjele, Denmark

⁴ Finnish Environment Institute, Helsinki, Finland

⁵ NASA Goddard Institute for Space Studies, New York, NY 10025, USA

⁶ LEPSE, Univ. Montpellier, INRAE, Montpellier SupAgro. Montpellier, France

⁷ Centre for Crop Science, Queensland Alliance for Agriculture and Food Innovation (QAAFI), University of Queensland, Toowoomba, Australia

⁸ University of Florence, Florence, Italy

⁹ Institute of BioEconomy, National Research Council, Florence, Italy

¹⁰ Centre for Agricultural Research, Martonvásár, Hungary

¹¹ Institute of Agricultural and Fisheries Research and Training (IFAPA), Cordoba, Andalusia, Spain

¹² Plant Production Systems Group, Wageningen University & Research, Wageningen, The Netherlands

¹³ CEIGRAM, Universidad Politécnica de Madrid, Madrid, Spain

¹⁴ DAEF, Universidad de Castilla-La Mancha, Toledo, Spain

¹⁵ Rothamsted Research, Harpenden, United Kingdom

¹⁶ University of Potsdam, Germany

¹⁷ Global Change Research Institute, The Czech Academy of Sciences, Brno, Czech Republic.

* e-mail: webber@zalf.de

Abstract: The simulated data set described in this paper was created with an ensemble of nine different crop models: HERMES (HE), Simplace<Lintul5,Slim3, FAO-56 ET0> (L5), SiriusQuality (SQ), MONICA (MO), Sirius2014 (S2), FASSET (FA), 4M (4M), SSM (SS), DSSAT-CSM IXIM (IX). Simulations were performed for grain maize (six models) and winter wheat (eight models) under diverse conditions over agricultural land areas of the EU-27 at a 25 x 25 km spatial resolution. Simulations were drawn from combinations of three representative concentration pathways and climate outputs from five general circulation models for time periods 2040-2069 and 2070-2099. Historical climate data was the basis for simulation years 1980-2010 and considered as a baseline. Simulation results can be used to analyze crop responses to changing climatic variables. This data paper describes the creation, motivation and format of the simulation results to enable reuse of the data set. It also offers some possible further uses of the dataset in other contexts.

Keywords: crop modelling, climate change impacts, European Union, maize, wheat, heat stress, drought.

1 BACKGROUND: The data set consists of simulations from historical (1980-2010) and scenario periods (2040-2069 and 2070-2099) for six grain maize (MZ) and eight winter wheat (WW) process-based models. The models were applied on a spatial grid of 25 km resolution across the EU-27 (Austria, Belgium, Bulgaria, Cyprus, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Luxembourg, Malta, Netherlands, Portugal, Romania, Spain, Sweden, United Kingdom, Czech Republic, Estonia, Hungary, Latvia, Lithuania, Poland, Slovenia, Slovak Republic). The original purpose of this data set was to analyze the drivers of historical yield variability at both the national and subnational (NUTS2) levels, as well as the drivers of yield change under climate change for wheat and maize across Europe (Webber et al. 2018a). These two important food security crops are interesting to compare, as they have contrasting photosynthesis pathways (C3 and C4), a major determining factor of crop response to a change in ambient CO₂ levels. The two crops also differ in their main growing season (autumn versus spring sown for winter wheat and maize, respectively). A detailed description of the simulations, results

and discussion from this analysis are reported in Webber et al. (2018a). The data set described in this paper can be accessed from doi: [10.4228/ZALF.DK.88](https://doi.org/10.4228/ZALF.DK.88).

2 DATA CREATION PROCESS: A multi-model ensemble was used in this study to capture uncertainty associated with modeled processes and associated parameterizations. The nine crop models in the ensemble were: HERMES (HE), Simplace<Lintul5,Slim3, FAO-56 ET0> (L5), SiriusQuality (SQ), MONICA (MO), Sirius2014 (S2), FASSET (FA), 4M (4M), SSM (SS), DSSAT-CSM IXIM (IX). Models were selected based on their ability to simulate heat and drought stresses, as well as the interest of the respective modelling groups to participate in the study. Six of the models were also able to simulate crop canopy temperature (FA, L5, HE, SS, SQ and S2; see Table 1) allowing for the interaction of high temperature, drought stress and CO₂. More detailed model descriptions are provided in key references (Table 1) and in the SI materials of Webber et al. (2018a, b). A common protocol was defined and used by all modelers to standardize the modeling procedure, climate and soil data inputs as well as crop management practices. The complete protocol is provided in the supplemental methods of Webber et al. (2018a). All models were applied to the same spatial extent of EU-27 (Fig. 1), for 8,084 grids cells of 25 x 25 km resolution where soil data indicated at least a 40 cm rooting depth (see Figure 1). Data for sowing, anthesis and harvest dates were sourced from Eurostat¹, aggregated to 13 environmental zones and resampled to the simulation grid cells. Soil data were sourced from the JRC European Soil Data Portal.²

Table 1: Overview of models and key settings, including crop(s) simulated (winter wheat, grain maize, or both), processes affected by elevated CO₂ (canopy temperature, transpiration and/or radiation use efficiency, RUE) and the approach to simulate canopy temperature (CT). The three CT approaches include: empirical (EMP), energy balance assuming neutral stability (EBN) or energy balance correcting for atmospheric stability conditions (EBSC). 'NA' indicates that CT was not simulated.

Model name (abrv)	Crop	Processes affected by CO ₂	CT	Key references
HERMES (HE)	both	canopy temperature, transpiration and RUE	EBN	Kersebaum 2017
Simplace<Lintul5,Slim3, FAO-56 ET0> (L5)	both	canopy temperature, transpiration and RUE	EBSC	Webber et al. 2016
SiriusQuality (SQ)	winter wheat	RUE	EBN	Martre and Dambreville 2018
MONICA (MO)	both	transpiration and RUE	NA	Nendel et al. 2011
Sirius2014 (S2)	winter wheat	RUE	EBN	Jamieson et al. 1998
FASSET (FA)	both	canopy temperature, transpiration, RUE	EMP	Olesen et al. 2002
4M (4M)	both	transpiration and RUE	NA	Fodor et al. 2014
SSM (SS)	winter wheat	canopy temperature, transpiration, RUE	EBN	Soltani et al. 2013
DSSAT-CSM IXIM (IX)	grain maize	transpiration and RUE	NA	Lizaso et al. 2017

¹ <https://ec.europa.eu/jrc/en/mars>

² <https://esdac.jrc.ec.europa.eu/resource-type/european-soil-database-soil-properties>




Figure 1. Maps of average yield (kg/ha) for winter wheat for time period 2 (2040-2069), assuming elevated CO₂ levels (429, 499 and 571 ppm, for RCP2.6, 4.5 and 8.5, respectively) for each GCM (rows) and RCP (columns). These maps are based on simulations carried out by the FA model under rainfed conditions, including heat and drought stress, corresponding to Treatment 6 (T6) (see Figure 3).

2.1 Climate inputs: Observed meteorological data for the baseline period (period “0”= 1980-2010) were extracted from the Crop Growth Monitoring System (CGMS) of the Joint Research Centre (JRC) archive³. The JRC site-specific daily weather data is based on more than 3000 sites across Europe. The data is representative of agricultural land use and is interpolated to a regular grid at a spatial resolution of 25 km. Climate scenario data were constructed using an enhanced delta method (Ruane et al. 2015) for two periods (periods: “2”= 2040-2069 and “3”= 2070-2099). For each period, three representative concentration pathways (RCP; 2.6, 4.5, or 8.5) were coupled with five GCMs (GFDL-CM3, GISS-E2-R, HadGEM2-ES, MIROC5, and MPI-ESM-ER). Only two GCMs were available for RCP 2.6 (HadGEM2-ES and MPI-ESM-MR), whereas all five GCMs were available for RCP 4.5 and RCP 8.5 (Fig. 1 and Fig. 2). For each climate scenario, GCM and time period combination, simulations were conducted twice: the first set with atmospheric CO₂ concentration set at ambient levels corresponding to the historical baseline period (360 ppm) and a second set with elevated CO₂ (Fig. 2). Elevated CO₂ concentrations were determined based on the associated RCP and time period (429 and 442 ppm for RCP2.6 time periods 2 and 3, 499 and 532 ppm for RCP4.5 time periods 2 and 3, 571 and 801 ppm for RCP8.5 and time periods 2 and 3, respectively). For each scenario and time period, concentrations were based on values listed in the 2013 IPCC report (IPCC 2013). From these values, a central-year concentration, based on projections for the middle of the range of years, was assigned to each time period (see McDermid et al. 2015). As RCP 2.6 and 4.5 consider CO₂ mitigation measures, the increase in CO₂ concentration from time period 2 to 3 for these two scenarios is less than that for RCP 8.5. In total, there were 49 combinations used in the study (see Figure 2). The original climate data used in the study is available from doi [10.4228/ZALF.DK.59](https://doi.org/10.4228/ZALF.DK.59). An updated climate dataset that corrects for temperatures at high elevations is available at doi [10.4228/ZALF.DK.94](https://doi.org/10.4228/ZALF.DK.94). A detailed description of the creation of the climate data can be found in Fronzek et al., 2018b.




Figure 2. Simulations were conducted for 49 sets of climate and CO₂ concentration data. Climate data consist of 48 sets corresponding to climate scenarios defined by combinations of global climate models (GCM), representative concentration pathways (RCP), time period (2= 2040-2069, 3=2070-2099), and CO₂ level. One simulation set was conducted for a baseline climate (period = 0, 1980-2010; CO₂=360ppm, *GCM=NA and RCP=NA correspond to the fact that no GCM or RCP is applicable for observational historical data used in this study)

2.2 Soil inputs: The original soil data was sourced from the JRC European Soil Data Portal⁴ at 1 km resolution and included textural classes, depth available to roots, total available water content (TAWC), bulk density (BD), and soil organic carbon (SOC). To select for grids under current agricultural land use within the EU-27, soil layers were resampled to match the 250 x 250 m resolution Corine 2006 Land Cover Map Version 17⁵. Since data for Greece was not available from this map, the Corine 2000 Land Cover v16 map was used for Greece. Further, only soils with a depth of 40 cm or greater were included.

³ <https://ec.europa.eu/jrc/en/mars>

⁴ <https://esdac.jrc.ec.europa.eu/resource-type/european-soil-database-soil-properties>

⁵ http://www.eea.europa.eu/data-and-maps/data/ds_resolveuid/a47ee0d3248146908f72a8fde9939d9d

These data were used to derive estimates of soil water at saturation, field capacity and permanent wilting point, needed as inputs for the crop models.

2.3 Phenology and crop management inputs: Phenology observations of sowing, emergence, anthesis and harvest or maturity dates from the JRC AgriCast4 database⁶ were used by the modelling groups to calibrate phenology parameters with the historical weather data. Modelling groups calibrated their respective phenology parameters (eg, thermal times) to match observed anthesis and maturity dates. The resulting parameters for each model were kept constant for subsequent scenario simulations, with the explicit assumption that there was no adaption in crop variety.

2.4 Treatments: For each of the 49 climate combinations, two crops (grain maize, MZ, and winter wheat, WW) and up to six treatments (Fig. 3) were simulated by the models, dependent on the respective model's ability to simulate both crops and each treatment. The treatments were numbered T1 to T6 and defined by combinations of irrigation status (full or rain), heat stress (on or off) and heat by drought interaction (on or off) (Fig. 3). The interaction of heat and drought stresses were estimated using simulated canopy temperatures. Models used different methods to simulate canopy temperature (Table 1), with detailed descriptions in Table 2 of the supplementary materials of Webber et al. 2018a. As treatments T3 and T6 used canopy temperature, they were only simulated by models that consider canopy temperature (FA, L5, HE, SS, SQ, and S2)

Treatments			
ID	T	Irrigation status	Production case (code)
T1	T_{air}	Full	Potential (Pot) with no heat stress
T2	T_{air}	Full	Heat-limited with T_{air} (HL.air)
T3	T_{canopy}	Full	Heat-limited with T_c (HL.can)
T4	T_{air}	Rain	Water-limited with no heat stress (WL)
T5	T_{air}	Rain	Water-heat-limited with T_{air} (WHL.air)
T6	T_{canopy}	Rain	Water-heat-limited with T_c (WHL.can)

Figure 3. Six treatments were simulated, defined by irrigation status (full irrigation (Full) or rainfed (Rain)), stress type considered (heat or drought) and whether stress interactions were considered by using air (T_{air}) or canopy temperatures (T_{canopy} , T_c). Treatments using canopy temperature (T3 and T6) were only simulated by 6 of the models (FA, L5, HE, SS, SQ, and S2).

3 FILE FORMAT AND ANNOTATION: All files in the data set are comma separated value (csv) files compressed into a gzip format. Each file comprises the outputs for all crops (MZ, WW), treatments (TrtNo), scenarios by GCMs combination (sce), CO₂ concentrations (CO₂), periods (period) and year carried out by a single model (Model) and for a single simulation grid and is named as "EU_HS_2digitModelCode_row_col.csv.gz". As an example, the file with model outputs from model 4M with input data associated with grid cell in row 32 and column 125 would be named EU_HS_4M_32_125.csv.gz (Table 2). Additionally a redundant identifier is defined for the combination of scenario, GCM, and CO₂ concentration (ClimPerCO₂_ID), which are further defined in the Supplemental Materials of Webber et al. 2018a. Definitions and units for the variables found in the headers of each file are listed in doi [10.4228/ZALF.DK.88](https://doi.org/10.4228/ZALF.DK.88) and given (from top to bottom) in the same order as they appear in the files (from left to right). Header and variables are the same in every file of the data set. Missing values in the files are denoted as "NA" (Table 2). Data was annotated using metadata standards defined by DataCite 4.1 (DataCite 2017).

⁶ <https://ec.europa.eu/jrc/en/mars>

Table 2: Example of the information available in the data set. Model 4M used as an example. File names are given in format 'EU_HS_4M_row_col.csv.gz'. First rows with bolded text are the headers. Model= 2-digit model code, row_col= grid cell identifier, Crop = MZ or WW, ClimPerCO2ID= unique identifier for 48 period gcm_rcp CO₂ levels, period= period identifier (0, 2, 3), sce = GCM/RCP combination, CO₂= CO₂ level, TrtNo= 2-digit simulation treatment identifier, Irrigation= irrigation status (Full, Rain), ProdCase= stress status(Pot, HL_air, see also Figure 3), Year= year of harvest, Yield =Grain yield**, AntDOY= anthesis julian day (YYYYJJJ), MatDOY= maturity Julian day (YYYYJJJ), GNumber= Grain number per unit area, Biom-an= anthesis total above ground biomass, Biom-ma= maturity total above ground biomass, MaxLAI =maximum LAI, WDrain= cumulative water drained beneath max rooted zone*, CumET= cumulative crop evapotranspiration*, SoilAvW= final soil water content in maximum rooted zone**, Runoff= cumulative water runoff*, Transp= cumulative crop transpiration*, Evap= cumulative soil water evaporation*, CroN-an = Crop N content***, Crop-ma= Crop N content**, GrainN= Grain N content*, ET0= cum. Reference crop evapotranspiration*, SowDOY= Julian day of year at sowing (YYYYJJJ), EmergDOY= Julian day at emergence (YYYYJJJ), TcMaxAve= average daily maximum canopy temperature*, TmaxAve= average daily maximum air temperature*. *between sowing and maturity, **at maturity, ***at anthesis.

Model	row_col	Crop	ClimPerCO2ID	period	sce	CO2	TrtNo	Irrigation	ProdCase	Year
4M	32_125	MZ	C01	0	0.0	360	T1	Full	Pot	1981
4M	32_125	MZ	C01	0	0.0	360	T1	Full	Pot	1982
Yield	AntDOY	MatDOY	GNumber	Biom.an	Biom.ma	MAXLAI	WDrain	CumET	SoilAvW	Runoff
16556	189	270	NA	9628	31194	5.91	0	164	2	0
12438	192	268	NA	7113	23531	4.40	0	198	11	0
Transp	Evap	CroN.an	CroN.ma	GrainN	Et0	SowDOY	EmergDOY	TcMaxAve	TMaxAve	
145	20	NA	NA	NA	1069	114	119	NA	27.3	
140	58	NA	NA	NA	941	114	121	NA	25.5	

4 OPPORTUNITIES FOR REUSE: The data set may serve as the basis for further analysis of maize and wheat average yields and inter-annual variability, as well as crop response to climate and elevated CO₂ across Europe. Beyond quantifying and identifying impacts on yield, the dataset could also be used to assess changing water use and water demand under both rainfed and irrigated conditions. As such, it could inform risk assessments and irrigation design studies. Further, the dataset could potentially be linked with remote sensing data to understand patterns observed in soil-water, surface temperature or leaf area related indices, and how they may be affected by climate change. It also has the potential to act as a benchmark for more detailed and localized adaptation studies by providing European scale trends against which regional changes can be compared.

ACKNOWLEDGMENTS

H.W., T.G and F.E. acknowledge support from the FACCE JPI MACSUR project through the German Federal Ministry of Food and Agriculture (2815ERA01J). Support from the SUSTAg project funded through the German Federal Ministry of Food and Agriculture is acknowledged by H.W. and F.E. (031B0170B). J.E.O. and M.J. were funded by Innovation Fund Denmark (5105-00001B). S.F. received financial support from the Academy of Finland (decision 277276) and the Finnish Ministry of Agriculture and Forestry (MMM) through FACCE-MACSUR. A.R. was supported by the National Aeronautics and Space Agency Science Mission Directorate (WBS 281945.02.03.06.79). M.Bi., R.F., M.M., and G.T. acknowledge financial support from the JPI FACCE MACSUR2 project, funded by the Italian Ministry for Agricultural, Food, and Forestry Policies (D.M. 24064/7303/15 of 26/Nov/2015). N.F.'s contribution was supported by the Széchenyi 2020 program, the European Regional Development Fund-"Investing in your future" and the Hungarian Government (GINOP-2.3.2-15-2016-00028). M.R.R. and A.Ro. acknowledge support from MINECO (APCIN2016-0005-00-00). A.Ro. acknowledges support from Comunidad de Madrid (Spain) and Structural Funds 2014-2020 (ERDF and ESF), project AGRISOST-CM S2018/BAA-4330. M.A.S. and P.S. received grant-aided support from the BBSRC Designing Future Wheat programme [BB/P016855/1]. We thank R.P. Rötter for climate scenario construction. The contributions of Andreas Enders and Gunther Kraus in facilitating the data transfer and processing are acknowledged and appreciated.

REFERENCES

DataCite Metadata Working Group. 2017. DataCite Metadata Schema Documentation for the Publication and Citation of Research Data. Version 4.1. DataCite e.V. doi: [10.5438/0014](https://doi.org/10.5438/0014).

- Fronzek, S., Webber, H., Rötter, R., Ruane, A. and Ewerts, F., 2018a. A daily time-step observed and scenario climate dataset on a European grid for crop modelling applications (version 1). doi: [10.4228/ZALF.DK.59](https://doi.org/10.4228/ZALF.DK.59).
- Fronzek, S., Webber, H., Rötter, R., Ruane, A. and Ewerts, F., 2018b. A daily time-step observed and scenario climate dataset on a European grid for crop modelling applications (version 2). doi: [10.4228/ZALF.DK.94](https://doi.org/10.4228/ZALF.DK.94).
- IPCC, 2013. Annex II: Climate System Scenario Tables [Prather, M., Flato, G., Friedlingstein, P., Jones, C., Lamarque, J.-F., Liao, H., Rasch, P. (eds.)]. In: Stocker, T. et al. (Editors), Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. doi: [10.1017/CBO9781107415324.030](https://doi.org/10.1017/CBO9781107415324.030).
- Jamieson, P. D., Semenov, M.A., Brooking I.R. and Francis, G. S. 1998. Sirius: a mechanistic model of wheat response to environmental variation. *European Journal. Agronomy*. 8, 161–179. doi: [10.1016/S1161-0301\(98\)00020-3](https://doi.org/10.1016/S1161-0301(98)00020-3).
- Lizaso, J., Ruiz-Ramos, M., Rodríguez, L., Gabaldon-Leal, C., Oliveir, J.A., Lorite, I.J., Rodríguez, A., Maddonni, G.A., Otegui, M.E., 2017. Modeling the response of maize phenology, kernel set, and yield components to heat stress and heat shock with CSM-IXIM. *Field Crops Res.* 214, 239–252. doi: [10.1016/j.fcr.2017.09.019](https://doi.org/10.1016/j.fcr.2017.09.019).
- Martre, P. and Dambreville, A., 2018. A model of leaf coordination to scale-up leaf expansion from the organ to the canopy. *Plant Physiol.* 176, 704–716. doi: [10.1104/pp.17.00986](https://doi.org/10.1104/pp.17.00986).
- McDermid, S. P., et al. The AgMIP Coordinated Climate-Crop Modeling Project (C3MP): Methods and Protocols. *Handbook of Climate Change and Agroecosystems*: 191-220. doi: [10.1142/9781783265640_0008](https://doi.org/10.1142/9781783265640_0008).
- Nendel, C., Berg, M., Kersebaum, K.C., Mirschel, W., Specka, X., Wegehenkel, M., Wenkel, K.O., Wieland, R., 2011. The MONICA model: Testing predictability for crop growth, soil moisture and nitrogen dynamics. *Ecological Modelling*, Elsevier, vol. 222(9), pages 1614-1625. doi: [10.1016/j.ecolmodel.2011.02.018](https://doi.org/10.1016/j.ecolmodel.2011.02.018).
- Olesen, J.E., Petersen, B.M., Berntsen, J., Hansen, S., Jamieson, P.D., Thomsen, A.G., 2002. Comparison of methods for simulating effects of nitrogen on green area index and dry matter growth in winter wheat. *Field Crops Res.* 74, 131–149 (2002). doi: [10.1016/S0378-4290\(01\)00204-0](https://doi.org/10.1016/S0378-4290(01)00204-0).
- Ruane A.C., Winter, J.M., McDermid, S.P. and Hudson, N.I., 2015. AgMIP Climate Data and Scenarios for Integrated Assessment. In: *Handbook of Climate Change and Agroecosystems*. Imperial College Press, p 45–78. doi: [10.1142/9781783265640_0003](https://doi.org/10.1142/9781783265640_0003) / doi: [10.1038/s41467-018-06525-2](https://doi.org/10.1038/s41467-018-06525-2).
- Soltani, A., Maddah, V., and Sinclair, T.R., 2013. SSM-wheat: a simulation model for wheat development, growth and yield. *Int. J. Plant Prod.* 7, 711–740. doi: [10.22069/ijpp.2013.1266](https://doi.org/10.22069/ijpp.2013.1266).
- Webber, H., Ewert, F., Kimball, B.A., Siebert, S., White, J.W., Wall, G.W., Ottman, M.J., Trawally, D.N.A., Gaiser, T., 2016. Simulating canopy temperature for modelling heat stress in cereals. *Environ. Model. Softw.* 77, 143–155. doi: [10.1016/j.envsoft.2015.12.003](https://doi.org/10.1016/j.envsoft.2015.12.003).
- Webber, H., Ewert, F., Olesen, J.E., Müller, C., Fronzek, S., Ruane, A.C., Bourgault, M., Martre, P., Ababaei, B., Bindi, M., Ferrise, R., Finger, R., Fodor, N., Gabaldón-Leal, C., Gaiser, T., Jabloun, M., Kersebaum, K.-C., Lizaso, J. I., Lorite, I.J., Manceau, L., Moriondo, M., Nendel, C., Rodríguez, A., Ruiz-Ramos, M., Semenov, M.A., Siebert, S., Stella, T., Stratonovitch, P., Trombi, G. and Wallach, D., 2018a. “Diverging importance of drought stress for maize and winter wheat in Europe”. *Nature Communications* 9(1): 1-11. doi: [10.1038/s41467-018-06525-2](https://doi.org/10.1038/s41467-018-06525-2).
- Webber, H., J. White, B. Kimball, Ewert, F., Asseng, S., Rezaei, E.E., and Martre, P., 2018b. “Physical robustness of canopy temperature models for crop heat stress simulation across environments and production conditions.” *Field Crops Research* 216: 75 - 88. doi: [10.1016/j.fcr.2017.11.005](https://doi.org/10.1016/j.fcr.2017.11.005).
- Webber, H., Ababaei, B., Bindi, M., Ferrise, R., Fodor, N., Jabloun, M., Kersebaum, K.-C., Lizaso, J.I., Manceau, L., Moriondo, M., Martre, P., Nendel, C., Rodríguez, A., Ruiz-Ramos, M., Semenov, M.A., Siebert, S., Stella, T., Stratonovitch, P., Trombi, G., Ewert, F., Ruane, A.C., Lorite, I.J., Gabaldón-Leal, C., Gaiser, T. and Olesen, J., 2019. Diverging importance of drought stress for maize and winter wheat in Europe: Simulation Dataset. doi: [10.4228/ZALF.DK.88](https://doi.org/10.4228/ZALF.DK.88).