



Diversification improves the performance of cereals in European cropping systems

Moritz Reckling, Johannes Albertsson, Anthony Vermue, Georg Carlsson, Christine Watson, Eric Justes, Göran Bergkvist, Erik Steen Jensen, Cairistiona Topp

► To cite this version:

Moritz Reckling, Johannes Albertsson, Anthony Vermue, Georg Carlsson, Christine Watson, et al.. Diversification improves the performance of cereals in European cropping systems. *Agronomy for Sustainable Development*, 2022, 42 (6), pp.118. 10.1007/s13593-022-00850-z . hal-03922945

HAL Id: hal-03922945

<https://hal.inrae.fr/hal-03922945>

Submitted on 11 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Diversification improves the performance of cereals in European cropping systems

Moritz Reckling^{1,2} · Johannes Albertsson³ · Anthony Vermue⁴ · Georg Carlsson³ · Christine A. Watson^{2,5} · Eric Justes^{4,6} · Göran Bergkvist² · Erik Steen Jensen³ · Cairistiona F. E. Topp⁵

Accepted: 17 November 2022 / Published online: 5 December 2022
© The Author(s) 2022

Abstract

In the face of climate change, cropping systems need to achieve a high performance, providing food and feed and adapting to variable environmental conditions. Diversification of cropping systems can support ecosystem services and associated biodiversity, but there is little evidence on which temporal field arrangement affects the performance of crop yields (productivity and stability), partly due to a lack of long-term data and appropriate indicators. The objectives of this study were to quantify the effect of cropping system diversification on yield stability, environmental adaptability, and the probability of diversified systems to outperform less diverse cereal-based systems in Europe. Spring and winter cereal yields were analyzed from long-term field experiments from Sweden, Scotland, and France. We investigated diversification through (i) introduction of perennial leys, (ii) increasing the proportion of ley in the rotation, (iii) varying the order in which crops are positioned in the rotation, (iv) introduction of grain legumes, and (v) introduction of cover crops. The results showed that cereal crops within cropping systems incorporating perennial leys outperformed systems without leys in 60–94% of the comparisons with higher probabilities at low fertilizer intensities. The yield stability of oat did not differ, but mean yields were 33% higher, when grown directly after the ley compared to oat grown two years later in the crop sequence under similar management. Durum wheat grown in a cropping system with grain legumes had higher yields in lower-yielding environmental conditions compared to rotations without legumes. Diversification with cover crops did not significantly affect yield stability. We conclude that diverse cropping systems can increase cereal productivity and environmental adaptability and are more likely to outperform less diverse systems especially when introducing perennial forage legumes into arable systems. Effects of diversification on cereal yield stability were inconsistent indicating that higher productivity is achievable without reducing yield variability. These novel findings can support the design of more diverse and high-performing cropping systems.

Keywords Coefficient of variation · Cover crops · Legumes · Ley · Oat · Rotation · Stability · Taylor's power law · Variability · Wheat

✉ Moritz Reckling
moritz.reckling@zalf.de

¹ Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany

² Department of Crop Production Ecology, Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden

³ Department of Biosystems and Technology, Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden

⁴ UMR AGIR, University of Toulouse, INRAE, INPT, Centre de Recherches INRAE Occitanie-Toulouse, Auzeville-Tolosane, France

⁵ Scotland's Rural College (SRUC), Edinburgh, UK

⁶ CIRAD, Peryst Departement, Montpellier, France

1 Introduction

Agricultural cropping systems will experience increased perturbations due to climate change (Ray et al. 2015; Tigchelaar et al. 2018). The projected climate effects include increases in temperature and changes in patterns of precipitation and an increased frequency and severity of extreme weather events such as drought, heat stress, and flooding (Lobell et al. 2011). The increase in climate variation, alongside other factors, has resulted in decreased yield stability of major crop species in large parts of Europe and other regions of the world (Döring and Reckling 2018; Ray et al. 2015), and yield stability of crops is predicted to decrease

even more under future climate change scenarios (Tigchelaar et al. 2018).

Cropping systems need to be designed to enhance crop yields while maintaining or increasing yield stability under varying climatic conditions (Liu et al. 2019). Besides the choice of the evaluated outputs (e.g., yield stability), the factors influencing these outputs (e.g., rotational effects and climatic conditions) need to be considered in the design of diversified cropping systems. While most studies use simulation modeling to assess cropping systems under various conditions (Bergez et al. 2010), few use empirical data, e.g., where cropping systems are assessed over long time periods taking the role of crop rotations into account (Marini et al. 2020). Crop rotation principles are important in the design and assessment of farming systems because they describe the sequence and frequency of crops grown and the interactions within a system. Assessing yield stability of crops in different cropping systems requires yield data over long time periods. Long-term experiments (LTE) that compare different cropping systems provide such data (Fig. 1), i.e., crop yields over long time periods under different conditions and rotations (Johnston and Poulton 2018). The assessment of such experiments can provide fundamental knowledge on the role of crop rotation diversification in affecting yield stability, the probability of diversified systems to outperform less diverse systems, and environmental adaptability. As detailed in Section 2.2.4, the latter is assessed based on crop yields of individual cropping systems against the mean yields of a range of cropping systems. Such knowledge could support

the design of future cereal-based cropping systems that are better able to deal with the increasing climate variability.

There are several hundred LTEs available worldwide, and these are mainly analyzed for single research objectives comparing few treatments. It is only recently that the yield data from LTEs have been used to assess yield stability for separate LTEs (see Reckling et al. (2021) for a review) and also in combined analyses using several international LTEs (Marini et al. 2020; Reckling et al. 2018). Data from LTEs incorporate changes in climate and changes associated with the management such as the impacts of crop diversification on soil structure and soil carbon that affect crop performance.

Incorporating perennial leys and grain legumes into cropping systems increases the yield of subsequent crops under most conditions compared to systems without legumes and perennial crops (Angus et al. 2015; Preissel et al. 2015; St-Martin et al. 2017). This is due to a positive pre-crop effect (residual and break crop effect) that enhances main crop yields by 20% on a global average (Zhao et al. 2022). Cover crops can significantly reduce nutrient leaching and also affect the yield of subsequent crops (Hauggaard-Nielsen et al. 2012; Plaza-Bonilla et al. 2016). While the effect of these diversification strategies (incorporating leys with different lengths, grain legumes, cover crops, and crop sequences) on yield has been observed in separate analyses, the effect of these strategies on yield stability, environmental adaptability, and the probability of diversified systems to outperform less diverse systems has not been investigated.

Fig. 1 Tulloch is a Scottish LTE established in 1991 with a 6-year crop rotation. In the stocked system, crops are partly grazed by sheep and receive farm-yard manure (picture: C. Watson/SRUC).



While stability analysis was originally used to assess the stability of crop genotypes across environments, the analysis of yield stability of cropping systems, especially in relation to climate change, has gained importance (Lobell et al. 2011; Tigchelaar et al. 2018). To assess yield stability, environmental adaptability, and the probability of systems to outperform other systems, different regression- and variance-based indicators have been proposed (see Reckling et al. (2021) for an overview).

There are a number of types of indicators that can help to assess cropping systems in relation to yield stability, environmental adaptability, and the probability of diversified systems to outperform less diverse systems: (i) Static and variance-based yield stability indicators include the relatively simple coefficient of variation (CV), which is one of the most frequently used indicators in agronomic and ecological research (Ray et al. 2015). Applying the CV implies the assumption that the standard deviation increases linearly with the mean. However, under certain conditions, the unguarded interpretation of the CV of crop yield data may be misleading, especially when the crop yield data spans a large numeric range (Döring et al. 2015; Döring and Reckling 2018). (ii) Another static yield stability indicator is the POLAR (Power Law Residuals) (Döring et al. 2015) that estimates yield stability of crops independent of differences in mean yields between cropping systems. (iii) Dynamic indicators can be expressed as a measure of environmental adaptability. A regression-based indicator following Finlay and Wilkinson (1963) is used in this study to assess the interaction between the yield performances of the single crop within a cropping system in relation to the mean yield of this crop over all cropping systems. (iv) The probability that one cropping system outperforms another system can be estimated to account for the variance and also the mean yield between systems (Piepho 1998).

The objective of this study was to assess the effect of cropping system diversification strategies across different European climates on cereal yield stability, environmental adaptability, and the probability that diversified systems outperform less diverse systems. We investigated diversification through (i) integration of perennial leys, (ii) increasing proportion (length, i.e., number of years) of the perennial ley relative to the entire crop rotation, (iii) varying the order in which crops are positioned in the rotation, (iv) integration of grain legumes, and (v) integration of cover crops. We used cereal yield data from five LTEs from Sweden, Scotland, and France.

2 Materials and methods

2.1 Characteristics of long-term experiments

Long-term experiments (LTEs) containing cereals in different cropping systems from Sweden, France, and Scotland

with different bio-physical conditions and experimental designs (Table 1) were used for the analyses of yield stability.

2.1.1 Swedish long-term experiments

Annual cereal yield data (winter wheat and oats) was used from three Swedish LTEs located at Lanna, Stenstugu, and Säby and established in 1965, 1968, and 1969, respectively. The time periods used for this study were 1971–2014 (44 years), 1974–2014 (41 years), and 1975–2014 (40 years) for Lanna, Stenstugu, and Säby, respectively (Table 1). The experimental design included three different crop rotations (A, with grass-clover ley; B, with grass ley; and C, without ley; Table S1) and four levels of N fertilization (N0, N1, N2, and N3; Table S1) with amounts of 0 kg ha⁻¹ year⁻¹, 38–55 kg ha⁻¹ year⁻¹, 73–100 kg ha⁻¹ year⁻¹, and 105–145 kg ha⁻¹ year⁻¹, respectively, and depending on crop. The design has no spatial replicates but all crops in the crop rotations are present each year. Each crop rotation × N level combination is hereafter referred to as a cropping system. All nitrogen was applied as a mineral fertilizer (see Persson et al. (2008) for further details).

In all cropping systems, the crop sequence was the same for the first 4 years of the rotation (Table S1): winter/spring oilseed rape (*Brassica napus* L.) or white mustard (*Sinapis alba* L.), winter wheat (*Triticum aestivum* L.), spring oat (*Avena sativa* L.), and spring barley (*Hordeum vulgare* L.). The grass-clover ley of crop rotation A consisted of red clover (*Trifolium pratense* L.) and timothy (*Phleum pratense* L.) at Lanna and red clover, timothy, and alfalfa (*Medicago sativa* L.) at Stenstugu and Säby. The grass-clover ley contained > 30% legumes in the biomass in the mixture in most years. However, there was a large variation in the legume percentage between years and rotation systems but there was no detailed data available. The grass ley of crop rotation B consisted of timothy and meadow fescue (*Festuca pratensis* L.) at all sites. Spring wheat (*Triticum aestivum* L.) and a black fallow was used in crop rotation C instead of first and second year leys in rotations A and B (Table S1).

To avoid confounding effects of previous management at the experimental sites, the first 6-year rotation cycle was excluded from all analyses. Due to poor drainage at Säby, the winter survival of winter wheat was poor, and thus, the complete winter wheat dataset was excluded from the analysis. Spring barley was excluded because it was managed differently in the three crop rotations (Table S1).

2.1.2 Scottish long-term experiments

Tulloch, the Scottish LTE, was established in 1991 in the North-East of Scotland. The time periods used for this study were 1994–2017 (24 years) (Table 1). The 6-year crop rotations

Table 1 Details about the long-term experiments.

LTE name	Coordinates	Country	Annual crops analyzed	Time period	Clay (%)	Silt (%)	Sand (%)	Organic matter (%)	Mean temperature (C°) ^c	Precipitation (mm) ^c
Lanna ^a	58° 20' N, 13° 07' E	Sweden	Winter wheat, oat	1971–2014	36.5	51.2	12.3	4.2	6.9	667
Stenstugu ^a	57° 36' N, 18° 26' E	Sweden	Winter wheat, oat	1974–2014	21.6	31.9	46.5	2.8	7.4	792
Säby ^a	59° 49' N, 17° 42' E	Sweden	Oat	1975–2014	24.0	55.2	20.8	4.0	6.3	617
Auzeville ^b	43° 31' N, 1° 30' E	France	Durum wheat	2005–2016	25.6	36.8	37.6	1.5	13.8	669
Tulloch	57° 11' N, 2° 16' W	Scotland	Oat	1994–2017	13.0	17.0	70.0	9.0	7.9	816

^aSoil texture (recalculated) and organic matter from Persson et al. (2008)^bSoil texture and organic matter (recalculated) from Plaza-Bonilla et al. (2016)^cMean daily temperature and mean annual total precipitation during the experimental periods

have each course of the rotation present in each year, and there are two spatial replicates of each rotation (see Watson et al. (2011) for further details). The T50 crop rotation (see Table S2) consisted of 3 years of grass-clover ley (*Lolium perenne* L., *P. pratense* L., and *Trifolium repens* L.), followed by spring oats (*A. sativa* L.), swede (*Brassica napus* L.), and then, barley with undersown ley. The T67 system had 4 years of grass-clover ley followed by spring oats and then spring oats undersown with the grass-clover ley mixture. These rotations were grazed by sheep and received farm-yard manure as described in Table S2. In 2007, the T67 rotation was converted to a stockless rotation. The first 4 years of the oat data were excluded from the analysis as they did not follow 4 years of grass-clover.

2.1.3 French long-term experiment

The French LTE is located at Auzeville and was established in 2004. The time periods used for this study was 2005–2016 (12 years) (Table 1). The LTE has a split-plot design with crop rotation (R1 and R2; Table S3) as the main plot and cover crops as sub-plots (with or without; Table S3). There are no spatial replicates, but all crops are present each year.

Four cropping systems resulting from a combination of two 3-year crop rotations (Table S3) with or without cover crops were used in the present study. The R1 rotation consisted of durum wheat (*Triticum turgidum* L.) followed by sorghum (*Sorghum bicolor* L.) and then sunflower (*Helianthus annuus* L.). Crop rotation R2 had durum wheat followed by sunflower and then winter pea (*Pisum sativum* L.) during 2004–2010 and winter faba bean (*Vicia faba* L.) during 2011–2016 (Table S3). A vetch-oat (*Vicia sativa* L.–*A. sativa* L.) mixture was used as a cover crop in crop rotation R2 while a combination of vetch and other crop species was used in rotation R1 (see Table S3 for details). Analyses were performed for the grain yields from durum wheat, since cereals were the target crop types for this study and it was the only crop that was present in both crop rotations with the same pre-crops. Further information regarding the LTE can be found in Plaza-Bonilla et al. (2016).

2.2 Statistical analyses

2.2.1 Probability method (PM)

The probability of one system outperforming another system is a method described by Piepho (1998) that compares single crops, e.g., cereals in two cropping systems directly with each other in terms of differences in yield and their variance and co-variance (Piepho 1998) and is calculated as

$$\Pr(D_j > 0) = \Phi[\delta/\sigma_D] \quad (1)$$

where Φ is the cumulative distribution function of the standard normal distribution, δ is the estimated mean difference between systems, and $\sigma^2 D_j$ is the variance of a difference D_j in a randomly selected environment following Piepho (1998).

There is a connection between the probability of one system outperforming another and the variance-based stability indicators mentioned above (CV and POLAR) through accounting for the mean, the variance, and covariance of the yield data. The probability indicator was used in this study to calculate the likelihood that one system, e.g., with perennial ley, “outyields” the system without perennial ley in more than 50% of the cases. For the Swedish LTE, each cropping system was considered as a system, and all analyses were performed separately for each site and crop combination. The environmental variance model described by Piepho (1999) was used to calculate estimates of variances and mean yields for the probability calculations. Probability calculations were not performed for the French and Scottish LTEs due to the limited availability of data.

2.2.2 Coefficient of variation (CV)

The yield stability of cropping systems was determined by the coefficient of variation (CV). The CV is defined as the standard deviation σ divided by the mean μ and is calculated as:

$$CV = \sigma/\mu \cdot 100\% \quad (2)$$

For the Swedish LTEs, the dataset of each cropping system was divided into one subset for each crop rotation cycle. Hence, every CV value comprised yield data from 6 years, except for the seventh crop rotation cycle at Stenstugu and Säby, which only comprised yield data from 5 and 4 years, respectively. The winter wheat grain yield data from crop rotation cycle (subset) five at Stenstugu were excluded from the CV analyses due to missing data. The dataset from the French and Scottish LTE was also divided in subsets. The CV values were analyzed statistically using the MIXED procedure in SAS and Tukey’s test ($\alpha = 0.05$) was used as a post hoc test. The dataset of the Swedish LTEs was analyzed by comparing the mean CV values between crop rotations (A, B, and C) within each site \times crop \times N level combination. Rotation cycle (subset) and crop rotation were considered fixed factors in these analyses. To assess the effect of two different crop rotations on the durum wheat yield stability, the dataset of the French LTE was analyzed by comparing mean CV values between the R1 rotations with grain legumes (with and without cover crop) and the R2 rotations without grain legumes (with and without cover crop), since the interaction between the systems was not significant. To assess the effect of cover crops on the durum wheat yield

stability, mean CV values of cropping systems with cover crops (R1 with and R2 without grain legumes) and without cover crops (R1 with and R2 without grain legumes) were compared. Rotation cycle (subset), crop rotation, and cover crop were considered as fixed factors in the French analysis. The Scottish dataset was also analyzed in two different ways. To assess the yield stability of spring oat that was directly following the grass-clover ley (oat position 1) with the yield stability of spring oat separated from the grass-clover ley by other crops in the rotation (oat position 2), CV values from the Scottish T50 rotation were compared for the period 1994–2017. CV values of spring oats in the 3-year ley (T50) and 4-year ley (T67) rotations were compared without any statistical analyses because of an insufficient number of crop rotations cycles.

2.2.3 Power Law Residuals (POLAR)

The Power Law Residuals (POLAR) is an index of yield stability that is independent of the mean yield in contrast to the CV (Döring et al. 2015; Reckling et al. 2018). POLAR is based on Taylor’s power law (TPL), which states that the logarithm of the sample variance (σ^2) is a linear function of the logarithm of the sample mean (μ) across different subsets of data. Using the same subsets of data as described for the calculation of the CV, means ($\hat{\mu}$) and variances ($\hat{\sigma}^2$) were calculated resulting in pairs (with index i) consisting of a mean and a variance. Following TPL, a linear regression is calculated for \log_{10} of the variance over the \log_{10} of the mean. With $v_i = \log(\hat{\sigma}^2_i)$ and $m_i = \log(\hat{\mu}_i)$, the linear regression is $v = a + bm$. The residuals u_i from this regression line (the POLAR values) are then calculated according to Döring et al. (2015) as:

$$u_i = v_i - (a + bm_i) \quad (3)$$

A low POLAR value corresponds to a high stability, whereas a high value corresponds to a low stability. For the POLAR calculations, all yield data from the LTEs were divided into the same subsets as for the CV analyses, and for each subset, a mean and a variance were calculated. Subsequently, a linear regression was calculated for \log_{10} of the variance against the \log_{10} of the mean using all yield data subsets from all LTEs following (Döring et al. 2015; Reckling et al. 2018). The yield data from spring oat and winter wheat from Lanna and Stenstugu, spring oat from Säby and the Scottish LTE, and durum wheat for the French LTE were used for the calculation of the linear regression. The number of values for the regression was the same as for the calculated CVs. The POLAR values (residuals) obtained from this regression line were compared and statistically analyzed as described for the CV values.

2.2.4 Environmental adaptability (FW)

In this regression analyses, annual yields of individual cropping systems were plotted against the mean annual yields of all cropping systems included in the comparison (Finlay and Wilkinson 1963; Piepho 1998) to be interpreted as an indicator of environmental adaptability. The performance of an individual cropping system is evaluated by comparing the slope of the regression lines (FW, Finlay-Wilkinson regression coefficient b_i) independent of the intercept. A positive slope value greater than one ($b > 1$) indicates higher environmental adaptability. Higher environmental adaptability means that favorable environmental conditions result in higher yields. A slope value lower than one ($b < 1$) indicates a lower environmental adaptability. The slopes within each site \times crop \times N level combination of the Swedish LTE were compared. In the French LTE, the slopes of the R1 rotations (with and without cover crop) and the R2 rotations (with and without cover crop) were compared and the slopes of cropping systems with cover crops (R1 and R2) and without cover crops (R1 and R2). The slopes of the Scottish crop rotations T50 and T67 at first oat position were compared using data between 1995 and 2006, and the slopes of the two different oat positions were compared using data from the T50 crop rotation between years 1994 and 2017. Following Litell et al. (2006), all comparisons of slopes were performed using the MIXED procedure in SAS.

3 Results and discussion

3.1 Diversification through perennial legume crops

Winter wheat yields in the Swedish cropping system with the perennial grass-clover ley (A) outperformed winter wheat yields in the system without the ley (C) across all sites and N levels with a probability of 64–94% in the Swedish LTEs (Table 2 for the site Lanna, S4 for Stenstugu, and S5 for Säby). Winter wheat in the system with perennial grass ley (B) also outperformed the wheat in the system without ley (C) but only with a probability of 55–79% (Table 2). Diversification with perennial grass-clover ley (A) and grass ley (B) did not consistently affect winter wheat yield stability quantified with the CV (yield dependent indicator) and POLAR (yield independent indicator) compared to wheat in the system without perennial ley (C) (Table 3). In Stenstugu in the 0 N treatment, yield stability was significantly higher with a CV of 23% for winter wheat in the systems with perennial ley (small CV value indicates higher stability) compared to the system without ley with a CV of 29% (Table 3). There was no significant difference in the environmental adaptability, although the slope of wheat in the

systems with perennial crops had consistently higher values than 1 indicating higher environmental adaptation (Table 3).

Similar to the probability analysis for winter wheat, grain yields of oats in the Swedish crop rotations with perennial leys (A and B) outperformed the oat yield in the system without a perennial crop (C) by 55–95%, except at the Säby site at N level 3 (Table 2). Oat yield stability (CV and POLAR) was not affected by the presence of perennial leys, except for N level 3 at Säby. At this site and N level, the CV of oat yields was significantly lower (indicating higher stability) without a perennial crop (C) compared to the system with a perennial grass ley (B) (Table 4) but not for the yield-independent POLAR coefficient. In contrast to winter wheat, the presence of perennial crops affected the response of the oat yield to the environmental conditions. At N level 0 and all sites, the slope values of the environmental adaptability coefficient for crop rotations including grass-clover leys and grass leys (A, B) were significantly higher than without (C) indicating higher environmental adaptability (Table 4, Fig. 2). For fertilization levels N1 and N2 at Stenstugu, the coefficient for rotation A was significantly higher than rotation C indicating higher environmental adaptability (Table 4). At Säby, a higher environmental adaptability was found for N2 and N3 for systems with perennial leys.

Our results partly support our first hypothesis that cropping systems with perennial leys (in the three Swedish LTEs) outperformed those without leys in terms of winter wheat and oat yields (Table 2). Sanford et al. (2021) found that systems with a greater proportion of perennial crops had the highest long-term stability. In our study, the magnitude of differences in yield was similar for winter wheat and oat and across the three sites but decreased from 92 to 64% with increasing nitrogen fertilizer application rate (Tables 2, S4, S5). Thus, the greatest impact of crop diversification was found in the low-input systems. In such systems, many factors can be limited including nitrogen, and the residues from the perennial crops provide nutrients and better growing conditions for the following cereals compared to other pre-crops. Forage legumes fix more dinitrogen when N is limiting and contribute to better soil conditions of the following crops (Iannetta et al. 2016). It has already been shown for the Swedish LTEs investigated here that the systems with perennial grass and grass-clover increased soil organic carbon (Persson et al. 2008) and that the yields of the cereals were greater than in the system without leys (Bergkvist and Båth 2015). Macholdt et al. (2020) found differences in yield stability (indicated as the risk of yield falling below a threshold) between less and more diverse systems to be larger in low-input treatments. On the other hand, crops in such low-input systems had the highest production risk (probability of yield loss) compared to fertilized systems (Macholdt et al. 2020). We also found an indication that non-fertilized systems were less stable than fertilized systems

Table 2 Probability of the i th treatment combination (crop rotation \times N level) outperforming the i' th treatment combination for winter wheat and oat grain yields during 1971–2014 at the site Lanna.

System i	System i'																								
	Winter wheat												Oat												
	A0	B0	C0	A1	B1	C1	A2	B2	C2	A3	B3	C3	A0	B0	C0	A1	B1	C1	A2	B2	C2	A3	B3	C3	
A0	-	0.80	0.92	0.02	0.03	0.08	0.00	0.00	0.00	0.01	0.01	0.01	-	0.73	0.92	0.01	0.03	0.05	0.01	0.01	0.01	0.01	0.02	0.01	0.03
B0	0.20	-	0.75	0.01	0.01	0.02	0.00	0.00	0.00	0.01	0.01	0.01	0.27	-	0.76	0.01	0.01	0.02	0.00	0.00	0.00	0.01	0.01	0.01	0.02
C0	0.08	0.25	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.24	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
A1	0.98	0.99	1.00	-	0.80	0.82	0.02	0.12	0.25	0.10	0.14	0.14	0.99	0.99	1.00	-	0.80	0.82	0.10	0.13	0.22	0.12	0.11	0.18	0.18
B1	0.97	0.99	1.00	0.20	-	0.55	0.04	0.04	0.04	0.08	0.08	0.08	0.97	0.99	1.00	0.20	-	0.57	0.05	0.03	0.07	0.08	0.06	0.11	0.11
C1	0.92	0.98	1.00	0.18	0.45	-	0.03	0.04	0.01	0.06	0.07	0.06	0.95	0.98	1.00	0.18	0.43	-	0.04	0.04	0.04	0.06	0.05	0.09	0.09
A2	1.00	1.00	1.00	0.98	0.96	0.97	-	0.71	0.80	0.35	0.44	0.51	0.99	1.00	1.00	0.90	0.95	0.96	-	0.59	0.75	0.32	0.32	0.43	0.43
B2	1.00	1.00	1.00	0.88	0.96	0.96	0.29	-	0.69	0.24	0.31	0.33	0.99	1.00	1.00	0.87	0.97	0.96	0.41	-	0.74	0.27	0.22	0.35	0.35
C2	1.00	1.00	1.00	0.75	0.96	0.99	0.20	0.31	-	0.21	0.26	0.23	0.99	1.00	1.00	0.78	0.93	0.96	0.25	0.26	-	0.16	0.15	0.23	0.23
A3	0.99	0.99	1.00	0.90	0.92	0.94	0.65	0.76	0.79	-	0.56	0.64	0.98	0.99	1.00	0.88	0.92	0.94	0.68	0.73	0.84	-	0.50	0.64	0.64
B3	0.99	0.99	1.00	0.86	0.92	0.93	0.56	0.69	0.74	0.44	-	0.56	0.99	0.99	1.00	0.89	0.94	0.95	0.68	0.78	0.85	0.50	-	0.64	0.64
C3	0.99	0.99	1.00	0.86	0.92	0.94	0.49	0.67	0.77	0.36	0.44	-	0.97	0.98	1.00	0.82	0.89	0.91	0.57	0.65	0.77	0.36	0.36	-	-

A0, B0, C0, etc. refer to three different crop rotations (A, with grass-clover ley; B, with grass ley; and C, without ley) and four levels of N fertilization (0, 1, 2, and 3) with amounts of 0 kg ha⁻¹ year⁻¹, 38–55 kg ha⁻¹ year⁻¹, 73–100 kg ha⁻¹ year⁻¹, and 105–145 kg ha⁻¹ year⁻¹, respectively, and depending on crop (see Table S1). Probability calculations according to Piepho (1998) and Piepho (1999) using the environmental variance model. The probability that crop rotation A is outperforming crop rotation B and C within the same N level is italicized. The probability that crop rotation B is outperforming crop rotation C within the same N level is in italic presentation

Table 3 Mean winter wheat grain yields, mean CV values, and Finlay-Wilkinson (FW) regression coefficient b_i for each crop rotation and N level combinations at sites Lanna and Stenstugu.

Crop rotation	N level	Lanna				Stenstugu			
		Grain yield (Mg DM/ha \pm SEM)	CV	FW b_i	POLAR	Grain yield (Mg DM/ha \pm SEM)	CV	FW b_i	POLAR
A	0	2.81 \pm 0.11	20.2	1.00	− 0.20	3.05 \pm 0.13	23.5 b	1.07	0.01
B	0	2.33 \pm 0.10	23.2	1.00	− 0.15	2.46 \pm 0.11	23.0 b	0.97	− 0.14
C	0	2.04 \pm 0.10	23.7	1.00	− 0.24	2.18 \pm 0.11	29.2 a	0.96	0.04
<i>P</i> value			0.7086	0.9979	0.8653		0.0017	0.4356	0.0850
A	1	4.18 \pm 0.13	19.2	1.07	− 0.06	4.44 \pm 0.12	16.6	1.08	− 0.12
B	1	3.67 \pm 0.11	17.9	0.93	− 0.18	3.93 \pm 0.11	16.6	0.99	− 0.19
C	1	3.59 \pm 0.12	18.4	1.00	− 0.21	3.86 \pm 0.11	16.0	0.93	− 0.22
<i>P</i> value			0.8657	0.4277	0.2648		0.8833	0.1432	0.5278
A	2	5.02 \pm 0.14	16.7	1.05	− 0.07	5.36 \pm 0.15	14.8	0.98	− 0.13
B	2	4.75 \pm 0.13	17.2	0.99	− 0.06	4.96 \pm 0.15	16.4	1.01	− 0.11
C	2	4.52 \pm 0.13	17.7	0.95	− 0.14	4.91 \pm 0.15	16.7	1.01	− 0.07
<i>P</i> value			0.9173	0.4354	0.7939		0.5893	0.8602	0.8492
A	3	5.25 \pm 0.18	19.2	1.03	0.05	5.54 \pm 0.20	19.5	0.98	0.12
B	3	5.07 \pm 0.18	20.3	1.01	0.09	5.40 \pm 0.21	21.8	1.06	0.19
C	3	4.94 \pm 0.17	20.5	0.96	0.08	5.37 \pm 0.19	20.1	0.96	0.13
<i>P</i> value			0.8522	0.7590	0.9087		0.5724	0.1287	0.7295

A, B, and C refer to three different crop rotations (A, with grass-clover ley; B, with grass ley; and C, without ley) and four levels of N fertilization (0, 1, 2, and 3) with amounts of 0 kg ha^{−1} year^{−1}, 38–55 kg ha^{−1} year^{−1}, 73–100 kg ha^{−1} year^{−1}, and 105–145 kg ha^{−1} year^{−1}, respectively, and depending on crop (see Table S1). Means comprise seven crop rotation cycles for all sites except Stenstugu which only comprise six rotation cycles. For CV and FW b_i , within columns per N level, means followed by different letters are significantly different

(Tables 3 and 4) but we did not find a systematic effect of diversification through leys on yield stability of winter wheat and oat. However, there was clear indication of higher environmental adaptability of the cropping systems with perennial leys, especially for oat in low-input systems across all sites (Fig. 2). The residual nitrogen from roots and crop residues of forage legumes might be one of the reasons to support crop growth of spring oat when nitrogen is limiting (without nitrogen fertilization) especially under favorable climatic conditions (with a high nitrogen demand for crop growth). Under unfavorable conditions, e.g., water limitation in dry years, we found only slightly higher yields of spring oat (probably factors other than nitrogen were limiting). In the analysis by Marini et al. (2020), they found that the more diverse systems performed even better under these poor conditions and concluded that this increases their potential for adapting to changes in climate.

3.2 Diversification through the length of the perennial ley crops

The period of perennial leys can vary and affect the subsequent crop production. At the Scottish site, the length of the ley phase had no effect on the mean yield and yield stability of the oats as indicated by the CV and the POLAR and no significant difference in the adaption to the environmental

conditions according to the FW regression (Table 5). Thus, diversification through increasing the length of the perennial ley had no significant effect on yield regardless of the environmental conditions. This is supported by the Finlay and Wilkinson analysis showing no difference in adaption to environmental conditions although the 3-year ley had a slope < 1 indicating lower adaptability than the 4-year ley with a slope > 1 (Table 5). The difference between the systems (both already very diverse with perennial grass-clover) was probably too small to impact yield variation. However, the longer ley phase in the rotation might have positively affected the nitrogen balance due to more crop residues and higher N input via N₂ fixation (Iannetta et al. 2016). The higher N input of the longer ley phase is also reflected in a higher N content of oat grain following the 4-year ley compared with the 3-year ley (Watson et al. 2011).

3.3 Diversification through the crop sequence

The place of a crop within a cropping sequence can differ and affect the productivity. Oat yields in the three-year perennial ley rotation in Scotland were 33% higher when following directly after the ley compared to oat grown 2 years later in the crop sequence (second oat, following the first oat and the swede crop; Table 5). Even though the CV and POLAR value of the first oat position was numerically

Table 4 Mean oat grain yields, mean CV values, Finlay-Wilkinson (FW) regression coefficient b_i and POLAR coefficients for each crop rotation and N level combination at sites Lanna, Stenstugu and Säby.

Crop rotation	N level	Lanna				Stenstugu				Säby			
		Grain yield (Mg DM/ha \pm SEM)	CV	FW b_i	POLAR	Grain yield (Mg DM/ha \pm SEM)	CV	FW b_i	POLAR	Grain yield (Mg DM/ha \pm SEM)	CV	FW b_i	POLAR
A	0	1.94 \pm 0.10	30.8	1.09 a	0.01	2.02 \pm 0.10	28.9	1.09 a	-0.07	2.24 \pm 0.15	41.1	1.25 a	0.29
B	0	1.68 \pm 0.10	35.8	1.04 a	0.06	1.66 \pm 0.10	36.4	1.07 a	0.06	1.94 \pm 0.12	37.2	0.98 b	0.13
C	0	1.36 \pm 0.07	32.4	0.80 b	-0.11	1.31 \pm 0.08	33.2	0.79 b	-0.13	1.57 \pm 0.10	36.0	0.77 c	0.04
P value			0.3026	0.0068	0.1951		0.1545	0.0030	0.2494		0.6407	0.0001	0.1083
A	1	3.22 \pm 0.12	21.2	1.05	-0.07	3.41 \pm 0.14	26.5	1.06 a	0.11	3.42 \pm 0.15	23.8	1.00	0.01
B	1	2.89 \pm 0.11	22.2	0.98	-0.07	3.22 \pm 0.14	27.3	1.05 a	0.11	3.35 \pm 0.16	26.6	1.07	0.07
C	1	2.82 \pm 0.11	22.8	0.98	-0.09	2.93 \pm 0.12	26.4	0.89 b	0.05	3.09 \pm 0.14	26.2	0.93	-0.03
P value			0.7635	0.5979	0.9583		0.7902	0.0017	0.3619		0.6132	0.1571	0.6504
A	2	4.01 \pm 0.12	16.5	1.03	-0.18	4.34 \pm 0.18	25.7	1.06 a	0.22	4.04 \pm 0.17	22.0	0.94 b	-0.06
B	2	3.88 \pm 0.12	17.3	1.01	-0.18	4.26 \pm 0.17	25.0	1.02 ab	0.19	4.00 \pm 0.21	25.8	1.13 a	0.00
C	2	3.61 \pm 0.11	18.9	0.96	-0.12	4.04 \pm 0.15	24.4	0.92 b	0.14	3.92 \pm 0.17	24.7	0.93 b	0.06
P value			0.4333	0.7373	0.7254		0.7598	0.0327	0.2924		0.4949	0.0080	0.7178
A	3	4.23 \pm 0.15	21.9	1.02	0.11	4.76 \pm 0.20	25.5	1.02	0.25	4.11 \pm 0.20	25.4 ab	0.99 ab	0.07
B	3	4.24 \pm 0.14	20.4	0.95	0.04	4.67 \pm 0.20	27.1	1.02	0.31	4.03 \pm 0.22	30.9 a	1.10 a	0.25
C	3	4.07 \pm 0.16	21.7	1.03	0.06	4.53 \pm 0.19	27.3	0.96	0.31	4.19 \pm 0.18	23.7 b	0.91 b	0.07
P value			0.6938	0.3454	0.6794		0.3775	0.3186	0.4277		0.0374	0.0340	0.1277

A, B, C, etc. refer to three different crop rotations (A, with grass-clover ley; B, with grass ley; and C, without ley) and four levels of N fertilization (0, 1, 2, and 3) with amounts of 0 kg ha⁻¹ year⁻¹, 38–55 kg ha⁻¹ year⁻¹, 73–100 kg ha⁻¹ year⁻¹, and 105–145 kg ha⁻¹ year⁻¹, respectively, and depending on crop (see Table S1). Means comprise seven crop rotation cycles for all sites except for the winter wheat crop at Stenstugu which only comprise six rotation cycles. For CV and FW b_i , within columns per N level, means followed by different letters are significantly different.




Fig. 2 Regression lines of individual cropping systems (A, with grass-clover ley; B, with grass ley; and C, without ley) over the overall cropping systems year means for oat at N level 0 (0 kg N

ha⁻¹ year⁻¹) for the three sites, **A** Lanna LTE, **B** Stenstugu LTE, and **C** Säby LTE. Comparisons between regression lines are reported in Table 4.

lower (indicating a higher stability) than position two, no significant difference was found with either the CV or the POLAR method. Equally, there was no difference in the response of the first and second oat to different environmental conditions, as shown by FW regression (Table 5).

The positive effects on soil fertility are more important to the crop directly following the ley due to carry-over and residual effects (Persson et al. 2008). According to the analysis of the Swedish LTEs by Bergkvist and Båth (2015), the positive effect of the leys preceding the oat did not increase continuously over time. In such systems, yield stability may be affected when the processes of the pre-crop are more dominant in particular years, then in others, but there was no difference in the response of the first and second oat to the environmental adaptability (Table 5).

3.4 Diversification through the integration of grain legumes

Durum wheat grown in the French cropping system with grain legumes (pea and faba bean) in the rotation (no inter-cropping) tended to yield higher especially in years with yields below average (< 5500 kg/ha) and had relatively low yields in high yielding years (> 6500 kg/ha) compared to durum wheat grown in a cropping system without legumes (Fig. 3). Indeed, the system with the grain legume had a slope < 1 indicating lower environmental adaptability than the system without the legume (Table 6). Yield stability was not significantly different between the two systems probably due to the limited dataset (CV 13% vs 18% and a POLAR of -0.35 vs 0.02) (Table 6).

Table 5 Mean grain yields of oat, CV, POLAR, and Finlay-Wilkinson (FW) in the Scottish LTE for rotations with different length and positions of the oat in the crop sequence.

System	Grain yield (Mg DM/ha \pm SE)	CV (%)	POLAR (coefficient)	FW b_1 (regression coefficient)
Length of the ley ¹				
3-year ley	5.26 \pm 0.21	14.1	− 0.20	0.95
4-year ley	5.47 \pm 0.23	14.2	− 0.17	1.02
Oat position ²				
First	4.83 \pm 0.24	21.4 a	0.07 a	0.99a
Second	3.22 \pm 0.24	37.1 a	0.36 a	1.01a
P value		0.052	0.3036	0.8809

¹Length of the ley: 3-year and 4-year ley and the oat at first oat position in the rotation (data from 1995 to 2006). No statistical analyses were performed for CV, POLAR, and FW

²Oat position: first and second oat position in crop rotations with 3-year ley at the Scottish LTE (data from 1994 to 2017). For the CV, POLAR, and FW columns, means followed by different letters are significantly different.

The positive effects of the grain legumes on the durum wheat yield (in lower yielding years) could be a result of the positive pre-crop effect of grain legumes (Angus et al. 2015; Preissel et al. 2015). This process encompasses the N and non-N-related preceding crop effects (Chalk 1998) that are difficult to separate empirically. While the “nitrogen effect” comprises the provision of N to the subsequent crops, the other benefits include the “break-crop effect” that occurs when a disease cycle is broken, benefits to soil organic matter and structure and phosphorus mobilization (Watson et al. 2017). These effects occur especially in cereal-dominated cropping systems such as the systems without legumes in

this study. There are meta-analyses on the pre-crop effect on mean yield, with cereal yields being 1.46 t ha^{−1} in temperate Europe (Preissel et al. 2015) and 1.2 t ha^{−1} in Australia, Europe, and North America (Angus et al. 2015) higher after grain legumes than after cereal pre-crops. This is one of the first studies exploring the pre-crop effect on yield stability.


3.5 Diversification through cover crops

Diversification with cover crops in the rotations had no significant effect on the mean yield of durum wheat in the French LTE (Table 6). Cover crops did not affect yield stability of durum wheat significantly (CV 14% vs 17% and POLAR − 0.26 vs − 0.07, Table 6). The FW coefficient in the system with cover crops was < 1, indicating low environmental adaptability, while it was > 1 in the system without cover crops (Fig. 3). However, the difference between the systems was not significant, maybe due to the few observations available for the analysis.

Cover crops are primarily grown to reduce nitrate leaching, increase soil organic carbon, or reduce soil erosion (Plaza-Bonilla et al. 2016). Although the effect of cover crops on yield stability has not been investigated so far, improved soil conditions could result in higher yields in bad years (indicated by our study) and improve yield stability in the long-term. Longer datasets are needed to draw robust conclusions related to the impact of cover cropping on yield stability.

3.6 Diverging impacts of diversification

Our results provide diverging indications for the effects of cropping system diversification on cereal yield stability,

**Fig. 3** Regression lines of individual cropping systems over the overall cropping systems year means for durum wheat at the French LTE for rotations with and without grain legumes (left) and rotations with

and without cover crops (right). Comparisons between regression lines are reported in Table 6.

Table 6 Mean grain yield of durum wheat, CV, POLAR, and Finlay-Wilkinson (FW) for the rotations with and without grain legumes and with and without cover crops in the French LTE.

Systems	Grain yield (Mg DM/ha \pm SE)	CV (%)	POLAR (coefficient)	FW b_i (regression coefficient)
Integration of grain legumes				
With grain legume	5.58 \pm 0.15	13.2 a	− 0.35 a	0.77 b
Without grain legume	5.43 \pm 0.20	17.8 a	0.02 a	1.23 a
<i>P</i> value		0.2295	0.1553	0.0216
Integration of cover crops				
With cover crop	5.53 \pm 0.16	13.8 a	− 0.26 a	0.88 a
Without cover crop	5.48 \pm 0.20	17.3 a	− 0.07 a	1.12 a
<i>P</i> value		0.3601	0.4306	0.1372

Data from 2005 to 2016. For the CV, POLAR, and FW columns, means followed by different letters are significantly different.

environmental adaptability, and the probability of diversified systems outperforming less diverse systems. St-Martin et al. (2017) have drawn a similar conclusion related to cereal yield stability and environmental adaptability when analyzing three cropping systems in a single LTE. In contrast Macholdt et al. (2020) found in another LTE that winter barley grown in cropping sequences dominated by cereals had lower yield stability and environmental adaptability and greater production risks compared with winter barley grown in cropping systems with higher crop diversity and additional organic matter inputs. Using seven long-term experiments across a wide latitudinal gradient in Europe, Marini et al. (2020) found that growing multiple crop species (including legumes) in a rotation always provided higher yields for both winter and spring cereals (average + 860 and + 390 kg ha^{−1} per year, respectively) compared with continuous cereal cropping. Yield gains in diverse rotations were especially higher in (low-yielding) years with high temperature and low precipitation. Similarly, we found that all diversification measures increased cereal yields in years with below average yields (including durum wheat in cropping systems with cover crops and grain legumes in France). In high-yielding years, only cropping systems with perennial legume crops consistently increased cereal yields. The inclusion of grain legumes tended to reduce yields of cereals in high yielding environments. While winter cereals yielded more in diverse rotations consistently across the period of the LTE, the yield gain increased over time since establishment of the LTE in spring cereals (Marini et al. 2020). Globally and also in Europe, cropping system diversification uses very different approaches (Hufnagel et al. 2020), which range from diversifying simple maize-based systems (Bowles et al. 2020) to systems with a large spatial and genetic diversity (Ditzler et al. 2021). This makes comparisons difficult.

4 Conclusion

We conclude that diversification affected the performance of cereals within cropping systems. While most but not all diversification measures increased the productivity of cereals in long-term experiments, the effects on yield stability and environmental adaptability were inconsistent. For the five diversification measures tested, we conclude (i) diversification through perennial grass and legume crop mixtures outperformed systems without leys across the three sites for wheat and oat. We found a higher environmental adaptability of the cropping systems with perennial leys, especially in low-input systems. (ii) Diversification through the length of the perennial ley increased the yield of oats after a longer period of the ley, but did not affect yield stability or environmental adaptability. (iii) Diversification through changing the position of oats in the crop sequence increased the yield by 33% when it followed the ley directly compared to the crop grown 2 years later. (iv) Diversification through the integration of grain legumes indicated increased yields of durum wheat in lower-yielding years compared to the system without the grain legume. (v) Diversification through cover crops did not affect yield stability in the studied LTE.

We conclude that the four methods used for analyzing the yield data, were a strength of our study, since it allowed detecting diverging evidence of the cropping system performance. While the probability method was providing the most conclusive results by combining aspects of productivity and stability, there is a risk of making unjustified conclusions about yield stability if using only one method. The existing dataset could be extended, i.e., adding more experiments with additional treatments to analyzing further effects of diversification.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s13593-022-00850-z>.

Acknowledgements The authors thank Jan Eric Englund (SLU) for valuable advice on statistical analyses. We also thank Birgitta Båth, Robin Walker, and staff of INRAE for providing yield data from the Swedish, Scottish, and French LTEs.

Authors' contributions Conceptualization and methodology: MR, JA, CFET, and AV; writing—original draft: MR and JA; writing—review and editing: MR, JA, CFET, ESJ, GC, CW, EJ, and GB; resources: JA, CFET, CW, AV, and EJ.

Funding Open Access funding enabled and organized by Projekt DEAL. The work was funded by the FACCE ERA-Net Plus project Climate-CAFE (grant PTJ-031A544 and BB/M018148/1), the SusCrop-ERA-NET project LegumeGap (grant 031B0807B), and the PRIMA project Biodiversify (grant 01DH20014). Tulloch, The Scottish LTE is supported by the Rural and Environment Science and Analytical Services Division of the Scottish Government (SRUC-D3-1). CFET and CW were funded by the Scottish Government RESAS Strategic Research Programme. MR was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—420661662.

Data availability The datasets used in this study were sourced from the Swedish University of Agricultural Sciences (SLU), Sweden, Scotland's Rural College (SRUC), United Kingdom, and Centre INRAE Occitanie-Toulouse, France, under license for the current study. The datasets are not publicly available but may be obtained from the authors upon reasonable request and with the permission of the institutions mentioned above.

Code availability The code may be obtained from the authors upon reasonable request.

Declarations

Conflict of interest The authors declare no competing interests.

Ethics approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Angus JF, Kirkegaard JA, Hunt JR, Ryan MH, Ohlander L, Peoples MB (2015) Break crops and rotations for wheat. *Crop Pasture Sci* 66(6):523–552. <https://doi.org/10.1071/CP14252>
- Bergez JE, Colbach N, Crespo O, Garcia F, Jeuffroy MH, Justes E, Loyce C, Munier-Jolain N, Sadok W (2010) Designing crop management systems by simulation. *Eur J Agron* 32(1):3–9. <https://doi.org/10.1016/j.eja.2009.06.001>
- Bergkvist G, Båth B (2015) Nitrogen fertiliser dose influence the effect of two year rotational leys with grass or clover/grass on other crops in the rotation. *Asp Appl Biol* 128:133–139
- Bowles TM, Mooshammer M, Socolar Y, Calderón F, Cavigelli MA, Culman SW, Deen W, Drury CF, Garcia y Garcia A, Gaudin ACM, Harkcom WS, Lehman RM, Osborne SL, Robertson GP, Salerno J, Schmer MR, Strock J, Grandy AS (2020) Long-term evidence shows that crop-rotation diversification increases agricultural resilience to adverse growing conditions in North America. *One Earth* 2(3):284–293. <https://doi.org/10.1016/j.oneear.2020.02.007>
- Chalk PM (1998) Dynamics of biologically fixed N in legume-cereal rotations: a review. *Aust J Agric Res* 49(3):303–316
- Ditzler L, Apeldoorn DFv, Schulte RPO, Tittone P, Rossing WAH (2021) Redefining the field to mobilize three-dimensional diversity and ecosystem services on the arable farm. *Eur J Agron* 122:126197. <https://doi.org/10.1016/j.eja.2020.126197>
- Döring TF, Reckling M (2018) Detecting global trends of cereal yield stability by adjusting the coefficient of variation. *Eur J Agron* 99:30–36. <https://doi.org/10.1016/j.eja.2018.06.007>
- Döring TF, Knapp S, Cohen JE (2015) Taylor's power law and the stability of crop yields. *Field Crops Res* 183:294–302. <https://doi.org/10.1016/j.fcr.2015.08.005>
- Finlay K, Wilkinson G (1963) The analysis of adaptation in a plant-breeding programme. *Aust J Agric Res* 14(6):742–754. <https://doi.org/10.1071/AR9630742>
- Hauggaard-Nielsen H, Mundus S, Jensen ES (2012) Grass-clover undersowing affects nitrogen dynamics in a grain legume cereal arable cropping system. *Field Crops Res* 136:23–31. <https://doi.org/10.1016/j.fcr.2012.07.001>
- Hufnagel J, Reckling M, Ewert F (2020) Diverse approaches to crop diversification in agricultural research. A Review. *Agron Sustain Dev* 40(2):14. <https://doi.org/10.1007/s13593-020-00617-4>
- Iannetta PPM, Young M, Bachinger J, Bergkvist G, Doltra J, Lopez-Bellido RJ, Monti M, Pappa VA, Reckling M, Topp CFE, Walker RL, Rees RM, Watson CA, James EK, Squire GR, Begg GS (2016) A comparative nitrogen balance and productivity analysis of legume and non-legume supported cropping systems: the potential role of biological nitrogen fixation. *Front Plant Sci* 7(1700). <https://doi.org/10.3389/fpls.2016.01700>
- Johnston AE, Poulton PR (2018) The importance of long-term experiments in agriculture: their management to ensure continued crop production and soil fertility; the Rothamsted experience. *Eur J Soil Sci* 69(1):113–125. <https://doi.org/10.1111/ejss.12521>
- Litell R, Milliken G, Stroup W, Wolfinger R, Schabenberger O (2006) SAS for mixed models. SAS Institute, Inc., Cary
- Liu K, Johnson EN, Blackshaw RE, Hossain Z, Gan Y (2019) Improving the productivity and stability of oilseed cropping systems through crop diversification. *Field Crops Res* 237:65–73. <https://doi.org/10.1016/j.fcr.2019.03.020>
- Lobell DB, Schlenker W, Costa-Roberts J (2011) Climate trends and global crop production since 1980. *Science* 333(6042):616–620. <https://doi.org/10.1126/science.1204531>
- Macholdt J, Styczen ME, Macdonald A, Piepho H-P, Honermeier B (2020) Long-term analysis from a cropping system perspective: yield stability, environmental adaptability, and production risk of winter barley. *Eur J Agron* 117:126056. <https://doi.org/10.1016/j.eja.2020.126056>
- Marini L, St-Martin A, Vico G, Baldoni G, Berti A, Blecharczyk A, Małacka-Jankowiak I, Morari F, Sawinska Z, Bommarco R (2020) Crop rotations sustain cereal yields under a changing climate. *Environ Res Lett* 15. <https://doi.org/10.1088/1748-9326/abc651>

- Persson T, Bergkvist G, Kätterer T (2008) Long-term effects of crop rotations with and without perennial leys on soil carbon stocks and grain yields of winter wheat. *Nutr Cycl Agroecosyst* 81(2):193–202. <https://doi.org/10.1007/s10705-007-9144-0>
- Piepho HP (1998) Methods for comparing the yield stability of cropping systems. *J Agron Crop Sci* 180(4):193–213. <https://doi.org/10.1111/j.1439-037X.1998.tb00526.x>
- Piepho H-P (1999) Stability analysis using the SAS System. *Agron J* 91(1):154–160. <https://doi.org/10.2134/agronj1999.00021962009100010024x>
- Plaza-Bonilla D, Nolot J-M, Passot S, Raffailac D, Justes E (2016) Grain legume-based rotations managed under conventional tillage need cover crops to mitigate soil organic matter losses. *Soil Tillage Res* 156:33–43. <https://doi.org/10.1016/j.still.2015.09.021>
- Preissel S, Reckling M, Schläfke N, Zander P (2015) Magnitude and farm-economic value of grain legume pre-crop benefits in Europe: a review. *Field Crops Res* 175:64–79. <https://doi.org/10.1016/j.fcr.2015.01.012>
- Ray DK, Gerber JS, MacDonald GK, West PC (2015) Climate variation explains a third of global crop yield variability. *Nat Commun* 6:5989. <https://doi.org/10.1038/ncomms6989>
- Reckling M, Döring TF, Bergkvist G, Stoddard FL, Watson CA, Seddig S, Chmielewski F-M, Bachinger J (2018) Grain legume yields are as stable as other spring crops in long-term experiments across northern Europe. *Agron Sustain Dev* 38(6):63. <https://doi.org/10.1007/s13593-018-0541-3>
- Reckling M, Ahrends H, Chen T-W, Eugster W, Hadasch S, Knapp S, Laidig F, Linstädter A, Macholdt J, Piepho H-P, Schiffrers K, Döring TF (2021) Methods of yield stability analysis in long-term field experiments. A review. *Agron Sustain Dev* 41(2):27. <https://doi.org/10.1007/s13593-021-00681-4>
- Sanford GR, Jackson RD, Booth EG, Hedtcke JL, Picasso V (2021) Perenniality and diversity drive output stability and resilience in a 26-year cropping systems experiment. *Field Crops Res* 263:108071. <https://doi.org/10.1016/j.fcr.2021.108071>
- St-Martin A, Vico G, Bergkvist G, Bommarco R (2017) Diverse cropping systems enhanced yield but did not improve yield stability in a 52-year long experiment. *Agric Ecosyst Environ* 247:337–342. <https://doi.org/10.1016/j.agee.2017.07.013>
- Tigchelaar M, Battisti DS, Naylor RL, Ray DK (2018) Future warming increases probability of globally synchronized maize production shocks. *Proc Natl Acad Sci U S A* 115(26):6644–6649. <https://doi.org/10.1073/pnas.1718031115>
- Watson C, Baddeley J, Edwards A, Rees R, Walker R, Topp C (2011) Influence of ley duration on the yield and quality of the subsequent cereal crop (spring oats) in an organically managed long-term crop rotation experiment. *Org Agric* 1(1):147–159. <https://doi.org/10.1007/s13165-011-0012-5>
- Watson C, Reckling M, Preissel S, Bachinger J, Bergkvist G, Kuhlman T, Lindström K, Nemecek T, Topp C, Vanhatalo A, Zander Z, Murphy-Bokern D, Stoddard F (2017) Grain legume production and use in European agricultural systems. *Adv Agron* 144(1):235–303. <https://doi.org/10.1016/bs.agron.2017.03.003>
- Zhao J, Chen J, Beillouin D, Lambers H, Yang Y, Smith P, Zeng Z, Olesen JE, Zang H (2022) Global systematic review with meta-analysis reveals yield advantage of legume-based rotations and its drivers. *Nat Commun* 13(1):4926. <https://doi.org/10.1038/s41467-022-32464-0>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.