



**HAL**  
open science

## Fate of isoxaflutole and its diketonitrile metabolite under conventional and conservation tillage in an irrigated continuous-maize field

Lionel Alletto, Yves Coquet, Christophe Labat

### ► To cite this version:

Lionel Alletto, Yves Coquet, Christophe Labat. Fate of isoxaflutole and its diketonitrile metabolite under conventional and conservation tillage in an irrigated continuous-maize field. Diffuse inputs into the groundwater: Monitoring - Modelling - Management. Agriculture and water management in the light of future challenges, Jan 2007, Graz, Austria. hal-03934035

**HAL Id: hal-03934035**

**<https://hal.inrae.fr/hal-03934035v1>**

Submitted on 11 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Fate of isoxaflutole and its diketonitrile metabolite under conventional and conservation tillage in an irrigated continuous-maize field

L. Alletto<sup>1,2</sup>, Y. Coquet<sup>2</sup> & C. Labat<sup>2</sup>



<sup>1</sup> **École d'ingénieurs de Purpan, Agronomy department, Toulouse, France.**

<sup>2</sup> **UMR INRA/INA PG Environment and Arable Crops, Thiverval-Grignon, France.**



## **Introduction**

Context of the study

The cropping systems

Objectives of the study

## **Materials and methods**

The experimental site: localisation and soil characteristics

Isoxaflutole (IFT) properties

Sampling procedure: soil

Sampling procedure: water

## **Results and discussion**

General data

Persistence of isoxaflutole

Water and herbicide leaching

## **Conclusion**

## Context

National context: generalization of water resources pollution by pesticides...

Economic context: Midi-Pyrénées region = 2<sup>nd</sup> region for maize production in France

Agronomic context: Typical maize production management in the region:

- 80 % of the production is in continuous maize with more than 60 % irrigated.
- Tillage usually included a mouldboard ploughing (30-cm depth) at the end of the winter
- Soils are unprotected during the inter-crop (from November to May)

Environmental context:

⇒ In the region, this system of production has generated several environmental problems (nitrate, atrazine)

⇒ Now atrazine is forbidden: there is a need for development of new strategies to control weeds in continuous maize systems



Context



Scientific



**Conventional tillage (CT)**

/

**Conservation tillage (MT)**

**vs. Pesticide ?**


→ Organic carbon content  $\nearrow$  surface → pesticides sorption  $\nearrow$  (Locke et al., 1997)  
Desorption: very few studies... tend to increase under MT (Ding et al., 2002)

→ Degradation: highly contrasted results !


→ Runoff and erosion: MT are efficient to reduce erosion, but runoff depends on climatic conditions (Fawcett et al. 1994)

→ Leaching: contrasted results but for no-tillage systems leaching of pesticides increases (Watts & Hall, 1996)

**There is a need to evaluate and/or design new cropping systems to both maintain weed control efficiency and limit environmental impacts**



**Conventional tillage**



**The cropping systems**

# The cropping systems: soil surface differences



+



**Conventional tillage (CT)**




+



**Conservation tillage (MT)**

# The cropping systems: subsurface differences

## Conventional tillage (CT)



(Manichon, 1982; Roger-Estrade et al. ,2004)

## Conservation tillage (MT)



→ Consequences on water dynamics and solutes transport



## Objectives of the study

Evaluate the effects of tillage practices (conventional vs. conservation tillage ; inter-crop with or without cover-crop) on:

1- Isoxaflutole degradation and formation of diketonitrile



2- Leaching potential of isoxaflutole and its diketonitrile metabolite








# The experimental site: localisation and soil characteristics



## Legend


-  Garden house
-  Instrumented soil profiles

-  CT – without cover-crop
-  CT – with cover-crop
-  MT – without cover-crop
-  MT – with cover-crop



- **Agricultural field of 15 ha**
- **Continuous maize production**
- **« Boulbènes » soils (Gleyic Luvisol)**
- **Irrigation with a centre pivot**
- **Conservation tillage (MT) on 3 ha since 2000**

The experimental site: localisation and soil characteristics




Main features of the Boulbènes soil types:

- Loamy soils with unstable structure
- Low organic matter level
- High sensitivity to crusting
- Hydromorphic profile

	Horizon	Prof.	pH	Clay	Silt	Sand	OC	CaCO <sub>3</sub>
		(m)				(g kg <sup>-1</sup> )		
<b>CT</b>	LA1	0-0.10	7.2	282	538	166	8.18	9
	LA2	0.10-0.28	7.2	279	560	144	8.91	12
	Btg1	0.28-0.55	7.5	394	489	108	3.64	18
<b>55-80cm</b>	Btg2	0.55-0.80	6.9	450	439	103	3.11	0
	<b>MT</b>	LA1	0-0.10	7.3	265	569	145	8.72
	"LA2"	0.10-0.28	7.2	276	559	149	8.35	9
	Btg1	0.28-0.55	7.2	387	476	123	4.72	13
	Btg2	0.55-0.80	6.7	447	402	144	4.39	0

## Isoxaflutole (IFT) properties

- Proherbicide, isoxazoles



- Annual grasses and broadleaves weeds, roots uptake

- Pre-emergence of maize (75 g ha<sup>-1</sup>)

- Inhibitor of the biosynthesis of carotenoids


- IFT: low solubility** in water (6.2 mg L<sup>-1</sup>), **rapidly degraded** ( $DT_{50}$  : 1.4-3 j), **good retention** on organic compounds ( $K_{OC}$  : 122 L kg<sup>-1</sup>)

- Degradation: formation of the **diketonitrile (DKN)** with a **higher solubility** (300 mg L<sup>-1</sup>), a **lower retention** ( $K_{OC}$  : 92 L kg<sup>-1</sup>) and a **higher persistence** ( $DT_{50}$  : 8-16 j)



## Sampling procedure: soil

- Before treatment: sampling of soil from surface to 80-cm depth to control initial concentration of IFT and DKN
- Treatment day: control of the variability of the treatment with fibreglass paper
- Sampling of soil from surface to a maximum of 30-cm depth
- Sampling time:  $t_{ini}$ ,  $t_0$ ,  $t_2$ ,  $t_3$ ,  $t_5$ ,  $t_7$ ,  $t_{11}$ ,  $t_{14}$ ,  $t_{21}$ ,  $t_{28}$
- Storage of frozen samples (-18°C) until analysis
- Analysis by HPLC-MS/MS



## Sampling procedure: water

- Ceramic cups: 4 by soil pit at 20 and 70 cm-depth
- Fibreglass wick lysimeters: 2 by soil pit at 40 cm (25x25 cm). Fibreglass wick length: 70 cm
- Sampling of soil water with ceramic cups and fibreglass wick lysimeters
- Storage of frozen samples (-18°C) until analysis
- Analysis by HPLC-MS/MS







Materials-methods





## Persistence of isoxaflutole




● : IFT } in soil samples  
○ : DKN } (in % of applied IFT)

1 IFT  $DT_{50}$  was short (<1 d)  
No detection of IFT 7 DAT

2 No effect of tillage on IFT  $DT_{50}$   
But... a longer persistence under  
MT with cover crop: Sorption ?

→ IFT hydrolysis was catalysed by retention on solid phase (Rice et al., 2004)  
But...  
→ Sorption of IFT on organic matter was found to increase  $DT_{50}$  (Rouchaud et al., 2002)

3 Increase of DKN with rainfall events





## Water and herbicide leaching

1 Cumulative water drainage  $\approx 7$  L under MT\_Bare soil and  $< 7$  L under MT\_Cover-crop

2 Cumulative leaching of DKN reached about 8 % of applied dose under MT\_Bare soil and  $< 2$  % under MT\_Cover-crop

→ Lower initial water content in the soil profile under cover-crop → increase sorption  
 → Quantity and/or nature of the residues *vs.* sorption and degradation of DKN...

■ Water drainage (mL)

● : DKN concentration in water (in % of applied IFT)

○ : Cumulative loss of herbicide

1 Cumulative water drainage  $\approx 14$  L under CT\_Bare soil & CT\_Cover-crop

2 Cumulative leaching of DKN reached about 15 % of applied dose at the end of the growing season

3 Maximum DKN concentration was  $32 \mu\text{g L}^{-1}$  64 DAT (but in a low volume)

- Tillage practices had no effect on the in-field degradation of isoxaflutole
- Residues on soil surface seemed to slow down degradation rate of IFT

*Effect of interception and retention ?*

- Migration in soil was faster and more important under conventional technique (data not shown)
- Water drainage was two times higher under conventional technique
- Herbicide leaching was between 2 and 7 times lower under conservation technique, with the lowest leaching under the cover-crop plot

*Retention processes and/or degradation were modified by tillage and residues management ?*

*Thank you.*

