



HAL
open science

Phylogenomics and a revised tribal classification of subfamily Dipterocharpoideae (Dipterocharpaceae)

Tijana Cvetković, Damien Hinsinger, Daniel Thomas, Jan Wieringa, Elango Velautham, Joeri Strijk

► **To cite this version:**

Tijana Cvetković, Damien Hinsinger, Daniel Thomas, Jan Wieringa, Elango Velautham, et al.. Phylogenomics and a revised tribal classification of subfamily Dipterocharpoideae (Dipterocharpaceae). *Taxon*, 2022, 71 (1), pp.85-102. 10.1002/tax.12648 . hal-03935018

HAL Id: hal-03935018

<https://hal.inrae.fr/hal-03935018>

Submitted on 31 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Phylogenomics and a revised tribal classification of subfamily Dipteroocarpoideae (Dipteroocarpaceae)

Tijana Cvetković,^{1,2}  Damien D. Hingsinger,^{3,4}  Daniel C. Thomas,⁵ Jan J. Wieringa,⁶  Elango Velautham⁵ & Joeri S. Strijk^{3,7} 

1 *Biology Centre CAS, Institute of Parasitology, Branišovská 1160/31, 37005 České Budějovice, Czech Republic*

2 *Biodiversity Genomics Team, Guangxi Key Laboratory of Forest Ecology and Conservation, Guangxi University, Daxuedonglu 100, Nanning, Guangxi, 530005, China*

3 *Alliance for Conservation Tree Genomics, Pha Tad Ke Botanical Garden, P.O. Box 959, 06000 Luang Prabang, Lao P.D.R.*

4 *Université Paris-Saclay, INRAE, Etude du Polymorphisme des Génomes Végétaux (EPGV), 91000 Evry, France*

5 *Singapore Botanic Gardens, 1 Cluny Road, Singapore, 259569, Singapore*

6 *Naturalis Biodiversity Center, Darwinweg 2, 2333 CR, Leiden, the Netherlands*

7 *Institute for Biodiversity and Environmental Research, Universiti Brunei Darussalam, Jalan Tungku Link, BE1410, Brunei Darussalam*

Address for correspondence: Joeri S. Strijk, jstrijk@actg.science

DOI <https://doi.org/10.1002/tax.12648>

Abstract Dipteroocarpoideae, the largest subfamily in the Meranti family (Dipteroocarpaceae) are an ecologically dominant group of trees throughout much of wet tropical Asia. Increasing anthropogenic pressures on this economically important tree family make it essential to resolve their complex evolutionary relationships and understand the distribution of genetic diversity throughout the family and distribution range. Dipteroocarpaceae have been the focal group in a wide range of studies, owing to their economic value, importance in historical biogeography and key role in the evolution of the Asian tropical forest biome. Despite this, persistent taxonomic and evolutionary questions remain, ranging from questions on the geographic origin, sequence of dispersal and the identification of diagnostic characters to circumscribe proper evolutionary groups. Here we present a comprehensive phylogenomic hypothesis for Dipteroocarpoideae, based on the analyses of plastome and nuclear cistron (NRC) data, and provide an in-depth review on the validity of morphological characters underlying the new tribal classification proposed here for the subfamily. Phylogenomic relationships were inferred using maximum likelihood and Bayesian approaches. Estimates of origin and onset of diversification in major clades and lineages were reconstructed using plastome, nuclear and combined datasets. Results of the separate and combined genomic datasets partly corroborate elements of previous classification systems (with improved support at all levels for major clades) but provide strong support for revising the tribal classification of the subfamily into four main clades: Dipteroocarpeae (*Dipterocarpus*), Dryobalanopseae (*Dryobalanops*), Shoreeae (*Hopea*, *Neobalanocarpus*, *Parashorea*, and all parts of a polyphyletic *Shorea*) and Vaterieae (including all other presently accepted Dipteroocarpoideae genera). Multi-fossil-dated divergence time estimation suggests Vaterieae first originated in the Late Cretaceous, followed by Dipteroocarpeae, with subsequent rise of the Dryobalanopseae and Shoreeae in the Eocene. Diversification of all tribes commenced before the Early Miocene. Our results provide strong support for the position of *Neobalanocarpus heimii*, *Parashorea* and (sub-)sections of the genera *Anisoptera*, *Hopea*, *Shorea* and *Vatica*. Hypotheses on the origin of *Neobalanocarpus heimii* by intergeneric hybridisation between *Anthoshorea* (maternally inherited) and *Hopea* (paternally inherited) species were corroborated. Finally, our study provides support for future revisionary changes: (1) the elevation to generic rank of sections in *Shorea*; and (2) revising the infrageneric classification of *Hopea* as all (sub-)sections were recovered as not monophyletic.

Keywords classification; Dipteroocarpaceae; Dipteroocarpoideae; genome skimming; historical diversification; molecular dating; nuclear ribosomal cistrons; phylogenomics; plastomes; tropical Asian rainforest

Supporting Information may be found online in the Supporting Information section at the end of the article.

INTRODUCTION

Dipteroocarpaceae includes nearly 700 species with their centre of diversity in Malesia (Symington, 1943; Ashton, 1982; Whitten & al., 1987; Appanah & Turnbull, 1998; Takhtajan, 2009). They are a major and often dominant structural com-

ponent of Asian tropical lowland forests (Ghazoul, 2016) and the most important native tree family for timber harvesting in tropical Asia (Tsumura & al., 2011; FAO, 2014; ITTO, 2019). Due to their ecological importance, economic value and increasing anthropogenic pressures on Dipteroocarpaceae species and their habitats (IUCN, 2020), studies contributing to

Article history: Received: 28 Jul 2020 | returned for (first) revision: 22 Oct 2020 | (last) revision received: 9 May 2021 | accepted: 25 May 2021 | published online: 7 Jan 2022 | **Associate Editor:** Eric H. Roalson | © 2022 The Authors.

TAXON published by John Wiley & Sons Ltd on behalf of International Association for Plant Taxonomy.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

our understanding of the phylogenetic relationships and the distribution of genetic diversity across the family's range are an essential part of the scientific foundation required to establish conservation and restoration priorities.

Dipterocarpaceae have traditionally been classified into three subfamilies: Dipterocarpoideae in Asia, Monotoideae in Africa and South America, and Pakaraimoideae in South America (Maguire & al., 1977; Ashton, 1982; Kostermans, 1985; Londoño & al., 1995; Morton, 1995; Maury-Lechon & Curtet, 1998). Recent molecular phylogenetic studies have suggested that the monospecific genus *Pakaraimaea* (Pakaraimoideae), previously classified in Dipterocarpaceae, may be more closely related to Cistaceae (e.g., Heckenhauer & al., 2017), and the genus was included in this family in APG IV (2016). Heckenhauer & al. (2017) pointed out that the position of *Pakaraimaea* among Cistaceae is not supported by its morphology and ecology (placed among Tiliaceae, close to *Schoutenia* by Kostermans, 1978 and Takhtajan, 1980), and their limited sampling was not sufficient to confirm its position with certainty. The phylogenetic position of Monotoideae is still unclear, and either Sarcolaenaceae (endemic to Madagascar) or Monotoideae have been proposed as sister to Dipterocarpoideae (Takhtajan, 1980, 2009; APG III, 2009; APG IV, 2016; Heckenhauer & al., 2017).

Various classifications have been proposed for the family (see Maury-Lechon & Curtet, 1998; Ghazoul, 2016; and Heckenhauer & al., 2017, for overviews). In his seminal revision of Malesian Dipterocarpaceae, Ashton (1982) recognized 13 genera in Asian Dipterocarpoideae and subdivided the subfamily into two tribes: Dipterocarpeae (*Anisoptera*, *Cotylelobium*, *Dipterocarpus*, *Stemonoporus*, *Upuna*, *Vateria*, *Vateriopsis*, *Vatica*), characterized by valvate sepals in fruits, solitary vessels, scattered resin canals, and $n = 11$; and Shoreeae (*Dryobalanops*, *Hopea*, *Neobalanocarpus*, *Parashorea*, *Shorea*) characterized by imbricate sepals, grouped vessels, resin canals in tangential bands, and $n = 7$ (Ashton, 1982; Cao & al., 2006; Gamage & al., 2006). Takhtajan (1980, 2009) proposed the classification into four tribes and 13 genera: Dipterocarpeae (*Anisoptera*, *Cotylelobium*, *Dipterocarpus*, *Stemonoporus*, *Vateria*, *Vateriopsis*, *Vatica*), Dryobalanopseae (*Dryobalanops*), Parashoreae (*Parashorea*), and Shoreeae (*Hopea*, *Neobalanocarpus*, *Shorea*, *Upuna*).

Most molecular phylogenetic studies aiming to resolve tribal as well as inter- and infrageneric relationships had taxonomically restricted sampling or employed only standardized low-variable plastome and nuclear DNA loci (Kajita & al., 1998; Kamiya & al., 1998, 2005; Dayanandan & al., 1999; Morton & al., 1999; Gamage & al., 2006; Tsumura & al., 1996, 2011; Yulita & al., 2005; Cao & al., 2006; Heckenhauer & al., 2017). Two recent phylogenomic studies used RADseq-derived SNP (single-nucleotide polymorphism) and plastome data to gain insight into phylogenetic relationships and floral evolution in Shoreeae (Heckenhauer & al., 2018, 2019). The results of these studies have provided valuable information on relationships but also highlighted incongruences in the deeper classification, still largely based on morphological characters.

At tribal level, the circumscriptions of the two tribes by Ashton (1982), Dipterocarpeae (*Anisoptera*, *Cotylelobium*, *Dipterocarpus*, *Stemonoporus*, *Upuna*, *Vateria*, *Vateriopsis*, *Vatica*) and Shoreeae (*Dryobalanops*, *Hopea*, *Neobalanocarpus*, *Parashorea*, *Shorea*), were largely corroborated, with the exception of the phylogenetic position of *Dipterocarpus*, which remained unclear and poorly supported in previous molecular phylogenetic studies (e.g., Tsumura & al., 1996; Kajita & al., 1998; Kamiya & al., 1998; Gamage & al., 2003, 2006; Indrioko & al., 2006; Heckenhauer & al., 2017). Morphological characters alone were shown to be unable to resolve the position of *Dryobalanops*, resulting in either inclusion in Shoreeae or as separate tribe (Heckenhauer & al., 2017).

Generic circumscription in Shoreeae has been problematic, and persisting uncertainty in the morphology-based taxonomy of the tribe has been hypothesized to either be the result of a considerable overlap of the morphospaces of the large genera *Shorea* and *Hopea*, intergeneric hybridization, or the presence of ancestral polymorphisms (Ashton, 1982; Murawski & al., 1994; Bawa, 1998; Kamiya & al., 2005; Cao & al., 2006). Molecular data including both cpDNA and RADseq-derived SNPs have corroborated previous studies and showed that *Shorea* sensu Ashton is paraphyletic and that the monophyletic genera *Parashorea* and *Hopea* are nested within (see Ashton, 1982; Heckenhauer & al., 2018, 2019).

The large genus *Shorea* (ca. 200 spp.: Ashton, 2004; ca. 360 spp.: Heckenhauer & al., 2018) is predominantly Malesian and has gone through a remarkable radiation in Borneo (136 species including 96 endemics reported by Ashton, 2004). Previous classifications have segregated *Shorea* into multiple genera based on timber colour, wood anatomy, and morphological features of the flower and embryo, as well as leaf epidermal characters: *Anthoshorea* (White Meranti), *Richetia* (Yellow Meranti) (Heim, 1892; Symington, 1943; Ashton, 1963, 1982; Meijer & Wood, 1964, 1976; Maury, 1978; Maury-Lechon, 1979a,b; Kostermans, 1983, 1984, 1992; Appanah & Turnbull, 1998; Maury-Lechon & Curtet, 1998), *Rubroshorea* (Red Meranti) and *Shorea* s.str. (Balau, Selangan Batu in Borneo). Symington (1943) classified the main genera *Shorea*, *Pentacme* and *Parashorea* into wood groups: Balau, Red, White, and Yellow Meranti. Maury (1978) and Kostermans (1984) introduced *Doona* and *Pentacme* besides the recognized genera *Anthoshorea*, *Shorea*, *Rubroshorea* and *Richetia*. Ashton (1982) recognized 11 sections in *Shorea*: *S.* sect. *Anthoshorea* (corresponding to White Meranti), sect. *Richetioides* (Yellow Meranti), sect. *Shorea*, sect. *Pentacme* and sect. *Neohopea* (Balau), and sect. *Brachypterae*, sect. *Doona*, sect. *Mutica*, sect. *Ovalis*, sect. *Pachycarpae* and sect. *Rubella* and (Red Meranti).

Geographic distribution of meranti groups varies from widespread (i.e., *Shorea* sect. *Shorea* and sect. *Anthoshorea* occur from Sri Lanka to Malesia) to locally endemic (i.e., Red Meranti sections are restricted to the biogeographic region of western Malesia, while *S.* sect. *Pachycarpae* and sect. *Rubella* are found only in Borneo and the Philippines) (Symington, 1943; Ashton, 1982; Kamiya & al., 1998, 2005). Heckenhauer

& al. (2018, 2019) concluded that the current infrageneric classification should be abandoned as most sections and subsections are non-monophyletic, and either support recognizing a single wide circumscription of *Shorea* (*Shorea* sensu Ashton) or to recognize *Anthoshorea*, *Doona*, *Richetia*, *Rubroshorea* and *Shorea* s.str. at generic level.

Infrageneric relationships in some other groups have also remained problematic, and Maury-Lechon & Curtet (1998) emphasized the mixed taxa of *Vatica* and *Cotylelobium* have remained poorly understood. *Sunaptea* is placed among *Vatica*, but morphological and anatomical characters in embryos, fruit-seeds and seedlings would suggest a close relationship with *Cotylelobium*.

Hybridization events may have contributed to problematic aspects of the current classification. Nuclear- and plastome-based phylogenies have indicated hard incongruence for the phylogenetic placement of *Parashorea* within *Shorea* (Heckenhauer & al., 2017, 2018, 2019), and a putative hybrid origin of the monotypic genus *Neobalanocarpus* (Shoreeae) has been hypothesized, based on conflicts between phylogenies derived from the nuclear *PgiC* gene and those derived from plastome fragments (Kamiya & al., 2005). However, *Neobalanocarpus* was not included in the recent studies by Heckenhauer & al. (2018, 2019), and the results based on limited DNA data need further corroboration by more extensive phylogenomic analyses.

To address some of these enduring conflicts between historically morphology-based classification systems and lacking genomic data, we employed previously released (Cvetković & al., 2017, 2019) and newly sequenced plastome and nuclear ribosomal cistron (NRC) data for 126 species of Dipterocarpaceae. Our main objectives were: (i) to test the monophyly of the proposed two tribes in Asian Dipterocarpaceae (Ashton, 1982): Dipterocarpeae (*Anisoptera*, *Cotylelobium*, *Dipterocarpus*, *Stemonoporus*, *Upuna*, *Vateria*, *Vateriopsis*, *Vatica*) and Shoreeae (*Dryobalanops*, *Hopea*, *Neobalanocarpus*, *Parashorea*, *Shorea*) and clarify the uncertain phylogenetic position of *Dipterocarpus* (Heckenhauer & al., 2017); (ii) to test previous hypotheses on the paraphyly of *Shorea* sensu Ashton and validity of proposed genus-level segregates (Heckenhauer & al., 2019); (iii) to test the monophyly of proposed subsections in *Vatica*; (iv) to test previous hypotheses of ancient hybridization events in the evolution of *Neobalanocarpus* (Kamiya & al., 2005) and *Parashorea* (Heckenhauer & al., 2019) and check for hard phylogenetic incongruence in other groups; (v) to gain insight into molecular divergence age estimates of the main clades and the tempo of diversification of Southeast Asian Dipterocarpaceae.

■ MATERIALS AND METHODS

Sampling. — Leaf material was collected during field work with collected materials frozen in liquid nitrogen or silica gel-dried. Herbarium material in the collections of Naturalis Biodiversity Center (L) were also sampled. Vouchers were

deposited in our herbarium (BGT, Brunei Darussalam), Singapore Botanic Gardens herbarium (SING) and Naturalis Biodiversity Center (L, WAG).

New plastome and NRC data was generated for 141 accessions, for 126 species in Dipterocarpaceae with a focus on Southeast Asian taxa (*Anisoptera* 4, *Cotylelobium* 2, *Dipterocarpus* 26, *Dryobalanops* 4, *Hopea* 23, *Neobalanocarpus* 1, *Parashorea* 4, *Shorea* 50, *Vatica* 13). In addition, new genomic data (both the plastomes and NRC sequences) for 23 species in two outgroup families, Malvaceae s.l. and Thymelaeaceae, was added (Cvetković & al., 2021). Finally, we retrieved plastome and ITS data from GenBank to further extend our dataset (Dipterocarpaceae: 6 species; outgroups: 18 species; see Appendix 1).

DNA extraction, sequencing, and phylogenomic analyses. — Total genomic DNA was extracted from frozen and silica-dried leaf material using the Plant Genomic DNA Kit (Tiangen Biotech, Beijing, China), following Hinsinger & Strijk (2015). The NEBNext Ultra II DNA Library Prep Kit (Ipswich, Massachusetts, U.S.A.) was used for construction of 350-bp paired-end libraries at Novogene (Beijing, China). Sequencing was performed on an Illumina HiSeq2500 platform (San Diego, California, U.S.A.) by Novogene (Beijing, China), with a read length of 2×150 bp.

Plastome and NRC assembly, annotation and phylogenomic analyses were performed following Cvetković & al. (2019), with DNA matrix assemblies constructed using ECuADOR v.2.0 (Armijos Carrion & al., 2020) (suppl. Appendices S1–S3). Model selection for molecular evolution was performed with ModelTest-NG v.0.1.5 (GTR+I+G4 chosen for plastome, confirmed by both the Akaike information criterion [AIC] and the corrected AIC [AICc]; GTR+I+G4 for combined plastome-NRC datasets, confirmed by the Bayesian information criterion [BIC], AIC and AICc; TIM2+I+G4 for NRC, confirmed by the BIC, AIC and AICc) (Darriba & al., 2020). Likelihood phylogenetic inference was performed using RAxML-NG v.0.9.0 (Kozlov & al., 2019) following Cvetković & al. (2019).

In addition to the likelihood analyses, we performed a coalescent method (ASTRAL) to analyse the rate variation and signal in the plastid protein-coding genes. Coding genes were extracted from assembled plastomes with Geneious R11 v.11.0.4 (<http://www.geneious.com>) (Kearse & al., 2012). Extracted genes were aligned individually using MAFFT v.7.475 with the FFT-NS-i algorithm and the “--adjustdirectionaccurately” option (suppl. Appendix S4). A maximum likelihood (ML) tree was then built for each gene with IQ-TREE v.1.6.12 (Chernomor & al., 2016), including 1000 replicates for both ultrafast bootstrap and SH-aLRT. Substitution model for each gene was automatically chosen by IQ-TREE, then used for tree building. Resulting ML trees for individual genes were then concatenated and analyzed using ASTRAL v.5.7.4 (Mirarab & al., 2014) with default parameters.

Fossil calibration and molecular divergence time estimation. — Molecular divergence age estimation was performed using four calibrations (Fig. 1). The crown age of the Malvales divergence from Brassicales (102.7 Ma; Magallón & al., 2015)




Fig 1. Chronogram of Diptero-carpoideae based on plastome and nuclear sequences (combined dataset) plus outgroups inferred by BEAST 2. Node ages (in Ma) shown at nodes, with the 95% highest posterior density intervals (HPD; blue bars). All nodes with posterior probability (PP) 1, except nodes indicated with blue circles (PP = 0.79–0.99) or grey circles (PP = 0.33–0.69). Overlay with revised tribal classification: A1: Vateriae; A2: Diptero-carpeae; A3: Dryobalanopseae; A4: Shoreae; A5: Doona + Anthoshorea + Neobalanocarpus + Hopea clade; A6: *Shorea* sect. *Doona*; A7: *S.* sect. *Anthoshorea*; A8: Richetioides + Parashorea + *Shorea* + Rubroshorea clade; A9: *S.* sect. *Richetioides*; A10: *S.* sect. *Shorea*; A11: *S.* sect. *Rubroshorea*. Fossils used in this study (red circles): I, the crown age of Malvales divergence from Brassicales (Magallón & al., 2015) (102.7 Ma); II, stem age for the ancestral node leading to Sterculioideae (Hernández-Gutiérrez & Magallón, 2019) (78.89 Mr); III, *Bombacacidites anne* (66–56 Ma) (Van Der Hammen, 1954); IV, *Malvaciphyllum macondicus* (61.6–56 Ma) (Carvalho & al., 2011). Geological time scale shown in millions of years.




Fig 1. Continued.

was set as a root prior with a normal distribution (95% CI: 100.9–104 Ma, mean 102.7 Ma; node I in Fig. 1). A lognormal prior (95% CI: 76.5–86.5 Ma, mean 80.9 Ma; node II in Fig. 1) was assigned to a stem age for the ancestral node leading to Sterculioideae (81.8 Ma; Hernández-Gutiérrez & Magallón, 2019). *Bombacacidites anne*, a pollen fossil from the Mid-

to Late Palaeocene found in Colombia, was assigned to the crown *Bombax* (66–56 Ma) (Van Der Hammen, 1954). To this node we applied a lognormal prior with a mean of 59.5 Ma (95% CI: 56.0–65.4 Ma; node III in Fig. 1). *Malvaciphyllum macondicus*, a leaf imprint found in Colombian mid- to late Palaeocene deposits was used for the crown eumalvaceae

(61.6–56 Ma) (Carvalho & al., 2011). For the node we used a lognormal prior with a mean of 58.6 Ma (95% CI: 56.0–63.1 Ma; node IV in Fig. 1).

We performed divergence time estimation using plastome, NRC and combined datasets (suppl. Appendices S5–S7) in BEAST v.2.6.1 (Drummond & Rambaut, 2007; Suchard & Rambaut, 2009) on the CIPRES Science Gateway (Miller & al., 2010). We used an uncorrelated relaxed clock with rates obtained from a lognormal distribution (UCLN) (Drummond & al., 2006) and the Yule model (Yule, 1925) as tree prior. The distribution of the two hyperparameters (uclMean.c: Mean = 10.0; uclStdev.c: Mean = 0.333) with an exponential distribution were set following Areces-Berazain & Ackerman (2016). For detailed calibration of the priors used in this study see suppl. Table S1.

Here, we have applied the same settings from Cvetković & al. (2021), with the following modifications: two independent MCMC (Drummond & al., 2002) runs were conducted for 515 million generations for the plastome dataset, 200 million generations for the NRC dataset and 470 million generations for the combined dataset, each with the first 80% of tree samples discarded as burn-in. Effective sample size values (ESS), assessed in Tracer v.1.7.1 (Rambaut & al., 2018), were over 150 for plastome, NRC and combined datasets.

■ RESULTS

Plastome and NRC size. — The alignment lengths for the plastome, NRC, and combined datasets were 203,064, 6459 and 209,523 bp, respectively. The reconstructed plastome lengths of Dipterocarpaceae species ranged from 118,091 bp (*Dipterocarpus sublamellatus* BGT4422) to 156,123 bp (*Parashorea smythiesii* BGT4413). The typical plastome organization was confirmed: approximately 85 kbps, 20 kbps and 24 kbps, and overall GC of 35%, 32% and 43% in LSC (large single copy), SSC (small single copy) regions and two IR (inverted repeats) copies, respectively. The NRC lengths ranged from 5769 bp (*Shorea smithiana* BGT4355) to 5832 bp (*Hopea polyalthioides* BGT4365 and *Hopea cf. polyalthioides* BGT4360). We presented in detail both the plastome and NRC characteristics of outgroups used elsewhere (Cvetković & al., 2021).

Phylogenomic analyses. — The best-scoring ML tree derived from the plastome data recovered three well-supported clades, representing the Malvales (sub-)families Thymelaeaceae, Malvaceae s.l. and Dipterocarpaceae subfamily Dipterocarpoideae (BS = 100; suppl. Figs. S1–S4).

In Dipterocarpoideae, four main clades were recovered, all with high bootstrap support (BS = 100): Vaterieae (clade A1), Dipterocarpeae (clade A2), Dryobalanopseae (clade A3) and Shoreeae (clade A4) (suppl. Figs. S1–S4).

(1) Clade Vaterieae – Genera *Anisoptera*, *Cotylelobium* and *Vatica* were all strongly supported and recovered as monophyletic (BS = 100; A1, suppl. Figs. S1–S4). *Anisoptera* was recovered as sister to the *Cotylelobium* + *Vatica* clade (BS = 100). The two sections present in our *Anisoptera* sampling

(*A. sect. Glabrae*, sect. *Anisoptera*) were supported as monophyletic (BS = 100; A1, suppl. Fig. S2). However, monophyly of *Vatica* sections was not supported with representatives of *V. sect. Sunapteae* distributed throughout the *Vatica* clade (BS = 100; A1, suppl. Figs. S1–S3).

(2) Clade Dipterocarpeae was retrieved as monophyletic (BS = 99; A2, suppl. Figs. S1–S4).

(3) Clade Dryobalanopseae was retrieved as monophyletic (BS = 100; A3, suppl. Figs. S1–S4).

(4) Clade Shoreeae: *Hopea*, *Neobalanocarpus*, *Parashorea* and *Shorea* were retrieved as the monophyletic tribe Shoreeae (A4, suppl. Figs. S1–S4). *Shorea* was retrieved as paraphyletic with *Hopea*, *Neobalanocarpus* and *Parashorea* nested within (suppl. Figs. S1–S4).

Clade A5 contains species assigned to *Doona* (A6), *Anthoshorea* (A7), *Neobalanocarpus* and *Hopea*; *Doona* (A6) is sister to an *Anthoshorea* + *Neobalanocarpus* + *Hopea* clade (BS = 100; A5, suppl. Figs. S1–S4). *Hopea* was retrieved as monophyletic (BS = 100); but neither of the two *Hopea* sections (sect. *Hopea*, sect. *Dryobalanoides*) was retrieved as monophyletic group (A5, suppl. Fig. S2). *Neobalanocarpus heimii* was recovered as sister to *Hopea* (BS = 100; A5, suppl. Figs. S1–S3).

Clade A8 is composed of species assigned to *Richetia* (A9), *Parashorea*, *Shorea* s.str. (A10), and *Rubroshorea* (A11); all of these clades are strongly supported. Most sections (*Shorea* sect. *Brachyptereae*, sect. *Mutica*, sect. *Pachycarpae*, sect. *Shorea*) within *Shorea* s.str. and *Rubroshorea* are not supported as monophyletic. *Parashorea* was retrieved as sister to a clade containing *Shorea* s.str. + *Rubroshorea* (BS = 100; A8, suppl. Figs. S1–S3).

Neobalanocarpus heimii was recovered as nested in *Hopea* using the NRC dataset (BS = 93; B10, suppl. Figs. S3, S4), which is incongruent with the position retrieved using the plastome data (i.e., as sister to *Hopea*).

The backbone of Shoreeae received moderate support (B4, BS = 80, suppl. Figs. S3, S4), but some strongly supported clades were recovered: *Richetia* (B5; BS = 100), *Shorea* s.str. (B7; BS = 100), *Rubroshorea* (B8; BS = 99), *Parashorea* (BS = 76), *Anthoshorea* (B9; BS = 100) and *Hopea* (B10; BS = 99). A sister relationship between *Anthoshorea* (including a part of *Shorea* sect. *Mutica*) and *Hopea* was weakly supported (BS = 56).

The RAxML tree constructed using our combined dataset was largely congruent with the topology recovered using the plastome data (suppl. Figs. S5, S6).

Moreover, the four main clades and the placement of genera in Dipterocarpoideae in the ML analyses were additionally supported by Bayesian analysis (for details about the posterior probability values, see Figs. 1 and 2, suppl. Fig. S7).

In addition, the phylogenomic relationships in the ML analyses were confirmed by the concatenated individual species tree inferred by a coalescent method (ASTRAL) (suppl. Fig. S8), but not by the concatenated individual gene tree (suppl. Fig. S9).




Fig 2. Chronograms of main sections in Diptercarpoideae based on plastome (A) and NRC (B) sequences plus outgroups inferred by BEAST 2. Node ages (in Ma) shown at nodes; 95% highest posterior density (HPD) intervals as black bars. All nodes with posterior probability (PP) 1, except nodes with blue circles (PP = 0.83–0.98) and grey circles (PP = 0.56–0.66). Overlay with revised tribal classification: A1: Vateriae; A2: Diptercarpeae; A3: Dryobalanopseae; A4: Shoreae; A5: Shoreae + Anthoshorea + Neobalanocarpus + Hopea clade; A6: Shoreae sect. *Doona*; A7: S. sect. *Anthoshorea*; A8: *Richettioides* + Parashorea + Rubroshorea clade; A9: S. sect. *Richettioides*; A10: S. sect. *Shorea*; A11: S. sect. *Shorea*; B1: Vateriae; B2: Diptercarpeae; B3: Dryobalanopseae; B4: Shoreae; B5: sect. *Richettioides*; B6: Shoreae + Rubroshorea + Parashorea + Anthoshorea + Neobalanocarpus + Hopea clade; B7: S. sect. *Shorea*; B8: S. sect. *Rubroshorea*; B9: S. sect. *Anthoshorea*; B10: Neobalanocarpus + Hopea clade. Geological time scale shown in millions of years.

Dating analyses. — An Early Upper Cretaceous origin for all three Malvacean families was retrieved using each of our three datasets (plastome, NRC, combined) [ca. 95 Ma for Malvaceae s.l.; Thymelaeaceae and Dipterocarpaceae, subfamily Dipterocarpoideae] (Figs. 1, 2, suppl. Fig. S7, suppl. Table S1).

Tribe Vateriae first originated in the Late Upper Cretaceous (ca. 62/76/52 Ma, plastome, NRC and combined datasets, respectively), followed by tribe Dipterocarpeae (ca. 56/71/47 Ma, as before), and tribes Dryobalanopseae and Shoreeae in the Eocene (ca. 48/62/41 Ma, as before). The longest period of relative stasis between origin and onset of diversification occurred in tribe Dryobalanopseae (ca. 30 Myr; plastome data), and tribe Dipterocarpeae (ca. 40–50 Myr; NRC and combined data). The shortest intervals were detected in tribe Shoreeae (ca. 4–7 Myr; all datasets). Plastome and NRC data show divergence of all generic precursors occurring as early as the Early Miocene. Onset of species diversification of all tribes was initiated before the Late Miocene (combined dataset, Fig. 1) or Early Miocene (plastome and NRC data, Fig. 2, suppl. Fig. S7).

Among genera in the tribe Vateriae, *Anisoptera* and *Vatica* diversified in the Late Oligocene and the Miocene. *Cotylelobium* diverged much later at the end stages of the Miocene (ca. 8.52–6.82 Ma for plastome and NRC datasets, respectively) or Early Pleistocene (ca. 2.23 Ma, combined data) (Figs. 1, 2, suppl. Fig. S7, suppl. Table S1). These estimates are similar to those recovered in tribe Shoreeae, where plastome data suggest a species diversification from the Late Oligocene–Early Miocene (27–21 Ma) (in *Shorea* s.str., *Rubroshorea*, *Anthoshorea*, *Parashorea*, and *Hopea*), except for *Richetioides* (Middle Miocene; 14.05 Ma). As before, nuclear data suggest slightly older age estimates (incl. *Richetioides*, 22–31 Ma; NRC data).

The position of *Neobalanocarpus* varied in our recovered phylogenetic trees, with divergence estimated to have started around 27.07 Ma (Oligocene; 23.34–34.69, plastome), 39.67 Ma (Late Eocene; 28.85–51.46, NRC) and 22.03 Ma (Late Oligocene; 17.76–26.21, combined data) (see Figs. 1, 2, suppl. Fig. S7).

■ DISCUSSION

A revised tribal classification for Dipterocarpoideae and phylogenetic affinities of *Dipterocarpus*. — Our analyses of plastome data resolved the backbone of the Dipterocarpoideae phylogenetic placement and clarified the previously unresolved positions of *Dipterocarpus* and *Dryobalanops* (Heckenhauer & al., 2017). The current tribal classifications of Dipterocarpoideae recognize either the two tribes Dipterocarpeae and Shoreeae (Ashton, 1982; Cao & al., 2006), or the four tribes Dipterocarpeae, Dryobalanopseae, Parashoreae, and Shoreeae (Takhtajan, 2009), which is not supported by our results. Tribe circumscription here is in partial agreement with that outlined by Brandis (1895), Ashton (1982), Kamiya

& al. (2005), Takhtajan (2009), and Heckenhauer & al. (2018, 2019). Our study recovers the following four strongly supported tribes, and we propose to restructure the tribal classification of Dipterocarpoideae accordingly:

Tribe **Dipterocarpeae** Rchb., Handb. Nat. Pfl.-Syst.: 304. 1837 – Type: *Dipterocarpus* C.F.Gaertn.

Emergent or canopy trees, columnar but hardly buttressed with untidy globose crowns; prominently lenticellate orange-brown massively flaky bark; large leaf buds; amplexicaul bud scales; furnished stipules with diverse species-defining indumenta; plicate venation resulting in corrugation of their coriaceous leaves; thickly geniculate and often long petioles with often complex rings of vascular bundles and resin canals; variously thickened calyx ornamentations (tubercules, simple or folded wings); large flowers bearing a tubular calyx united at base into a smooth, angled, tuberculate or flanged tube enclosing but free from the ovary; two aliform, valvate sepals all along their development; stamens (15–40) are elongate orange anthers and stout tapering connectival appendages; dispersed resin canals in the wood and the largest stamens are the unique characteristic for *Dipterocarpus*; chromosome number $n = 11$ (summarized in Heckenhauer & al., 2017). From India and Sri Lanka to SE Asia.

One genus: *Dipterocarpus* C.F.Gaertn.

Tribe **Dryobalanopseae** Baill., Hist. Pl. 4: 210, 213. 1873 – Type: *Dryobalanops* C.F.Gaertn.

Emergent or canopy; fibres are with bordered pits; scattered resin canals; solitary vessels; fruit sepal is thickened. The subvalvate sepals in fruit (imbricate at first, after only retaining some traces of imbrication) is a unique characteristic for *Dryobalanops*. Chromosome number $n = 7$. Native in W. Malesia.

Note: The name Dryobalanopseae was incorrect and superfluous when published, since it included *Dipterocarpus*, but it was validly published with the correct name on page 213 and the description on 210, and is available for a tribe containing *Dryobalanops* and not *Dipterocarpus*; see Art. 19 Note 3 of the ICN (Turland & al., 2018).

One genus: *Dryobalanops* C.F.Gaertn.

Tribe **Shoreeae** Miq., Fl. Ned. Ind. 1(2): 503. 1859 – Type: *Shorea* Roxb. ex C.F.Gaertn.

= [unranked] Hopeae Korth., Verh. Nat. Gesch. Ned. Bezitt., Bot.: 55. 1840.

– “Parashoreae” Takht., Flowering Pl.: 269. 2009, nom. nud.
– Based on *Parashorea* Kurz.

Emergent or canopy, understorey trees; resin canals in tangential bands; thickened sepal base; fruit sepals imbricate at the incassate-cupped base of the ripe fruit; 3 strata in pollen exine: absent tiloid structure of exine; T and Y columellae shape-type; always grouped vessels with cellular divisions; radial canal formation; 2 or 3 incassate bases of sepals (and accrescent sepals) in fruits; free bases of fruit sepals; frigid pericarp tissue; circular fruit equatorial section; embryo cotyledons “covering-piled”; inferior or median-inferior hypocotyl; bilobed seedling

cotyledons; 4 root-xylem poles; uni- to tri-lacunar cotyledonary vascular bundles; stomatal types in first leaves paracytic, or para-cyclocytic, or anomo-cyclocytic; elongate stomata, sunken in the epiderm; chromosome number $n = 7$ (summarized in Appanah & Turnbull, 1998). From India and Sri Lanka to Malaysia.

Eight genera: *Anthoshorea* Pierre (synonym of *Shorea*), *Doona* Thwaites (synonym of *Shorea*), *Hopea* Roxb., *Neobalanocarpus* P.S.Ashton, *Parashorea* Kurz, *Richetioides* F.Heim (synonym of *Shorea*), *Rubroshorea* (synonym of *Shorea*), *Shorea* Roxb. ex C.F.Gaertn.

Notes: Hopeae, described by Korthals (1839–1842), would have had priority over Shoreeae but was described without explicit rank, which is mandatory before 1887 (Art. 37.1–37.3 of the *ICN*, Turland & al., 2018).

Shoreeae has also been spelled as ‘Shoreae’. Given this name has a non-Latin base (named after Sir John Shore), but has been Latinized, where the stem is the single syllable Shore, it seems proper to follow the original spelling of Miquel, who also used ‘Shoreeae’. In case we consider this name to have alternative possible genitives, Art. 18.1 (via Art. 17.1) even obliges us to do so. In botanical Latin, double ee’s are often avoided, but many recent tribal names based on generic names ending on -ea have also regularly been formed including the double ee. Since *Parashorea* is derived from *Shorea*, a tribe named after it should be spelled with the double ee as well, but so far this name lacks a formal description and is not in alignment with our classification superfluous.

Tribe **Vaterieae** Miq., Fl. Ned. Ind. 1(2): 502. 1859 – Type: *Vateria* L. [see note].

= Anisoptereae Miq., Fl. Ned. Ind. 1(2): 500. 1859 – Type: *Anisoptera* Korth.

Emergent or canopy trees; pollen grains tricolpate and lack endexine; universal presence of intercellular resin canals; valvate base of sepals in calyx of ripe fruit (imbricate at first, after only retain some traces of imbrication); solitary vessels, scattered resin canals; pericarp thickenings; 2 strata in pollen exine; tilioid structure of exine; columellae shape-type V and U; solitary vessels with cellular divisions of canal formation oblique; 0 or 5 incrassate bases of sepals (and accrescent sepals) in fruits; fused bases of fruit sepals; rigid to soft pericarp tissue; circular to 3-symmetric fruit equatorial section; cotyledons in embryo neither covering nor piled; hypocotyl apical or median; seedling cotyledons entire; 6, 8 or 10 root-xylem poles; 3- to multilacunar cotyledonary vascular bundles; stomatal types in first leaves anisocytic or anisocytic; elongate stomata; and sunken in the epiderm (imbricate) or round and raised above the epiderm (valvate); chromosome number $n = 11$ (summarized in Appanah & Turnbull, 1998). From the Seychelles through India and Sri Lanka to SE Asia.

Note: Miquel (1859) seems to have taken up the name from Korthals (1839), who had described the unranked name Vaterieae for a group containing *Vateria* and *Retinodendron* Korth. Subsequently, Blume (1852) used Vaterieae, but as a subfamily name (“subord.”) that included both *Vateria* and

Vatica. Miquel was the first to explicitly use the term at tribal level, and provided a new description. Since in this work he only had to deal with some species of *Vatica* (*Vateria* does not occur in Indonesia), it is not immediately apparent here that it is based on *Vateria* instead of *Vatica*. Since Vaterieae is only the correct tribe name when it is based on the generic name *Vateria*, and it likely is a classification following previous works of Korthals and Blume, we consider *Vateria* the type of the tribe, and hence Vaterieae the correct spelling.

Seven genera: *Anisoptera* Korth., *Cotylelobium* Pierre, *Stemonoporus* Thwaites, *Upuna* Symington, *Vateria* L., *Vateriopsis* F.Heim, *Vatica* L.

Resolving power of genomic data and morphological traits, and taxonomic areas requiring additional investigation.

— In this revised setup, the chromosome number of $x = 11$ is considered a synapomorphy of Dipterocarpeae, and imbricate flowers as a synapomorphy of Shoreeae (Indrioko & al., 2006). Unique morphological characters of *Dipterocarpus* were summarized in Heckenbauer & al. (2017, 2018). *Stemonoporus*, *Upuna*, *Vateria*, and *Vateriopsis* were not included in this study and will need to be included in an expanded survey. Monophyly of *Stemonoporus* and *Vateriopsis* was confirmed by both morphology (Ashton, 1982) and molecular phylogenetics (Dayanandan & al., 1999; Gamage & al., 2003, 2006; Heckenbauer & al., 2017). However, phylogenetic placement of *Upuna* and *Vateria* remains unresolved (Heckenbauer & al., 2017), and additional genomic data is needed for resolving their placement.

New insights on a hybrid origin of *Neobalanocarpus*.

— Various affinities of *Neobalanocarpus heimii* have been suggested using morphological and anatomical data: *N. heimii* was hypothesized to be closely related to *Hopea* sect. *Hopea* on the basis of the inflorescence, fruit embryo and germination mode (Ashton, 1982; Kamiya & al., 1998; Yulita & al., 2005) and *Doona* based on wood anatomy (Parameswaran & Gotwald, 1979). *Neobalanocarpus heimii* shares morphological characters with both *Anthoshorea* and *Hopea* (urceolate corolla and stamens with an acicular connective appendage [Dayanandan & al., 1999], and a linear anther in the flower and sub-equal short woody fruit sepals [Kamiya & al., 2005]).

The placement of *Neobalanocarpus heimii* as sister to *Hopea* (plastome data, Fig. 2A, suppl. Figs. S3A, S4A, S7A, S8) is in concordance with Gamage & al. (2006) and Tsumura & al. (2011); in NRC-derived phylogenetic trees, *Neobalanocarpus* is nested within the *Hopea* clade (Fig. 2B, suppl. Figs. S3B, S4B, S7B), which is not only incongruent with the results of the plastome data, but also with the results from phylogenetic inference based on the nuclear *PigC* gene that indicates that the genus is nested within *Anthoshorea* (White Meranti; Kamiya & al., 2005). Kamiya & al. (2005) hypothesized this incongruence is a likely indicator of an ancient hybridization event involving ancestors of *Anthoshorea* as paternal progenitor and ancestors of the *Hopea* crown group as maternal progenitor. This hypothesis for a hybrid origin is also corroborated by an irregular behaviour during meiosis

in *Neobalanocarpus* (Jong & Lethbridge, 1967; Kamiya & al., 2005). The strong support for inclusion of *Neobalanocarpus* in *Hopea* (Fig. 2B, suppl. Figs. S3B, S4B, S7B) was unexpected and may indicate an additional level of complexity not previously recovered. As the NRC reads in our study are derived from the same read pool as the plastome reads, the latter indicating the expected relationship as sister to *Hopea*, lab artefacts such as sample mix-ups are unlikely. NRC copies can homogenize to either maternal or paternal parent after hybridization (concerted evolution; see Álvarez & Wendel, 2003; Nieto Feliner & Rosselló, 2007). The phylogenetic signal presented by the NRC data may be the result of an additional hybridization event with a species in *Hopea*. This would have occurred after the hybridization event between the *Anthoshorea* crown group species and the ancestor of the *Hopea* crown group that gave rise to *Neobalanocarpus*. Additional nuclear data is clearly required to further disentangle this complex pattern of reticulation.

Paraphyly of *Shorea* and validity of generic segregates of *Shorea*. — Paraphyly of *Shorea* as indicated in previous studies (Kamiya & al., 1998; Heckenhauer & al., 2018, 2019) is corroborated by our results: *Hopea*, *Neobalanocarpus* and *Parashorea* are clearly nested within *Shorea* sensu Ashton (1982).

Hopea is consistently retrieved as sister to *Anthoshorea* (Heckenhauer & al., 2018, 2019, this study), but there are some inconsistencies in the placement of *Parashorea*. Previous analyses of plastome data recovered *Parashorea* as sister to a *Shorea* s.str. + *Rubroshorea* clade, while RADseq-derived SNP data indicated a sister relationship to *Richetia* (Heckenhauer & al., 2019). Our results, with extended taxon sampling in the generic segregates of *Shorea*, corroborate the phylogenetic position in the plastome phylogenetic analysis; the backbone of tribe Shoreeae in the NRC data-derived phylogenetic tree was moderately supported. Heckenhauer & al. (2019) hypothesized that the incongruent placement in the plastome and nDNA phylogenetic analyses may indicate ancient hybridization, and this hypothesis remains plausible given the signals from the extended plastome and NRC data.

Sections of *Shorea* recognized by Maury (1978) and Kostermans (1984) including *S.* sect. *Anthoshorea*, sect. *Doona*, sect. *Richetia*, sect. *Rubroshorea*, and sect. *Shorea* [*Shorea* s.str.] are resolved as separate, well-supported clades in both whole-plastome and NRC phylogenetic analyses, while the monophyly of most subsections (*S.* subsect. *Brachyptereae*, subsect. *Mutica*, subsect. *Pachycarpae*, subsect. *Shorea*) within *S.* sect. *Shorea* s.str. and sect. *Rubroshorea* is not supported, in agreement with recent studies based on plastome and RADseq-derived SNP data by Heckenhauer & al. (2018, 2019). Accessions of species in the genus *Pentacme* (Maury, 1978; Maury-Lechon, 1979a,b) and *Shorea* sect. *Neohopea* and sect. *Rubella* (Ashton, 1977, 1980, 1982) were not included in this study, preventing us from clear conclusions about their placement and relationships within Shoreeae.

Sectional non-monophyly in *Hopea* and *Vatica*. — *Hopea* was retrieved as a monophyletic group in our study

(as in Yulita & al., 2005; Cao & al., 2006; Gamage & al., 2006; Tsumura & al., 2011), but all sections and subsections were non-monophyletic (but *H.* sect. *Dryobalanoides* was monophyletic in Yulita & al., 2005). High levels of morphological diversity within sections and subsections (e.g., variability in ovary, leaf, bark and floral characters; Ashton, 2004) further support this (Heckenhauer & al., 2018; this study).

Monophyly of the genera *Anisoptera*, *Cotylelobium* and *Vatica* among Vateriae was confirmed by our study (as in Kajita & al., 1998; Gamage & al., 2003, 2006; Indrioko & al., 2006). Two monophyletic sections in *Vatica* (Cao & al., 2006) were not retrieved here. In addition, two entries of *V.* sect. *Sunaptea* were placed among sect. *Vatica* in our study, resolving the previously doubtful position of this group (Maury-Lechon & Curtet, 1998).

Dating analyses. — Here we focused on species in subfamily Dipterocarpoideae and present in detail the origin and divergence of outgroups used elsewhere (Cvetković & al. 2021). Results obtained with our combined dataset (51.78 [45.91–55.77] Ma) partly confirm results from Heckenhauer & al. (2017: 54.9 Ma [39.3–71.6 Ma]) but provide improved phylogenetic resolution. Major clades in Heckenhauer & al. (2017) showed wider range age estimates. A key difference is the position of *Dipterocarpus* forming a monophyletic clade with the rest of species belonging to Vateriae, in contrast to all our analyses that recovered four tribes in Dipterocarpoideae, including a well-supported Dipterocarpeae. The placement of *Neobalanocarpus* in their study is compatible with our plastome dataset; however, our NRC-based results present an additional previously undetected hybridization event with species in *Hopea*. We agree with Heckenhauer & al. (2017) that calibration remains difficult in the group, despite the large numbers of reported fossils for the family (see discussion further below).

Dipterocarpaceae are an ancient group that evolved slowly and dispersed widely. Evidence for significant increases in speciation rates is absent (Strijk & al., unpub. data) and most extant species are relatively young. Despite many studies in this family, little is known about how this pace of speciation has affected the distribution of genomic diversity or even when this has taken place exactly. A Gondwanan origin is still the more widely accepted theory (i.e., allopatrically evolved from Monotoideae on Deccan and Mascarene Plates) (Ducousso & al., 2004; Dutta & al., 2009; Rust & al., 2010; Beimforde & al., 2011; Shukla & al., 2013; Ghazoul, 2016; Kooyman & al., 2019), but remains contentious. Fossils have been reported from E Africa (Bancroft, 1935), Alaska (Wolfe, 1977) and S and SE Asia (discussed below). Fossils found in the Neogene sediments of India and Nepal (e.g., Prasad, 1990, 1994; Prakash & al., 1994; Prasad & Awasthi, 1996; Khan & Bera, 2010) are absent from the Palaeogene sediments (Bande & Prakash, 1986; Awasthi & Srivastava, 1990; Khan & Bera, 2010; Srivastava & Mehrotra, 2010; Shukla & al., 2013; Kooyman & al., 2019). The Palaeogene fossils in SE Asia (Muller, 1981; Dutta & al., 2011) are presented by several studies to support a SE Asian origin of Dipterocarpaceae

(Lakhanpal, 1970, 1974; Awasthi, 1996; Sasaki, 2006; Shukla & al., 2012, 2013).

The Indian plate (with Madagascar) separated from E Africa 158–160 Ma (Middle Jurassic) moving 15–25 cm/yr (Briggs, 2003; Rust & al., 2010; Ghazoul, 2016). Initial colliding with Eurasia commenced in the early Cenozoic (55–65 Ma; with final suturing 42–55 Ma) (Briggs, 2003), enabling biotic interchange between endemics from both sides (Rust & al., 2010). Nowadays, SE Asia is recognized as having one of the world's most complex and least understood geological histories (Sun & al., 2000). In the Early Palaeogene, the Indian subcontinent witnessed enormous floral diversification as a result of a global rise in temperature and changes in latitudinal position (Zachos & al., 2001; Rana & al., 2004; Sahni & al., 2006; Garg & al., 2008; Shukla & al., 2013; Paul & al., 2015). Modern-type, broad-leaf tropical Dipterocarpaceae forests were spread across the Indian subcontinent from the Eocene (52 Ma) (Van Aarssen & al., 1994; Anderson & Muntean, 2000; Dutta & al., 2009; Mallick & al., 2009; Rust & al., 2010; Rudra & al., 2014; Paul & al., 2015). Our results suggest an early Eocene diversification for Dipterocarpoideae (51.78 Ma), and the origin of all its main tribes during the Middle Eocene (ca. 41–52 Ma). This is remarkable since the origin of the clade has been hypothesized to be of Cretaceous origin (ca. 96 Ma), with a lag in initial divergence for c. 40 Ma (Eocene diversification for Dipterocarpoideae, 51.78 Ma). In contrast, the Legumes have been recently assessed as originating and diversifying around the Cretaceous-Palaeogene Mass extinction event (Vanneste & al., 2014; Koenen & al., 2021), a window in time that the Dipterocarps seem to have passed through without any evidence of major diversification.

Arid climate, uplifting of the Himalaya-Tibetan plateau, monsoon initiation and intensification, and Northern Hemisphere glaciations are thought to have caused complete extirpation of the family in most of parts in the Indian subcontinent during the Cenozoic (Milanković, 1941; Quade & al., 1989; Ruddiman & Kutzbach, 1989; Kroon & al., 1991; Molnar & al., 1993; Derry & France-Lanord, 1996; Ramstein & al., 1997; Dynesius & Jansson, 2000; An & al., 2001; Zheng & al., 2004; Cliff & al., 2008; Boos & Kuang, 2010; Shukla & al., 2013; Rudra & al., 2014). In this study, all main sections of Dipterocarpoideae diversified during the Neogene – probably due to irregular annual and mass flowering effects as a result of extended periods of drought (Heckenhauer & al., 2017, 2018). The increased drought during the Neogene is considered to be one of the driving forces behind adaptation and diversification of the family in SE Asia.

Dipterocarpaceae fossils suggest a floristic link between Chinese and Indian palaeofloras and the presence of massive vegetation exchanges between Indian Gondwana and Laurasia (Sun & Wang, 2005; Jacques & al., 2011, 2013, 2015; Feng & al., 2013). Members of the family became adapted to a seasonal climate from the Late Eocene to the Early Miocene (Wolfe, 1994a,b; Zachos & al., 2001; Ho & al., 2003; Mosbrugger & al., 2005; Shi & Li, 2010; Shi & al., 2014a,b; Jacques & al., 2015), and gradually disappeared from SE China

from the Late Miocene (Feng & al., 2013; Jiang & al., 2013; Huang & al., 2016; Liu & Quan, 2017). The family was almost or completely absent from mainland China during the last glacial maximum (LGM) (Ni & al., 2010; Jacques & al., 2015).

In contrast, the SW Chinese Miocene palaeofloras have a different floristic composition without clear Indian affinities (Jiang & al., 2013; Jacques & al., 2015; Huang & al., 2016; Liu & Quan, 2017; Cvetković & al., 2019). The migration route is thought to be through SE Asia, linking India with SE China as opposed to SW China (Jacques & al., 2015). Yunnan has also been affected by a period of aridification since the Miocene (Xia & al., 2009; Jacques & al., 2011; Sun & al., 2011; Xing & al., 2012; Jiang & al., 2013; Huang & al., 2016; Liu & Quan, 2017). The existence and position of the “savanna corridor” across Sundaland during the Neogene and Quaternary, and in which way it affected diversification of the family, remain controversial to this day (Morley & Flenley, 1987; Heaney, 1991; Morley, 2000, 2012; Sun & al., 2000; Gathorne-Hardy & al., 2002; Meijaard, 2003; Bird & al., 2005; Cannon & al., 2009; Wurster & al., 2010; Slik & al., 2011; Iwanaga & al., 2012; Kamiya & al., 2012; Ohtani & al., 2013; Raes & al., 2014). The absence of migration pathways and dispersal capabilities could explain the patterns of narrow endemism in Dipterocarpaceae (Ashton, 1982), but the historical interplay between dispersal and local extirpation, species ecology and adaptive radiations in the family remain less well understood (Kooyman & al., 2019).

■ CONCLUSIONS

Our plastome and NRC datasets confirm some results from previous studies, but also provide novel insights into the tribal classification of Dipterocarpoideae and present strong support for a new tribal classification for the group. Our data resolves the poorly understood phylogenetic relationships of *Dipterocarpus*, establishes non-monophyly of sections in *Hopea*, *Shorea* and *Vatica*, and re-assesses hybridization of *Neobalanocarpus* and *Parashorea*, revealing a previously undetected event. In our study we have focused on the use of a single (extended) nuclear region but in order to fully corroborate this signal, more extensive data, especially from the nuclear genome, is needed. This will aid in ruling out other potential explanations such as incomplete lineage sorting, chloroplast capture and other organism-level processes that can cause phylogenomic discordance (Spooner & al., 2020).

Parashorea is closely related to *Shorea*, and *Neobalanocarpus heimii*'s interspecific hybridisation (Kamiya & al., 2005; Gamage & al., 2006) of *Anthoshorea* (maternally inherited) and *Hopea* (paternally inherited) is supported by its placement between the *Anthoshorea* and *Hopea* clades in our plastome-based results, and in the *Hopea* clade using NRC data.

We propose here the recognition of four tribes in Dipterocarpoideae: Vateriae (including all genera of presently accepted Dipterocarpaceae except for *Dipterocarpus*: *Anisoptera*, *Cotylelobium*, *Stemonoporus*, *Upuna*, *Vateria*, *Vateriopsis*,

Vatica), Dipterocarpeae (*Dipterocarpus*), Dryobalanopseae (*Dryobalanops*), and Shoreae (*Hopea*, *Neobalanocarpus*, *Parashorea*, *Shorea* s.str. and several generic segregates of *Shorea* s.l. including *Anthoshorea*, *Doona*, *Richetioides*, and *Rubroshorea*).

Our molecular results are consistent with previous hypotheses that several *Shorea* sections including *S.* sect. *Anthoshorea*, sect. *Doona*, sect. *Richetia*, sect. *Rubroshorea* and sect. *Shorea* are distinct and could be elevated to the genus rank (Maury, 1978; Maury-Lechon, 1979a,b; Heckenhauer & al., 2018, 2019); this could also resolve the paraphyly of *Shorea* sensu Ashton (1982) in which the genera *Hopea*, *Neobalanocarpus* and *Parashorea* are nested. Most *Shorea* subsections were shown to be non-monophyletic, indicating that the infra-generic classification at this rank needs to be revised.

■ AUTHOR CONTRIBUTIONS

TC, DDH and DCT performed the experiments and data analyses; TC, DDH, DCT, JJW, EV and JSS collected samples; JSS designed the experiments and provided the funding. All authors contributed to editing, writing and reviewing the manuscript, and approved the final version for submission. The authors alone are responsible for the content and writing of the paper. — TC, <https://orcid.org/0000-0002-5981-0134>; DDH, <https://orcid.org/0000-0001-7459-7610>; JJW, <https://orcid.org/0000-0003-0566-372X>; JSS, <https://orcid.org/0000-0003-1109-7015>

■ ACKNOWLEDGEMENTS

We thank the horticultural staff of Singapore Botanic Gardens, Xishuangbanna Tropical Botanical Garden and Naturalis Biodiversity Center (L, WAG) for their kind assistance in our sampling efforts. This work was supported by the China Scholarship Council (2016GXZS80 to TC), grants from Guangxi University and funding through the Bagui Scholarship team funding (C33600992001 to JSS) and China Postdoctoral Science Foundation Grants (2015M582481 and 2016T90822 to DDH).

■ LITERATURE CITED

- Álvarez, I. & Wendel, J.F. 2003. Ribosomal ITS sequences and plant phylogenetic inference. *Molec. Phylogen. Evol.* 29: 417–434. [https://doi.org/10.1016/s1055-7903\(03\)00208-2](https://doi.org/10.1016/s1055-7903(03)00208-2)
- An, Z.S., Kutzbach, J.E., Prell, W.L. & Porter, S.C. 2001. Evolution of Asian monsoons and phased uplift of the Himalaya–Tibetan plateau since Late Miocene times. *Nature* 411: 62–66. <https://doi.org/10.1038/35075035>
- Anderson, K.B. & Muntean, J.V. 2000. The nature and fate of natural resins in the geosphere. Part X. Structural characteristics of the macromolecular constituents of modern Dammar resin and Class II ambers. *Geochem. Trans.* 1, 1. <https://doi.org/10.1186/1467-4866-1-1>
- APG III 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* 161: 105–121. <https://doi.org/10.1111/j.1095-8339.2009.00996.x>
- APG IV 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot. J. Linn. Soc.* 181: 1–20. <https://doi.org/10.1111/boj.12385>
- Appanah, S. & Turnbull, J.M. 1998. *A review of dipterocarps: Taxonomy, ecology and silviculture*. Bogor: Center for International Forestry Research.
- Areces-Berzain, F. & Ackerman, J.D. 2016. Phylogenetics, delimitation and historical biogeography of the pantropical tree genus *Thespesia* (Malvaceae, Gossypieae). *Bot. J. Linn. Soc.* 181: 171–198. <https://doi.org/10.1111/boj.12414>
- Armijos Carrion, A.D., Hinsinger, D.D. & Strijk, J.S. 2020. ECuADOR—Easy Curation of Angiosperm Duplicated Organellar Regions, a tool for cleaning and curating plastomes assembled from next generation sequencing pipelines. *PeerJ* 8: e8699 <https://doi.org/10.7717/peerj.8699>
- Ashton, P.S. 1963. Taxonomic notes on Bornean Dipterocarpaceae. *Gard. Bull. Singapore* 20: 229–284.
- Ashton, P.S. 1977. Dipterocarpaceae. Pp. 166–196 in: Abeywickrama, B.A. & Dassanayake, M.D. (eds.), *A revised handbook to the flora of Ceylon*, vol. 1(2): Boca Raton: CRC Press.
- Ashton, P.S. 1980. Dipterocarpaceae. Pp. 364–423 in: Dassanayake, M.D. & Fosberg, F.R. (eds.), *A revised handbook to the Flora of Ceylon*, vol. 1. Rotterdam: Balkema.
- Ashton, P.S. 1982. *Flora Malesiana*, ser. I, *Spermatophyta, flowering plants*, vol. 9(2), *Dipterocarpaceae*. The Hague, etc.: Martinus Nijhoff. <https://doi.org/10.5962/bhl.title.40744>
- Ashton, P.S. 2004. Dipterocarpaceae. Pp. 63–388 in: Soepadmo, E., Saw, L.G. & Chung, R.C.K. (eds.), *Tree flora of Sabah and Sarawak*. Kuala Lumpur: Government of Malaysia.
- Awasthi, N. 1996. *Dipterocarpus* in the Indian subcontinent: Past, present and future. Pp. 138–156 in: Appanah, S. & Khoo, K.C. (eds.), *Proceedings of fifth round-table conference on dipterocarps*, Chiang Mai, 7–10 November 1994. Kepong: Forest Research Institute.
- Awasthi, N. & Srivastava, R. 1990. Some new carbonized woods from the Neogene of Kerala coast and their bearing on Palaeoclimate. *Palaeobotanist* 38: 285–292.
- Baillon, H. 1873. *Histoire des plantes*, vol. 4. Paris: Librairie Hachette. <https://doi.org/10.5962/bhl.title.40796>
- Bancroft, H. 1935. Some fossil dicotyledonous wood from Mount Elgon, East Africa. I. *Amer. J. Bot.* 22: 164–183. <https://doi.org/10.2307/2436007>
- Bande, M.B. & Prakash, U. 1986. The Tertiary flora of Southeast Asia with remarks on its paleoenvironment and phytogeography of the Indo-Malayan region. *Rev. Palaeobot. Palynol.* 49: 203–233.
- Bawa, K.S. 1998. Conservation of genetic resources in the Dipterocarpaceae. Pp. 45–55 in: Appanah, S. & Turnbull, J.M. (eds.), *A review of dipterocarps: Taxonomy, ecology and silviculture*. Bogor: Center for International Forestry Research.
- Beimforde, C., Schäfer, N., Dörfelt, H., Nascimbene, P.C., Singh, H., Heinrichs, J., Reitner, J., Rana, R.S. & Schmidt, A.R. 2011. Ectomycorrhizas from a Lower Eocene angiosperm forest. *New Phytol.* 192: 988–996. <https://doi.org/10.1111/j.1469-8137.2011.03868.x>
- Bird, M.I., Taylor, D. & Hunt, C. 2005. Palaeoenvironments of insular Southeast Asia during the last glacial period: A savanna corridor in Sundaland? *Quatern. Sci. Rev.* 24: 2228–2242. <https://doi.org/10.1016/j.quascirev.2005.04.004>
- Blume, C.L. 1852. *Museum botanicum Lugduno-Batavum; sive, Stirpium exoticarum novarum vel minus cognitarum ex vivis aut siccis brevis expositio et descriptio*, vol. 2. Lugduni-Batavorum [Leiden]: impensis auctoris.
- Boos, W.R. & Kuang, Z. 2010. Dominant control of the south Asian monsoon by orographic insulation versus plateau heating. *Nature* 463: 218–222. <https://doi.org/10.1038/nature08707>
- Brandis, D. 1895. An enumeration of the Dipterocarpaceae, based chiefly upon the specimens preserved at the Royal Herbarium and Museum, Kew, and the British Museum; with Remarks on the genera and species. *J. Linn. Soc., Bot.* 31: 1–148. <https://doi.org/10.1111/j.1095-8339.1895.tb00803.x>

- Briggs, J.C.** 2003. The biogeographic and tectonic history of India. *J. Biogeogr.* 30: 381–388. <https://doi.org/10.1046/j.1365-2699.2003.00809.x>
- Cannon, C.H., Morley, R.J. & Bush, A.B.G.** 2009. The current refugial rainforests of Sundaland are unrepresentative of their biogeographic past and highly vulnerable to disturbance. *Proc. Natl. Acad. Sci. U.S.A.* 106: 11188–11193. <https://doi.org/10.1073/pnas.0809865106>
- Cao, C.P., Gailing, O., Siregar, I., Indrioko, S. & Finkeldey, R.** 2006. Genetic variation at AFLPs for the Dipterocarpaceae and its relation to molecular phylogenies and taxonomic subdivisions. *J. Pl. Res.* 119: 553–558. <https://doi.org/10.1007/s10265-006-0005-8>
- Carvalho, M.R., Herrera, F.A., Jaramillo, C.A., Wing, S.L. & Callejas, R.** 2011. Paleocene Malvaceae from northern South America and their biogeographical implications. *Amer. J. Bot.* 98: 1337–1355. <https://doi.org/10.3732/ajb.1000539>
- Chernomor, O., Haeseler, A. von & Minh, B.Q.** 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Syst. Biol.* 65: 997–1008. <https://doi.org/10.1093/sysbio/syw037>
- Clift, P.B., Hodges, K.V., Heslop, D., Hannigan, R., Long, H.V. & Calves, G.** 2008. Correlation of Himalayan exhumation rates and Asian monsoon intensity. *Nature, Geosci.* 1: 875–880. <https://doi.org/10.1038/ngeo351>
- Cvetković, T., Hinsinger, D.D. & Strijk, J.S.** 2017. The first complete chloroplast sequence of a major tropical timber tree in the Meranti family: *Vatica odorata* (Dipterocarpaceae), *Mitochondrial DNA, B*, 2(1): 52–53. <https://doi.org/10.1080/23802359.2016.1275837>
- Cvetković, T., Hinsinger, D.D. & Strijk, J.S.** 2019. Exploring evolution and diversity of Chinese Dipterocarpaceae using next-generation sequencing. *Sci. Rep.* 9: 11639. <https://doi.org/10.1038/s41598-019-48240-y>
- Cvetković, T., Areces-Berazain, F., Hinsinger, D.D., Thomas, D.C., Wieringa, J.J., Ganesan, S.K. & Strijk, J.S.** 2021. Phylogenomics resolves deep subfamilial relationships in Malvaceae s.l. *G3 Genes, Genomes, Genetics* 11(7), jkab136. <https://doi.org/10.1093/g3journal/jkab136>
- Darriba, D., Posada, D., Kozlov, A.M., Stamatakis, A., Morel, B. & Flouri, T.** 2020. ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Molec. Biol. Evol.* 37: 291–294. <https://doi.org/10.1093/molbev/msz189>
- Dayanandan, S., Ashton, P.S., Williams, S.M. & Primack, R.B.** 1999. Phylogeny of the tropical tree family Dipterocarpaceae based on nucleotide sequences of the chloroplast RBCL gene. *Amer. J. Bot.* 86: 1182–1190. <https://doi.org/10.2307/2656982>
- Derry, L.A. & France-Lanord, C.** 1996. Neogene Himalayan weathering history and river $^{87}\text{Sr}/^{86}\text{Sr}$: Impact on the marine Sr record. *Earth Planet. Sci. Lett.* 142: 59–74. [https://doi.org/10.1016/0012-821X\(96\)00091-X](https://doi.org/10.1016/0012-821X(96)00091-X)
- Drummond, A.J. & Rambaut, A.** 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *B. M. C. Evol. Biol.* 7: 214. <https://doi.org/10.1186/1471-2148-7-214>
- Drummond, A.J., Nicholls, G.K., Rodrigo, A.G. & Solomon, W.** 2002. Estimating mutation parameters, population history and genealogy simultaneously from temporally spaced sequence data. *Genetics* 161: 1307–1320. <https://doi.org/10.1093/genetics/161.3.1307>
- Drummond, A.J., Ho, S.Y.W., Phillips, M.J. & Rambaut, A.** 2006. Relaxed phylogenetics and dating with confidence. *PLoS Biol.* 4: e88. <https://doi.org/10.1371/journal.pbio.0040088>
- Ducousso, M., Bena, G., Bourgeois, C., Buyck, B., Eyssartier, G., Vincelette, M., Rabevohitra, R., Randrihasipara, L., Dreyfus, B. & Prin, Y.** 2004. The last common ancestor of Sarcolaenaceae and Asian dipterocarp trees was ectomycorrhizal before the India–Madagascar separation, about 88 million years ago. *Molec. Ecol.* 13: 231–236. <https://doi.org/10.1046/j.1365-294x.2003.02032.x>
- Dutta, S., Mallick, M., Bertram, N., Greenwood, P.F. & Mathews, R.P.** 2009. Terpenoid composition and class of Tertiary resins from India. *Int. J. Coal Geol.* 80: 44–50. <https://doi.org/10.1016/j.coal.2009.07.006>
- Dutta, S., Tripathi, S.M., Mallick, M., Mathews, R.P., Greenwood, P.F., Rao, M.R. & Summons R.E.** 2011. Eocene out-of-India dispersal of Asian dipterocarps. *Rev. Palaeobot. Palynol.* 166: 63–68. <https://doi.org/10.1016/j.revpalbo.2011.05.002>
- Dynesius, M. & Jansson, R.** 2000. Evolutionary consequences of changes in species' geographical distributions driven by Milankovitch climate oscillations. *Proc. Natl. Acad. Sci. U.S.A.* 97: 9115–9120. <https://doi.org/10.1073/pnas.97.16.9115>
- FAO** 2014. *State of the World's forests: Enhancing the socioeconomic benefits from forests*. Rome: FAO.
- Feng, X., Tang, B., Kodrul, T.M. & Jin, J.** 2013. Winged fruits and associated leaves of *Shorea* (Dipterocarpaceae) from the late Eocene of South China and their phytogeographic and paleoclimatic implications. *Amer. J. Bot.* 100: 574–581. <https://doi.org/10.3732/ajb.1200397>
- Gamage, T.W., de Silva, M., Yoshida, A., Szmjdt, A.E. & Yamazaki, T.** 2003. Molecular phylogeny of Sri Lankan Dipterocarpaceae in relation to other Asian Dipterocarpaceae based on chloroplast DNA sequences. *Tropics* 13: 79–87. <https://doi.org/10.3759/tropics.13.79>
- Gamage, D., de Silva, M.P., Inomata, N., Yamazaki, T. & Szmjdt, A.E.** 2006. Comprehensive molecular phylogeny of the subfamily Dipterocarpoideae (Dipterocarpaceae) based on chloroplast DNA sequences. *Genes Genet. Syst.* 81: 1–12. <https://doi.org/10.1266/ggs.81.1>
- Garg, R.K.-A., Khowaja-Ateequzzaman, Prasad, V., Tripathi, S.K.M., Singh, I.B., Jauhri, A.K. & Bajpai, S.** 2008. Age-diagnostic dinoflagellate cysts from the lignite-bearing sediments of the Vastan lignite mine, Surat District, Gujarat, western India. *J. Palaeontol. Soc. India* 53: 99–105.
- Gathorne-Hardy, F.J., Syaukani, R.G.D., Eggleton, P. & Jones, D.T.** 2002. Quaternary rainforest refugia in south-east Asia: Using termites (Isoptera) as indicators. *Biol. J. Linn. Soc.* 75: 453–466. <https://doi.org/10.1046/j.1095-8312.2002.00031.x>
- Ghazoul, J.** 2016. *Dipterocarp biology, ecology, and conservation*. Oxford: Oxford University Press.
- Heaney, L.R.** 1991. A synopsis of climatic and vegetational change in Southeast Asia. *Clim. Change* 19: 53–61. <https://doi.org/10.1007/BF00142213>
- Heckenhauer, J., Samuel, R., Ashton, P.S., Turner, B., Barfuss, M.H.J., Jang, T.-S., Tensch, E.M., Mccann, J., Salim, K.A., Attanayake, A.M.A.S. & Chase, M.W.** 2017. Phylogenetic analyses of plastid DNA suggest a different interpretation of morphological evolution than those used as the basis for previous classifications of Dipterocarpaceae (Malvales). *Bot. J. Linn. Soc.* 20: 31–39. <https://doi.org/10.1093/botlinnean/box044>
- Heckenhauer, J., Samuel, R., Ashton, P.S., Kamariah, A.S. & Paun, O.** 2018. Phylogenomics resolves evolutionary relationships and provides insights into floral evolution in the tribe Shoreae (Dipterocarpaceae). *Molec. Phylog. Evol.* 127: 1–13. <https://doi.org/10.1016/j.ympev.2018.05.010>
- Heckenhauer, J., Paun, O., Chase, M.W., Ashton, P.S., Kamariah, A.S. & Samuel, R.** 2019. Molecular phylogenomics of the tribe Shoreae (Dipterocarpaceae) using whole plastid genomes. *Ann. Bot. (Oxford)* 123: 857–865. <https://doi.org/10.1093/aob/mcy220>
- Heim, F.** 1892. Sur un nouveau genre de Diptérocarpacées: *Vateriopsis seychellarum* Heim; *Vateria seychellarum* Dyer, in Baker. *Bull. Soc. Bot. France.* 39: 149–154.
- Hernández-Gutiérrez, R. & Magallón, S.** 2019. The timing of Malvales evolution: Incorporating its extensive fossil record to inform about lineage diversification. *Molec. Phylog. Evol.* 140: 106606. <https://doi.org/10.1016/j.ympev.2019.106606>
- Hinsinger, D.D. & Strijk, J.S.** 2015. Complete chloroplast genome sequence of *Castanopsis concinna* (Fagaceae), a threatened species

- from Hong Kong and South-Eastern China. *Mitochondrial DNA*, A, 2: 28: 65–66. <https://doi.org/10.3109/19401736.2015.1110800>
- Ho, K.-S., Chen, J.-C., Lo, C.-H. & Zhao, H.-L.** 2003. ⁴⁰Ar–³⁹Ar dating and geochemical characteristics of late Cenozoic basaltic rocks from the Zhejiang-Fujian region, SE China: Eruption ages, magma evolution and petrogenesis. *Chem. Geol.* 197: 287–318. [https://doi.org/10.1016/S0009-2541\(02\)00399-6](https://doi.org/10.1016/S0009-2541(02)00399-6)
- Huang, Y., Jia, L., Wang, Q., Mosbrugger, V., Utescher, T., Su T. & Zhou, Z.** 2016. Cenozoic plant diversity of Yunnan: A review. *Pl. Diversity* 38: 271–282. <https://doi.org/10.1016/j.pld.2016.11.004>
- Indrioko, S., Gailing, O. & Finkeldey, R.** 2006. Molecular phylogeny of Dipterocarpaceae in Indonesia based on chloroplast DNA. *Pl. Syst. Evol.* 261: 99–115. <https://doi.org/10.1007/s00606-006-0435-8>
- ITTO** 2019. Annual review and assessment of the world timber situation 2019. International Tropical Timber Association. https://www.itto.int/annual_review/ (accessed 10 Sep 2019).
- IUCN** 2020. The IUCN Red List of Threatened Species. Version 2020-1. <https://www.iucnredlist.org> (accessed 10 Sep 2019).
- Iwanaga, H., Teshima, K.M., Khatab, I.A., Inomata, N., Finkeldey, R., Siregar, I.Z., Siregar, U.J. & Szmidt, A.E.** 2012. Population structure and demographic history of a tropical lowland rainforest tree species *Shorea parvifolia* (Dipterocarpaceae) from Southeastern Asia. *Ecol. Evol.* 2: 1663–1675. <https://doi.org/10.1002/ece3.284>
- Jacques, F.M.B., Shi, G.L. & Wang, W.M.** 2011. Reconstruction of Neogene zonal vegetation in South China using the Integrated Plant Record (IPR) analysis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 307: 272–284. <https://doi.org/10.1016/j.palaeo.2011.05.025>
- Jacques, F.M.B., Shi, G.L. & Wang, W.M.** 2013. Neogene zonal vegetation of China and the evolution of the winter monsoon. *Bull. Geosci.* 88: 175–193. <https://doi.org/10.3140/bull.geosci.1359>
- Jacques, F.M.B., Shi, G., Su, T. & Zhou, Z.** 2015. A tropical forest of the middle Miocene of Fujian (SE China) reveals Sino-Indian biogeographic affinities. *Rev. Palaeobot. Palynol.* 216: 76–91. <https://doi.org/10.1016/j.revpalbo.2015.02.001>
- Jiang, W., Cheng, Y., Yang, X. & Yang, S.** 2013. Chinese loess plateau vegetation since the last glacial maximum and its implications for vegetation restoration. *J. Appl. Ecol.* 50: 440–448. <https://doi.org/10.1111/1365-2664.12052>
- Jong, K. & Lethbridge, A.** 1967. Cytological studies in the Dipterocarpaceae. I. Chromosome numbers of certain Malaysian genera. *Notes Roy. Bot. Gard. Edinburgh* 27: 175–184.
- Kajita, T., Kamiya, K., Nakamura, K., Tachida, H., Wickneswari, R., Tsumura, Y., Yoshimaru, H. & Yamazaki, T.** 1998. Molecular phylogeny of Dipterocarpaceae in Southeast Asia based on nucleotide sequences of *matK*, *trnL* intron, and *trnL-trnF* intergenic spacer region in chloroplast DNA. *Molec. Phylog. Evol.* 10: 202–209. <https://doi.org/10.1006/mpev.1998.0516>
- Kamiya, K., Harada, K., Ogino, K., Kajita, T., Yamazaki, T., Lee, H.-S. & Ashton, P.S.** 1998. Molecular phylogeny of dipterocarpaceae species using nucleotide sequences of two non-coding regions in chloroplast DNA. *Tropics* 7: 195–207. <https://doi.org/10.3759/tropics.7.195>
- Kamiya, K., Harada, K., Hidenori, T. & Ashton, P.S.** 2005. Phylogeny of *PggiC* gene in *Shorea* and its closely related genera (Dipterocarpaceae), the dominant trees in Southeast Asian tropical rain forests. *Amer. J. Bot.* 92: 775–788. <https://doi.org/10.3732/ajb.92.5.775>
- Kamiya, K., Nanami, S., Kenzo, T., Yoneda, R., Diway, B., Chong, L., Azani, M.A., Majid, N.M., Lum, S.K.Y., Wong, K.-M. & Harada, K.** 2012. Demographic history of *Shorea curtisii* (Dipterocarpaceae) inferred from chloroplast DNA sequence variations. *Biotropica* 44: 577–587. <https://doi.org/10.1111/j.1744-7429.2011.00834.x>
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. & Drummond, A.** 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Khan, M.A. & Bera, S.** 2010. Record of fossil fruit wing of *Shorea* Roxb. from the Neogene of Arunachal Pradesh. *Curr. Sci.* 98: 1573–1574.
- Koenen, E.J.M., Ojeda, D.I., Bakker, F.T., Wieringa, J.J., Kidner, C., Hardy, O.J., Pennington, R.T., Herendeen, P.S., Bruneau, A. & Hughes, C.E.** 2021. The origin of the legumes is a complex paleopolyploid phylogenomic tangle closely associated with the Cretaceous–Paleogene (K–Pg) mass extinction event. *Syst. Biol.* 70: 508–526. <https://doi.org/10.1093/sysbio/syaa041>
- Kooyman, R.M., Morley, R.J., Crayn, D.M., Joyce, E.M., Rossetto, M., Slik, J.W.F., Strijk, J.S., Su, T., Yap, J.-Y.S. & Wilf, P.** 2019. Origins and assembly of Malaysian rainforests. *Annual Rev. Ecol. Evol. Syst.* 50: 119–143. <https://doi.org/10.1146/annurev-evolsys-110218-024737>
- Korthals, P.W.** 1839–1842. *Verhandelingen over de natuurlijke geschiedenis der Nederlandsche overzeesche bezittingen*, vol. 2. Leiden: In commissie bij S. en J. Luchtmans en C.C. van der Hoek. <https://doi.org/10.5962/bhl.title.114730>
- Kostermans, A.J.G.H.** 1978. *Pakaraimaea dipterocarpacea* Maguire and Ashton belongs to Tiliaceae and not to Dipterocarpaceae. *Taxon* 27: 357–359. <https://doi.org/10.1111/j.1469-8137.2006.01860.x>
- Kostermans, A.J.G.H.** 1983. The Ceylonese species of *Shorea* Roxb. (Dipterocarpaceae). *Bot. Jahrb. Syst.* 104: 183–201. <https://doi.org/10.1007/s10265-010-0348-z>
- Kostermans, A.J.G.H.** 1984. Monograph of the genus *Doona* Thwaites (Dipterocarpaceae). *Bot. Jahrb. Syst.* 104: 425–454.
- Kostermans, A.J.G.H.** 1985. Family status for the Monotoideae Gilg and the Pakaraimoideae Ashton, Maguire and De Zeeuw (Dipterocarpaceae). *Taxon* 34: 426–435. <https://doi.org/10.2307/1221209>
- Kostermans, A.J.G.H.** 1992. *A handbook of the Dipterocarpaceae of Sri Lanka*. Colombo: Wildlife Heritage Trust of Sri Lanka.
- Kozlov, A.M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A.** 2019. RAXML-NG: A fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35: 4453–4455. <https://doi.org/10.1093/bioinformatics/btz305>
- Kroon, D., Steens, T. & Troelstra, S.R.** 1991. Onset of monsoonal related upwelling in the western Arabian Sea as revealed by planktonic foraminifers. Pp. 257–263 in: Stewart, M.J. (ed.), *Proceedings of the Ocean Drilling Program*, vol. 117, *Scientific results: Oman margin/Neogene package*. College Station, TX: Ocean Drilling Program. <https://doi.org/10.2973/odp.proc.sr.117.126.1991>
- Lakhanpal, R.N.** 1970. Tertiary floras of India and their bearing on the historical geology of the region. *Taxon* 19: 675–694. <https://doi.org/10.2307/1219280>
- Lakhanpal, R.N.** 1974. Geological history of the Dipterocarpaceae. Pp. 30–39 in: Lakhanpal, R.N. (ed.), *Symposium on origin and phytogeography of angiosperms*. Lucknow: Birbal Sahni Institute of Palaeobotany.
- Liu, Y.C. & Quan, C.** 2017. Late Cenozoic climates of low-latitude East Asia: A paleobotanical example from the Baise Basin of Guangxi, southern China. *Palaeoworld* 26: 572–580. <https://doi.org/10.1016/j.palwor.2016.08.001>
- Londoño, A.C., Alvarez, E., Forero, E. & Morton, C.M.** 1995. A new genus and species of Dipterocarpaceae from the Neotropics. I. Introduction, taxonomy, ecology and distribution. *Brittonia* 47: 225–236. <https://doi.org/10.2307/2807116>
- Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L.L. & Hernández-Hernández, T.** 2015. A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New Phytol.* 207: 437–453. <https://doi.org/10.1111/nph.13264>
- Mallick, M., Dutta, S., Greenwood, P.F. & Bertram, N.** 2009. Pyrolytic and spectroscopic studies of Eocene resin from Vastan Lignite Mine, Cambay Basin, western India. *J. Geol. Soc. India* 74: 16–22. <https://doi.org/10.1007/s12594-009-0098-5>

- Maguire, B., Ashton, P.S., De Zeeuw, C., Giannasi, D.E. & Niklas, K.J.** 1977. Pakaraimoideae, Diptercarpaceae of the Western Hemisphere. *Taxon* 26: 341–385. <https://doi.org/10.1002/j.1996-8175.1977.tb04186.x>
- Maury, G.** 1978. *Diptercarpaceae: Du fruit à la plantule*. Dissertation. Université Paul Sabatier, Toulouse, Paris, France.
- Maury-Lechon, G.** 1979a. Conséquences taxonomiques de l'étude des caractères des fruits/germinations, embryons et plantules des Diptercarpacées. Pp. 81–106 in: Maury-Lechon, G. (ed.), *Diptercarpacées: Taxonomie, phylogénie, écologie*. Mémoires du Muséum National d'Histoire Naturelle, Série B, Botanique, n.s., 26. Paris: Editions du Muséum.
- Maury-Lechon, G.** 1979b. Interprétation phylogénique des caractères des pollens, fruits-germinations et plantules des Diptercarpacées. Pp. 139–144 in: Maury-Lechon, G. (ed.), *Diptercarpacées: Taxonomie, phylogénie, écologie*. Mémoires du Muséum National d'Histoire Naturelle, Série B, Botanique, n.s., 26. Paris: Editions du Muséum.
- Maury-Lechon, G. & Curtet, L.** 1998. Biogeography and evolutionary systematics of Diptercarpaceae. Pp. 5–44 in: Appanah, S. & Turnbull, J.M. (eds.), *A review of diptercarps: Taxonomy, ecology and silviculture*. Bogor: Center for International Forestry Research.
- Meijaard, E.** 2003. Mammals of south-east Asian islands and their Late Pleistocene environments. *J. Biogeogr.* 30: 1245–1257. <https://doi.org/10.1046/j.1365-2699.2003.00890.x>
- Meijer, W. & Wood, G.H.S.** 1964. Diptercarps of Sabah (North Borneo). *Sabah Forest Rec.* 4: 1–344.
- Meijer, W. & Wood, G.H.S.** 1976. *Keys to diptercarps of Sabah*. Bogor: BIOTROP.
- Milanković, M.** 1941. *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem*. Royal Serbian Academy Special Publication 133. Belgrade: Königlich Serbische Akademie.
- Miller, M.A., Pfeiffer, W. & Schwartz, T.** 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Pp. 45–52 in: *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, Louisiana, 14 Nov 2010. Piscataway: IEEE. <https://doi.org/10.1109/GCE.2010.5676129>
- Miquel, F.A.W.** 1859. *Flora von Neederlandsch Indië*, vol. 1(2). Amsterdam: C. G. van der Post; Utrecht: C. van der Post. <https://doi.org/10.5962/bhl.title.93>
- Mirarab, S., Reaz, R., Bayzid, Md.S., Zimmermann, T., Swenson, M.S. & Warnow, T.** 2014. ASTRAL: Genome-scale coalescent-based species tree estimation. *Bioinformatics* 30: i541–i548. <https://doi.org/10.1093/bioinformatics/btu462>
- Molnar, P., England, P. & Martinrod, J.** 1993. Mantle dynamics, uplift of the Tibetan Plateau, and the Indian monsoon. *Rev. Geophys.* 31: 357–396. <https://doi.org/10.1029/93RG02030>
- Morley, R.J.** 2000. *Origin and evolution of tropical rain forests*. John Wiley & Sons.
- Morley, R.J.** 2012. A review of the Cenozoic palaeoclimate history of Southeast Asia. Pp. 79–114 in: Gower, D.J., Johnson, K., Richardson, J., Rosen, B., Rüber, L. & Williams, S. (eds.), *Biotic evolution and environmental change in Southeast Asia*. Cambridge, U.K.; Cambridge University Press.
- Morley, R.J. & Flenley, J.R.** 1987. Late Cainozoic vegetational and environmental changes in the Malay archipelago. Pp. 50–59 in: Whitmore, T.C. (eds.), *Biogeographical evolution of the Malay archipelago*. Oxford: Clarendon Press.
- Morton, C.M.** 1995. A new genus and species of Diptercarpaceae from the Neotropics. II. Stem anatomy. *Brittonia* 47: 237–247. <https://doi.org/10.2307/2807117>
- Morton, C.M., Dayanandan, S. & Dissanayake, D.** 1999. Phylogeny and biosystematics of *Pseudomonotes* (Diptercarpaceae) based on molecular and morphological data. *Pl. Syst. Evol.* 216: 197–205. <https://doi.org/10.1007/BF01084398>
- Mosbrugger, V., Utescher, T. & Dilcher, D.L.** 2005. Cenozoic continental climatic evolution of Central Europe. *Proc. Natl. Acad. Sci. U.S.A.* 102: 14964–14969. <https://doi.org/10.1073/pnas.0505267102>
- Muller, J.** 1981. Fossil pollen records of extant angiosperms. *Bot. Rev. (Lancaster)* 47: 1–142. [https://doi.org/10.1663/0006-8101\(2004\)070\[0425:FPROEA\]2.0.CO;2](https://doi.org/10.1663/0006-8101(2004)070[0425:FPROEA]2.0.CO;2)
- Murawski, D.A., Jaun, G. & Bawa, K.S.** 1994. The effect of selective logging on mating patterns of *Shorea megistophylla* (Diptercarpaceae) from Sri Lanka. *Conservation Biol.* 8: 997–1002. <https://doi.org/10.1046/j.1523-1739.1994.08040997.x>
- Ni, J., Yu, G., Harrison, S.P. & Prentice, I.C.** 2010. Palaeovegetation in China during the late Quaternary: Biome reconstructions based on a global scheme of plant functional types. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 289: 44–61. <https://doi.org/10.1016/j.palaeo.2010.02.008>
- Nieto Feliner, G. & Rosselló, J.A.** 2007. Better the devil you know? Guidelines for insightful utilization of nrDNA ITS in species-level evolutionary studies in plants. *Molec. Phylog. Evol.* 44: 911–919. <https://doi.org/10.1016/j.ympev.2007.01.013>
- Ohtani, M., Kondo, T., Tani, N., Ueno, S., Lee, L.S., Ng, K.K.S., Muhammad, N., Finkeldey, R., Na'iem, M., Indrioko, S., Kamiya, K., Harada, K., Diway, B., Khoo, E., Kawamura, K. & Tsumura, Y.** 2013. Nuclear and chloroplast DNA phylogeography reveals Pleistocene divergence and subsequent secondary contact of two genetic lineages of the tropical rainforest tree species *Shorea leprosula* (Diptercarpaceae) in South-East Asia. *Molec. Ecol.* 22: 2264–2279. <https://doi.org/10.1111/mec.12243>
- Parameswaran, N. & Gotwald, H.** 1979. Problematic taxa in the Diptercarpaceae: Their anatomy and taxonomy. *Mém. Mus. Natl. Hist. Nat., B, Bot.* 26: 69–75.
- Paul, S., Sharma, J., Singh, B.D., Saraswati, P.K. & Dutta, S.** 2015. Early Eocene equatorial vegetation and depositional environment: Biomarker and palynological evidences from a lignite-bearing sequence of Cambay Basin, western India. *Int. J. Coal Geol.* 149: 77–92. <https://doi.org/10.1016/j.coal.2015.06.017>
- Prakash, U., Vaidyanathan, L. & Tripathi, P.P.** 1994. Plant remains from the Tipam sandstones of northeast India with remarks on the palaeoecology of the region during the Miocene. *Palaeontographica Abt. B. Paläophytol.* 231: 113–146.
- Prasad, M.** 1990. Some more leaf-impressions from the lower Siwalik sediments of Koilabas, Nepal. *Palaeobotanist* 37: 299–305.
- Prasad, M.** 1994. Siwalik (Middle Miocene) leaf-impressions from the foothills of Himalayas, India. *Terestr. Resources* 15: 53–90.
- Prasad, M. & Awasthi, N.** 1996. Contribution to the Siwalik flora from Surai Khola sequence, western Nepal and its palaeoecological and phytogeographical implications. *Paleobotanist* 43: 1–42.
- Quade, J., Cerling, T.E. & Bowman, J.E.** 1989. Development of Asian monsoon revealed by marked ecological shift during the latest Miocene in northern Pakistan. *Nature* 342: 163–166. <https://doi.org/10.1038/342163a0>
- Raes, N., Cannon, C.H., Hijmans, R.J., Piessens, T., Saw, L.G., Van Welzen, P.C. & Slik, J.W.F.** 2014. Historical distribution of Sundaland's Diptercarp rainforests at Quaternary glacial maxima. *Proc. Natl. Acad. Sci. U.S.A.* 111: 16790–16795. <https://doi.org/10.1073/pnas.1403053111>
- Rambaut, A., Drummond, A.J., Xie, D., Baele, G. & Suchard, M.A.** 2018. Posterior summarisation in Bayesian phylogenetics using Tracer 1.7. *Syst. Biol.* 67: 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ramstein, G., Fluteau, F., Besse, J. & Jousaume, S.** 1997. Effect of orogeny, plate motion and land-sea distribution on Eurasian climate change over the past 30 million years. *Nature* 386: 789–795. <https://doi.org/10.1038/386788a0>
- Rana, R.S., Kumar, K. & Singh, H.** 2004. Vertebrate fauna from the subsurface Cambay Shale (Lower Eocene), Vastan lignite mine, Gujarat, India. *Curr. Sci. India* 87: 1726–1733.
- Ruddiman, W.F. & Kutzbach, J.E.** 1989. Forcing of late Cenozoic Northern Hemisphere climate by plateau uplift in southern Asia

- and the American west. *J. Geophys. Res.* 94: 18409–18427. <https://doi.org/10.1029/JD094iD15p18409>
- Rudra, A., Dutta, S. & Raju, S.V.**, 2014. Molecular composition and paleobotanical origin of Eocene resin from northeast India. *J. Earth System Sci.* 123(5): 935–941. <https://doi.org/10.1007/s12040-014-0443-y>
- Rust, J., Singh, H., Rana, R.S., McCann, T., Singh, L., Anderson, K., Sarkar, N., Nascimbene, P.C., Stebner, F., Thomas, J.C., Kraemer, M.S., Williams, C.J., Engel, M.S., Sahni, A. & Grimaldi, D.** 2010. Biogeographic and evolutionary implications of a diverse paleobiota in amber from the early Eocene of India. *Proc. Natl. Acad. Sci. U.S.A.* 107: 18360–18365. <https://doi.org/10.1073/pnas.1007407107>
- Sahni, A., Saraswati, P.K., Rana, R.S., Kumar, K., Singh, H., Alimohammadian, H., Sahni, N., Rose, K.D., Singh, L. & Smith, T.** 2006. Temporal constraints and depositional palaeoenvironments of the Vastan lignite, sequence, Gujarat: Analogy for the Cambay Shale hydrocarbon source rock. *J. Petrol. Geol.* 15: 1–20.
- Sasaki, S.** 2006. Ecology and physiology of Dipterocarpaceae. Pp. 3–22 in: Suzuki, K., Ishii, K., Sakurai, S. & Sasaki, S. (eds.), *Plantation technology in tropical forest science*. Tokyo: Springer.
- Shi, G. & Li, H.** 2010. A fossil fruit wing of *Dipterocarpus* from the middle Miocene of Fujian, China and its palaeoclimatic significance. *Rev. Palaeobot. Palynol.* 162: 599–606. <https://doi.org/10.1016/j.revpalbo.2010.08.001>
- Shi, G.L., Dutta, S., Paul, S., Wang, B. & Jacques, F.M.B.** 2014a. Terpenoid compositions and botanical origins of Late Cretaceous and Miocene amber from China. *PLoS ONE* 9(10): e111303. <https://doi.org/10.1371/journal.pone.0111303>
- Shi, G.L., Jacques, F.M.B. & Li, H.M.** 2014b. Winged fruits of *Shorea* (Dipterocarpaceae) from the Miocene of Southeast China: Evidence for the northward extension of dipterocarps during the Mid-Miocene Climatic Optimum. *Rev. Palaeobot. Palynol.* 200: 97–107. <https://doi.org/10.1016/j.revpalbo.2013.09.003>
- Shukla, A., Guleria, J.S. & Mehrotra, R.C.** 2012. Fossil record of a *Shorea* Roxb. fruit wing from the Early Miocene sediments of Kachchh, Gujarat and its bearing on palaeoclimatic interpretation. *J. Earth System Sci.* 121: 195–201. <https://doi.org/10.1007/s12040-012-0142-5>
- Shukla, A., Mehrotra, R.C. & Guleria, J.S.** 2013. Emergence and extinction of Dipterocarpaceae in western India with reference to climate change: Fossil wood evidences. *J. Earth System Sci.* 122: 1373–1386. <https://doi.org/10.1007/s12040-013-0341-8>
- Slik, J.W.F., Aiba, S.-I., Bastian, M., Brearley, F.Q., Cannon, C.H., Eichhorn, K.A.O., Fredriksson, G., Kartawinata, K., Laumonier, Y., Mansor, A., Marjokorpi, A., Meijaard, E., Morley, R.J., Nagamasu, H., Nilus, R., Nurtjahya, E., Payne, J., Permana, A., Poulsen, A.D., Raes, N., Riswan, S., van Schaik, C.P., Sheil, D., Sidiyasa, K., Suzuki, E., Van Valkenburg, J.L.C.H., Webb, C.O., Wich, S., Yoneda, T., Zakaria, R. & Zweifelu, N.** 2011. Soils on exposed Sunda shelf shaped biogeographic patterns in the equatorial forests of Southeast Asia. *Proc. Natl. Acad. Sci. U.S.A.* 108: 12343–12347. <https://doi.org/10.1073/pnas.1103353108>
- Spooner, D.M., Ruess, H., Ellison, S., Senalik, D. & Simon, P.** 2020. What is truth: Consensus and discordance in next-generation phylogenetic analyses of *Daucus*. *J. Syst. Evol.* 58: 1059–1070. <https://doi.org/10.1111/jse.12678>
- Srivastava, G. & Mehrotra, R.C.** 2010. Tertiary flora of north-east India vis-à-vis movement of the Indian plate. *Mem. Geol. Soc. India* 75: 123–130.
- Suchard, M.A. & Rambaut, A.** 2009. Many-core algorithms for statistical phylogenetics. *Bioinformatics* 25: 1370–1376. <https://doi.org/10.1093/bioinformatics/btp244>
- Sun, B.N., Wu, J.Y., Liu, C.Y.S., Ding, S.-T., Li, X.-C., Xie, S.-P., Yana, D.-F. & Lin, Z.-C.** 2011. Reconstructing Neogene vegetation and climates to infer tectonic uplift in western Yunnan, China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 304: 328–336. <https://doi.org/10.1016/j.palaeo.2010.09.023>
- Sun, X.J. & Wang, P.X.** 2005. How old is the Asian monsoon system? Palaeobotanical records from China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 222: 181–222. <https://doi.org/10.1016/j.palaeo.2005.03.005>
- Sun, X., Li, X., Luo, Y. & Chen, X.** 2000. The vegetation and climate at the last glaciation on the emerged continental shelf of the South China Sea. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 160: 301–316. [https://doi.org/10.1016/S0031-0182\(00\)00078-X](https://doi.org/10.1016/S0031-0182(00)00078-X)
- Symington, C.F.** 1943. *Forester's manual of dipterocarps*. Malayan Foresters Records 16. Kuala Lumpur: Penerbit Universiti Malaysia.
- Takhtajan, A.** 1980. Outline of the classification of flowering plants (Magnoliophyta). *Bot. Rev. (Lancaster)* 46: 225–359. <https://doi.org/10.1007/BF02861558>
- Takhtajan, A.** 2009. *Flowering plants*. New York: Springer Science & Business Media.
- Tsumura, Y., Kawahara, T., Wickneswari, R. & Yoshimura, K.** 1996. Molecular phylogeny of Dipterocarpaceae in Southeast Asia using RFLP of PCR-amplified chloroplast genes. *Theor. Appl. Genet.* 93(1–2): 22–29. <https://doi.org/10.1007/BF00225722>
- Tsumura, Y., Kado, T., Yoshida, K., Abe, H., Ohtani, M., Taguchi, Y., Fukue, Y., Tani, N., Ueno, S., Yoshimura, K., Kamiya, K., Harada, K., Takeuchi, Y., Diway, B., Finkeldey, R., Na'iem, M., Indrioko, S., Kit, K., Ng, S., Muhammad, N. & Lee, S.L.** 2011. Molecular database for classifying *Shorea* species (Dipterocarpaceae) and techniques for checking the legitimacy of timber and wood products. *J. Pl. Res.* 124: 35–48. <https://doi.org/10.1007/s10265-010-0348-z>
- Turland, N.J., Wiersema, J.H., Barrie, F.R., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Kusber, W.-H., Li, D.-Z., Marhold, K., May, T.W., McNeill, J., Monro, A.M., Prado, J., Price, M.J. & Smith, G.F. (eds.)** 2018. *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017*. Regnum Vegetabile 159. Glashütten: Koeltz Botanical Books. <https://doi.org/10.12705/Code.2018>
- Van Aarssen, B.G.K., De Leeuw, J.W., Collinson, M., Boon, J.J. & Goth, K.** 1994. Occurrence of polycadinene in fossil and recent resins. *Geochim. Cosmochim. Acta* 58(1): 223–229. [https://doi.org/10.1016/0016-7037\(94\)90459-6](https://doi.org/10.1016/0016-7037(94)90459-6)
- Van Der Hammen, T.** 1954. El desarrollo de la flora Columbiana en los periodos geologicos I: Maectrichtiano hasta terciario mas inferior (Una investigación Palinológica de la formación de Guaduas y equivalentes). *Bol. Geol.* 1: 49–106.
- Vanneste, K., Baele, G., Maere, S. & Van de Peer, Y.** 2014. Analysis of 41 plant genomes supports a wave of successful genome duplications in association with the Cretaceous–Paleogene boundary. *Genome Res.* 24: 1334–1347. <https://doi.org/10.1101/gr.168997.113>
- Whitten, A.J., Mustafa, M. & Henderson, G.S.** 1987. *The ecology of Sulawesi*. Yogyakarta: Gadjah Mada UP.
- Wolfe, J.A.** 1977. Paleogene floras from the Gulf of Alaska region. *U.S. Profess. Pap. U.S. Geol. Surv.* 997: 1–108.
- Wolfe, J.A.** 1994a. Tertiary climatic changes at middle latitudes of western North America. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 108: 195–205. [https://doi.org/10.1016/0031-0182\(94\)90233-X](https://doi.org/10.1016/0031-0182(94)90233-X)
- Wolfe, J.A.** 1994b. An analysis of Neogene climates in Beringia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 108: 207–216. [https://doi.org/10.1016/0031-0182\(94\)90234-8](https://doi.org/10.1016/0031-0182(94)90234-8)
- Wurster, C.M., Bird, M.I., Bull, I.D., Creed, F., Bryant, C., Dungait, J.A.J. & Paz, V.** 2010. Forest contraction in north equatorial Southeast Asia during the Last Glacial Period. *Proc. Natl. Acad. Sci. U.S.A.* 107: 15508–15511. <https://doi.org/10.1073/pnas.1005507107>
- Xia, K., Su, T., Liu, Y.S., Xing, Y.W., Jacques, F.M.B. & Zhou, Z.K.** 2009. Quantitative climate reconstructions of the late Miocene Xiaolongtan megafloora from Yunnan, southwest China. *Palaeogeogr.*

- Palaeoclimatol. Palaeoecol.* 276: 80–86. <https://doi.org/10.1016/j.palaeo.2009.02.024>
- Xing, Y., Utescher, T., Jacques, F.M.B., Su, T., Liu, Y.C., Huang, Y. & Zhou, Z. 2012. Paleoclimatic estimation reveals a weak winter monsoon in southwestern China during the late Miocene: Evidence from plant macrofossils. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 358–360: 19–26. <https://doi.org/10.1016/j.palaeo.2012.07.011>
- Yule, U. 1925. A mathematical theory of evolution, based on the conclusions of Dr. J.C. Willis, F.R.C. *Philos. Trans., Ser. B* 213: 21–87. <https://doi.org/10.1098/rstb.1925.0003>
- Yulita, K.S., Bayer, R.J. & West, J.G. 2005. Molecular phylogenetic study of *Hopea* and *Shorea* (Dipterocarpaceae): Evidence from the *trnL-trnF* and internal transcribed spacer regions. *Pl. Spec. Biol.* 20: 167–182. <https://doi.org/10.1111/j.1442-1984.2005.00136.x>
- Zachos, J., Pagani, M., Sloan, L., Thomas, E. & Billups, K. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to Present. *Science* 292: 686–693. <https://doi.org/10.1126/science.1059412>
- Zheng, H.B., Powell, C.M., Rea, D.K., Wang, J.L. & Wang, P.X. 2004. Late Miocene and mid-Pliocene enhancement of the East Asian monsoon as viewed from the land and sea. *Global Planet. Change* 41: 147–155. <https://doi.org/10.1016/j.gloplacha.2004.01.003>

Appendix 1. Accession list and voucher information for samples used in this study.

Species name; location data (country, province); voucher information; GenBank accession numbers of cpDNA (complete plastid genome), NRC (nuclear ribosomal cistrons) or ITS (internal transcribed spacer). Newly generated sequences, 151 plastome and 164 NRC, are indicated with *; missing information is indicated by –.

Anisoptera costata Korth., Singapore, *Strijk 4412* (SING), MZ901761*, –, *Anisoptera costata* Korth., Brunei, *Strijk 2509* (BGT), MZ901760*, MZ782371*, *Anisoptera marginata* Korth., Singapore, *Strijk 4401* (SING), MZ901762*, MZ782372*, *Anisoptera megistocarpa* Slooten, Singapore, *Strijk 4398* (SING), MZ901763*, MZ782373*, *Anisoptera megistocarpa* Slooten, Singapore (MacRitchie Reservoir), *Strijk 4429* (SING), MZ901764*, MZ782374*, *Anisoptera scaphula* (Roxb.) Kurz, Singapore (Yishun Park), *Strijk 4306* (SING), MZ901765*, MZ782375*, *Cotylelobium lanceolatum* Craib, Singapore, *Strijk 4374* (SING), MZ901766*, MZ782383*, *Cotylelobium melanoxylon* (Hook.f.) Pierre, Brunei, *Strijk 2100* (BGT), MZ901767*, MZ782384*, *Dipterocarpus alatus* Roxb., Singapore (Yishun Park), *Strijk 4300* (SING), MZ901912*, MZ889121*, *Dipterocarpus baudii* Korth., Singapore, *Strijk 4356* (SING), MZ901768*, MZ782386*, *Dipterocarpus caudatus* Foxw., Singapore, *Strijk 4278* (SING), MZ901769*, MZ782387*, *Dipterocarpus chartaceus* Symington, Singapore, *Strijk 4383* (SING), MZ901770*, MZ782388*, *Dipterocarpus confertus* Slooten, Singapore, *Strijk 4416* (SING), –, MZ782389*, *Dipterocarpus cornutus* Dyer, Singapore (Yishun Park), *Strijk 4299* (SING), –, MZ782390*, *Dipterocarpus costulatus* Slooten, Singapore (Yishun Park), *Strijk 4302* (SING), –, MZ782391*, *Dipterocarpus dyeri* Pierre, Singapore, *Strijk 4381* (SING), –, MZ782392*, *Dipterocarpus elongatus* Korth., Singapore (Chestnut Nature Park), *Strijk 4331* (SING), MZ901771*, MZ782393*, *Dipterocarpus fagineus* Vesque, Singapore, *Strijk 4405* (SING), MZ901772*, MZ782394*, *Dipterocarpus gracilis* Blume, Singapore (Yishun Park), *Strijk 4303* (SING), MZ901773*, MZ782396*, *Dipterocarpus gracilis* Blume, China (Qin Xiu Shan), *Strijk 3593* (BGT), –, MZ782395*, *Dipterocarpus grandiflorus* Blanco, Singapore, *Strijk 4392* (SING), MZ901774*, MZ782397*, *Dipterocarpus grandiflorus* Blanco, Singapore (MacRitchie Reservoir), *Strijk 4424* (SING), MZ901775*, MZ782398*, *Dipterocarpus humeratus* Slooten, Singapore, *Strijk 4380* (SING), MZ901776*, MZ782399*, *Dipterocarpus kerrii* King, Singapore, *Strijk 4272* (SING), MZ901777*, MZ782400*, *Dipterocarpus kunstleri* King, Singapore, *Strijk 4273* (SING), –, MZ782401*, *Dipterocarpus oblongifolius* Blume, Singapore, *Strijk 4290* (SING), MZ901778*, MZ782402*, *Dipterocarpus palembanicus* Slooten, Singapore, *Strijk 4335* (SING), MZ901779*, MZ782403*, *Dipterocarpus retusus* Blume, China (Qin Xiu Shan), *Strijk 3592* (BGT), –, MZ782404*, *Dipterocarpus rigidus* Ridl., Singapore, *Strijk 4397* (SING), MZ901780*, MZ782405*, *Dipterocarpus semivestitus* Slooten, Singapore, *Strijk 4414* (SING), MZ901781*, MZ782406*, *Dipterocarpus sp.*, China (Qin Xiu Shan), *Strijk 3590* (BGT), –, MZ782412*, *Dipterocarpus sublamellatus* Foxw., Singapore (MacRitchie Reservoir), *Strijk 4422* (SING), MZ901782*, MZ782407*, *Dipterocarpus tempehes* Slooten, Singapore (MacRitchie Reservoir), *Strijk 4421* (SING), MZ901784*, MZ782409*, *Dipterocarpus tempehes* Slooten, Singapore, *Strijk 4271* (SING), MZ901783*, MZ782408*, *Dipterocarpus tuberculatus* Roxb., Singapore, *Strijk 4379* (SING), MZ901785*, MZ782410*, *Dipterocarpus validus* Blume, Singapore, *Strijk 4395* (SING), –, MZ782411*, *Dipterocarpus zeylanicus* Thw., Singapore (Yishun Park), *Strijk 4301* (SING), MZ901786*, MZ782413*, *Dryobalanops aromatica* C.F.Gaertn., Singapore, *Strijk 4289* (SING), MZ901787*, MZ782414*, *Dryobalanops beccarii* Dyer, Singapore (Yishun Park), *Strijk 4315* (SING), MZ901788*, MZ782415*, *Dryobalanops lanceolata* Burck, Singapore (Yishun Park), *Strijk 4309* (SING), MZ901789*, MZ782416*, *Dryobalanops oblongifolia* Dyer, Singapore, *Strijk 4283* (SING), MZ901790*, MZ782417*, *Hopea auriculata* Foxw., Singapore (Yishun Park), *Strijk 4297* (SING), MZ901805*, MZ782420*, *Hopea beccariana* Burck, Singapore, *Strijk 4435* (SING), MZ901806*, MZ782421*, *Hopea coriacea* Burck, Singapore, *Strijk 4367* (SING), MZ901808*, MZ782423*, *Hopea dryobalanoides* Miq., Singapore (Yishun Park), *Strijk 4296* (SING), MZ901809*, MZ782424*, *Hopea dryobalanoides* Miq., –, –, MH791329, *Hopea exalata* W.T.Lin, Y.Y.Yang & Q.S.Hsue, China (Yunnan), *Strijk 4625* (BGT), MZ901810*, MZ782425*, *Hopea ferrea* Laness., Singapore, *Strijk 4364* (SING), MZ901811*, MZ782426*, *Hopea ferruginea* Parisi, Singapore, *Strijk 4280* (SING), MZ901812*, MZ782427*, *Hopea ferruginea* Parisi, Singapore (MacRitchie Reservoir), *Strijk 4425* (SING), MZ901813*, MZ782428*, *Hopea glaucescens* Symington, Singapore, *Strijk 4362* (SING), MZ901814*, MZ782429*, *Hopea griffithii* Kurz, Singapore (MacRitchie Reservoir), *Strijk 4419* (SING), MZ901816*, MZ782431*, *Hopea griffithii* Kurz, Singapore, *Strijk 4363* (SING), MZ901815*, MZ782430*, *Hopea helferi* (Dyer) Brandis, Singapore (Yishun Park), *Strijk 4308* (SING), MZ901817*, MZ782432*, *Hopea johorensis* Symington, Singapore, *Strijk 4359* (SING), MZ901818*, MZ782433*, *Hopea kerangasensis* P.S.Ashton, Singapore, *Strijk 4361* (SING), MZ901819*, MZ782434*, *Hopea mengarawan* Miq., Singapore (Chestnut Nature Park), *Strijk 4330* (SING), MZ901821*, MZ782436*, *Hopea mengarawan* Miq., Singapore, *Strijk 4282* (SING), MZ901820*, MZ782435*, *Hopea nervosa* King, Singapore (Yishun Park), *Strijk 4293* (SING), MZ901822*, MZ782437*, *Hopea nutans* Ridl., Singapore (Yishun Park), *Strijk 4294* (SING), MZ901823*, MZ782438*, *Hopea odorata* Roxb., Singapore, *Strijk 4270* (SING), MZ901824*, MZ782439*, *Hopea pachycarpa* (F.Heim) Symington, Singapore, *Strijk 4368* (SING), MZ901825*, MZ782440*, *Hopea polyalthioides* Symington, Singapore, *Strijk 4365* (SING), MZ901826*, MZ782441*, *Hopea cf. polyalthioides* Symington, Singapore, *Strijk 4360* (SING), MZ901807*, MZ782422*, *Hopea pubescens* Ridl., Singapore, *Strijk 4333* (SING), MZ901827*, MZ782442*, *Hopea sangal* Korth., Singapore (Yishun Park), *Strijk 4312* (SING), MZ901828*, MZ782443*, *Hopea semicuneata* Symington, Singapore, *Strijk 4366* (SING), MZ901829*, MZ782444*, *Hopea subulata* Symington, Singapore, *Strijk 4378* (SING), MZ901830*, MZ782445*, *Hopea wightiana* Wall., Singapore (Yishun Park), *Strijk 4295* (SING), MZ901831*, MZ782446*, *Neobalanocarpus heimii* (King) P.S.Ashton, Singapore, *Strijk 4279* (SING), MZ901832*, MZ782449*, *Neobalanocarpus heimii* (King) P.S.Ashton, –, –, MH746730, AY026657, *Parashorea chinensis* Wang Hsie, China (Yunnan), *Strijk 3594* (BGT), <https://doi.org/10.5061/dryad.1512pn3> (previous study), *Parashorea chinensis* Wang Hsie, China (Yunnan), *Strijk 4551* (BGT), MZ901833*, MZ782450*, *Parashorea macrophylla* Wyatt-Sm. ex P.S.Ashton, –, –, MH791330, *Parashorea malaanonan* (Blanco) Merr., Singapore, *Strijk 4386* (SING), MZ901834*, MZ782451*, *Parashorea smythiesii* Wyatt-Sm. ex P.S.Ashton, Singapore, *Strijk 4413* (SING), MZ901835*, MZ782452*, *Parashorea stellata* Slooten (1927, non Kurz 1870), Singapore, *Strijk 4402* (SING), MZ901836*, MZ782453*, *Shorea acuminata* Dyer, Singapore (Yishun Park), *Strijk 4323* (SING), MZ901837*, MZ782458*, *Shorea agamii* P.S.Ashton, Singapore (Yishun Park), *Strijk 4324* (SING), MZ901838*, MZ782459*, *Shorea amplexicaulis* P.S.Ashton, Singapore, *Strijk 4436* (SING), MZ901839*, MZ782460*, *Shorea argentifolia* Symington, Singapore (Yishun Park), *Strijk 4320* (SING), MZ901840*, MZ782461*, *Shorea atrinervosa* Symington,

Appendix 1. Continued.

Singapore, *Strijk 4431* (SING), –, MZ782462*, *Shorea balanocarpoides* Symington, Singapore (Yishun Park), *Strijk 4319* (SING), MZ901841*, MZ782463*, *Shorea bentongensis* Foxw., Singapore (Yishun Park), *Strijk 4318* (SING), MZ901842*, MZ782464*, *Shorea bracteolata* Dyer, Singapore, *Strijk 4384* (SING), MZ901843*, MZ782465*, *Shorea confusa* P.S.Ashton, Singapore, *Strijk 4382* (SING), MZ901845*, MZ782467*, *Shorea curtisii* Dyer ex King, Singapore, *Strijk 4291* (SING), MZ901846*, MZ782468*, *Shorea faguetiana* F.Heim, Singapore (Yishun Park), *Strijk 4321* (SING), MZ901847*, MZ782469*, *Shorea falciferoides* subsp. *falciferoides* Foxw., Singapore (The Gallop Arboretum), *Strijk 4352* (SING), MZ901848*, MZ782470*, *Shorea flaviflora* G.H.S.Wood ex P.S.Ashton, Singapore, *Strijk 4275* (SING), MZ901849*, MZ782471*, *Shorea gibbosa* Brandis, Singapore, *Strijk 4343* (SING), MZ901850*, MZ782472*, *Shorea glauca* King, Singapore, *Strijk 4344* (SING), –, MZ782473*, *Shorea gratissima* Dyer, Singapore (Yishun Park), *Strijk 4317* (SING), MZ901851*, MZ782474*, *Shorea guiso* (Blanco) Blume, Singapore (The Gallop Arboretum), *Strijk 4353* (SING), MZ901852*, MZ782475*, *Shorea hemsleyana* (King) King ex Foxw., Singapore, *Strijk 4341* (SING), MZ901853*, MZ782476*, *Shorea hypochra* Hance, Singapore (Yishun Park), *Strijk 4304* (SING), MZ901854*, MZ782477*, *Shorea cf. johorensis* Foxw., Singapore (MacRitchie Reservoir), *Strijk 4420* (SING), MZ901844*, MZ782466*, *Shorea johorensis* Foxw., Singapore (Yishun Park), *Strijk 4325* (SING), MZ901855*, MZ782478*, *Shorea kudatensis* G.H.S.Wood ex Meijer, Singapore, *Strijk 4385* (SING), MZ901856*, MZ782479*, *Shorea kunstleri* King, Singapore, *Strijk 4342* (SING), MZ901857*, MZ782480*, *Shorea laevis* Ridl., Singapore, *Strijk 4276* (SING), MZ901858*, MZ782481*, *Shorea lepidota* (Korth.) Blume, Singapore, *Strijk 4345* (SING), MZ901859*, MZ782482*, *Shorea leprosula* Miq., Singapore, *Strijk 4346* (SING), MZ901860*, MZ782483*, *Shorea macrantha* Brandis, Singapore, *Strijk 4399* (SING), MZ901861*, MZ782484*, *Shorea macrophylla* (de Vr.) P.S.Ashton, Singapore, *Strijk 4372* (SING), –, MZ782485*, *Shorea macroptera* Dyer, Singapore (MacRitchie Reservoir), *Strijk 4472* (SING), MZ901864*, MZ782487*, *Shorea macroptera* Dyer, Singapore, *Strijk 4339* (SING), MZ901863*, MZ782486*, *Shorea materialis* Ridley, Singapore (Yishun Park), *Strijk 4307* (SING), –, MZ782488*, *Shorea maxima* (King) Symington, Singapore, *Strijk 4357* (SING), MZ901865*, MZ782489*, *Shorea maxwelliana* (King) Symington, Singapore (Yishun Park), *Strijk 4322* (SING), MZ901866*, MZ782490*, *Shorea mecistopteryx* Ridl., Singapore, *Strijk 4430* (SING), MZ901867*, MZ782491*, *Shorea multiflora* (Burck) Symington, Singapore, *Strijk 4348* (SING), MZ901868*, MZ782492*, *Shorea ochrophloia* Strugnell ex Symington, Singapore (The Gallop Arboretum), *Strijk 4351* (SING), MZ901869*, MZ782493*, *Shorea ovalis* (Korth.) Blume, Singapore, *Strijk 4434* (SING), MZ901870*, MZ782494*, *Shorea pachyphylla* Ridl. ex Symington, –, –, MH841940, *Shorea parvifolia* Dyer, Singapore (The Gallop Arboretum), *Strijk 4354* (SING), MZ901871*, MZ782495*, *Shorea pauciflora* King, Singapore (MacRitchie Reservoir), *Strijk 4423* (SING), MZ901873*, MZ782497*, *Shorea pauciflora* King, Singapore, *Strijk 4337* (SING), MZ901872*, MZ782496*, *Shorea peltata* Symington, Singapore, *Strijk 4400* (SING), MZ901874*, MZ782498*, *Shorea platycarpa* F.Heim, Singapore, *Strijk 4371* (SING), MZ901875*, MZ782499*, *Shorea platyclados* Slooten ex Foxw., Singapore, *Strijk 4396* (SING), MZ901876*, MZ782500*, *Shorea resinosa* Foxw., Singapore, *Strijk 4274* (SING), MZ901877*, MZ782501*, *Shorea robusta* Gaertn., Singapore, *Strijk 4417* (SING), MZ901878*, MZ782502*, *Shorea roxburghii* G.Don, Singapore, *Strijk 4390* (SING), MZ901879*, MZ782503*, *Shorea roxburghii* G.Don, China (Yunnan), *Strijk 4627* (BGT), MZ901880*, MZ782504*, *Shorea singkawang* (Miq.) Burck, Singapore, *Strijk 4415* (SING), MZ901881*, MZ782505*, *Shorea smithiana* Symington, Singapore (The Gallop Arboretum), *Strijk 4355* (SING), MZ901882*, MZ782506*, *Shorea stenoptera* Burck., Singapore (Yishun Park), *Strijk 4316* (SING), MZ901883*, MZ782507*, *Shorea sumatrana* (Slooten ex Thorenaar) Desch, Singapore, *Strijk 4286* (SING), MZ901884*, MZ782508*, *Shorea symingtonii* G.H.S.Wood, Singapore, *Strijk 4340* (SING), MZ901885*, MZ782509*, *Shorea xanthophylla* Symington, Singapore, *Strijk 4373* (SING), MZ901886*, MZ782510*, *Shorea zeylanica* (Thwaites) P.S.Ashton, –, –, MH841939, *Vatica bella* Slooten, Singapore (Yishun Park), *Strijk 4310* (SING), MZ901791*, MZ782518*, *Vatica flavida* Foxw., Singapore, *Strijk 4377* (SING), MZ901792*, MZ782519*, *Vatica lobata* Foxw., Singapore (Yishun Park), *Strijk 4313* (SING), MZ901793*, MZ782520*, *Vatica maingayi* Dyer, Singapore (Yishun Park), *Strijk 4314* (SING), MZ901794*, MZ782521*, *Vatica maritima* Slooten, Singapore, *Strijk 4375* (SING), MZ901795*, MZ782522*, *Vatica nitens* King, Singapore, *Strijk 4404* (SING), MZ901796*, MZ782523*, *Vatica odorata* (Griff.) Symington, Singapore, *Strijk 4376* (SING), MZ901797*, MZ782524*, *Vatica odorata* (Griff.) Symington, Singapore (MacRitchie Reservoir), *Strijk 4426* (SING), MZ901798*, MZ782525*, *Vatica odorata* (Griff.) Symington, China (Yunnan), *Strijk 1594* (BGT), KX966283, –, *Vatica perakensis* King, Singapore, *Strijk 4403* (SING), MZ901799*, MZ782526*, *Vatica rassak* (Korth.) Blume, Singapore, *Strijk 4292* (SING), MZ901800*, MZ782527*, *Vatica ridleyana* Brandis, Singapore, *Strijk 4358* (SING), MZ901801*, MZ782528*, *Vatica stapfiana* (King) Slooten, Singapore, *Strijk 4287* (SING), MZ901802*, MZ782529*, *Vatica umbonata* (Hook.f.) Burck, Singapore, *Strijk 4288* (SING), MZ901803*, MZ782530*, *Vatica yeechongii* Saw, Singapore (Yishun Park), *Strijk 4311* (SING), MZ901804*, MZ782531*, **Outgroups:** *Abelmoschus esculentus* (L.) Moench, U.S.A. (Austin), –, NC035234, KP222461, *Adansonia digitata* L., Gabon (Nyanga, Mayumba), *Wieringa 8465* (WAG8001190), MZ901888*, MZ782370*, *Adansonia digitata* L., Singapore, *Strijk 4394* (SING), MZ901887*, MZ782369*, *Abelmoschus esculentus* (L.) Moench, U.S.A. (Austin), –, NC035234, KP222461, *Althaea officinalis* L., –, KY085914, EF679733, *Aquilaria sinensis* (Lour.) Gilg, China (Hainan), –, NC029243, GQ891956, *Arabidopsis arenosa* (L.) Lawalrée, –, LT161918, DQ528871, *Arabidopsis thaliana* L., U.S.A., –, NC000932, DQ528813, *Arabis alpina* L., –, HF934132, AF137559, *Bombacopsis glabra* (Pasq.) A.Robyns, China (Yunnan), *Strijk 4531* (BGT), MZ901889*, MZ782376*, *Bombax ceiba* L., China (Yunnan), *Strijk 4596* (BGT), MZ901890*, MZ782377*, *Bombax ceiba* L., China (Yunnan), –, MG569974, HQ658377, *Bombax malabaricum* DC., China (Yunnan), *Strijk 4592* (BGT), MZ901891*, MZ782378*, *Ceiba insignis* (Kunth) P.E.Gibbs & Semir, China (Qin Xiu Shan), *Strijk 4467* (BGT), MZ901892, MZ782379*, *Chorisia speciosa* A.St.-Hil., China (Yunnan), *Strijk 4595* (BGT), MZ901892*, MZ782380*, *Cola micrantha* K.Schum., Cameroon, –, *Wieringa 5862* (WAG8003629), MZ901894*, MZ782381*, *Colona floribunda* (Kurz) Craib, China (Yunnan), *Strijk 4483* (BGT), MZ901895*, MZ782382*, *Dicranolepis disticha* Planch., Cameroon, *Wieringa 5821* (WAG8003655), MZ901896*, MZ782385*, *Durio zibethinus* L., –, MG138151, MF629775, *Edgeworthia chrysantha* Sieb. & Zucc, China (Hangzhou), *Strijk 4048* (BGT), MZ901897*, MZ782418*, *Eriolaena spectabilis* (DC.) Planch. ex Mast., China (Yunnan), *Strijk 4474* (BGT), MZ901898*, MZ782419*, *Firmiana major* (W.W.Sm.) Hand.-Mazz., China (Sichuan), –, MG229069, AF460186, *Gossypium barbadense* L., Japan (Kyoto), –, NC008641, GU935141, *Gossypium herbaceum* L., –, JF317353, GQ166629, *Gossypium hirsutum* L., Coker310FR –, NC007944, KC404827, *Gossypium turneri* Fryxell, –, NC026835, U12726, *Lavatera punctata* All., Italy, *Wieringa 5952* (WAG8003579), MZ901899*, MZ782447*, *Microcos paniculata* L., China (Yunnan), *Strijk 4575* (BGT), MZ901900*, MZ782448*, *Peddiea africana* Harv., Cameroon, *Wieringa 5900* (WAG8003696), MZ901901*, MZ782454*, *Pterospermum kingtungense* C.Y.Wu ex H.H.Hsue, China (Yunnan), *Strijk 4497* (BGT), MZ901902*, MZ782455*, *Reevesia thyrsoides* Lindl., China (Yunnan), *Strijk 4519* (BGT), MZ901903*, MZ782456*, *Scaphopetalum longipedunculatum* Mast., Gabon, *Sosef 2028* (WAG8004058), MZ901904*, MZ782457*, *Sterculia lanceifolia* Roxb., China (Yunnan), *Strijk 4602* (BGT), MZ901905*, MZ782511*, *Sterculia lanceolata* Cav., China (Yunnan), *Strijk 4537* (BGT), MZ901906*, MZ782512*, *Sterculia micrantha* Chun & H.H.Hsue, China (Yunnan), *Strijk 4482* (BGT), MZ901907*, MZ782513*, *Theobroma cacao* L., –, NC014676, JQ228377, *Theobroma cacao* f. *pentagonum* (Bernoulli) Cuatrec., China (Yunnan), *Strijk 4534* (BGT), MZ901908*, MZ782514*, *Theobroma sp.*, China (Yunnan), *Strijk 4535* (BGT), MZ901909*, MZ782515*, *Tilia amurensis* Kom., China (Heilongjiang), –, NC028588, KF445432, *Tilia mandshurica* Rupr. & Maxim., China (Heilongjiang), –, KT894773, KX062302, *Tilia oliveri* Szyszyl., China (Chongqing), –, NC028590, KF897525, *Urena lobata* L., Gabon, *Wieringa 8287* (WAG8001343), MZ901911*, MZ782517*, *Urena lobata* L., China (Yunnan), *Strijk 4494* (BGT), MZ901910*, MZ782516*.