



**HAL**  
open science

## Growing energy cover crops in double cropping: which ecosystem services or environmental impacts associated?

Camille Launay, Louise Hermet, Sylvain Frédéric, Sabine Houot, Julie Constantin

### ► To cite this version:

Camille Launay, Louise Hermet, Sylvain Frédéric, Sabine Houot, Julie Constantin. Growing energy cover crops in double cropping: which ecosystem services or environmental impacts associated?. XVII. Congress of the European Society for Agronomy, Aug 2022, Potsdam, Germany. pp.67-68. hal-03937907

**HAL Id: hal-03937907**

**<https://hal.inrae.fr/hal-03937907>**

Submitted on 13 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# XVII. Congress of the European Society for Agronomy

August 29 – September 2, 2022  
Potsdam, Germany



Diversification & Digitalisation  
Trends that Shape Future Agriculture

## BOOK OF ABSTRACTS

# **Book of abstracts: XVII. Congress of the European Society for Agronomy**

August 29 – September 02, 2022 – Potsdam, Germany

## **Congress chair:**

**Claas Nendel**, ZALF and University of Potsdam, Germany

## **Scientific committee:**

**Pietro Barbieri**, Bordeaux Sciences Agro, France

**Benjamin Dumont**, University of Liège, Belgium

**Nadja El Benni**, Agroscope, Switzerland

**Victoria Gonzalez-Dugo**, Institute of Sustainable Agriculture, Spain

**Kathrin Grahmann**, ZALF Germany

**Hans-Peter Kaul**, BOKU, Austria

**Zita Kriaučiūnienė**, Vytautas Magnus University, Lithuania

**Edith LeCadre-Barthélémy**, Institut Agro of AGROCAMPUS OUEST, Rennes, France

**Evelin Loit**, EMÜ, Estonia

**James Lowenberg-DeBoer**, Harper Adams University, Newport, Shropshire, UK

**Jørgen E. Olesen**, Aarhus University, Denmark

**Ingrid Öborn**, SLU, Sweden

**Michele Perniola**, University of Basilicata, Italy

**Daniel Plaza-Bonilla**, University of Lleida-Agrotecnio, Spain

**Vera Potopová**, ČZU, Czech Republic

**Xavier Reboud**, INRAE, Dijon, France

**Moritz Reckling**, ZALF, Germany

**Francelino Rodrigues**, Lincoln Agritech Ltd, New Zealand

**Mariana Rufino**, Lancaster University, UK

**Julie Ryschawy**, INRAE, France

**Stanislaw Samborski**, Warsaw University of Life Sciences, Poland

**Henrik G. Smith**, University Lund, Sweden

**Wopke van der Werf**, WUR, the Netherlands

**Sebastian Vogel**, ATB, Germany

**Christine Watson**, SRUC, United Kingdom

**Željka Zgorelec**, University of Zagreb, Croatia

## **Layout concept:**

Leibniz Centre for Agricultural Landscape Research (ZALF)

This book of abstracts will be published only electronically: [www.esa-congress-potsdam2022.de](http://www.esa-congress-potsdam2022.de)

Leibniz Centre for Agricultural Landscape Research (ZALF)

Eberswalder Straße 84 | 15374 Müncheberg, Germany

T +49 (0)33432 | 82 200

E [zalf@zalf.de](mailto:zalf@zalf.de)

W [www.leibniz-zalf.de](http://www.leibniz-zalf.de)

© ZALF 2022



## **Growing energy cover crops in double cropping: which ecosystem services or environmental impacts associated?**

Camille Launay<sup>1</sup>; Louise Hermet<sup>2</sup>; Sylvain Frédéric<sup>3</sup>; Sabine Houot<sup>4</sup>; Dr. Julie Constantin<sup>1</sup>

<sup>1</sup> Université de Toulouse, INRAE, UMR AGIR; <sup>2</sup> ARVALIS - Institut du végétal; <sup>3</sup> GRDF; <sup>4</sup> ECOSYS, INRAE, AgroParisTech, Université Paris-Saclay

Key-words: anaerobic digestion, nutrient recycling, carbon storage, modeling

Energy cover crops are currently being developed in France and Europe to provide biogas through anaerobic digestion and thus contribute to renewable energy production. These crops have the double advantage of (1) not competing with food crops, as they do not occupy the soil at the same time, and (2) providing the ecosystem services expected from a cover crop (e.g. reduction of nitrate leaching, reduction of erosion, carbon storage).

However, if agronomic issues related to their management and methanogenic potential have been addressed (Molinuevo-Salces et al. 2013\*; Wannasek et al. 2019\*\*), issues related to their environmental impact have been very little explored, particularly coupled with digestate return to the field. We focused on ecosystem services related to nitrogen recycling (reduction of nitrate leaching, reduction of nitrous oxide and ammonia emissions) and soil carbon storage. We also studied whether energy cover crops could compete with food production by inducing nitrogen or water stress on the following main crops, which could lead to yield loss.

We conducted a two-year field experiment in southwestern France. Two main crop sequences were implemented: wheat [*Triticum turgidum* subsp. *Durum* (Desf.) Husn.] - Winter barley [*Hordeum vulgare* L.] - winter peas [*Pisum sativum* L.] and wheat [*Triticum turgidum* subsp. *Durum* (Desf.) Husn.] – Sunflower [*Helianthus annuus* L.] - grain sorghum [*Sorghum bicolor* (L.)]. Summer energy cover crops (sorghum [*Sorghum bicolor* (L.) Moench] – vetch [*Vicia benghalensis* L.] mixture) were grown on the short fallow period of the first rotation and winter energy cover crops (rye [*Secale cereale* L.] – fababean [*Vicia faba* L.] mixture) on the long fallow period of the second crop sequence. These energy cover crops were exported from the plot at harvest to produce biogas and fertilized with digestate produced during anaerobic digestion. They were compared to the same unfertilized and non-exported cover crop and to a soil left bare. For the long fallow period, the comparison was also made with a white mustard, growing over the autumn and plowed into the soil earlier. The biomass and yield of cover and main crops, ammonia volatilization and soil water and nitrogen content were regularly monitored. This experiment was then simulated with the STICS crop model in order to assess unmeasured variables (i.e. nitrate leached, nitrous oxide emissions carbon stored over 30 years, crop stress indices) and to extend the simulated period from 2020 to 2050.

About competition with food crops, our study shows that in 2021, energy cover crops did not decrease the yield of the main crops. Repeating the trial over several climatic years using the model will allow us to confirm whether this result is isolated or generic. Secondly, with regard



to ecosystem services, we observed a decrease in leaching during long cover crop period that was lower than with fast-growing mustard, and a decrease in nitrous oxide emissions compared to bare soil. Conversely, spreading digestate increased ammonia emissions, particularly when spread on the crop and not buried in the soil as done before sowing. According to biomass produced and incorporated to the soil (including digestate) in the different treatments, we expect a similar carbon storage that would be verified through simulation results. A greenhouse gas (GHG) balance will allow us to assess if the reduction of direct and indirect N<sub>2</sub>O emissions and increased in carbon storage compensate for higher ammonia emissions. Finally, the GHG balance of these energy cover crops, if negative, and the assessment of potential other environmental impacts would make possible to ensure that the gas produced is renewable.

\* <https://doi.org/10.1016/j.biombioe.2013.10.008>

\*\* <https://doi.org/10.1016/j.eja.2019.125934>