



**HAL**  
open science

## A multispecies, intraspecific functional traits data set on fish species from the Bay of Biscay, France

Romane Rozanski, David Eme, Anne Boiron Leroy, Marta M Rufino, Camille Albouy

► **To cite this version:**

Romane Rozanski, David Eme, Anne Boiron Leroy, Marta M Rufino, Camille Albouy. A multispecies, intraspecific functional traits data set on fish species from the Bay of Biscay, France. *Ecology*, 2023, pp.e3924. 10.1002/ecy.3924 . hal-03946746

**HAL Id: hal-03946746**

**<https://hal.inrae.fr/hal-03946746>**

Submitted on 19 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**DATA PAPER**

# A multispecies, intraspecific functional traits data set on fish species from the Bay of Biscay, France

Romane Rozanski<sup>1,2,3</sup>  | David Eme<sup>2,4</sup>  | Anne Boiron Leroy<sup>5</sup>  |  
 Marta M. Rufino<sup>6,7</sup>  | Camille Albouy<sup>1,2,3</sup> 

<sup>1</sup>Ecosystems and Landscape Evolution, Department of Environmental Systems Science, Institute of Terrestrial Ecosystems, ETH Zürich, Zürich, Switzerland

<sup>2</sup>DECOD (Ecosystem Dynamics and Sustainability), IFREMER, INRAE, Institut Agro – Agrocampus Ouest, Nantes, France

<sup>3</sup>Unit of Land Change Science, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

<sup>4</sup>RiverLY Research Unit, National Research Institute for Agriculture Food and Environment (INRAE), Villeurbanne, France

<sup>5</sup>PMMLT, IFREMER Station de La Tremblade, La Tremblade, France

<sup>6</sup>Division of Modeling and Management of Fisheries Resources, IPMA – Portuguese Institute for the Sea and the Atmosphere, Lisbon, Portugal

<sup>7</sup>Faculty of Sciences, CEAUL, Centre of Statistics and its Applications of the University of Lisbon, Lisbon, Portugal

**Correspondence**

Romane Rozanski  
 Email: rrozanski@ethz.ch

**Handling Editor:** William K. Michener

**Abstract**

The global biodiversity crisis due to anthropogenic pressures jeopardizes marine ecosystem functioning and services. Community responses to these environmental changes can be assessed through functional diversity, a biodiversity component related to organism–environment interactions, and estimated through biological traits related to organism functions (locomotion, feeding mode, and reproduction). Fish play a key role in marine systems functioning and supply proteins for billions of humans worldwide, yet most of the knowledge is limited to several commercial species and little is known about the intraspecific variability of their functional traits. The data provided here consist of 867 records of individuals from 85 species of ray-finned (Actinopterygii) and cartilaginous (Chondrichthyes) fish sampled in the Bay of Biscay (Atlantic, France) between autumn 2017 and 2019. We provided for each individual the taxonomic classification, 16 ecomorphological measures (5 directly made on fresh individuals and 11 realized using individual pictures) that were converted into nine ecomorphological traits classically documented in the literature (biomass, protrusion, oral gape shape, surface and position, eye size and position, body transversal shape and surface, pectoral fin position and caudal peduncle throttling) and eight life history traits obtained from FishBase (maximum length, average depth, depth range, trophic level, reproduction mode, fertilization mode, parental care, vertical position in the water column). These traits document several functions such as dispersion, feeding mode, habitat use, position in the food web, and reproduction. To improve the development of new traits, we provided a picture of each individual with an ROI file containing the different morpho-anatomical measures made using “ImageJ” software and an R function to extract them. In addition, we provided the metadata from each sampling site (years, dates, stations, sampling hours, strata, gears, latitudes, longitudes, and depths) and environmental variables measured in situ (conductivity, salinity, water temperature, water density, and air temperature). This data set accounting for the intraspecific variability

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Ecology* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

among 85 fish species is of interest to better understand the effects of environmental forcing in a global change context as in the Bay of Biscay, a highly fished transition zone harboring mixed assemblages of boreal, temperate, and subtropical fish species that are susceptible to display variability in functional trait to adapt to changing conditions. The data set is freely available without copyright restrictions; users should cite this paper in research products (publications, presentations, reports, etc.) derived from the data set.

#### KEYWORDS

Actinopterygii, Atlantic Ocean, ecomorphological traits, functional diversity, intraspecific traits variability, life history traits

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### DATA AVAILABILITY STATEMENT

A compressed file entitled “Fish\_pictures\_&\_ROI\_files.zip” containing the 867 pictures of fish individuals (JPEG format) and the 867 ROI files is too large to be included as Supporting Information and these items are only available in Figshare at <https://doi.org/10.6084/m9.figshare.19410818.v2>. Remaining data and code are provided as Supporting Information and are also available in Figshare at <https://doi.org/10.6084/m9.figshare.19410818.v2>.

#### ORCID

Romane Rozanski  <https://orcid.org/0000-0002-5558-6622>

David Eme  <https://orcid.org/0000-0001-8790-0412>

Anne Boiron Leroy  <https://orcid.org/0000-0002-9557-9611>

Marta M. Rufino  <https://orcid.org/0000-0002-0734-7491>

Camille Albouy  <https://orcid.org/0000-0003-1629-2389>

#### SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Rozanski, Romane, David Eme, Anne Boiron Leroy, Marta M. Rufino, and Camille Albouy. 2022. “A Multispecies, Intraspecific Functional Traits Data Set on Fish Species from the Bay of Biscay, France.” *Ecology* e3924. <https://doi.org/10.1002/ecy.3924>