



**HAL**  
open science

## Transgenerational epigenetics in quail

Chloé Cerutti, Sophie Leroux, Paul Terzian, Joanna Lledo, David Gourichon, Rémy-Félix Serre, Céline Vandecasteele, Christophe Klopp, Carole Iampietro, Christine Gaspin, et al.

► **To cite this version:**

Chloé Cerutti, Sophie Leroux, Paul Terzian, Joanna Lledo, David Gourichon, et al.. Transgenerational epigenetics in quail: whole genome DNA methylation analysis. Genotoul Biostat Bioinfo Day, Dec 2022, Castanet-Tolosan, France. hal-03957296

**HAL Id: hal-03957296**

**<https://hal.inrae.fr/hal-03957296>**

Submitted on 26 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Transgenerational epigenetics in quail : whole genome DNA methylation analysis

Chloé Cerutti<sup>1</sup>, Sophie Leroux<sup>1</sup>, Paul Terzian<sup>2</sup>, Joanna Lledo<sup>3</sup>, David Gourichon<sup>4</sup>, Rémy-Félix Serre<sup>3</sup>, Céline Vandecasteele<sup>3</sup>, Christophe Klopp<sup>2</sup>, Carole Iampietro<sup>3</sup>, Christine Gaspin<sup>2</sup>, Denis Milan<sup>1</sup>, Cécile Donnadiou<sup>3</sup>, Frédérique Pitel<sup>1</sup> and Guillaume Devailly<sup>1</sup>

<sup>1</sup> GenPhySE, Université de Toulouse, INRAE, ENVT, 31326, Castanet Tolosan, France  
{chloe.cerutti, sophie.leroux1, denis.milan, frederique.pitel, guillaume.devailly}@inrae.fr

<sup>2</sup> MIAT, UR875 INRAE, 31326, Castanet-Tolosan, France

{paul.terzian, christophe.klopp, christine.gaspin}@inrae.fr

<sup>3</sup> GeT-PlaGe - Génome et Transcriptome - Plateforme Génomique, US1426 GET - INRAE, 31326, Castanet-Tolosan, France

{joanna.lledo, remy-felix.serre, celine.vandecasteele, carole.iampietro, cecile.donnadiou}@inrae.fr

<sup>4</sup> PEAT, UE1295 - INRAE Centre Val de Loire, 37380, Nouzilly, France

{david.gourichon}@inrae.fr

## Abstract

Prenatal environment influence on adult phenotype is partially mediated by epigenetics phenomena. One study highlighted significant effects of the *in-ovo* injection of endocrine disruptors or DNA methyltransferase inhibitor on quail development, significantly reducing their weight [1]. Recently, an increasing number of studies highlighted the transmission of epigenetics marks between generations following an environmental exposure. However, there is much debate about their transmission beyond the exposed individuals. Recent studies revealed that non-genetics inheritance was probably present in avian species. In one of them [2], fertilized eggs were divided into two groups: one group injected with an endocrine disruptor, Genistein, and a non-injected control group. After three generations without any other injection, several traits were impacted by the ancestor treatment including reproduction and behavior. This pilot study highlighted the potential existence of transgenerational transmission of environmental effects in quails. To better understand the transgenerational transmission of these environmental effects, we generated DNA methylation data from blood samples of the third generation using WGBS (Whole Genome Bisulfite Sequencing) and ONT (Oxford Nanopore Technologies). To analyse the DNA methylation states, we developed a bioinformatics pipeline in order to detect differentially methylated cytosines (DMCs) due to transgenerational phenomena. We detected thousands of DMCs between both groups (Table 1). In addition to these analyses, these data allowed us to perform a comparative analysis between the WGBS and ONT sequencing technologies (Fig. 1).

	measured CpGs sites	epiline DMCs detected
WGBS	5 881 815	12 872
ONT	7 537 007	22 325

Table 1. Number of measured CpGs sites and epiline DMCs detected from WGBS and ONT methylation data. A CpG site refers to a dinucleotide CG.




Figure 1. Heatmap of methylation rate of 10,000 common epigenetic differentially methylated CpGs (DMCs) detected from WGBS and ONT. The red and blue colors refer to the epi- and epi+ groups, respectively. The violet and green colors refer to the female (F) and male (M) quails, respectively. The diverging color gradient refers to a low (blue) or high (red) methylation rate.

## Acknowledgements

This work was supported by FEDER (European Funding for Regional Economic Development), the PhD scholarship to CC from Région Occitanie and INRAE Animal Genetics Division. We are grateful to the genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, <https://doi.org/10.15454/1.5572369328961167E12>) for providing computing and storage resources.

## References

- [1] C. Cerutti, S. Leroux, D. Gourichon, Y. Labrune, I. David, T. Zerjal, V. Coustham, G. Devailly, F. Pitel. *Short communication: Effects of in-ovo injection of endocrine disruptors and methyltransferase inhibitor on quail growth and egg-laying performances*. Vol. 16, pp. :1751-7311, 2022.
- [2] S. Leroux, D. Gourichon, C. Leterrier, Y. Labrune, V. Coustham, S. Rivière, T. Zerjal, J.L. Coville, M. Morisson, F. Minvielle, F. Pitel. *Embryonic environment and transgenerational effects in quail*. Vol. 49, pp. :1297-9686, 2017.