



HAL
open science

Effects of pathogen reproduction system on the evolutionary and epidemiological control provided by deployment strategies for two major resistance genes in agricultural landscapes

Marta Zaffaroni, Loup Rimbaud, Jean-françois Rey, Julien Papaïx, Frédéric Fabre

► To cite this version:

Marta Zaffaroni, Loup Rimbaud, Jean-françois Rey, Julien Papaïx, Frédéric Fabre. Effects of pathogen reproduction system on the evolutionary and epidemiological control provided by deployment strategies for two major resistance genes in agricultural landscapes. *Evolutionary Applications*, 2024, 17 (1), pp.e13627. 10.1111/eva.13627 . hal-03984706v2

HAL Id: hal-03984706

<https://hal.inrae.fr/hal-03984706v2>

Submitted on 31 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Effects of pathogen reproduction system on the evolutionary and epidemiological control provided by deployment strategies for two major resistance genes in agricultural landscapes

Marta Zaffaroni^{1,2}  | Loup Rimbaud³  | Jean-François Rey²  | Julien Papaix²  | Frédéric Fabre¹ 

¹INRAE, Bordeaux Sciences Agro, SAVE, Villenave d'Ornon, France

²INRAE, BioSP, Avignon, France

³INRAE, Pathologie Végétale, Montfavet, France

Correspondence

Marta Zaffaroni, INRAE, Bordeaux Sciences Agro, SAVE, Villenave d'Ornon 33882, France.

Email: marta.zaffaroni@inrae.fr

Funding information

ANR COMBINE, Grant/Award Number: ANR-22-CE32-0004; Ecophyto II APR Levier Territoriaux MEDEE, Grant/Award Number: No.SIREPA4621

Abstract

Resistant cultivars are of value for protecting crops from disease, but can be rapidly overcome by pathogens. Several strategies have been proposed to delay pathogen adaptation (*evolutionary control*), while maintaining effective protection (*epidemiological control*). Resistance genes can be (i) combined in the same cultivar (pyramiding), (ii) deployed in different cultivars sown in the same field (mixtures) or in different fields (mosaics), or (iii) alternated over time (rotations). The outcomes of these strategies have been investigated principally in pathogens displaying pure clonal reproduction, but many pathogens have at least one sexual event in their annual life cycles. Sexual reproduction may promote the emergence of superpathogens adapted to all the resistance genes deployed. Here, we improved the spatially explicit stochastic model *landsepi* to include pathogen sexual reproduction, and we used the improved model to investigate the effect of sexual reproduction on evolutionary and epidemiological outcomes across deployment strategies for two major resistance genes. Sexual reproduction favours the establishment of a superpathogen when single mutant pathogens are present together at a sufficiently high frequency, as in mosaic and mixture strategies. However, sexual reproduction did not affect the strategy recommendations for a wide range of mutation probabilities, associated fitness costs, and landscape organisations.

KEYWORDS

deployment strategy, disease control, durable resistance, evolutionary epidemiology, sexual reproduction, simulation modelling

1 | INTRODUCTION

The deployment of resistant cultivars in agricultural landscapes is a relatively low-input and cost-effective way to protect crops

from plant pathogens. However, resistant cultivars have often been rapidly overcome by pathogens, especially when a single resistant cultivar is widely cultivated over a large geographic area (García-Arenal & McDonald, 2003; McDonald & Linde, 2002;

Julien Papaix and Frédéric Fabre contributed equally to the article.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Evolutionary Applications* published by John Wiley & Sons Ltd.

Parlevliet, 2002). Ultimately, this may result in recurrent cycles of resistance deployment followed by rapid pathogen adaptation, often described as boom-and-bust cycles (McDonald & Linde, 2002). Several strategies have been proposed to promote a more durable management of resistant cultivars. These strategies involve increasing cultivated host genetic diversity (McDonald, 2010, 2014; Zhan et al., 2015) with the aim of confronting pathogens with eco-evolutionary challenges to prevent or delay their adaptation to plant resistance (evolutionary control), while maintaining effective disease protection (epidemiological control). Plant breeders can stack resistance sources in the same cultivar by pyramiding (Fuchs, 2017; McDonald & Linde, 2002), or farmers can alternate resistances over time by rotating cultivars in the same field (Curl, 1963). Host genetic diversity can also be introduced spatially. Resistant cultivars can be combined within the same field in cultivar mixtures (Mundt, 2002; Wolfe, 1985) or cultivated in different fields in landscape mosaics (Burdon et al., 2014; Zhan et al., 2015).

Given the multitude of deployment options, it is not straightforward to compare deployment strategies for identification of the optimal deployment strategy in a given epidemiological context. In addition, evolutionary and epidemiological control may not necessarily be correlated: any strategy designed to control the emergence of resistance-adapted pathogens in agro-ecosystems may potentially come into conflict with epidemiological control (Burdon et al., 2014; Papaix et al., 2018; Rimbaud, Papaix, Barrett, et al., 2018). Finally, particularly for airborne plant pathogens, which often disperse over large distances, deployment strategies are more likely to be effective if implemented across landscapes at large spatial scales, rendering experimental testing logistically demanding (but see Djian-Caporalino et al., 2014; Koller et al., 2018; Lohaus et al., 2000; Zhu et al., 2000). In this context, many mathematical models have been developed to facilitate assessments of the variation of evolutionary and epidemiological outcomes across different resistance deployment strategies (reviewed by Rimbaud et al., 2021). These models have been used to unravel the effects of resistance deployment strategies on pathogen epidemiology and evolution and to compare these strategies in a given epidemiological context.

Most of the models reviewed by Rimbaud et al. (2021) include only selection and/or mutation as evolutionary forces. This approach is suitable for the simulation of pathogens with purely clonal reproduction systems. Under the hypothesis of a purely clonal reproduction system, new pathogen variants are either already present (possibly at low frequency) at the beginning of the simulated period, either introduced through migration or generated by mutation. However, some pathogens are not purely clonal and their life cycles include at least one sexual event per cropping season (mixed reproduction system), with some even reproducing exclusively by sexual means (purely sexual reproduction system). Of the 43 plant pathogens analysed by McDonald and Linde (2002), only 17 have exclusively clonal reproduction, the other 26 pathogens presenting at least one sexual reproduction event during their life cycle. The genetic recombination occurring during sexual reproduction can

efficiently create gene combinations that would be accessible only through sequential mutation events in a purely clonal reproduction system. Several authors have argued that pathogens with mixed reproduction system have the highest potential for evolving and breaking down the resistances deployed in agriculture (McDonald & Linde, 2002; Stam & McDonald, 2018). Genetic recombination first creates many new variants of the pathogen (Halkett et al., 2005; Tibayrenc & Ayala, 2002). The populations of the fittest variants then expand rapidly through clonal reproduction, potentially breaking down the resistance, (i.e. increasing the frequency of pathogen strains adapted to the resistance genes present). Genetic recombination can, therefore, have a major impact on the evolutionary and epidemiological outcomes of resistance deployment strategies (Arenas et al., 2018; Stam & McDonald, 2018). It has been shown that even low rates of recombination in pests and pathogens have profound implications for policies concerning drug and pesticide resistance (Halkett et al., 2005). Similarly, by mixing the genotypes of parental individuals, recombination can favour the emergence of the generalist superpathogens able to overcome pyramided cultivars (McDonald & Linde, 2002; Uecker, 2017). However, the ability of recombination to favour the emergence of superpathogens also depends on subtle interactions between mutation and recombination rates on the one hand, and pathogen population size on the other (Althaus & Bonhoeffer, 2005). Indeed, recombination can generate variants accumulating infectivities, but it can also break down such genetic combinations (Hadany & Beker, 2003).

Despite the potentially major impact of the pathogen reproduction system on the epidemiological and evolutionary control provided by resistance deployment strategies, this impact has been little studied and is poorly understood (Mundt, 2018). Genetic recombination is considered in only three (Cr  t   et al., 2020; Sapoukhina et al., 2009; Xu, 2012) of the 69 models reviewed by Rimbaud et al. (2021) and in a recent study by Saubin et al. (2021). These studies considered pathogens with mixed reproduction systems, but they did not compare purely clonal reproduction with mixed reproduction systems, all other things being equal. It is, therefore, difficult to assess the impact of reproduction system on the epidemiological and evolutionary control provided by resistance deployment strategies from the data currently available. In addition, these works focused on just one or two resistance deployment strategies, preventing a global assessment of all possible spatiotemporal deployment options. They highlighted the role of the fitness cost of resistance in superpathogen persistence (Xu, 2012), and in the efficacy of rotation (Cr  t   et al., 2020) and mixture (Sapoukhina et al., 2009; Xu, 2012) strategies. In addition, Saubin et al. (2021) assessed the impact of ploidy on resistance durability, revealing that resistance durability was greater, but more variable, for diploid pathogens.

Here, we investigated the effect of pathogen sexual reproduction on the evolutionary and epidemiological control achieved with four main categories of deployment strategies (rotation, pyramiding, mixture and mosaic). We adapted the *landsepi* model (Rimbaud, Papaix, Rey, et al., 2018), which simulates the spread of epidemics

across an agricultural landscape and the evolution of a pathogen in response to the deployment of host resistance, to include pathogen sexual reproduction. We then used this model to compare the resistance deployment strategies considered for situations in which two major resistance genes conferring immunity are deployed. The new model is flexible enough to vary resistance deployment strategy and pathogen life cycle, making it possible to compare pathogens with different reproduction systems (purely clonal vs. mixed). We parameterised the model to simulate grapevine downy mildew, which is caused by the oomycete *Plasmopara viticola*. However, our general conclusions are likely to have broader implications to other pathosystems.

2 | MATERIALS AND METHODS

2.1 | Model overview

The model used in this study is an adapted version of that presented by Rimbaud, Papaix, Rey, et al. (2018), which simulates the clonal reproduction, spread and evolution of a pathogen in an agricultural landscape over multiple cropping seasons. Here, we introduce between-season sexual reproduction to address the issue of pathogens with mixed reproduction systems. Multiple clonal reproduction events occur during the life cycle of these pathogens, with a final sexual reproduction event at the end of the host cropping season. We split the modelled cropping season into two different time periods: (i) within the cropping season, when multiple clonal reproduction events take place and (ii) the period between cropping seasons, when a single sexual reproduction event may take place. Below, we describe only the major changes between cropping seasons, the modifications within cropping seasons being only minor. The entire model is described in Note S2.

2.2 | Landscape and resistance deployment strategies

We considered agricultural landscapes randomly generated with a T-tessellation algorithm (Papaix et al., 2014) in which four cultivars were randomly allocated to fields: a susceptible cultivar (SC) initially infected with a pathogen not adapted to any resistance, two resistant cultivars, each carrying a single major resistance gene (RC_1 and RC_2), and one resistant cultivar carrying both resistance genes (RC_{12}). The cropping ratio φ represents the proportion of fields in the landscape (hereafter the candidate fields) cultivated with resistant cultivars, which are planted according to one of the following strategies:

1. Mosaics: RC_1 and RC_2 are cultivated in the equal proportions of the candidate fields;
2. Mixtures: both RC_1 and RC_2 are cultivated in all the candidate fields, in equal proportions within each field;

3. Rotations: RC_1 and RC_2 are cultivated alternately in candidate fields during three consecutive cropping seasons each;
4. Pyramiding: RC_{12} is cultivated in all candidate fields.

The remaining proportion $1-\varphi$ of fields are planted with the susceptible cultivar.

A cultivar carrying a major resistance gene is assumed to be immune to disease (i.e. pathogen infection probability is equal to 0), unless the pathogen has acquired the corresponding infectivity gene. At the beginning of each simulation, the pathogen population is composed exclusively of non-adapted pathogens (denoted "WT" here for "wild-type"). During the simulation, a WT can acquire infectivity gene $g \in \{1, 2\}$ through a single mutation, with a probability τ , or, alternatively, through sexual reproduction with another individual pathogen carrying such an infectivity gene. Pathogen adaptation leads to resistance breakdown, i.e., a complete restoration of pathogen infectivity on resistant hosts. The acquisition of infectivity may be penalised by a fitness cost θ (Brown, 2015; Laine & Barrès, 2013; Thrall & Burdon, 2003). This fitness cost is associated in the model to a lower infection probability for mutant pathogens on hosts not carrying the corresponding resistance gene. In other words, mutant pathogens pay a fitness cost for their unnecessary virulences on a given host. For the superpathogen, the fitness costs are multiplicative on the susceptible cultivar. Note that we assumed the same mutation probability and fitness cost for all major genes and infectivity genes, respectively. Here, a pathogen genotype is represented by a set of binary variables indicating whether it carries infectivity genes able to overcome cultivar resistance genes. There are four possible pathogen genotypes: wild-type, unable to break down the resistance conferred by any resistance gene ("00"), single mutant " SM_1 " (or " SM_2 "), able to break down to the first (or second) resistance gene ("10" and "01", respectively), and superpathogen "SP", able to break down both resistance genes ("11"). The relative infection probabilities of these pathogens on the different cultivars are summarised in Table 1.

TABLE 1 Plant-pathogen interaction matrix.

	Host genotype v			
	SC	RC_1	RC_2	RC_{12}
Pathogen genotypes p				
WT	1	0	0	0
SM_1	$1-\theta$	1	0	0
SM_2	$1-\theta$	0	1	0
SP	$(1-\theta)^2$	$1-\theta$	$1-\theta$	1

Note: The matrix gives the coefficient by which the infection probability is multiplied. The value of this coefficient reflects the relative infection probabilities for the wild-type (WT) and adapted (single mutants SM_1 and SM_2 , and SP) pathogen genotypes on the susceptible (SC) and resistant cultivars carrying a single major resistance gene (RC_1 and RC_2), or their combination (RC_{12}). θ is the fitness cost of infectivity with respect to the major resistance genes considered. This matrix corresponds to a simplified version of a more general interaction matrix (Table S2 in Note S7), for which simulations were also performed.

2.3 | Demogenetic dynamics within the cropping season

The demogenetic dynamics of the host–pathogen interaction within the cropping season are based on a compartmental model with a discrete time step, schematically reported in Figure 1. Below, $H_{i,v,t}$, $L_{i,v,p,t}$, $I_{i,v,p,t}$, $R_{i,v,p,t}$, and $P_{i,p,t}$ denote the numbers of healthy, latent, infectious and removed individuals, and of pathogen propagules, respectively, in the field $i=1, \dots, J$, for cultivar $v=1, \dots, V$, pathogen genotype $p=1, \dots, P$ at time step $t=1, \dots, T \times Y$ (Y is the number of cropping seasons and T the number of time steps per season). Note that, in this model, an "individual" is defined as a given amount of plant tissue, and is referred to as a "host" hereafter for the sake of simplicity. At the beginning of the cropping season, healthy hosts are contaminated with the primary inoculum generated at the end of the previous cropping season.

2.4 | Demogenetic dynamics between cropping seasons

The demogenetic dynamics of the host–pathogen interaction between cropping seasons is presented schematically in Figure 1. At the end of the cropping season, the crop is harvested and the leaves of the host plants fall to the ground, imposing a potential bottleneck on the pathogen population before the start of the next cropping season. The remaining hosts produce clonal or sexual propagules. Clonal propagules can mutate in the same way as they do during the cropping season. The production of propagules through sexual reproduction and the possibility of genetic recombination are detailed in the Section 2.4.1. The propagules produced during the period between cropping seasons, whether clonal or sexual, are uniformly released throughout the following cropping season, constituting the primary inoculum.




FIGURE 1 Model overview. Within-cropping season dynamics: healthy hosts can be contaminated by pathogen propagules (produced both at the end of the previous cropping season and within the current cropping season) and may become infected. Following a latent period, infectious hosts start producing propagules through clonal reproduction. These propagules may mutate and disperse across the landscape. At the end of the infectious period, infected hosts become epidemiologically inactive. Qualitative resistance prevents the infection of contaminated hosts, i.e. their transition to the latently infected state. Green boxes indicate healthy hosts contributing to host growth, as opposed to diseased plants (i.e. symptomatic, red boxes) or plants with latent infections (dark blue box). Between-cropping season dynamics: at the end of each cropping season, pathogens experience a bottleneck during the off-season period, and propagules are then produced (by clonal or sexual reproduction). Clonal propagules may mutate, whereas genetic recombination may occur during sexual reproduction. Propagules produced between host cropping seasons are gradually released during the following host cropping season. The parameters associated with epidemiological processes are indicated in grey and detailed in Table 2. The distributions used to simulate stochasticity in model transitions are indicated in red; \mathcal{B} : binomial, Γ : gamma, \mathcal{P} : Poisson, \mathcal{M} : multinomial, \mathcal{U} : uniform, \mathcal{B}_{ern} : Bernoulli. Host logistic growth is deterministic. The model's assumptions and equations are described in Notes S1 and S2.

2.4.1 | Pathogen sexual reproduction

We considered in the study heterothallic plant pathogen species that reproduce sexually only when mycelia of two mating types grow together in intimate proximity (Cohen & Rubin, 2012). Accordingly, sexual reproduction can only take place between lesions located on the same plant. Therefore, we assumed that in field i , the pool of infectious hosts associated with the same cultivar v undergoes sexual reproduction. Two parental infectious hosts, infected with pathogens Par_1 and Par_2 , respectively, are randomly sampled without replacement from the pool of infectious hosts. The $c = \{\text{Par}_1; \text{Par}_2\}$ pair produces $P_{v,c}^{\text{sex}}$ propagules, drawn from a Poisson distribution in which the expectation is the sum of the number r_{exp} of propagules produced by each of the parental infectious hosts:

$$P_{v,c}^{\text{sex}} \sim \text{Poisson}(2 \times r_{\text{exp}}) \quad (1)$$

The genotype of each propagule is then retrieved from the parental genotypes: the genotype at every locus g is randomly sampled from one of the two parents $\{\text{Par}_1; \text{Par}_2\}$. For example, assuming that parental infection Par_1 provides infectivity genes against resistance gene $g=1$ (corresponding to genotype "10") and parental infection Par_2 provides infectivity genes effective against resistance $g=2$ (genotype "01"), the resulting propagule genotype may be the same as that of one of the two parents (with probability 0.5), a SP genotype "11" (with probability 0.25), or a WT genotype "00" (with probability 0.25). This process is iterated for all the pairs $c = 1, \dots, C$ of infectious hosts associated with all the cultivars $v = 1, \dots, V$ in a given field i , resulting in a total number of sexual propagules:

$$P_i^{\text{sex}} = \sum_{v=1}^V \sum_{c=1}^C P_{v,c}^{\text{sex}} \quad (2)$$

2.5 | Propagule dispersal

Clonal and sexual propagules disperse similarly (no dispersal dimorphism) within the landscape according to a power-law dispersal kernel.

2.6 | Simulation plan and model outputs

2.6.1 | Model parameterisation for *Plasmopara viticola*

We parameterised the model to simulate epidemics of *Plasmopara viticola*, the causal agent of grapevine downy mildew, which has a mixed reproduction system (Gessler et al., 2011; Wong et al., 2001). Downy mildew is a real threat to grapevines in all vine-growing areas of the world, causing significant yield losses and leading to a massive use of pesticides (Gessler et al., 2011). In recent years, breeders have been developing programs for breeding resistance to grapevine

downy mildew, resulting in the creation of several resistant varieties, with the aim of lowering rates of fungicide application on grapevines. However, *P. viticola* has already been shown to have a high evolutionary potential, as demonstrated by the rapid emergence of fungicide resistance (Blum et al., 2010; Chen et al., 2007) and the breakdown of some of the resistances deployed (Delmas et al., 2016; Paineau et al., 2022; Peressotti et al., 2010). All the model parameters used in the simulations are listed in Table 2.

2.6.2 | Simulation plan

The model is used to assess evolutionary and epidemiological outputs for different deployment strategies. In addition to the four resistance deployment strategies considered (mosaic, mixture, rotation, pyramiding), we varied the cropping ratio of fields where resistance is deployed (φ , five values). We simulated different pathogen evolutionary potentials, by varying the mutation probability (τ , two levels) and the fitness cost (θ , three values). We explored the effect of the pathogen reproduction system by either having the pathogen reproduce sexually at the end of the cropping season (mixed reproduction system) or having no sexual reproduction event (purely clonal reproduction system). The abovementioned factors were explored with a complete factorial design of 240 parameter combinations (Table 2). Simulations were also performed with five different landscape structures (with about 155 fields and a total area of $2 \times 2 \text{ km}^2$, see Figure S11) and 50 replications in each landscape structure, resulting in a total of 250 replicates per parameter combination. The whole numerical design represents a total of 60,000 simulations. Each simulation was run for 50 cropping seasons of 120 days each. Trial simulations showed that this simulation horizon was sufficiently long to differentiate between deployment strategies in terms of their evolutionary and epidemiological performances. The simulations have been performed using the R package *landsepi* (v1.2.4, Rimbaud et al., 2022).

2.6.3 | Model outputs

At the end of a simulation run, the results were evaluated by considering evolutionary and epidemiological outputs. For evolutionary outputs, we determined the time point at which the generalist superpathogen SP was established in the resistant host population. We first studied SP establishment by defining E_{SP} a binary variable set to 1 if the SP becomes established before the end of a simulation run and 0 otherwise. Assuming that the SP became established, we then studied the time to establishment T_{SP} . This time corresponds to the time point at which the number of resistant host plants infected with SP exceeds a threshold above which extinction in a constant environment becomes unlikely. We also determined the time required for the two single mutants to become established (T_{SM_1} and T_{SM_2}). Finally, we monitored the size of the superpathogen population SP_{fi} and the maximum number of

TABLE 2 Summary of model parameters and numerical simulation plan (factors in bold are varied according to a complete factorial design).

Notation	Parameter	Value	Source
Simulation factors			
Y	Number of cropping seasons	50 years	Fixed
T	Number of time steps in a cropping season	120 days	Fixed
J	Number of fields in the landscape	[155; 154; 152; 153; 156]	Varied
V	Number of host cultivars	[2 ^a , 3 ^b]	Fixed
Initial conditions and seasonality (same value for all cultivars)			
C_v^0	Plantation host density of cultivar v (in pure crops)	1 m ⁻²	Fixed
C_v^{max}	Maximal host density of cultivar v (in pure crops)	20 m ⁻²	Fixed
δ_v	Host growth rate of cultivar v	0.1 day ⁻¹	[1]
Φ	Initial probability of infection of susceptible hosts	5 × 10 ⁻⁴	Fixed
λ	Off-season survival probability of pathogen spores	10 ⁻⁴	Fixed
Pathogen aggressiveness components			
e_{max}	Maximal expected infection probability	0.9	[2, 3]
Γ_{exp}	Expected latent period duration	7 days	See Note S3
Γ_{var}	Variance of the latent period duration	8 days	See Note S3
Y_{exp}	Expected infectious period duration	14 days	See Note S3
Y_{var}	Variance of infectious period duration	22 days	See Note S3
r_{exp}	Expected propagule production rate	2 day ⁻¹	See Note S3
Sexual reproduction			
R	Pathogen reproduction system	[Purely clonal, mixed]	Varied
p_{inh}	Probability of a sexual propagule inheriting the genotype at locus g from parent Par_1 genotype	0.5	Fixed
Pathogen dispersal			
$g(\cdot)$	Dispersal kernel	Power-law function	See Note S2
μ_{mean}	Mean dispersal distance	20 m	[4]
a	Scale parameter	5	Fixed
b	Width of the tail	3.5	[5, 6, 7]
Contamination of healthy hosts			
$\pi(\cdot)$	Contamination function	Sigmoid	See Note S2
σ	Related to the position of the inflection point	3	[4]
κ	Related to the position of the inflection point	5.33	[4]
Host-pathogen genetic interaction			
G	Total number of major genes	2	Fixed
τ	Mutation probability	[10⁻⁷; 10⁻⁴]^c	Varied
θ	Cost of infectivity	[0; 0.25; 0.5]^d	Varied
Landscape organisation			
	Resistance deployment strategy	Mixture; MOsaic; PYramiding; ROTation	Varied
α	Level of spatial aggregation	0	Fixed
φ	Cropping ratio of fields in which resistance is deployed	[0.17; 0.33; 0.5; 0.67; 0.83]	Varied

Source: [1] Bove and Rossi (2020), [2] Bove et al. (2019), [3] Boso and Kassemeyer (2008), [4] Rimbaud, Papaix, Rey, et al. (2018), [5] Frantzen and Van den Bosch (2000), [6] Papaix et al. (2014), [7] Grosdidier et al. (2018).

^aPyramiding.

^bMixture, mosaic, rotation.

^cCorresponding to a relatively low and relatively high mutation probability.

^dCorresponding to a null, relatively low and relatively high cost of infectivity.

heterogeneous parental pairs HP_{t^f} (i.e. parental pairs involving SM_1 and SM_2) in the landscape after the bottleneck. In a given field and for a given host cultivar, the maximum number of heterogeneous parental pairs was calculated as the minimum between the population size of SM_1 and SM_2 after harvest at t^f ; which gives, for the whole landscape: $HP_{t^f} = \sum_i^J \sum_v^V [\min(SM_{1;i,v,t^f}, SM_{2;i,v,t^f})]$. For epidemiological output, we assessed the area under the disease progress curve (AUDPC) to measure disease severity over the whole landscape, averaged across all the simulated cropping seasons. AUDPC is normalised by dividing by mean disease severity in a fully susceptible landscape; its value therefore ranges from 0 (i.e. no disease) to 1 (i.e. disease severity identical to that in a fully susceptible landscape).

2.7 | Statistical analysis

We first used a classification tree to determine how the factors of interest and their interactions affected the binary evolutionary output E_{SP} . We considered the following six factors as qualitative explanatory variables: resistance deployment strategy, cropping ratio, mutation probability and fitness cost of the infectivity genes, the pathogen reproduction system and landscape structure. In addition, for each combination of resistance deployment strategy, mutation probability, fitness cost and pathogen reproduction system, we fitted second-order polynomial regressions (or second-order logistic regressions) to assess the response of T_{SP} and AUDPC (or E_{SP}) to variations of cropping ratio. Note that fitting a second-order logistic regression was impossible for factor combinations that almost always or never led to SP establishment in the 250 replicates. In such cases, a second-order polynomial regression was fitted instead. Finally, for each combination of resistance deployment strategy, mutation probability, fitness cost, pathogen reproduction system and cropping ratio, we fitted local polynomial regressions to the temporal dynamics of the population of SP_{t^f} and HP_{t^f} .

Statistical analyses were performed with R (v4.0.5, R Core Team, 2021) software. The function *rpart* within the package *rpart* (v4.1.16, Therneau et al., 2022) was used to fit the classification and regression trees (we set a minimum number of values in any terminal node equal to 3% the total number of values). The function *geom_smooth* within the package *ggplot2* (v3.3.6, Wickham et al., 2022) was used to fit second-order logistic (method="glm", formula= $y \sim \text{poly}(x, 2)$, family="binomial"), second-order polynomial (method="lm", formula= $y \sim \text{poly}(x, 2)$) and local polynomial (method="loess", formula= $y \sim x$) regressions.

3 | RESULTS

The SP became established before the end of the 50-year simulation in 69.2% of the 60,000 simulations. In these 41,504 simulations, the mean time to SP establishment was 4.87 years, and the 2.5th and 97.5th percentiles were 0.6 and 33.44 years, respectively.

For the 60,000 simulations performed, the AUDPC ranged from 15% (i.e., mild epidemics) to 99% (i.e., severe epidemics). Below, we determine the roles of the principal factors driving such variability in output.

3.1 | Factors affecting superpathogen establishment

We constructed a classification tree for identifying parameter combinations leading to SP establishment (E_{SP}) (Figure 2a). E_{SP} was dependent principally on the mutation probability, the resistance deployment strategy and the fitness cost. At high mutation probabilities, the SP almost invariably became established in the pathogen population, regardless of the other factors, except for mosaic strategy at high fitness costs. In that setting, the SP became established in less than one over seven simulations. At low mutation probabilities, specific combinations of these factors determined whether or not the SP became established. For example, the SP was never established in conditions in which the resistance genes were pyramided in the same cultivar. The SP became established in less than one over two simulations when resistance genes were deployed in (i) mosaic and rotation, for high fitness costs ($\theta=0.5$); (ii) mixture and mosaic, for fitness costs below 0.5 and purely clonal reproduction. For the remaining parameter combinations, the SP became established in more than one over two simulations. The pathogen reproduction system had a secondary influence on SP establishment. However, for mixture, mosaic and rotation strategies with a low or no fitness cost, the SP almost always became established for pathogens with a mixed reproduction system, whereas the proportion of simulations in which the SP became established was substantially lower for pathogens with a clonal reproduction system, particularly for mixture and mosaic strategies.

At low mutation probabilities, SP establishment was a highly stochastic event in mixture, mosaic and rotation strategies; it occurred in 24% to 95% of the simulations, depending on the values of the other factors (Figure 2a). To get insight on the mechanisms behind SP establishment for this subset of factors, we hypothesised that the probability of SP establishment increases with the time interval between the establishment of the two single mutants $|T_{SM_1} - T_{SM_2}|$. This hypothesis was based on the rationale that, as SP and single mutants are in competition for hosts to infect, longer intervals would favour the establishment of the SP as one of the two resistant hosts remains an empty ecological niche for longer. It can, therefore, be infected by the SP if it emerges through mutation or recombination. This hypothesis holds only for the mosaic and mixture strategies, as the two resistant hosts must be deployed at the same time, excluding *de facto* the rotation strategies from the subsequent analysis. Therefore, using the function *glm* within the package *stats* (v3.6.2, R Core Team, 2022), we fitted a logistic regression to assess the relationship between E_{SP} and the time elapsed between the establishment of the two single mutants $|T_{SM_1} - T_{SM_2}|$. As expected, the probability of SP establishment increased sharply with $|T_{SM_1} - T_{SM_2}|$,




FIGURE 2 (a) Classification tree for the binary output E_{SP} . The number and proportion of simulations (out of the 60,000 performed) associated with each end node are indicated. Orange bars indicate the proportion of simulations in which the SP became established before the end of the simulation, whereas blue bars indicate the proportion of simulations in which this was not the case. The factors identified by the tree are the mutation probability for infectivity genes, the resistance deployment strategy (MIxture, MOsaic, ROTation and PYramiding), the fitness cost of infectivity genes and the pathogen reproduction system (purely clonal or mixed). (b) Relationship between the time elapsed between the establishment of the two single mutants (SM_1 and SM_2) and the probability of superpathogen establishment ($pr(E_{SP} = 1)$) for the MIxture and MOsaic strategies. Logistic regression was used to fit relationships to simulation outputs corresponding to the combination of parameters highlighted in brackets under the final nodes of the tree. Confidence intervals are delimited by the 2.5th and 97.5th percentiles.

whatever the fitness cost. Moreover, the probability of SP establishment was systematically higher for mixtures than for mosaics (Figure 2b). Finally, a specific feature of rotation strategies may also

favour the emergence of the SP regardless of the pathogen reproduction system. Indeed, a SP generated by mutation from a single mutant late in the season (i.e. when the ecological niche is no longer

empty) could still have an opportunity to establish itself in an empty niche if this event occurs shortly before the switch to a different variety in the rotation.

To deepen the analysis on the parameter combinations leading to SP establishment, we assessed the relationship between the variable E_{SP} and the cropping ratio for all combinations of resistance deployment strategy, fitness cost and pathogen reproduction system considered (Figure 3). We focused on low mutation probabilities, as shown in Figure 3 (but see Figure S1 for its analogous version with high mutation probability). The probability of E_{SP} generally increased with cropping ratio for mixture, mosaic and rotation strategies unless establishment is already certain at the lowest cropping ratio. However, for mixture strategies with null or high fitness costs, the probability of E_{SP} for pathogens undergoing purely clonal reproduction followed a U-shaped curve, with the lowest probability of E_{SP} achieved for an intermediate cropping ratio. The SP was never established in simulations based on pyramiding strategies. Furthermore, for mixture and mosaic strategies, the probability of E_{SP} was consistently lower for pathogens with clonal rather than mixed reproduction. In addition, the probability of E_{SP} was lower for mosaics than for mixtures in pathogens with a clonal reproduction system.

The effect of the pathogen reproduction system on the probability of E_{SP} can be explained by the demogenetic dynamics of the pathogen population after the bottleneck at the end of the cropping season. Contrasting dynamics were, indeed, observed across resistance deployment strategies and fitness costs, as illustrated in Figure 4 for intermediate cropping ratios. With mixture and mosaic strategies, the maximum number of heterogeneous parental pairs after the bottleneck HP_t was relatively high, at least during the first 10 cropping seasons. In this setting, sexual recombination between single mutants favoured the generation of SP propagules, which constituted the primary inoculum for the following season. Accordingly, the number of SP_t increased more rapidly, reaching a higher level for pathogens with mixed reproduction systems than for those with purely clonal reproduction, particularly if there was no fitness cost (for both mosaic and mixture strategies) or if the fitness cost was low (mixture strategy only). As a mirror effect, the number of HP_t stabilised at lower levels for pathogens with a mixed reproduction system. This effect disappeared at higher fitness costs. By contrast, the small number or absence of HP_t observed with the pyramiding and rotation strategies greatly decreased the likelihood of recombination between single mutants. Consequently, the production of SP propagules was not favoured by sexual reproduction in these strategies. Note that the trends in the demogenetic dynamics of SP_t and HP_t were similar for the other combinations of cropping ratios and mutation probabilities (Figures S2–S10).

3.2 | Factors affecting the time to superpathogen establishment

The mean time to SP establishment T_{SP} , estimated conditionally on SP establishment (i.e. for the subset of replicates such that $E_{SP}=1$), was

generally not influenced by the cropping ratio and the type of reproduction, except for mixture at high fitness cost and for mosaic and rotation at low fitness cost. In these settings, the T_{SP} decreased with cropping ratio and it was lower for pathogens with purely clonal reproduction systems (for mixture and mosaic only, Figure 3a,b). Furthermore, for mixture strategy, T_{SP} was lower for pathogens with mixed rather than purely clonal reproduction systems, for null fitness cost. Similarly, for mosaic strategy at high mutation probability, T_{SP} was lower for pathogens with mixed rather than purely clonal reproduction systems, for fitness costs that were low or zero (Figure S1B). Finally, our results showed that the variance of T_{SP} increased substantially with fitness cost, suggesting that, in these contexts, the mean time to SP establishment poorly reflected the underlying evolutionary dynamics.

3.3 | Factors affecting the mean area under the disease progress curve

In a fully susceptible landscape, the mean area under the disease progress curve, $AUDPC_0$ was 0.63 for both pathogen reproduction systems. This value implies that diseased hosts (those in an infectious or removed state, see Figure 1) accounted for a mean of 63% of the available host individuals over the entire period simulated. $AUDPC$ generally decreased with cropping ratio (Figure 3). At low mutation probability, the best epidemiological control (i.e. the lowest $AUDPC$) was obtained with the pyramiding strategy, which decreased $AUDPC$ by up to 85% at high cropping ratios, independently of the fitness cost incurred for pathogen adaptation. With the other strategies, the highest $AUDPC$ reductions achieved (for the 250 replicates) were 22% for mosaics, 32% for mixtures, 50% for rotation. These values were obtained at a high cropping ratio and fitness cost. By contrast, almost no epidemic control (i.e. $AUDPC \approx 1$) was observed for these strategies in the absence of a fitness cost. Finally, the pathogen reproduction system did not affect the $AUDPC$.

4 | DISCUSSION

We addressed the question of the effect of the type of pathogen reproduction system on the epidemiological and evolutionary control provided by plant resistance. Epidemiological control relates to plant health and the demographic dynamics of the pathogen, whereas evolutionary control relates to the durability of resistance and the genetic dynamics of the pathogen. Sexual reproduction principally favours the exchange of genes via recombination. We therefore studied the fate of the superpathogen during the deployment of two resistance genes.

4.1 | Effect of the pathogen reproduction system on evolutionary and epidemiological outputs

McDonald and Linde (2002) hypothesised that pathogens with mixed reproduction systems pose the greatest risk of genetic resistance




FIGURE 3 Probability of SP establishment (first row of each panel), time to SP establishment, given that the SP becomes established, (second row) and AUDPC (third row) at low ($\tau = 10^{-7}$) mutation probability and at zero ($\theta = 0$), low ($\theta = 0.25$) and high ($\theta = 0.5$) fitness cost (FC). Panels show the probability of E_{SP} , T_{SP} and AUDPC as a function of the cropping ratio for the two pathogen reproduction systems and the four deployment strategies considered. Curves are based on the fitting of logistic or second-order polynomial regressions to simulation outputs (represented by points, note that, in the first row of each panel, the points represent the proportion of $E_{SP} = 1$ among the 50 replicates); shaded envelopes delimited by the 2.5th and 97.5th percentiles.




FIGURE 4 Population size of the superpathogen SP_{it} (in blue) and maximum number of heterogeneous parental pairs HP_{it} (in orange) in the landscape after the annual bottleneck. The curves represent the population dynamics across resistance deployment strategies (MIxture, MOsaic, ROtation and PYramiding), fitness costs and reproduction systems, at low mutation probability ($\tau = 10^{-7}$) and intermediate cropping ratio ($\varphi = 0.5$). The curves are based on the fitting of local polynomial regressions and shaded envelopes delimited by the 2.5th and 97.5th percentiles. Note that the curves for pyramiding and rotation (at high fitness cost) overlap.

breakdown, because they benefit from the advantages of both reproduction systems. Between-cropping seasons, the occurrence of a single sexual reproduction event generates new pathogen genotypes that may combine mutations already present in the population. During the cropping season, clonal reproduction enables the fittest pathogen genotypes to invade the population rapidly. However, in tests of their risk model on 34 pathosystems, McDonald and Linde (2002) found no significant effects of the pathogen reproduction system on the risk of breakdown, which was instead affected by gene/genotype flow and mutation. Our results confirm the importance of mutation rate as a driver of pathogen evolution. Indeed, the SP was established in almost all simulations with a high mutation probability (except for mosaic at high fitness cost), regardless of the deployment strategy or pathogen reproduction system. This finding can be explained by the interplay between mutation probability and population size (Althaus & Bonhoeffer, 2005; Christiansen et al., 1998). Indeed, we can estimate the probability that at least one SP appears from the WT population infecting the susceptible fields within five growing seasons (17 generations of clonal reproduction

each) as $1 - (1 - \tau^2)^{N_{\text{mean}} \times 17 \times 5}$, where τ is the mutation probability and N_{mean} is the mean pathogens population on susceptible cultivars. In our settings, N_{mean} ranges from $[2.7 \times 10^6; 1.3 \times 10^7]$, respectively, for the highest and the lowest cropping ratios. Therefore, at high mutation probability ($\tau = 10^{-4}$), the SP will emerge in [90–99] of 100 simulations during the first five cropping seasons. Accordingly, the SP almost surely pre-exists in the pathogen population as standing genetic variation for the remaining 45 years of simulation (McDonald et al., 2002).

Our results also show that the effect of sexual reproduction on the likelihood of the generalist SP becoming established depends on the resistance deployment strategy. This finding goes a step further than the analysis presented by McDonald and Linde (2002), who did not consider the effect of deployment strategies, shading light on a topic of great importance for plant pathology and resistance durability (Mundt, 2018). Our simulations suggest that recombination favours the establishment of the SP only when heterogeneous pairs of single mutant parents are potentially abundant after crop harvest. This is the case for the mosaic and mixture strategies (Figure 4).

For these strategies, populations of single mutant pathogens can increase in size on their specific hosts, with recombination subsequently occurring on susceptible hosts during sexual reproduction, potentially generating SP propagules between two cropping seasons. The timing of sexual reproduction is also a key element explaining why SP establishment is favoured by a mixed reproduction system. Indeed, the SP propagules generated by recombination during the off-season emerge right at the start of the following cropping season, when most hosts are healthy, favouring SP establishment in this empty ecological niche. By contrast, for pathogens with purely clonal reproduction, the SP is generated by mutation from a single mutant when the population is large enough. This event probably occurs late during the cropping season when the competition between the SP and the two single mutants for the infection of healthy hosts is much stronger. Accordingly, we found that the probability of SP establishment increased when the competition with the single mutants is lower, in particular when only one single mutant pathogen is established on a resistant host and the second host is free from disease (Figure 2b).

By contrast, sexual reproduction does not favour the establishment of the SP in pyramiding and rotation strategies, because heterogeneous pairs of single mutants are scarce in these conditions (Figure 4), as the cultivars carrying the single resistance genes are not deployed at all, or not deployed simultaneously. Similar results were reported in the context of the resistance to xenobiotics (Althaus & Bonhoeffer, 2005; Taylor & Cunniffe, 2022). In particular, sexual reproduction in fungi increases the frequency of the double-resistant strain adapted to a mixture of fungicides (as for the SP here) only when the frequency of single-resistant strains is significantly higher than that of double-resistant or avirulent strain (Taylor & Cunniffe, 2022).

4.2 | No deployment strategy is universally optimal

Consistent with the findings of previous comparisons of deployment strategies (Djidjou-Demasse et al., 2017; Lof & van der Werf, 2017; Rimbaud, Papaix, Barrett, et al., 2018; Sapoukhina et al., 2009), our results confirm that no one strategy is universally optimal. Instead, the strategy used should be adapted to the pathosystem and production situation, and a decision must be taken as to whether to prioritise epidemiological or evolutionary outputs. With this in mind, given that pre-adapted pathogens were assumed to be initially absent, the order of magnitude of the mutation probability relative to pathogen population size is a key factor. Conversely, the pathogen reproduction system had no effect on strategy recommendations for various fitness costs, mutation probabilities and cropping ratios. Similarly Taylor and Cunniffe (2022) showed that sexual reproduction did not affect recommendations for the management of fungicides mixtures.

At low mutation probabilities, a SP will emerge by mutation from the wild-type at most 1 in every 10,000 simulation runs during the 17×50 generations. It explains the better performance of

pyramiding over all other strategies (Leach et al., 2001). Pyramiding strategies ensure both epidemiological and evolutionary control of the targeted disease, as reported by Djian-Caporalino et al. (2014), Rimbaud, Papaix, Barrett, et al. (2018). In particular, the decrease in disease severity is proportional to the cropping ratio of the pyramided variety in the landscape as the dilution effect is maximal in this setting (Keesing & Ostfeld, 2021). For the other strategies, the probability of SP establishment generally increases with cropping ratio, as higher cropping ratios favour the development of large populations of single mutants, in turn favouring the emergence of the SP. However, for mixture strategies with null or high fitness costs and pathogens with purely clonal reproduction, the relationship between cropping ratio and the probability of SP establishment is U-shaped. Among the mechanisms underlying this relationship, the competition between single mutants (SM_1 and/or SM_2) and SP and thus the relative order of their establishment should be a major driver. Indeed, in this setting, the SP must appear from a single mutation of one of SM_1 or SM_2 , and colonise a host not already occupied by SM_1 or SM_2 , to establish. At small cropping ratio, the two single mutants quickly appear in susceptible fields, but their probability to disperse to their corresponding resistant host is small because such hosts are scarce in the landscape. At the opposite, at high cropping ratio, the population size of the wild-type is small, which reduces the chances that two single mutants emerge within a short time frame, but they can infect their corresponding resistant host more easily. Overall it results that, at both small and large cropping ratio, the time between the establishment of the two SMs is relatively long (top row of Figure S14). Ultimately, this gives more time to the SP, that will emerge by mutation from the first SM established, to establish on the other resistant host (bottom row of Figure S14). At intermediate cropping ratio, instead, the population size of the wild-type is high enough to allow appearance of the two single mutants quickly one after the other (top row of Figure S14), and the proportion of resistant fields is high enough to allow their establishment before colonisation of a SP. This reduces the probability of SP establishment at intermediate compared to low and high cropping ratios (bottom row of Figure S14). The processes leading to the non-monotonic response of the time elapsed between SMs establishment and cropping ratio also apply for mosaic strategy (top row in Figure S14). However, in this case, contrary to the mixture strategy, the SP, which will for example emerge from SM_1 in fields planted with RC_1 , must additionally disperse to another field sown with the RC_2 . The probability of such successful event is weak for low cropping ratio whatever the fitness cost and increases with the cropping ratio (bottom row of Figure S14).

At high mutation probabilities, the SP becomes established a mean of 1.5 years after the beginning of a simulation run for pyramiding strategies (Figure S1D). There is no dilution effect at work during most of the 50-year time frame considered, and epidemiological and evolutionary control disappear. In this setting, the strategies delaying SP establishment for the longest were mosaic, at high fitness costs (Figure S1B). Higher fitness costs in this strategy slowed SP establishment through disruptive selection. This

mechanism exploits fitness costs to favour local host specialisation of the pathogen, limiting the likelihood of a generalist SP emerging (Barrett et al., 2009). Despite generally providing the best evolutionary control, the mosaic strategy was the worst strategy (in comparisons with rotation and mixture) in our conditions for epidemiological control. One key reason for this is the high probability of autoinfections, 0.87 on average, a consequence of our choice of large field sizes (mean of 160m × 160m) relative to short mean pathogen dispersal distances (20m). The frequent infection events resulting from propagules produced in the same field favours the mixture strategy over the mosaic strategy (Mundt, 2002). Like us, Djidjou-Demasie et al. (2017) also found that pyramiding and mosaic strategies provided similar levels of epidemiological control if the probability of autoinfection was high. In their study, frequent between-field infections and high rates of mutation were required for mosaic strategies to outperform pyramiding.

Crucially, our results highlight the need for knowledge about mutation probability, pathogen population size and cost of infectivity to guide the choice of a deployment strategy. In particular, the ratio between mutation probability and pathogen population size will guide the choice as to whether or not to use a pyramiding strategy, as it greatly affects the establishment of a SP (Figure S15 in Note S6). Unfortunately, there has been little quantitative characterisation of these parameters (Laine & Barrès, 2013). Point mutations are the simplest evolutionary events conferring virulence to a resistance gene. Such events occur once every 10^5 to 10^7 propagules per generation (Stam & McDonald, 2018). However, many other mutational events *sensu lato* (e.g. complete or partial gene deletion, insertion of transposable elements) increase the overall mutation probability conferring virulence (Daverdin et al., 2012; Paineau et al., 2023). Similarly, despite census population sizes of plant fungi are likely very large at field scale (McDonald et al., 2002), estimates of effective population size that will effectively contribute to the epidemics are lacking. The cost of infectivity, instead, has a monotonic influence: the higher the cost, the higher the levels of evolutionary and epidemiological control achieved. Such costs are not pervasive among plant-pathogenic fungi and vary with host genotype and abiotic environment (Laine & Barrès, 2013). For example, substantial sporulation costs have been reported in rusts (Bahri et al., 2009; Thrall & Burdon, 2003) but no such costs evidenced for grapevine downy mildew (Delmas et al., 2016; Toffolatti et al., 2012). Finally, we assumed a complete restoration of pathogen infectivity on resistant hosts. However, several modelling studies considered that the adapted pathogens experience a fitness cost on all host genotypes (Clin et al., 2021, 2022; Lo Iacono et al., 2012; Sapoukhina et al., 2009). This assumption leads to a slightly different plant-pathogen interaction matrix (Table S2 in Note S7). We investigated to which extent the structure of this matrix impacts our results (Note S7). Firstly, our results on the effect of pathogen reproduction system were robust to the presence (or absence) of a fitness cost for adapted pathogens on all host genotypes (Figures S16 and S17). Furthermore, our results indicated that the structure of

the plant–pathogen interaction matrix did not substantially affect evolutionary and epidemiological outputs for fitness costs $\theta \leq 0.25$ (Figures S18 and S19). However, for the highest considered fitness cost, the AUDPC could be considerably lower when the adapted pathogens experience a fitness cost on all host genotypes, especially at high cropping ratio. This is a direct consequence of the globally smaller fitness of adapted pathogens in a landscape composed of many resistant hosts, as compared with the scenario where they pay a fitness cost only for their unnecessary virulence.

4.3 | Further perspectives

The ecoevolutionary model presented here represents a solid foundation for further investigations of the effects of other mechanisms linked to the sexual reproduction of pathogens. For example, we assume that all the sexual propagules emerge in the cropping season immediately following their production, but specialised reproductive structures can survive in the soil for many years (up to 5 years for *P. viticola*, Caffi et al., 2010). This feature may impact the outputs of deployment strategies, in particular rotations (Papavizas & Ayers, 1974). We also assume that sexual and clonal propagules have similar dispersal capacities. This may not always be the case, as shown for black sigatoka (Rieux et al., 2014) and grapevine downy mildew (Rossi & Caffi, 2012). Such dispersal dimorphism probably affects the effectiveness of resistance deployment strategies such as mixtures and mosaics (Papaix et al., 2018; Sapoukhina et al., 2010; Watkinson-Powell et al., 2020).

Furthermore, we focus here exclusively on qualitative resistance genes (i.e. major genes), but quantitative resistance is attracting increasing interest for use in pathogen control (Niks et al., 2015; Parlevliet, 2002). As the model can also handle quantitative resistances, it would be interesting to broaden our analysis in this direction. Recombination in a diverse pathogen population, as favoured by the partial effect of quantitative resistance on pathogens, might accelerate pathogen evolution towards higher levels of aggressiveness (Drenth et al., 2019; Frézal et al., 2018). Conversely, recombination, by breaking up blocks of co-adapted genes, may slow the adaptation of pathogens to quantitative resistance genes (McDonald & Linde, 2002).

Finally, we assumed that a given deployment strategy is implemented over the entire agricultural landscape from the beginning of a simulation. It would be interesting to investigate the effects of the gradual introduction of resistant cultivars in the landscape.

ACKNOWLEDGEMENTS

This work was funded by the MEDEE project of the Plan d'action national Ecophyto II: Axe 2 - APR Leviers Territoriaux (No.SIREPA 4621) and by the ANR COMBINE project (ANR-22-CE32-0004).

CONFLICT OF INTEREST STATEMENT

All authors declare that they have no conflicts of interest.

DATA AVAILABILITY STATEMENT

The code for reproducing the analysis, results and figures presented in this paper are openly available at <https://doi.org/10.57745/PACERT>.

ORCID

Marta Zaffaroni  <https://orcid.org/0000-0002-2951-8626>

Loup Rimbaud  <https://orcid.org/0000-0002-8098-9984>

Jean-François Rey  <https://orcid.org/0000-0003-3281-6701>

Julien Papaix  <https://orcid.org/0000-0003-2273-5334>

Frédéric Fabre  <https://orcid.org/0000-0001-8271-7678>

REFERENCES

- Althaus, C. L., & Bonhoeffer, S. (2005). Stochastic interplay between mutation and recombination during the acquisition of drug resistance mutations in human immunodeficiency virus type 1. *Journal of Virology*, 79(21), 13572–13578.
- Arenas, M., Araujo, N. M., Branco, C., Castelhana, N., Castro-Nallar, E., & Pérez-Losada, M. (2018). Mutation and recombination in pathogen evolution: Relevance, methods and controversies. *Infection, Genetics and Evolution*, 63, 295–306.
- Bahri, B., Kaltz, O., Leconte, M., de Vallavieille-Pope, C., & Enjalbert, J. (2009). Tracking costs of virulence in natural populations of the wheat pathogen, *Puccinia striiformis f.sp.tritici*. *BMC Evolutionary Biology*, 9, 26.
- Barrett, L. G., Kniskern, J. M., Bodenhausen, N., Zhang, W., & Bergelson, J. (2009). Continuum of specificity and virulence in plant host–pathogen interactions: Causes and consequences. *New Phytologist*, 183(3), 513–529.
- Blum, M., Waldner, M., & Gisi, U. (2010). A single point mutation in the novel *PvCesA3* gene confers resistance to the carboxylic acid amide fungicide mandipropamid in *Plasmopara viticola*. *Fungal Genetics and Biology*, 47(6), 499–510.
- Boso, S., & Kassemeyer, H. H. (2008). Different susceptibility of European grapevine cultivars for downy mildew. *Vitis*, 47(1), 39–49.
- Bove, F., Bavaresco, L., Caffi, T., & Rossi, V. (2019). Assessment of resistance components for improved phenotyping of grapevine varieties resistant to downy mildew. *Frontiers in Plant Science*, 10, 1–10.
- Bove, F., & Rossi, V. (2020). Components of partial resistance to *Plasmopara viticola* enable complete phenotypic characterization of grapevine varieties. *Scientific Reports*, 10, 1–12.
- Brown, J. K. (2015). Durable resistance of crops to disease: A Darwinian perspective. *Annual Review of Phytopathology*, 53, 513–539.
- Burdon, J. J., Barrett, L. G., Rebetzke, G., & Thrall, P. H. (2014). Guiding deployment of resistance in cereals using evolutionary principles. *Evolutionary Applications*, 7(6), 609–624.
- Caffi, T., Rossi, V., & Bugiani, R. (2010). Evaluation of a warning system for controlling primary infections of grapevine downy mildew. *Plant Disease*, 94(6), 709–716.
- Chen, W.-J., Delmotte, F., Cervera, S. R., Douence, L., Greif, C., & Corio-Costet, M.-F. (2007). At least two origins of fungicide resistance in grapevine downy mildew populations. *Applied and Environmental Microbiology*, 73(16), 5162–5172.
- Christiansen, F. B., Otto, S. P., Bergman, A., & Feldman, M. W. (1998). Waiting with and without recombination: The time to production of a double mutant. *Theoretical Population Biology*, 53(3), 199–215.
- Clin, P., Grogard, F., Andrivon, D., Mailleret, L., & Hamelin, F. M. (2022). Host mixtures for plant disease control: Benefits from pathogen selection and immune priming. *Evolutionary Applications*, 15(6), 967–975.
- Clin, P., Grogard, F., Mailleret, L., Val, F., Andrivon, D., & Hamelin, F. M. (2021). Taking advantage of pathogen diversity and immune priming to minimize disease prevalence in host mixtures: A model. *Phytopathology*, 111(7), 1219–1227.
- Cohen, Y., & Rubin, A. E. (2012). Mating type and sexual reproduction of *Pseudoperonospora cubensis*, the downy mildew agent of cucurbits. *European Journal of Plant Pathology*, 132, 577–592.
- Crété, R., Pires, R. N., Barbetti, M. J., & Renton, M. (2020). Rotating and stacking genes can improve crop resistance durability while potentially selecting highly virulent pathogen strains. *Scientific Reports*, 10, 1–17.
- Curl, E. A. (1963). Control of plant diseases by crop rotation. *The Botanical Review*, 29(4), 413–479.
- Daverdin, G., Rouxel, T., Gout, L., Aubertot, J.-N., Fudal, I., Meyer, M., Parlange, F., Carpezat, J., & Balesdent, M.-H. (2012). Genome structure and reproductive behaviour influence the evolutionary potential of a fungal phytopathogen. *PLoS Pathogens*, 8, e1003020.
- Delmas, E. L., Fabre, F., Jolivet, J., Mazet, I. D., Richart Cervera, S., Delière, L., & Delmotte, F. (2016). Adaptation of a plant pathogen to partial host resistance: Selection for greater aggressiveness in grapevine downy mildew. *Evolutionary Applications*, 9, 709–725.
- Djian-Caporalino, C., Palloix, A., Fazari, A., Marteu, N., Barbary, A., Abad, P., Sage-Palloix, A.-M., Mateille, T., Risso, S., Lanza, R., Taussig, C., & Castagnone-Sereno, P. (2014). Pyramiding, alternating or mixing: Comparative performances of deployment strategies of nematode resistance genes to promote plant resistance efficiency and durability. *BMC Plant Biology*, 14(1), 1–13.
- Djidjou-Demasse, R., Moury, B., & Fabre, F. (2017). Mosaics often outperform pyramids: Insights from a model comparing strategies for the deployment of plant resistance genes against viruses in agricultural landscapes. *New Phytologist*, 216, 239–253.
- Drenth, A., McTaggart, A. R., & Wingfield, B. D. (2019). Fungal clones win the battle, but recombination wins the war. *IMA Fungus*, 10(1), 1–6.
- Frantzen, J., & Van den Bosch, F. (2000). Spread of organisms: Can travelling and dispersive waves be distinguished? *Basic and Applied Ecology*, 1(1), 83–92.
- Frézal, L., Jacqua, G., & Neema, C. (2018). Adaptation of a fungal pathogen to host quantitative resistance. *Frontiers in Plant Science*, 9, 1554.
- Fuchs, M. (2017). Pyramiding resistance-conferring gene sequences in crops. *Current Opinion in Virology*, 26, 36–42.
- García-Arenal, F., & McDonald, B. A. (2003). An analysis of the durability of resistance to plant viruses. *Phytopathology*, 93(8), 941–952.
- Gessler, C., Pertot, I., & Perazzolli, M. (2011). *Plasmopara viticola*: A review of knowledge on downy mildew of grapevine and effective disease management. *Phytopathologia Mediterranea*, 50(1), 3–44.
- Grosdidier, M., Ios, R., Husson, C., Cael, O., Scordia, T., & Marçais, B. (2018). Tracking the invasion: Dispersal of *Hymenoscyphus fraxineus* airborne inoculum at different scales. *FEMS Microbiology Ecology*, 94, fiy049.
- Hadany, L., & Beker, T. (2003). On the evolutionary advantage of fitness-associated recombination. *Genetics*, 165(4), 2167–2179.
- Halkett, F., Simon, J. C., & Balloux, F. (2005). Tackling the population genetics of clonal and partially clonal organisms. *Trends in Ecology & Evolution*, 20(4), 194–201.
- Keesing, F., & Ostfeld, R. S. (2021). Dilution effects in disease ecology. *Ecology Letters*, 24(11), 2490–2505.
- Koller, T., Brunner, S., Herren, G., Hurni, S., & Keller, B. (2018). Pyramiding of transgenic pm3 alleles in wheat results in improved powdery mildew resistance in the field. *Theoretical and Applied Genetics*, 131(4), 861–871.
- Laine, A.-L., & Barrès, B. (2013). Epidemiological and evolutionary consequences of life-history trade-offs in pathogens. *Plant Pathology*, 62, 96–105.
- Leach, J. E., Vera Cruz, C. M., Bai, J., & Leung, H. (2001). Pathogen fitness penalty as a predictor of durability of disease resistance genes. *Annual Review of Phytopathology*, 39(1), 187–224.

- Lo Iacono, G., van den Bosch, F., & Paveley, N. (2012). The evolution of plant pathogens in response to host resistance: Factors affecting the gain from deployment of qualitative and quantitative resistance. *Journal of Theoretical Biology*, 304, 152–163.
- Lof, M. E., & van der Werf, W. (2017). Modelling the effect of gene deployment strategies on durability of plant resistance under selection. *Crop Protection*, 97, 10–17.
- Lohaus, G., Hussmann, M., Pennewiss, K., Schneider, H., Zhu, J.-J., & Sattelmacher, B. (2000). Solute balance of a maize (*Zea mays* L.) source leaf as affected by salt treatment with special emphasis on phloem retranslocation and ion leaching. *Journal of Experimental Botany*, 51(351), 1721–1732.
- McDonald, B. A. (2010). How can we achieve durable disease resistance in agricultural ecosystems? *New Phytologist*, 185, 3–5.
- McDonald, B. A. (2014). Using dynamic diversity to achieve durable disease resistance in agricultural ecosystems. *Tropical Plant Pathology*, 39, 191–196.
- McDonald, B. A., & Linde, C. (2002). Pathogen population genetics, evolutionary potential, and durable resistance. *Annual Review of Phytopathology*, 40, 349–379.
- McDonald, B. A., Suffert, F., Bernasconi, A., & Mikaberidze, A. (2002). How large and diverse are field populations of fungal plant pathogens? The case of *Zymoseptoria tritici*. *Evolutionary Applications*, 15(9), 1360–1373.
- Mundt, C. C. (2002). Use of multiline cultivars and cultivar mixtures for disease management. *Annual Review of Phytopathology*, 40(1), 381–410.
- Mundt, C. C. (2018). Pyramiding for resistance durability: Theory and practice. *Phytopathology*, 108(7), 792–802.
- Niks, R. E., Qi, X., & Marcel, T. C. (2015). Quantitative resistance to biotrophic filamentous plant pathogens: Concepts, misconceptions, and mechanisms. *Annual Review of Phytopathology*, 53(1), 1110–1146.
- Paineau, M., Mazet, I., Wiedemann-Merdinoglu, S., Fabre, F., & Delmotte, F. (2022). The characterization of pathotypes in grapevine downy mildew provides insights into the breakdown of rpv3, rpv10, and rpv12 factors in grapevines. *Phytopathology*, 112(11), 2329–2340.
- Paineau, M., Minio, A., Mestre, P., Fabre, F., Mazet, I. D., Couture, C., Legeai, F., Dumartinet, T., Cantu, D., & Delmotte, F. (2023). An effector deletion leads to the breakdown of partial grapevine resistance to downy mildew. *bioRxiv*.
- Papaix, J., Rimbaud, L., Burdon, J. J., Zhan, J., & Thrall, P. H. (2018). Differential impact of landscape-scale strategies for crop cultivar deployment on disease dynamics, resistance durability and long-term evolutionary control. *Evolutionary Applications*, 11(5), 705–717.
- Papaix, J., Touzeau, S., Monod, H., & Lannou, C. (2014). Can epidemic control be achieved by altering landscape connectivity in agricultural systems? *Ecological Modelling*, 284, 35–47.
- Papavizas, G. C., & Ayers, W. A. (1974). *Aphanomyces species and their root diseases in pea and sugarbeet*. Technical Report. Agricultural Research Service, U.S. Dept. of Agriculture.
- Parlevliet, J. E. (2002). Durability of resistance against fungal, bacterial and viral pathogens; present situation. *Euphytica*, 124(2), 147–156.
- Peressotti, E., Wiedemann-Merdinoglu, S., Delmotte, F., Bellin, D., Di Gaspero, G., Testolin, R., Merdinoglu, D., & Mestre, P. (2010). Breakdown of resistance to grapevine downy mildew upon limited deployment of a resistant variety. *BMC Plant Biology*, 10(1), 1–11.
- R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- R Core Team. (2022). *The R stats package*. R Package Version 4.3.0.
- Rieux, A., Soubeyrand, S., Bonnot, F., Klein, E. K., Ngando, J. E., Mehl, A., Ravigne, V., Carlier, J., & De Bellaire, L. (2014). Long-distance wind-dispersal of spores in a fungal plant pathogen: Estimation of anisotropic dispersal kernels from an extensive field experiment. *PLoS One*, 9(8), e103225.
- Rimbaud, L., Fabre, F., Papaix, J., Moury, B., Lannou, C., Barret, L. G., & Thrall, P. H. (2021). Models of plant resistance deployment. *Annual Review of Phytopathology*, 59(1), 125–152.
- Rimbaud, L., Papaix, J., Barrett, L. G., Burdon, J. J., & Thrall, P. H. (2018). Mosaics, mixtures, rotations or pyramiding: What is the optimal strategy to deploy major gene resistance? *Evolutionary Applications*, 11(10), 1791–1810.
- Rimbaud, L., Papaix, J., Rey, J.-F., Barrett, L. G., & Thrall, P. H. (2018). Assessing the durability and efficiency of landscape-based strategies to deploy plant resistance to pathogens. *PLoS Computational Biology*, 14(4), e1006067.
- Rimbaud, L., Papax, J., Rey, J.-F., Zaffaroni, M., & Gaussen, J.-L. (2022). Landsepi: Landscape epidemiology and evolution. R Package Version 1.2.4.
- Rossi, V., & Caffi, T. (2012). The role of rain in dispersal of the primary inoculum of *plasmopara viticola*. *Ecology and Epidemiology*, 102(2), 158–165.
- Sapoukhina, N., Durel, C. E., & Le Cam, B. (2009). Spatial deployment of gene-for-gene resistance governs evolution and spread of pathogen populations. *Theoretical Ecology*, 2, 229–238.
- Sapoukhina, N., Tyutyunov, Y., Sache, I., & Arditi, R. (2010). Spatially mixed crops to control the stratified dispersal of airborne fungal diseases. *Ecological Modelling*, 221(23), 2793–2800.
- Saubin, M., De Mita, S., Zhu, X., Sudret, B., & Halkett, F. (2021). Impact of ploidy and pathogen life cycle on resistance durability. *Peer Community Journal*, 1, e8.
- Stam, R., & McDonald, B. A. (2018). When resistance gene pyramids are not durable—The role of pathogen diversity. *Molecular Plant Pathology*, 19(3), 521–524.
- Taylor, N., & Cunniffe, N. J. (2022). Optimal resistance management for mixtures of high-risk fungicides: Robustness to the initial frequency of resistance and pathogen sexual reproduction. *Phytopathology*, 113, 55–69.
- Therneau, T., Atkinson, B., & Ripley, B. (2022). Recursive partitioning and regression trees. R Package Version 4.1.16.
- Thrall, P. H., & Burdon, J. J. (2003). Evolution of virulence in a plant host-pathogen metapopulation. *Science*, 299(5613), 1735–1737.
- Tibayrenc, M., & Ayala, F. J. (2002). The clonal theory of parasitic protozoa: 12 years on. *Trends in Parasitology*, 18(9), 405–410.
- Toffolatti, S. L., Venturini, G., Maffi, D., & Vercesi, A. (2012). Phenotypic and histochemical traits of the interaction between *Plasmopara viticola* and resistant or susceptible grapevine varieties. *BMC Plant Biology*, 12(1), 124.
- Uecker, H. (2017). Evolutionary rescue in randomly mating, selfing, and clonal populations. *Evolution*, 71(4), 845–858.
- Watkinson-Powell, B., Gilligan, C. A., & Cunniffe, N. J. (2020). When does spatial diversification usefully maximize the durability of crop disease resistance? *Phytopathology*, 110(11), 1808–1820.
- Wickham, H., Chang, W., Henry, L., Pedersen, T. L., Takahashi, K., Wilke, C., Woo, K., Yutani, H., Dunnington, D., & RStudio. (2022). Create elegant data visualisations using the grammar of graphics. R Package Version 3.3.6.
- Wolfe, M. (1985). The current status and prospects of multiline cultivars and variety mixtures for disease resistance. *Annual Review of Phytopathology*, 23(1), 251–273.
- Wong, F. P., Burr, H. N., & Wilcox, W. F. (2001). Heterothallism in *Plasmopara viticola*. *Plant Pathology*, 50, 427–432.
- Xu, X. (2012). Super-races are not likely to dominate a fungal population within a life time of a perennial crop plantation of cultivar mixtures: A simulation study. *BMC Ecology*, 12, 1–10.
- Zhan, J., Thrall, P. H., Papaix, J., Xie, L., & Burdon, J. J. (2015). Playing on a Pathogen's weakness: Using evolution to guide sustainable

plant disease control strategies. *Annual Review of Phytopathology*, 53, 19–43.

Zhu, Y., Chen, H., Fan, J., Wang, Y., Li, Y., Chen, J., Fan, J., Yang, S., Hu, L., Leung, H., Mew, T. W., Teng, P. S., Wang, Z., & Mundt, C. C. (2000). Genetic diversity and disease control in rice. *Nature*, 406(6797), 718–722.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Zaffaroni, M., Rimbaud, L., Rey, J.-F., Papaïx, J., & Fabre, F. (2024). Effects of pathogen reproduction system on the evolutionary and epidemiological control provided by deployment strategies for two major resistance genes in agricultural landscapes. *Evolutionary Applications*, 17, e13627. <https://doi.org/10.1111/eva.13627>