



HAL
open science

Natural streambank structure assessment in mountain rivers: an approach combining ecology and hydromorphology

Rousset Juliette, Piton Guillaume, Didier Marie, François Adeline, Evette André

► **To cite this version:**

Rousset Juliette, Piton Guillaume, Didier Marie, François Adeline, Evette André. Natural streambank structure assessment in mountain rivers: an approach combining ecology and hydromorphology. EGU General Assembly 2023, European Geoscience Union, Apr 2023, Vienna, Austria. pp.3/EGU23-5403, <10.5194/egusphere-egu23-5403>. <hal-04007459>

HAL Id: hal-04007459

<https://hal.inrae.fr/hal-04007459v1>

Submitted on 28 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

EGU23-5403

<https://doi.org/10.5194/egusphere-egu23-5403>

EGU General Assembly 2023

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



Natural streambank structure assessment in mountain rivers: an approach combining ecology and hydromorphology

Rousset Juliette¹, Piton Guillaume², Didier Marie¹, François Adeline¹, and Evette André¹

¹Univ. Grenoble Alpes, INRAE, UR LESSEM, 2 rue de la papeterie BP 76, 38402 Saint-Martin-d'Hères, France

²Univ. Grenoble Alpes, INRAE, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

Due to the steep relief of the mountain massifs, the valleys are strategic locations for the establishment of human activities. Over last decades, the floodplain of alpine rivers has been highly anthropized. However, mountain rivers have a strong erosive power that can threaten human infrastructures. To protect them, riverbanks are regularly protected with civil engineering works. Yet, this environment is home to a high level of biodiversity. Thus, soil water bioengineering techniques appear sustainable by allowing both to protect the anthropogenic assets and to welcome the rich riparian biodiversity and the associated ecological services. To implement these Nature-Based Solutions, it is necessary to draw inspiration from the biotic and abiotic components of natural riverbanks, which are still poorly understood at high altitude due to several pressures (climatic, hydrologic, hydraulic or morphologic). In order to understand the structure and functioning of natural models of alpine riverbanks, a protocol has been implemented on 21 mountain banks with mature vegetation and no signs of erosion. The sites were located in the Vanoise massif (France) on an altitudinal gradient (1331 – 2131 m) ranging from montane to subalpine belts with a wide range of channel slope (0,8 – 28,4 %). By focusing on the bank toe, biotic (vegetation cover and species biological features) and abiotic (altitude and hydrogeomorphology) components have been measured. First observations showed that the grain size distribution of the sediments forming the bank was coarser and more homogeneous than those forming the channel bed. For the step-pool channels, bank grain sizes were coarser than for plane-bed channels. Shields parameters for the 21 banks were well below the thresholds for sediment motion, and were therefore relatively stable. Furthermore, on all the sites the shear stresses were relatively high but according to the literature values, classic bioengineering structures could be implemented on 16 of the sites and would withstand a 100-year return period flood, which is consistent with our observation of mature, stable and healthy vegetation patches. The mineral component of the bank toe is closely linked to the woody plant structure. The vegetation cover was dominated by 12 species of willow and the green alder. Green alder cover increased with the altitude and the bed slope. Willow species cover varied with altitude and was less important for steep river. Thus, the natural models of banks were: (i) for the rivers with steep slope, models mixing mineral and plant units which both took part jointly in the stability of the bank; and (ii) for the rivers with lower slope the stability of the bank was mainly ensured by the vegetation. This work provided additional knowledge on the structure and functioning of the banks of stable and mature high-altitude rivers. Accordingly, these results will allow to design soil

water bioengineering techniques more adapted to mountain environments (plant composition and structure, mineral unit size).