



**HAL**  
open science

# Inorganic phosphate content does not affect oviposition preference in the invasive pest *Drosophila suzukii*

Laure Olazcuaga, Robin Guilhot, Jean-loup Claret, Nicolas O Rode

## ► To cite this version:

Laure Olazcuaga, Robin Guilhot, Jean-loup Claret, Nicolas O Rode. Inorganic phosphate content does not affect oviposition preference in the invasive pest *Drosophila suzukii*. *Journal of Applied Entomology*, In press, 10.1111/jen.13107 . hal-04014886

**HAL Id: hal-04014886**

**<https://hal.inrae.fr/hal-04014886>**

Submitted on 4 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Inorganic phosphate content does not affect oviposition preference in the invasive pest *Drosophila suzukii*

Laure Olazcuaga<sup>1,2</sup>  | Robin Guilhot<sup>1</sup>  | Jean-Loup Claret<sup>1</sup>  | Nicolas O. Rode<sup>1</sup> 

<sup>1</sup>CBGP, INRAE, CIRAD, IRD, Institut Agro, Univ Montpellier, Montpellier, France

<sup>2</sup>Department of Agricultural Biology, Colorado State University, Fort Collins, Colorado, USA

## Correspondence

Laure Olazcuaga, CBGP, University Montpellier, CIRAD, INRAE, Institut Agro, IRD, 755 avenue du Campus Agropolis CS 30016, 34988 Montpellier sur Lez cedex, France.  
Email: [olaz.laure@gmail.com](mailto:olaz.laure@gmail.com)

## Funding information

European Union program FEFER FSE IEJ 2014-2020; INRAE scientific department SPE, Grant/Award Number: AAP-SPE 2016; CeMEB LabEx/University of Montpellier, Grant/Award Number: ANR-10-LABX-04-01; National Science Foundation, Grant/Award Number: DEB-1930650

## Abstract

The broad variation in host use among polyphagous insects is well documented but still poorly understood. In numerous pest insects, the proximate mechanisms responsible for variation in oviposition preference among host plants remain to be elucidated. The invasive crop pest, *Drosophila suzukii*, attacks a wide range of host fruits. Females prefer ovipositing on particular fruit media (blackberry, cherry, blackcurrant) that are rich in phosphorus. As phosphorus is known to be involved in female reproduction in insect species such as *Drosophila*, it could drive oviposition preference in *D. suzukii*. Phosphorus is either present as inorganic or organic phosphate in fruits. As the absolute content in macromolecules associated with phosphate in fruits (i.e. proteins and carbohydrates) do not affect oviposition in *D. suzukii*, we tested for the effect of inorganic phosphate on oviposition preference. We measured the egg-laying preferences of *D. suzukii* in a choice environment containing 12 artificial media with increasing content in inorganic phosphate (monopotassium dihydrogen phosphate). In our assay, *D. suzukii* females did not prefer ovipositing in media with high inorganic phosphate content compared to media with lower inorganic phosphate content. As a confirmation, we verified the previous result of a higher female preference for media made of phosphorus-rich fruits (blackberry, cherry, blackcurrant). The higher preference for phosphorus-rich fruits could be driven by macromolecules containing phosphorus (e.g. phospholipids) or by the presence of one or more molecules that do not contain phosphorus, but that happen to be correlated to fruit phosphorus content. Studying the proximate mechanisms driving host use will ultimately help improve the management of *D. suzukii* and other crop pests.

## KEYWORDS

chemical ecology, host choice, insect behaviour, oviposition site selection, spotted-wing drosophila

## 1 | INTRODUCTION

Polyphagous insect pests that infest many host plants are responsible for important economic losses in agriculture (Oerke, 2006).

Hence, elucidating the mechanisms that drive patterns of host use by polyphagous insects is an important goal in ecological entomology. Oviposition preference, one of the main traits that determines host use in polyphagous insects, varies widely among host fruits and

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Journal of Applied Entomology* published by Wiley-VCH GmbH.

among varieties within a given host fruit (Singer, 1986). Both the physical (skin thickness, fruit shape, and colour) and chemical (nutritional quality, presence of toxins) properties of fruits are thought to affect female oviposition preference (Renwick, 1989).

*Drosophila suzukii* is an insect pest, native to Asia, which invaded both Europe and North America in 2008 (Framout et al., 2017). This extremely polyphagous pest can infest fruits from at least 19 different plant families (Kenis et al., 2016). In particular, it infests soft fruits (strawberry, raspberry, blueberry, mulberry), stone fruits (cherry) and grape, which results in important economic losses for farmers (for example \$US 36.1 million in revenue loss between 2009 and 2011 for Californian raspberry producers; Farnsworth et al., 2017). Current control methods rely heavily on the use of broad-spectrum insecticides (Schetelig et al., 2018). However, as the timing of infestation coincides with ripening, pesticide use is restricted to protect consumers from exposure to chemical residues (Haviland & Beers, 2012). Given the known negative impacts of insecticides on human health and on the environment (Bourguet & Guillemaud, 2016), alternative pest management strategies are deeply needed. Notably, the development of push-pull strategies (e.g. trap-plants) requires a good understanding of the factors that drive oviposition preference in insect pests (Alkema et al., 2019).

The roles of certain physical (e.g. Burrack et al., 2013; Lee et al., 2011; Little et al., 2018; Little et al., 2019; Tait et al., 2020; Takahara & Takahashi, 2017) and chemical characteristics (e.g. Dweck et al., 2016; Karageorgi et al., 2017; Silva-Soares et al., 2017; Young et al., 2018; see Olazcuaga et al., 2019 for a review) in driving oviposition preference of *D. suzukii* are well known. For instance, oviposition preference for a given fruit increases as pH increases and as the force necessary to penetrate the skin decreases (Baena et al., 2022; Kinjo et al., 2013; Lee et al., 2015). The ratio of protein to carbohydrates (P:C ratio) also affects female oviposition preference with females preferring to lay eggs in media rich in carbohydrates relative to proteins (Silva-Soares et al., 2017; Young et al., 2018). However, the effects of specific molecules of the fruit on oviposition preference have been little investigated (see Olazcuaga et al., 2019 for an indirect approach). Olazcuaga et al. (2019) found that, in a choice environment, females preferred to lay eggs in media made from fruits rich in phosphorus (specifically, blackberry, cherry and blackcurrant), resulting in a positive correlation between oviposition preference and phosphorus content across those fruits. Although the existence of the positive correlation between oviposition preference and phosphorus content is not direct evidence of a causal link between phosphorus and preference, it suggests that the presence of large amounts of phosphorus might drive female oviposition preference in *D. suzukii*. Indeed, phosphorus is known to be important for female survival and reproduction in insects, including some *Drosophila* species (Bergwitz, 2012; King & Wilson, 1955; Markow et al., 1999). In both frugivorous and cactophilic *Drosophila*, the elemental composition of females is richer in phosphorus than that of males (Markow et al., 1999). The authors assumed that females may actively seek phosphorus-rich resources for feeding, as they might use phosphorus for RNA transcription during oogenesis. We

thus hypothesize that *D. suzukii* females could seek phosphorus-rich fruits for feeding and potentially for oviposition, resulting in a positive correlation between oviposition preference and the phosphorus content of fruits.

In plant organs such as fruits, leaves, seeds or roots, phosphorus is mostly present as inorganic phosphate, usually associated with potassium, or as organically bound phosphate (e.g. sugar phosphates, phospholipids, nucleotides; Lott et al., 2000; Marschner, 2012). Organically bound phosphate is unlikely to affect oviposition preference, as the absolute content of macromolecules associated with phosphate in fruits such carbohydrates or proteins do not affect oviposition preference in *D. suzukii* (Olazcuaga et al., 2019). We hypothesized that female oviposition might be driven by the inorganic phosphate content of fruits and tested whether oviposition preference of *Drosophila suzukii* does depend on inorganic phosphate content.

## 2 | MATERIALS AND METHODS

### 2.1 | Fly population and maintenance

To minimize phenotypic variation, we used an inbred line of *D. suzukii* (WT3 2.0; Paris et al., 2020). Inbred lines represent powerful tools to unravel potential drivers of oviposition preference of natural populations of *D. suzukii* (Karageorgi et al., 2017). Individuals were maintained in vials with standard laboratory fly food (GF, Backhaus et al., 1984) for several discrete generations in a growth chamber (15 days of larval development, 6 days of adult maturation and 24 h to lay eggs; 21°C ( $\pm 2^\circ\text{C}$ ), 65% relative humidity and 16:8 (L:D) light cycle). We performed oviposition preference assays under the same environmental conditions.

### 2.2 | Preparation of media with increasing content in inorganic phosphate

To test whether the oviposition preference of *D. suzukii* depends on the inorganic phosphate content, we supplemented a minimum fly food medium with increasing amounts of monopotassium dihydrogen phosphate ( $\text{KH}_2\text{PO}_4$ ). Other sources of phosphorus (i.e. calcium or iron phosphate) are less likely to be used by plants due to their insolubility (Lindsay C. Campbell, pers. com.). The minimum medium was made by adding 1% (w/v) of agar, 6% of inactive malted brewer's yeast, 1.5% of yeast extract and an antimicrobial solution (consisting of 6 mL of 99% propionic acid, 10 mL of 96% ethanol and 0.1% of methylparaben sodium salt) to 1000 mL of sterile deionized water (Table S1). Since  $\text{KH}_2\text{PO}_4$  is a good buffer, we adjusted the pH of each medium to 4 (i.e. the pH of fruit media used in Olazcuaga et al., 2019 and in the present assay) using hydrochloric acid (HCl).

To maximize the ability to detect an effect of the inorganic phosphate content on oviposition preference, we varied the inorganic

phosphate content from 0 to 4.7 g/kg medium. This range corresponds to a monopotassium dihydrogen phosphate ranging from 0 to 6.6 g/kg medium and phosphorus content ranging from 0 to 1.5 g/kg medium, several times larger than the range of phosphorus content observed in the 12 artificial fruit media used in Olazcuaga et al., 2019 (0 to 0.30 g/kg medium).

### 2.3 | Standardization of the age of the flies

Following Olazcuaga et al., 2019, we used groups of 6-day old flies for the oviposition experiment (i.e. sexually mature females; Emiljanowicz et al., 2014). The artificial media used for the experiment were freshly made and flies were not previously exposed to fruit media, two factors known to influence oviposition site selection in *D. sukuzii* (Elsensohn, Aly, et al., 2021; Elsensohn, Schal, & Burrack, 2021; Hoffmann, 1985; Tait et al., 2020). To maximize the number of replicates and minimize the stress of handling adults with CO<sub>2</sub> to control sex ratio, we did not control or measure the sex ratio of the adults in each group of 20 individuals. As we used a very large number of replicates (see below), our oviposition preference assay represents that of a population with an average sex ratio of 50:50.

### 2.4 | Oviposition preference assay

Following Olazcuaga et al., 2019, oviposition preference was measured in oviposition arenas (choice assay) with 12 compartments. We randomly distributed each of 12 different inorganic phosphate content media across the 12 compartments of each oviposition arena.

In each arena, we placed a group of 20 flies aged 6 days, as described above. After 24 h, we counted the number of eggs laid in each compartment. This method is a direct approach to measuring phosphate preference and is, therefore, more powerful than other types of indirect approaches (e.g. measuring preference in arenas containing 12 puree fruits where the phosphate content would have been equalized among the fruits), thus avoiding potential false negative results. We replicated this assay 129 times. In addition, to confirm the existence of a positive correlation between oviposition preference and phosphorus content of fruit media, we performed the same protocol as Olazcuaga et al. (2019) using the 12 artificial fruit media that vary in phosphorus content. To estimate the number of replicate arenas for this confirmatory assay (hereafter 'fruit assay'), we performed a power analysis with simulations based on the subsampling of the original data (see Sup. Mat.). We found that an assay with six fruit arenas provided enough power to detect a significant effect of phosphorus content on oviposition preference in more than 99% of the simulations. Hence, we used 20 fruit arenas for the fruit assay. The position of 129 test arenas and 20 fruit arenas was randomized within the growth chamber.

## 2.5 | Statistical analyses

To investigate whether oviposition preference depends on inorganic phosphate, we tested for the effect of the inorganic phosphate content on the number of eggs laid in each compartment using a Generalized Linear Mixed Model with a log link (Poisson distribution for count data). The model included inorganic phosphate content as a fixed effect (continuous variable). To control for potential correlations in the number of eggs among compartments across and within arenas, we added the position of the compartment within the arena and arena identity as random effects (with 12 and 129 levels respectively). To account for overdispersion, we added an observation-level random effect (1548 levels). We tested for the effect of inorganic phosphate content using a Likelihood Ratio Test (LRT; Bates et al., 2015). To quantify the proportion of variance in oviposition preference explained by inorganic phosphate content, we computed the likelihood-based coefficient of determination using the *rr2* package (Ives, 2019). In addition, we separately performed the same analysis on the data from the 20 fruit arenas using the phosphorus content of the 12 fruit media as estimated in Olazcuaga et al., 2019.

## 3 | RESULTS

### 3.1 | Testing for a causal effect of inorganic phosphate on oviposition preference


The number of eggs laid in each compartment did not increase with inorganic phosphate content (LRT  $\chi^2 = 2.94$ ,  $df = 1$ ,  $p$ -value = 0.09; Figure 1a). In contrast, the number of eggs laid in each compartment increased with the phosphorus content of the 12 fruit media in fruit arenas (LRT  $\chi^2 = 23.53$ ,  $df = 1$ ,  $p$ -value < 0.001; Figure 1b).

## 4 | DISCUSSION

We used an inbred line of *D. sukuzii* to test whether inorganic phosphate content affects oviposition preference in this invasive pest. Our results confirm previous findings of a greater oviposition preference for fruit media with the highest phosphorus content. However, females did not prefer to lay eggs on minimum media with a high inorganic phosphate content.

### 4.1 | Identifying molecules involved in oviposition preference

Two hypotheses may explain why we found that females prefer to oviposit in fruit media with high phosphorus content, but not in minimum media with high inorganic phosphate content. First, although the content in macromolecules associated with phosphorus such as carbohydrates or proteins do not drive oviposition preference



**FIGURE 1** Relationship between female oviposition preference and (a) increasing content in inorganic phosphate supplemented in standard medium or (b) increasing content in phosphorus in 12 different artificial fruit media. Large- and dark-filled circles with error bars represent the mean with 95% confidence intervals, small- and light-filled circles represent the observed data. The dotted red line represents the number of eggs predicted by the best model. The  $P$ -value and coefficient of determination of the effect of phosphate or phosphorus content on oviposition preference are given above each panel.

in *D. sukukii* (Olazcuaga et al., 2019), other macromolecules (e.g. phospholipids) might drive oviposition preference. Second, oviposition preference could be driven by one or more molecules that contain no phosphorus, but whose content may be correlated with phosphorus content across the 12 fruits studied. Some volatile compounds, such as linalool, are known to be involved in the oviposition choice of *D. sukukii* (Baena et al., 2022). However, whether the presence of these compounds correlates with phosphorus content remains to be determined.

## 4.2 | Deciphering mechanisms that may influence oviposition preference

Two proximate mechanisms could explain the oviposition preference for specific fruits (Visser, 1986). First, females might be attracted by olfactory compounds which are honest cues of the quality of the fruit substrate (olfaction-driven preference; Bruce

et al., 2005). In this scenario, oviposition preference would be an active choice made by females for oviposition (i.e. associated with a non-random flight targeted towards the best fruit substrate for either female feeding or egg development). Second, females might not be attracted by olfactory compounds, but may spend more time on fruits that are more nutritious for them (gustation-driven preference; Amrein & Thorne, 2005). If females randomly lay eggs when they feed, they might eventually lay more eggs on fruits they feed on the most. To dissociate these two mechanisms, using a preference experiment, Clymans et al. (2019) showed that *D. sukukii* females are attracted to fermentation volatiles when searching for food and to fruit volatiles when searching for an oviposition substrate. Similar experiments would allow for a better understanding and identification of the factors driving host preference in *D. sukukii*. Finally, the detection of specific volatile compounds could help to improve the design of traps to increase pest trapping efficiency and ultimately improve the prediction of potential damages (Burrack et al., 2015).

### 4.3 | Integration of the results in the context of natural populations

Although the experimental conditions differ from natural conditions, especially with the use of an inbred line and fruit purees, the results found in this study represent a strong basis to draw inferences on oviposition preference in natural populations for two reasons. First, fruit media may provide a faithful sensorial representation of fruits in the field. Indeed, we recently found that female oviposition preference was higher on fruit medium corresponding to the fruit from which the female originated than on fruit media corresponding to other fruits (Olazcuaga et al., 2022). This discrimination of *D. sukukii* females among different artificial fruit media suggests that these media do harbour the same cues as natural fruits. Second, the behaviour of an inbred line accurately represents that of a natural population. Indeed, oviposition preference on 12 artificial fruit media was consistent between the inbred line (this study) and a genetically diverse population used in a previous study (Olazcuaga et al., 2019).

## 5 | CONCLUSION

By addressing an important and long-standing question regarding the factors that drive oviposition preference in crop pests, our study proposes a straightforward methodology to identify chemical compounds that may determine oviposition preference in an insect pest. This methodology can be used to test for the effects of other chemical compounds on the oviposition preference of important insect pests.

### AUTHOR CONTRIBUTIONS

Conceptualization: L.O., N.O.R., R.G. Experimental design and data acquisition: L.O. and J-L.C with inputs of R.G. and N.O.R. Statistical analyses and writing: L.O. and N.O.R with inputs of R.G. and J-L.C.

### ACKNOWLEDGEMENTS

We are very grateful to Graeme D. Batten and Lindsay C. Campbell for insightful discussions on the chemical composition of fruits and Ruth A. Hufbauer for comments on the manuscript and for insightful discussions. We also thank two anonymous reviewers for their useful comments. L.O. acknowledges support from the European Union program FEFER FSE IEJ 2014-2020 (project CPADROL), the INRAE scientific department SPE (AAP-SPE 2016) and the US National Science Foundation (DEB-1930650 to Ruth Hufbauer). N.O.R. acknowledges support from the CeMEB LabEx/University of Montpellier (ANR-10-LABX-04- 01).

### CONFLICT OF INTEREST

The authors declare that they have no conflicting interest.

### DATA AVAILABILITY STATEMENT

The data and R scripts for our analyses are available at: <https://github.com/olazlaure/PhosphorusPreference2022>. They are archived on Figshare: <https://doi.org/10.6084/m9.figshare.22032500>.

### ORCID

Laure Olazcuaga  <https://orcid.org/0000-0001-9100-1305>

Robin Guilhot  <https://orcid.org/0000-0003-1441-8486>

Jean-Loup Claret  <https://orcid.org/0000-0002-1397-5638>

Nicolas O. Rode  <https://orcid.org/0000-0002-1121-4202>

### REFERENCES

- Alkema, J. T., Dicke, M., & Wertheim, B. (2019). Context-dependence and the development of push-pull approaches for integrated management of *Drosophila sukukii*. *Insects*, 10, 454.
- Amrein, H., & Thorne, N. (2005). Gustatory Perception and Behavior in *Drosophila melanogaster*. *Current Biology*, 15, R673–R684.
- Backhaus, B., Sulkowski, E., & Schlote, F. (1984). A semi-synthetic, general-purpose medium for *Drosophila melanogaster*. *Drosophila Information Service*, 60, 210–212.
- Baena, R., Araujo, E. S., Souza, J. P. A., Bischoff, A. M., Zarbin, P. H. G., Zawadneak, M. A. C., Araujo, E. S., Souza, J. P. A., Bischoff, A. M., Zarbin, P. H. G., Zawadneak, M. A. C., & Cuquel, F. L. (2022). Ripening stages and volatile compounds present in strawberry fruits are involved in the oviposition choice of *Drosophila sukukii* (Diptera: Drosophilidae). *Crop Protection*, 153, 105883.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48.
- Bergwitz, C. (2012). Dietary phosphate modifies lifespan in *Drosophila*. *Nephrology Dialysis Transplantation*, 27, 3399–3406.
- Bourguet, D., & Guillemaud, T. (2016). The hidden and external costs of pesticide use. In *Sustainable agriculture reviews* (pp. 35–120). Springer International Publishing.
- Bruce, T. J. A., Wadhams, L. J., & Woodcock, C. M. (2005). Insect host location: A volatile situation. *Trends in Plant Science*, 10, 269–274.
- Burrack, H. J., Asplen, M., Bahder, L., Collins, J., Drummond, F. A., Guédot, C., Asplen, M., Bahder, L., Collins, J., Drummond, F. A., Guedot, C., Isaacs, R., Johnson, D., Blanton, A., Lee, J. C., Loeb, G., Rodriguez-Saona, C., van Timmeren, S., Walsh, D., & McPhie, D. R. (2015). Multistate Comparison of Attractants for Monitoring *Drosophila sukukii* (Diptera: Drosophilidae) in Blueberries and Caneberries. *Environmental Entomology*, 44, 704–712.
- Burrack, H. J., Fernandez, G. E., Spivey, T., & Kraus, D. A. (2013). Variation in selection and utilization of host crops in the field and laboratory by *Drosophila sukukii* Matsumura (Diptera: Drosophilidae), an invasive frugivore. *Pest Management Science*, 69, 1173–1180.
- Clymans, R., Van Kerckvoorde, V., Bangels, E., Akkermans, W., Alhmedi, A., De Clercq, P., van Kerckvoorde, V., Bangels, E., Akkermans, W., Alhmedi, A., de Clercq, P., Beliën, T., & Bylemans, D. (2019). Olfactory preference of *Drosophila sukukii* shifts between fruit and fermentation cues over the season: Effects of physiological status. *Insects*, 10, 200.
- Dweck, H. K. M., Ebrahim, S. A. M., Khallaf, M. A., Koenig, C., Farhan, A., Stieber, R., Ebrahim, S. A. M., Khallaf, M. A., Koenig, C., Farhan, A., Stieber, R., Weißflog, J., Svatoš, A., Grosse-Wilde, E., Knaden, M., & Hansson, B. S. (2016). Olfactory channels associated with the *Drosophila* maxillary palp mediate short- and long-range attraction. *eLife*, 5, 1–19.
- Elsensohn, J. E., Aly, M. F. K., Schal, C., & Burrack, H. J. (2021). Social signals mediate oviposition site selection in *Drosophila sukukii*. *Scientific Reports*, 11, 3796.
- Elsensohn, J. E., Schal, C., & Burrack, H. J. (2021). Plasticity in oviposition site selection behavior in *Drosophila sukukii* (Diptera: Drosophilidae) in relation to adult density and host distribution and quality. *Journal of Economic Entomology*, 114, 1517–1522.

- Emiljanowicz, L. M., Ryan, G. D., Langille, A., & Newman, J. (2014). Development, reproductive output and population growth of the fruit fly pest *Drosophila suzukii* (Diptera: Drosophilidae) on artificial diet. *Journal of Economic Entomology*, *107*, 1392–1398.
- Farnsworth, D., Hamby, K. A., Bolda, M., Goodhue, R. E., Williams, J. C., & Zalom, F. G. (2017). Economic analysis of revenue losses and control costs associated with the spotted wing drosophila, *Drosophila suzukii* (Matsumura), in the California raspberry industry. *Pest Management Science*, *73*, 1083–1090.
- Fraimout, A., Debat, V., Fellous, S., Hufbauer, R. A., Foucaud, J., Pudlo, P., Debat, V., Fellous, S., Hufbauer, R. A., Foucaud, J., Pudlo, P., Marin, J. M., Price, D. K., Cattell, J., Chen, X., Deprá, M., François Duyck, P., Guedot, C., Kenis, M., ... Estoup, A. (2017). Deciphering the routes of invasion of *Drosophila suzukii* by means of ABC random forest. *Molecular Biology and Evolution*, *34*, 980–996.
- Haviland, D. R., & Beers, E. H. (2012). Chemical control programs for *Drosophila suzukii* that comply with international limitations on pesticide residues for exported sweet cherries. *Journal of Integrated Pest Management*, *3*, F1–F6.
- Hoffmann, A. A. (1985). Effects of Experience on Oviposition and Attraction in *Drosophila*: Comparing Apples and Oranges. *The American Naturalist*, *126*, 41–51.
- Ives, A. R. (2019). R2 for correlated data: phylogenetic models, LMMs, and GLMMs. *Systematic Biology*, *68*, 234–251.
- Karageorgi, M., Bracker, L. B., Lebreton, S., Minervino, C., Cavey, M., Siju, K. P., Bräcker, L. B., Lebreton, S., Minervino, C., Cavey, M., Siju, K. P., Grunwald Kadow, I. C., Gompel, N., & Prud'homme, B. (2017). Evolution of multiple sensory systems drives novel egg-laying behavior in the fruit pest *Drosophila suzukii*. *Current Biology*, *27*, 847–853.
- Kenis, M., Tonina, L., Eschen, R., van der Sluis, B., Sancassani, M., Mori, N., Tonina, L., Eschen, R., van der Sluis, B., Sancassani, M., Mori, N., Haye, T., & Helsen, H. (2016). Non-crop plants used as hosts by *Drosophila suzukii* in Europe. *Journal of Pest Science*, *89*, 735–748.
- King, R. C., & Wilson, L. P. (1955). Studies with radiophosphorus in *Drosophila*. V. The phosphorus balance of adult females. *Journal of Experimental Zoology*, *130*, 71–82.
- Kinjo, H., Kunimi, Y., Ban, T., & Nakai, M. (2013). Oviposition efficacy of *Drosophila suzukii* (Diptera: Drosophilidae) on different cultivars of blueberry. *Journal of Economic Entomology*, *106*, 1767–1771.
- Lee, J. C., Bruck, D. J., Curry, H., Edwards, D., Haviland, D. R., Van Steenwyk, R. A., Bruck, D. J., Curry, H., Edwards, D., Haviland, D. R., van Steenwyk, R. A., & Yorgey, B. M. (2011). The susceptibility of small fruits and cherries to the spotted-wing drosophila, *Drosophila suzukii*. *Pest Management Science*, *67*, 1358–1367.
- Lee, J. C., Dreves, A. J., Cave, A. M., Kawai, S., Isaacs, R., Miller, J. C., Dreves, A. J., Cave, A. M., Kawai, S., Isaacs, R., Miller, J. C., van Timmeren, S., & Bruck, D. J. (2015). Infestation of wild and ornamental noncrop fruits by *Drosophila suzukii* (Diptera: Drosophilidae). *Annals of the Entomological Society of America*, *108*, 117–129.
- Little, C. M., Chapman, T. W., & Hillier, N. K. (2018). Effect of color and contrast of highbush blueberries to host-finding behavior by *Drosophila suzukii* (Diptera: Drosophilidae). *Environmental Entomology*, *47*, 1242–1251.
- Little, C. M., Rizzato, A. R., Charbonneau, L., Chapman, T., & Hillier, N. K. (2019). Color preference of the spotted wing drosophila, *Drosophila suzukii*. *Scientific Reports*, *9*, 16051.
- Lott, J. N. A., Ockenden, I., Raboy, V., & Batten, G. D. (2000). Phytic acid and phosphorus in crop seeds and fruits: a global estimate. *Seed Science Research*, *10*(1), 11–33.
- Markow, T. A., Raphael, B., Dobberfuhr, D., Breitmeyer, C. M., Elser, J. J., & Pfeiler, E. (1999). Elemental stoichiometry of *Drosophila* and their hosts. *Functional Ecology*, *13*, 78–84.
- Marschner, P. (2012). *Marschner's mineral nutrition of higher plants* (P. Marschner, Ed.; 3rd ed.) Academic Press.
- Oerke, E.-C. (2006). Crop losses to pests. *The Journal of Agricultural Science*, *144*, 31–43.
- Olazcuaga, L., Foucaud, J., Deschamps, C., Loiseau, A., Claret, J.-L., Vedovato, R., Foucaud, J., Deschamps, C., Loiseau, A., Claret, J. L., Vedovato, R., Guilhot, R., Sévely, C., Gautier, M., Hufbauer, R. A., Rode, N. O., & Estoup, A. (2022). Rapid and transient evolution of local adaptation to seasonal host fruits in an invasive pest fly. *Evolution Letters*, *6*, 490–505.
- Olazcuaga, L., Rode, N. O., Foucaud, J., Facon, B., Ravigné, V., Ausset, A., Rode, N. O., Foucaud, J., Facon, B., Ravigné, V., Ausset, A., Leménager, N., Loiseau, A., Gautier, M., Estoup, A., & Hufbauer, R. A. (2019). Oviposition preference and larval performance of *Drosophila suzukii* (Diptera: Drosophilidae), spotted-wing *Drosophila*: Effects of fruit identity and composition. *Environmental Entomology*, *48*, 867–881.
- Paris, M., Boyer, R., Jaenichen, R., Wolf, J., Karageorgi, M., Green, J., Boyer, R., Jaenichen, R., Wolf, J., Karageorgi, M., Green, J., Cagnon, M., Parinello, H., Estoup, A., Gautier, M., Gompel, N., & Prud'homme, B. (2020). Near-chromosome level genome assembly of the fruit pest *Drosophila suzukii* using long-read sequencing. *Scientific Reports*, *10*, 11227.
- Renwick, J. A. A. (1989). Chemical ecology of oviposition in phytophagous insects. *Experientia*, *45*, 223–228.
- Schetelig, M. F., Lee, K.-Z., Otto, S., Talmann, L., Stökl, J., Degenkolb, T., Lee, K. Z., Otto, S., Talmann, L., Stökl, J., Degenkolb, T., Vilcinskis, A., & Halitschke, R. (2018). Environmentally sustainable pest control options for *Drosophila suzukii*. *Journal of Applied Entomology*, *142* (3–17), 3.
- Silva-Soares, N. F., Nogueira-Alves, A., Baldade, P., & Mirth, C. (2017). Adaptation to new nutritional environments: larval performance, foraging decisions, and adult oviposition choices in *Drosophila suzukii*. *BMC Ecology*, *17*, 1–13.
- Singer, M. C. (1986). The definition and measurement of oviposition preference in plant-feeding insects. In J. R. Miller & T. A. Miller (Eds.), *Insect-plant interactions*, Springer Series in Experimental Entomology (pp. 65–94). Springer.
- Tait, G., Park, K., Nieri, R., Crava, M. C., Mermer, S., Clappa, E., Park, K., Nieri, R., Crava, M. C., Mermer, S., Clappa, E., Boyer, G., Dalton, D. T., Carlin, S., Brewer, L., Walton, V. M., Anfora, G., & Rossi-Stacconi, M. V. (2020). Reproductive Site Selection: Evidence of an Oviposition Cue in a Highly Adaptive Dipteran, *Drosophila suzukii* (Diptera: Drosophilidae). *Environmental Entomology*, *49*, 355–363.
- Takahara, B., & Takahashi, K. H. (2017). Associative learning of color and firmness of oviposition substrates in *Drosophila suzukii*. *Entomologia Experimentalis et Applicata*, *162*, 13–18.
- Visser, J. H. (1986). Host odor perception in phytophagous insects. *Annual Review of Entomology*, *31*, 121–144.
- Young, Y., Buckiewicz, N., & Long, T. A. F. (2018). Nutritional geometry and fitness consequences in *Drosophila suzukii*, the spotted-wing drosophila. *Ecology and Evolution*, *1–10* (8), 2842.

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Olazcuaga, L., Guilhot, R., Claret, J.-L., & Rode, N. O. (2023). Inorganic phosphate content does not affect oviposition preference in the invasive pest *Drosophila suzukii*. *Journal of Applied Entomology*, *00*, 1–6. <https://doi.org/10.1111/jen.13107>