



Cytogenotoxicity screening of urban and rural marshes: An integrated in vivo approach coupling fish and plant-based tests adapted for low income countries

Elodie Pepey, Geneviève Conéjéro, Elia Njara Ndrianaivo, Jean-luc Verdeil,
Jean-michel Mortillaro, David Boggio, Modestine Raliniaina, Simon Pouil

► To cite this version:

Elodie Pepey, Geneviève Conéjéro, Elia Njara Ndrianaivo, Jean-luc Verdeil, Jean-michel Mortillaro, et al.. Cytogenotoxicity screening of urban and rural marshes: An integrated in vivo approach coupling fish and plant-based tests adapted for low income countries. *Environmental Toxicology and Chemistry*, 2023, 42 (6), pp.1266-1275. 10.1002/etc.5605 . hal-04028206v1

HAL Id: hal-04028206

<https://hal.inrae.fr/hal-04028206v1>

Submitted on 14 Mar 2023 (v1), last revised 5 Sep 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Elodie Pepey ORCID iD: 0000-0003-2624-0929

David Boggio ORCID iD: 0000-0003-2320-7250

Simon Pouil ORCID iD: 0000-0003-1531-0362

Cytogenotoxicity screening of urban and rural marshes: An integrated in vivo approach coupling fish and plant-based tests adapted for low income countries

Elodie Pepey^{1,2*}, Geneviève Conéjéro³, Elia Njara Ndrianaivo⁴, Jean-Luc Verdeil^{5,6}, Jean-Michel Mortillaro^{1,2,4}, David Boggio⁷, Modestine Raliniaina⁴, Simon Pouil⁸

¹ CIRAD, UMR ISEM, F-34398 Montpellier, France

² ISEM, University of Montpellier, CNRS, EPHE, IRD, Montpellier, France

³ BPMP, University of Montpellier, CNRS, INRAE, Montpellier SupAgro, Montpellier, France

⁴ FOFIFA DRZVP, Antananarivo, Madagascar

⁵ PHIV, CIRAD, UMR AGAP Institut, 34398, Montpellier, France

⁶ UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, 34398, Montpellier, France

⁷ CIRAD, AMO, DSI, F-34398 Montpellier, France

⁸ Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, 37830, USA

* Correspondence: elodie.pepey@cirad.fr

Acknowledgement:

Authors are most grateful to Marie-Charline Raivomanana and Falimalala Andrianarimandimby (FOFIFA), Mylène Lorcin (INRS), Marc Canonne (Cirad) and Marc Lartaud (Cirad) for their technical assistance and to Hugues de Verdal (Cirad) for his advice on the experimental procedures implemented for animal welfare. The authors are also grateful to the NGO APDRA Pisciculture Paysanne for technical and logistic assistance before or during field work in Madagascar. We acknowledge the imaging facility MRI, member of the national infrastructure France-BioImaging supported by the French National Research Agency (ANR-10-INBS-04). We are grateful to the CIRAD Water, Soil and Plant Analysis Laboratory (US Analyses, CIRAD, Montpellier, France) for the analyses of trace elements. This is the publication ISEM XX.

Data availability Statement

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/etc.5605.

This article is protected by copyright. All rights reserved.

The data and methodology supporting the results of the present study are available within the article. Regarding the cytogenotoxicity data, the nature of our study did not generate large data sets requiring the use of data repositories. However, the corresponding author (E. Pepey; elodie.pepey@cirad.fr) may be contacted for further inquiries regarding the data.

Author Contribution statement:

Elodie Pepey: Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Supervision, Writing - Review & Editing, Funding acquisition. **Geneviève Conéjéro:** Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Visualization, Writing - Review & Editing, Funding acquisition. **Elia Njara Ndrianaivo:** Investigation, Resources. **Jean-Luc Verdeil:** Methodology, Formal analysis, Writing - Review & Editing. **Jean-Michel Mortillaro:** Resources, Writing - Review & Editing. **David Boggio:** Software, Formal analysis. **Modestine Raliniaina:** Resources. **Simon Pouil:** Conceptualization, Formal analysis, Writing - Original Draft, Visualization, Writing - Review & Editing

Abstract

Effects of anthropogenic activities such as urbanisation, population growth and agriculture on water quality are major concerns particularly in low-income countries where water quality monitoring can be challenging. The purpose of this study was to evaluate the cytogenotoxic potential of water from urban and rural Malagasy marshes, coupling a fish (Nile tilapia *Oreochromis niloticus*) and a plant (*Allium cepa*) species as bioindicators. The fish and plants were exposed for 72 h to water sampled in the two locations investigated. DNA strand breaks were assessed using the comet assay on fish erythrocytes, while mitotic index and nucleolar alterations were estimated in cells of the plant root apex. Comet assays revealed significant DNA strand breaks to fish erythrocytes in both the marshes investigated while the mitotic index and nucleolar characteristics in the roots of *A. cepa* mainly highlighted potential cytotoxicity in the urban marsh. Our results demonstrate the advantages of coupling *in vivo* biological test systems to screen potential cytogenotoxicity of surface water in low-income countries where comprehensive datasets of aquatic contaminants are often lacking.

Keywords: *Allium cepa*; Comet assay; Madagascar; Mitotic index; Nucleolar alterations; *Oreochromis niloticus*

INTRODUCTION

Water availability and quality are among the most important concerns for human societies (McLennan, 2021). Water quality has deteriorated from urbanisation, population growth, industrialisation and agriculture. Today, poor water quality is widespread and has a major negative effect on many aspects of human society, economy, ecology and the environment (Kirschke et al., 2020). In low-income countries, concerns about the distribution and allocation of water resources and water pollution are increasing. So far, water monitoring data are lacking in less-developed countries, notably because of capacity challenges to define an enabling environment or prioritizing parameters as well as limiting financial and human means (Capps et al., 2015; Kirschke et al., 2020).

One of the main concerns in water quality deterioration is potential exposure to substances that are both cytotoxic and genotoxic. Cytogenotoxicity can only be evaluated by biological-oriented tests (Hemachandra & Pathiratne, 2017). Among the toxic effects induced by chemicals, main concerns are the mutagenic/genotoxic hazards which could lead to ecological and human health risks (Ohe et al., 2004). Coupling different *in vivo* models, such as plants and fishes, and different targeting endpoints with direct exposure to the water samples allows an integrated view of the cytogenotoxic risk for aquatic biomonitoring, as recommended by Orsière et al. (2005). Furthermore, several studies already shown that genotoxicity tests based on fish and plants are producing results showing good correlations with human and other mammalian test systems (e.g., Dos Reis et al., 2017; Leme & Marin-Morales, 2009, Vlastos et al., 2016) suggesting that they are appropriate models for assessing cytogenotoxicity potential of contaminated waters to humans.

This article is protected by copyright. All rights reserved.

The potential benefits of using fish as indicators of cytogenotoxicity are now widely recognized (de Lapuente et al., 2015). This is due to their ecological and economic importance and their similarity to mammals for which toxicological mechanisms are far better described (Martins & Costa, 2017). The comet assay consisting in quantifying DNA breaks by gel electrophoresis (Singh et al., 1988), applied to fish erythrocytes, is known to be a sensitive way to monitor aquatic environments affected by pollutants (de Lapuente et al., 2015; Frenzilli et al., 2009; Martins & Costa, 2017). Compared with other genotoxicity tests, the advantages of the comet assay are the detection of slight DNA damage, the low number of cells required, low cost, precision, ease of application, reproducibility, and short period of time to conduct the experiment. In plants, the *Allium cepa* assay, assessing chromosome damages and disturbances in the mitotic cycle, is well-known among plant cytogenotoxicity tests and is now used in many laboratories for a wide range of applications (Leme & Marin-Morales, 2009). As for the comet-assay test, the advantages of the *A. cepa* test are that it is fast, easy to use, inexpensive and requires minimal sample preparation (Barbério et al., 2013; Leme & Marin-Morales, 2009). For these reasons, all these trials can be conducted in remote areas from low-income countries because such tests require limited infrastructures to be performed (Hemachandra & Pathiratne, 2016; 2017).

The objective of this study was to investigate the cytogenotoxic potential of water sampled from two marshes in Madagascar, where urban population is growing at twice the rate of that in the rest of the world (+ 5% per year; Bastaraud et al., 2018). One of the two marshes is mainly used to supply water to the capital city of Antananarivo (> 1 300 000 inhabitants) while the second is mainly surrounded by agricultural land. Given the risk for aquatic organisms as well as the risk of contaminants entering the food chain and affecting human health, we chose to assess

in vivo exposure using Nile tilapia *Oreochromis niloticus*, an aquatic species, and, *A. cepa*, a terrestrial species, as bioindicators of cytogenotoxicity. These species are consumed by local populations (FAO, 2022) and are both present in the two locations.

MATERIALS AND METHODS

Study areas and sample collection

The investigated urban marsh (Masay) is an artificial basin built in 2004 measuring approximately 1 km² located in the northern part of the city of Antananarivo (18°52'51"S, 47°32'23"E) acting as a buffer basin for wastewater and runoff from the surrounding districts (Figure 1). The basin, opens into the Andriantany canal and the Ikopa river, collects rainwater and untreated domestic wastewater and industrial discharges from two watersheds and regulates excess water in order to prevent flooding. Masay marsh is surrounded by residential housing and agri-food industrial establishments and numerous car garages while uncontrolled landfills are also existing near the marsh. Locally, several unpublished studies highlighted the pollution of this marsh notably by heavy metals and other contaminants with sometimes illegal fishing for self-consumption and sale on local markets. For the purpose of comparison, water was also collected in a rural marsh located along the Viliahazo river (18°48'5"S, 47°40'13"E) which is used for agriculture, mostly for rice and cassava cultivation (Figure 1) and fishing. There is also a significant market gardening activity in the area allowing the supply of vegetables to the city of Antananarivo, an agriculture activity requiring much more pesticides compared to other cropping systems (Razanakoto et al., 2021).

The investigation was carried out in the rainy season, February 2019. Unfiltered surface water was collected from the urban and rural marshes at a depth of 50 cm. For technical reasons, the samples of water were kept oxygenated using air

This article is protected by copyright. All rights reserved.

bubbling and stored for three days at room temperature in the dark in separate 250-L food-grade plastic drums prior to the experiment. Ceretti et al. (2010) found that storage materials were not associated with the genotoxic properties of the water even after 10 days of storage at 40 °C. Thus, it is reasonable to assume that our storage conditions had no effect on water toxicity.

Water analysis

The physical-chemical parameters of the water under each experimental condition were measured at the beginning of the experiment. The temperature and the conductivity of the water were determined using a combined tester (HI 98311, HANNA Instruments Co.) while pH was measured using a portable digital pH probe (HI 98128, HANNA Instruments Co.). Dissolved oxygen (DO) was measured using a DO meter (HI 9146, HANNA Instruments Co.) and the biochemical oxygen demand after five days (BOD5) was analysed as described by Aminot & K  rouel (2004).

After the storage period, 100 mL sub-samples of water originating from each location were taken from each drum and filtered using 0.2   m PES filters and kept in 50 mL Falcon   tubes with trace element grade nitric acid (pH < 2) for the analysis of trace metals. All the samples were kept at room temperature in the dark prior to analysis.

As potential cytogenotoxic agents (Tchounwou et al., 2012), the dissolved and particulate trace metals (i.e. As, Co, Cr, Cu, Fe, Mn, Ni, Pb and Zn) were analysed. In the water samples kept in Falcon   tubes, trace elements were directly analysed by ICP-MS (ICP-MS X series 2 Thermo Scientific). Trace metals on the filters were digested using hydrofluoric and perchloric acids following method ISO 14869-1. All quality assurance procedures (e.g. blanks; reference materials: SLRS6 and CRM 7004

(Metranal 34)) were applied in each analytical method. Only the elements recovered from reference materials in the 80-120% range.

Ethical statement

The study was carried out in accordance with the recommendations of Directive 2010-63-EU concerning the protection of animals used for scientific purposes. The protocols were evaluated and approved by the ethical committee C2EA – 36 (Comité d'éthique en expérimentation animale Languedoc-Roussillon) and authorized by the French ministry of higher education and research (APAFiS n° 2020031711429384 #2020-01 v2.universities).

Bioassay assembly

Thirty-two adult Nile tilapia (fresh weight: 127 ± 48 g) were obtained from a local fish farm. Fish were randomly placed in four 150 L glass aquariums ($n = 5$ per aquarium) filled with dechlorinated tap water and equipped with air-lift filters and hiding places. Fish were acclimated for 10 days under a natural light and natural photoperiod and were fed with a commercial diet (PGR32, Aqua Madagascar Innovation). Feeding was stopped 48 hours before the bioassays began. After the acclimation period, the water in the two aquariums was replaced by water collected from the urban and rural marshes, while the two other aquariums were used as controls. While dechlorinated tap water was used as a negative control, a further set of fish were exposed to ethyl methanesulfonate (EMS) at 10 mg L^{-1} and used as positive controls, similarly to a previous genotoxicity investigation in Nile tilapia (Könen & Çavaş, 2008). After the 3-days exposure, five fish from each condition, including controls, were anesthetised with Eugenol (0.5 mL L^{-1}) and blood was sampled from the ventral tail vein using a heparinized syringe.

For the *A. cepa* test, we followed the protocol initially proposed by Fiskesjö (1985). Briefly, the plants were first grown on plastic glasses in tap water. When the roots were at least one cm long, they were transferred to the water collected from the urban and rural marshes while, as for fish, dechlorinated tap water and EMS at 10 mg L⁻¹ were used as negative and positive controls, respectively. After 72 h of exposure, root tips from all experimental and control conditions (n = 4 bulbs per condition) were processed for microscopy. The test was conducted under laboratory conditions in a shaded area at room temperature.

Bioassay procedures

Comet assay using fish. The alkaline comet assay using Nile tilapia erythrocytes was performed according to Singh et al. (1988) and Osman et al. (2012) with some modifications. Before the assay, frosted microscope slides were pre-coated with 1% (w/v) of normal-melting agarose (NMA) and dried at room temperature for at least two days. Each fish blood puncture was immediately diluted with phosphate-buffered saline (PBS) at a ratio of 1:500 (v/v). Then, 40 µL of each diluted blood sample was added to 360 µL of 0.5% (w/v) low-melting agarose (LMA) and 150 µL of the mixture was immediately placed on the pre-coated slide and covered with a coverslip. The slides were incubated at 4 °C for 15 min to allow solidification then immersed in cold lysing buffer (2.5 M NaCl, 100 mM EDTA, 10 mM TRIS base, pH 10, 1% Triton X-100, 10% DMSO) at 4 °C overnight (Hemachandra & Pathiratne, 2016; Iturburu et al., 2018). Then, the slides were incubated at 4 °C in electrophoresis buffer (300 mM NaOH, 1 mM EDTA, pH ≥ 13) for 40 min, and electrophoresis was carried out at 0.8 V/cm and 300 mA at 4 °C for 40 min. A neutralisation step was then performed consisting of three successive 10-min incubation periods in 0.4 M Tris buffer (pH 7.5). Slides were rinsed in ultra-pure water for 5 min then immersed in ethanol (≥ 99.6 %) for 10 min to start dehydration before being dried at 45 °C for 1 h.

This article is protected by copyright. All rights reserved.

Finally, the slides were stained with 100 μ L of SYBR Gold (Thermo Fisher Scientific, Massachusetts, USA) diluted x10,000 with Tris/EDTA buffer (pH 8.0) and the comets were viewed under an epifluorescence microscope (Nikon Eclipse NI-E, objective Plan APO 0,75 NA 20x). For each fish, we analysed two slides each containing 200 cells (i.e. 400 cells per fish and approx. 2000 cells per condition). The images were analysed using the plug-in OpenComet with ImageJ software (Gyori et al., 2014) and the extent of DNA strand breaks was expressed as the percentage of DNA in the tail (Kumaravel & Jha, 2006).

Mitotic index and nucleolar characteristics in the plant. The 3-5 mm long root tips were cutted and were immediately transferred in a fixative solution (ethanol:glacial acetic acid 3:1, v/v) and left overnight (Pathiratne et al., 2015). After fixation, the root tips were transferred to 70% ethanol and stored at 4 °C until analysis. Samples were hydrolysed in 1 M HCl at 60 °C for 5 min (Creber et al., 1994) and washed three times with distilled water. The samples were processed with a ClearSee clearing protocol to enable visualisation of all the nuclei in the meristem region (Kurihara et al., 2015). After washing with a phosphate buffer, the root tips were immersed in a ClearSee solution (urea 25% w/v, sodium deoxycholate 15% w/v, xylitol, 10%, w/v) for 5 to 7 days under gentle shaking. The nuclei were then stained with a DAPI solution (5 μ g mL⁻¹) prepared in ClearSee solution for 60 min. After washing, five root tips were placed on a glass slide and gently squashed in a drop of ClearSee solution by applying very slight pressure to the coverslip. Pictures of the meristem tissues were acquired with a multiphotonic microscope, 880 NLO Zeiss (Iena, Germany) equipped with an Apochromatic 20x NA 1 objective, at 740 nm of excitation (band-pass emission, 405-495 nm; Plant Cell Imaging Platform PHIV, Montpellier Ressources Imagerie, ISO9001 certified). A set of confocal images was taken of the specimen in such a way that the area of the image along the x- and y-axes

This article is protected by copyright. All rights reserved.

remains the same but the distance from the objective (z-axis) differs in each image (z-stack). Z-stacks were performed on 100 μm with 1 μm between two optical slices (ZEN soft, Zeiss, Jena, Germany). As the distance between adjacent images is precisely controlled in the z-stack, it can be used to form a 3-dimensional image. This method allows us to obtain pictures on several intact cell layers. Some pictures were obtained with an 40X LD C-Apo 1.1NA W or an 63X Plan Apo 1.4 NA oil objective to observe the morphology of the nuclei.

The mitotic index was used to evaluate the cytotoxic potential in each of the experimental conditions. It was calculated as the percentage of cells undergoing mitotic stages (metaphase, anaphase, and telophase) in at least 1000 cells of meristem tissues observed per root apex and three root apex analysed by plant (i.e. approx. 3000 cells per plant and 10000-12000 cells per condition). In complement to the mitotic index, quantitative analysis of nucleoli has also been proposed as a sensitive biomarker for routine monitoring of cytotoxicity in plants (Arkhipchuk et al., 2000; Lima et al., 2019). In this study, nucleolar alterations were determined by counting the number of nucleoli per nucleus and expressed as a percentage of approximately 1000 nuclei of root meristem cells per condition (Singh et al., 1988). Cell countings were performed using the Cell Counter plugin in ImageJ-1.53a software.

Statistical analysis

The differences in tail DNA (in fish), mitotic index and nucleolar activity (in plant) among the different experimental conditions were evaluated by One-way ANOVA followed by Tukey's post hoc test. Data were arcsine transformed and average values were used in the statistical analysis. The normality and homogeneity of variances were graphically confirmed and evaluated by Shapiro-Wilk and Levene's tests. In addition, a non-parametric Spearman's correlation analysis was used with concentrations of trace elements in samples of water collected in the two locations

This article is protected by copyright. All rights reserved.

prior to the experiment. All statistical analyses were performed in R v.3.6.3 (R Development Core Team, 2022) using the packages *car* and *stats* while the package *ggplot2* was used to create graphics. The level of significance for statistical analyses was always set to $\alpha = 0.05$. If not specified, the results are expressed as mean $\pm$ SD.

RESULTS

DNA damage in fish and plants

Comet assay in fish. After 72 h of exposure, no mortality was recorded in control and experimental conditions. (i.e. water from the urban and rural marshes). Comet assays in blood cells (Figure 2) revealed significant differences in DNA strand breaks between conditions with high variability among individuals ($p < 0.001$). The genotoxicity detected in fish ($n = 5$) from the two marshes (mean $\pm$ SD: $43.7 \pm 25.8\%$, median: 42.1% for the urban marsh and mean $\pm$ SD: $39.2 \pm 26.6\%$, median: 36.3% for the rural marsh) was lower ($p < 0.001$) than that found in the positive control group (i.e., fish exposed to EMS, mean $\pm$ SD: $64.1 \pm 24.1\%$, median: 71.9%) but higher ($p < 0.001$) than the DNA strand breaks found in the negative group (i.e., fish kept in tap water, mean $\pm$ SD: $25.3 \pm 22.8\%$, median: 17.1%, Figure 2A). The DNA strand breaks to fish exposed to water from the two marshes was then compared to the damage detected in control conditions and confirmed by observations of DNA degradation in erythrocytes by epifluorescence microscopy (Figure 2B).

Mitotic index and nucleolar characteristics. Cytotoxicity was assessed by determining both the mitotic index and nucleolar alterations in plants (Figures 3 and 4). Significant differences were found in the mitotic index estimated for the cells of meristem tissues in the root apex of *A. cepa* between the experimental conditions ($p < 0.001$). Pairwise comparisons revealed that the mitotic index in the plants exposed to water sampled from the urban and rural marshes differed significantly ($p = 0.035$)

This article is protected by copyright. All rights reserved.

with values of 1.18 ± 0.75 ($n = 4$ plants) from that in plants exposed to water taken from the urban marsh and 3.02 ± 0.64 in plants exposed to water taken from the rural marsh. Interestingly, while the mitotic indices were similar ($p = 0.99$) in plants exposed to water taken from the rural and in plants exposed to tap water (i.e., negative controls), the mitotic indices of the plants exposed to water taken from the urban marsh were similar to the indices observed when plants were exposed to EMS (i.e., positive controls; $p = 0.345$, Figure 3). These results show that the mitotic index was only affected by exposure to water taken from the urban marsh. Significant differences were observed between the experimental conditions in the proportion of cells with ≥ 3 nucleoli per nucleus ($p < 0.001$). Nucleolar activity in plants exposed to water taken from the two marshes did not differ from the negative control ($p \geq 0.07$) but the proportion of cells with ≥ 3 nucleoli per nucleus were lower in the rural marsh than in the urban marsh (i.e. $1.4 \pm 0.7\%$ vs $12.0 \pm 6.0\%$; $p = 0.027$; Figure 4).

Correlations between cytogenotoxicity and concentrations of trace elements in the water

The results of water quality measurements and trace elements analysis are given in Table 1. Spearman's correlations revealed no significant correlations between cytogenotoxicity results (comet assay in fish; mitotic index and nucleoli activity in plant) and the concentrations of trace elements in water ($p \geq 0.13$). Among the trace elements we investigated, concentrations of the non-essential elements in the unfiltered water were always $< 1 \mu\text{g L}^{-1}$ (Table 1).

DISCUSSION

To the best of our knowledge, this is the first time cytogenotoxic effects of aquatic environments have been assessed in Madagascar. We combined different approaches for one species of fish (*O. niloticus*) and one species of plant (*A. cepa*) to

This article is protected by copyright. All rights reserved.

evaluate the levels of cytogenotoxic pollution in two marshes; one located in an urban area and one in a rural area of the Malagasy highlands. Such investigation is needed to highlight the interest of implementing ecotoxicity assessment tools in the management of surface water in Madagascar.

The alkaline in vivo comet assay can be used to detect DNA breaks in aquatic species following exposure to genotoxics (Frenzilli et al., 2009) and is recommended for use by regulatory agencies (Speit et al., 2015). The Nile tilapia is known to respond rapidly to environmental alterations (Vijayan et al., 1997) making it a good candidate for genotoxic studies while erythrocyte-based comet assay has already been successfully used for cytogenotoxicity assessments (Christofolletti et al., 2009; de Campos Ventura et al., 2008; Hemachandra & Pathiratne, 2016; Matsumoto et al., 2006). Cytogenotoxic effects can be detected very soon after exposure to contaminated waters. For instance, using the comet assay, cytogenotoxic effects have been detected in common carp *Cyprinus carpio* after 24 h of exposure to industrial plant effluents (SanJuan-Reyes et al., 2015) while Matsumoto et al. (2006) found genotoxicity and mutagenicity in Nile tilapia exposed 72 h to water contaminated with tannery effluents.

In the present study, we found that a greater cytogenotoxicity was induced by the contaminants in the water taken from the urban and rural marshes we investigated when compared to the negative control condition. According to de Souza Trigueiro et al. (2021), the median value of the percentage of DNA damage in the tail suggested that most of the damages were minimal or low for fish coming from the negative control condition while fish exposed to the water from the two marshes exhibited mostly high damages. Such genotoxic effects in fish can lead to several cellular and physiological alterations such as mutations changes in the lifespan and functions of circulating erythrocytes and cell death (Çavaş & Könen, 2007; da Silva et al., 2019; de

This article is protected by copyright. All rights reserved.

Souza Trigueiro et al., 2021). The slight changes observed in the genotoxic results between the two marshes combined with the absence of correlations between comet-assay results and concentrations of trace elements suggest that (1) the genotoxic agents involved are not trace elements (2) the genotoxic agents are present in both the urban and rural marshes although probably not present at the same concentrations in the marshes and/or (3) the genotoxic effects observed were not caused by the same contaminants in the two locations investigated in this study.

The *A. cepa* test is one of the most efficient, feasible and low-cost methods for the assessment of the cytogenotoxic effects of mixtures of contaminants in aquatic environments (Leme & Marin-Morales, 2009). In the *A. cepa* test, a decrease in the mitotic index in the root tip cells points to cytotoxicity, while induction of chromosomal abnormalities, micronuclei and nuclear buds in the root cells indicate cytogenotoxicity and can also be used to check for mutagenicity (Leme & Marin-Morales, 2009). In the present study, we combined the measurement of the mitotic index and observations of nucleolar abnormalities with a focus on the number of nucleoli per nucleus.

Nucleolar activity is sometimes considered to be a more sensitive parameter than the parameters mentioned above in the cytotoxicity studies (Lima et al., 2019). In *A. cepa*, some trace elements including Al, Cd, Cu, Pb have been reported to disrupt the synthesis of ribosomal RNAs but also to inhibit nucleolus reconstruction at the end of the telophase (Jiang et al., 2014; Qin et al., 2010; Wang et al., 2016). At the root apex, the nuclei cells mainly have two nucleoli, which sometimes merge at the end of the telophase (Jiang et al., 2014). Thus, observations of three or more nucleoli per nucleus in root meristem cells point to cytotoxicity (Lima et al., 2019). The 3D images obtained thanks to multiphoton microscopy allowed us to observe the nuclei in all their thickness and to count the exact number of nucleoli, which are not always on the

This article is protected by copyright. All rights reserved.

same plane in each nucleus, unlike what is visible in the 2-dimensional images commonly used for such measurements. Overall, the mitotic index and nucleolar activity measured on *A. cepa* indicated potential cytotoxicity of urban marsh water that primarily inhibited cell division, whereas no cytotoxicity of rural marsh water was found.

Cytogenotoxicity tests performed in fish and plants pointed to increasing toxicity effects ranked as follows: tap water (negative control) $\leq$ rural marsh $\leq$ urban marsh $\leq$ EMS (positive control), with several discrepancies between the comet assay performed in fish and the other cytotoxicity markers in plants. There are several possible explanations for these differences. Physical barriers in plants (i.e. cell walls and plasma membrane of the root) can prevent certain chemicals from entering, whereas in fish, chemicals present in the water can enter the body directly (Matsumoto et al., 2006). Furthermore, depending on the contaminants involved and their respective concentrations, cytogenotoxic effects may occur in fish but not in plants. Radić et al. (2013) found that although genotoxicity responses to the exposure of fertilizer factory effluents were mostly in agreement in common carp and duckweed *Lemna minor*, several discrepancies were observed. The authors linked these differences to differences in the sensitivity of model organisms or in experimental conditions.

The trace elements measured in the samples of water collected from the two marshes and from drinking water were within the Malagasy drinking water standards (MEAH, 2004) and do not exceed the WHO guidelines for drinking water quality (WHO, 2017). Nevertheless, both *in vivo* test systems (fish erythrocyte and *A. cepa* root-based tests) confirmed the cytogenotoxicity of the marshes' waters. The absence of correlations with the trace elements in water suggests that the cytogenotoxic effects are not due to only one of the elements measured. However, since our chemical

This article is protected by copyright. All rights reserved.

characterisation of these water samples was not comprehensive, it is difficult to conclude on the actual chemical contamination status as a complex mixture of chemicals at trace concentrations can have cytogenotoxic impacts. In addition to trace elements (Barbosa et al., 2010), a wide range of contaminants can have cytogenotoxic effects that are likely to be present in Madagascar, and potentially in our study locations.

Anthropogenic activities produce other contaminants of concern in Madagascar including persistent organic pollutants that are likely to have cytogenotoxic effects (Baqar et al., 2017). Through a screening of environmental pollutants in waste waters, Rasoazanany (2008) found that by comparison to the rural environment, the urban environment of Antananarivo is disrupted by the polluting elements including Cr and Pb. The nature of the pollutants occurring in an urban area and a rural area may differ. The agricultural practices used in cultivating vegetable crops in the vicinity of the rural marsh (Figure 1) included slash and burn, which may leach toxic pyrrolizidine alkaloids into the soil (Letsyo et al., 2021), and the use of pesticides could also explain the cytogenotoxic effects observed in this water body. A more comprehensive dataset of contamination levels of Malagasy water bodies is needed to fully explain our observations.

It has also been shown that the toxicity of surface water can vary with the season (Wu et al., 2004). Meteorological and hydrological factors may greatly affect water quality. In the present study, the experiment has been performed during the rainy season. We can thus hypothesise that rains may dilute pollutants dissolved in the water. Further investigations carried out during the dry season would be relevant to assess such changes. Indeed, several studies performed in tropical areas highlighted that induction of DNA damages in fish erythrocytes and gill tissues are higher in summer than in winter (Nagpure et al., 2015; Scalón et al., 2013). Our findings

This article is protected by copyright. All rights reserved.

suggest that, although specific priorities need to be considered depending on countries, cytogenotoxicity assessments in tropical areas should consider potential seasonal or time-specific changes especially in the context of global climate change.

CONCLUSION

All the bioassays we performed revealed cytogenotoxic potential, particularly in the marsh that received urban, industrial and artisanal waste from the capital city Antananarivo. We demonstrated the interest of using fish and plant tests combined to assess cytogenotoxicity. Indeed, while the comet assay performed on fish erythrocytes showed genotoxicity for both marshes, the tests on *A. cepa* showed differential cytotoxicity between the waters from the two sources. The present study provides scientific evidence for the urgent need to include ecotoxicity assessment tools in the management of surface water used for human activities in Madagascar. In addition, this study showed that *in vivo* test systems combining fish and plants are efficient in low income countries where comprehensive water quality monitoring programmes can be complex to achieve due to limited resources.

REFERENCES

- Aminot, A., & K  rouel, R. (2004). *Hydrologie des   cosyst  mes marins: param  tres et analyses*. Editions IFREMER.
- Arkhipchuk, V. V., Malinovskaya, M. V., & Garanko, N. N. (2000). Cytogenetic study of organic and inorganic toxic substances on *Allium cepa*, *Lactuca sativa*, and *Hydra attenuata* cells. *Environmental Toxicology: An International Journal*, 15(4), 338-344. [https://doi.org/10.1002/1522-7278\(2000\)15:4<338::AID-TOX10>3.0.CO;2-R](https://doi.org/10.1002/1522-7278(2000)15:4<338::AID-TOX10>3.0.CO;2-R)
- Baqar, M., Arslan, M., Sadeh, Y., Mahmood, A., Qadir, A., & Ahmad, S. R. (2017). Persistent organic pollutants in Pakistan: Potential threat to ecological

integrities in terms of genotoxicity and oxidative stress. *Human and Ecological Risk Assessment: An International Journal*, 23(6), 1249-1271. <https://doi.org/10.1080/10807039.2017.1292843>

Barbério, A. (2013). Bioassays with plants in the monitoring of water quality. In W. Elshorbagy & R. Chowdhury (Eds.), *Water Treatment* (pp. 317-334). InTech.

Barbosa, J. S., Cabral, T. M., Ferreira, D. N., Agnez-Lima, L. F., & De Medeiros, S.

B. (2010). Genotoxicity assessment in aquatic environment impacted by the presence of heavy metals. *Ecotoxicology and Environmental Safety*, 73(3), 320-325. <https://doi.org/10.1016/j.ecoenv.2009.10.008>

Bastarud, A., Rakotondramanga, J. M., Mahazosaotra, J., Ravaonindrina, N., & Jambou, R. (2018). Environmental factors and the microbial quality of urban drinking water in a low-income country: The case of Madagascar. *Water*, 10(10), 1450. <https://doi.org/10.3390/w10101450>

Capps, K. A., Bentsen, C. N., & Ramírez, A. (2016). Poverty, urbanization, and environmental degradation: urban streams in the developing world. *Freshwater Science*, 35(1), 429-435. <https://doi.org/10.1086/684945>

Çavaş, T., & Könen, S. (2007). Detection of cytogenetic and DNA damage in peripheral erythrocytes of goldfish (*Carassius auratus*) exposed to a glyphosate formulation using the micronucleus test and the comet assay. *Mutagenesis*, 22(4), 263-268.

Ceretti, E., Zani, C., Zerbini, I., Guzzella, L., Scaglia, M., Berna, V., Donato, F., Monarca, S., & Feretti, D. (2010). Comparative assessment of genotoxicity of mineral water packed in polyethylene terephthalate (PET) and glass bottles. *Water research*, 44(5), 1462-1470. <https://doi.org/10.1016/j.watres.2009.10.030>

Christofolletti, C. A., David, J. A. O., & Fontanetti, C. S. (2009). Application of the comet assay in erythrocytes of *Oreochromis niloticus* (Pisces): A

This article is protected by copyright. All rights reserved.

methodological comparison. *Genetics and Molecular Biology*, 32, 155-158.

<https://doi.org/10.1590/S1415-47572009005000020>

Creber, H. M. C., Davies, M. S., Francis, D., & Walker, H. D. (1994). Variation in DNA C value in natural populations of *Dactylis glomerata* L. *New Phytologist*, 128(3), 555-561. <https://doi.org/10.1111/j.1469-8137.1994.tb03001.x>

da Silva, G. S., de Matos, L. V., da Silva Freitas, J.O., de Campos, D. F., & de Almeida E Val, V.M.F. (2019). Gene expression, genotoxicity, and physiological responses in an Amazonian fish, *Colossoma macropomum* (CUVIER 1818), exposed to Roundup® and subsequent acute hypoxia. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 222, 49-58. <https://doi.org/10.1016/j.cbpc.2019.04.010>

de Campos Ventura, B., de Angelis, D. D. F., & Marin-Morales, M. A. (2008). Mutagenic and genotoxic effects of the Atrazine herbicide in *Oreochromis niloticus* (Perciformes, Cichlidae) detected by the micronuclei test and the comet assay. *Pesticide Biochemistry and Physiology*, 90(1), 42-51. <https://doi.org/10.1016/j.pestbp.2007.07.009>

de Lapuente, J., Lourenço, J., Mendo, S. A., Borràs, M., Martins, M. G., Costa, P. M., & Pacheco, M. (2015). The Comet Assay and its applications in the field of ecotoxicology: A mature tool that continues to expand its perspectives. *Frontiers in genetics*, 6, 180. <https://doi.org/10.3389/fgene.2015.00180>

de Souza Trigueiro, N. S., Gonçalves, B. B., Dias, F. C., de Oliveira Lima, E. C., Rocha, T. L., & Sabóia-Morais, S. M. T. (2021). Co-exposure of iron oxide nanoparticles and glyphosate-based herbicide induces DNA damage and mutagenic effects in the guppy (*Poecilia reticulata*). *Environmental Toxicology and Pharmacology*, 81, 103521. <https://doi.org/10.1016/j.etap.2020.103521>

- Dos Reis, G. B., Andrade-Vieira, L. F., de Campos Moraes, I., César, P. H. S., Marcussi, S., & Davide, L. C. (2017). Reliability of plant root comet assay in comparison with human leukocyte comet assay for assessment environmental genotoxic agents. *Ecotoxicology and Environmental Safety*, 142, 110-116. <https://doi.org/10.1016/j.ecoenv.2017.04.004>
- Dupuy, S., Defrise, L., Gaetano, R., & Burnod, P. (2017). *Antananarivo - Cartographie de l'occupation du sol 2017*, CIRAD Dataverse, v1. <https://doi:10.18167/DVN1/NHE34C>
- FAO (2022). *FishStatJ: Universal Software for Fishery Statistical Time Series*. Fisheries and aquaculture department. Statistics and Information Service.
- Fiskesjö, G. (1985). The *Allium* test as a standard in environmental monitoring. *Hereditas*, 102(1), 99-112. <https://doi.org/10.1111/j.1601-5223.1985.tb00471.x>
- Frenzilli, G. I. A. D. A., Nigro, M. A. R. C. O., & Lyons, B. P. (2009). The Comet assay for the evaluation of genotoxic impact in aquatic environments. *Mutation research/reviews in mutation research*, 681(1), 80-92. <https://doi.org/10.1016/j.mrrev.2008.03.001>
- Gyori, B. M., Venkatachalam, G., Thiagarajan, P. S., Hsu, D., & Clement, M. V. (2014). OpenComet: An automated tool for comet assay image analysis. *Redox biology*, 2, 457-465. <https://doi.org/10.1016/j.redox.2013.12.020>
- Hemachandra, C. K., & Pathiratne, A. (2017). Cytogenotoxicity screening of source water, wastewater and treated water of drinking water treatment plants using two *in vivo* test systems: *Allium cepa* root based and Nile tilapia erythrocyte based tests. *Water Research*, 108, 320-329. <https://doi.org/10.1016/j.watres.2016.11.009>
- Hemachandra, C. K., & Pathiratne, A. (2016). Combination of physico-chemical analysis, *Allium cepa* test system and *Oreochromis niloticus* erythrocyte based
- This article is protected by copyright. All rights reserved.

comet assay/nuclear abnormalities tests for cyto-genotoxicity assessments of treated effluents discharged from textile industries. *Ecotoxicology and environmental safety*, 131, 54-64.

<https://doi.org/10.1016/j.ecoenv.2016.05.010>.

Iturburu, F. G., Simoniello, M. F., Medici, S., Panzeri, A. M., & Menone, M. L.

(2018). Imidacloprid causes DNA damage in fish: Clastogenesis as a mechanism of genotoxicity. *Bulletin of environmental contamination and toxicology*, 100(6), 760-764. <https://doi.org/10.1007/s00128-018-2338-0>

Jiang, Z., Zhang, H., Qin, R., Zou, J., Wang, J., Shi, Q., Jiang, W., & Liu, D. (2014).

Effects of lead on the morphology and structure of the nucleolus in the root tip meristematic cells of *Allium cepa* L. *International journal of molecular sciences*, 15(8), 13406-13423. <https://doi.org/10.3390/ijms150813406>

Kirschke, S., Avellán, T., Bärlund, I., Bogardi, J. J., Carvalho, L., Chapman, D.,

Dickens, C. W. S., Irvine, K., Lee, S., Mehner, T., & Warner, S. (2020).

Capacity challenges in water quality monitoring: Understanding the role of human development. *Environmental monitoring and assessment*, 192(5), 1-16. <https://doi.org/10.1007/s10661-020-8224-3>

Könen, S., & Çavaş, T. (2008). Genotoxicity testing of the herbicide trifluralin and its commercial formulation Treflan using the piscine micronucleus test.

Environmental and molecular mutagenesis, 49(6), 434-438.

<https://doi.org/10.1002/em.20401>

Kumaravel, T. S., & Jha, A. N. (2006). Reliable Comet assay measurements for

detecting DNA damage induced by ionising radiation and chemicals. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 605(1-2), 7-16.

<https://doi.org/10.1016/j.mrgentox.2006.03.002>

- Kurihara, D., Mizuta, Y., Sato, Y., & Higashiyama, T. (2015). ClearSee: A rapid optical clearing reagent for whole-plant fluorescence imaging. *Development*, 142(23), 4168-4179. <https://doi.org/10.1242/dev.127613>
- Leme, D. M., & Marin-Morales, M. A. (2009). *Allium cepa* test in environmental monitoring: a review on its application. *Mutation research/reviews in mutation research*, 682(1), 71-81. <https://doi.org/10.1016/j.mrrev.2009.06.002>
- Letsyo, E., Adams, Z. S., Dzikunoo, J., & Asante-Donyinah, D. (2021). Uptake and accumulation of pyrrolizidine alkaloids in the tissues of maize (*Zea mays* L.) plants from the soil of a 4-year-old *Chromolaena odorata* dominated fallow farmland. *Chemosphere*, 270, 128669. <https://doi.org/10.1016/j.chemosphere.2020.128669>
- Lima, M. G. F., Rocha, L. C., Silveira, G. L., Alvarenga, I. F. S., & Andrade-Vieria, L. F. (2019). Nucleolar alterations are reliable parameters to determine the cytogenotoxicity of environmental pollutants. *Ecotoxicology and environmental safety*, 174, 630-636. <https://doi.org/10.1016/j.ecoenv.2019.03.033>
- Martins, M., & Costa, P. M. (2017). The comet assay in aquatic (eco) genotoxicology using non-conventional model organisms: Relevance, constraints and prospects. In M. L. Larramendy (Ed.). *Ecotoxicology and genotoxicology: non-traditional aquatic models* (pp. 1-32). Royal Society of Chemistry.
- Matsumoto, S. T., Mantovani, M. S., Malagutti, M. I. A., Dias, A. L., Fonseca, I. C., & Marin-Morales, M. A. (2006). Genotoxicity and mutagenicity of water contaminated with tannery effluents, as evaluated by the micronucleus test and comet assay using the fish *Oreochromis niloticus* and chromosome aberrations in onion root-tips. *Genetics and Molecular Biology*, 29, 148-158. <https://doi.org/10.1590/S1415-47572006000100028>

- McLennan, M. (2021). *The Global Risks Report 2021 16th Edition*. Cologne: World Economic Forum.
- MEAH (2004). *Décret n°2004-635 du 15 juin 2004 Portant modification du décret n°2003-941 du 09 Septembre 2003 relatif à la surveillance de l'eau, au contrôle des eaux destinées à la consommation humaine et aux priorités d'accès à la ressource en eau*. Ministère de l'Eau, de L'Assainissement et de l'Hygiène.
- Nagpure, N. S., Srivastava, R., Kumar, R., Dabas, A., Kushwaha, B., & Kumar, P. (2015). Assessment of pollution of river Ganges by tannery effluents using genotoxicity biomarkers in murrel fish, *Channa punctatus* (Bloch). *Indian Journal of Experimental Biology*, 53(7), 476-483.
- Ohe, T., Watanabe, T., & Wakabayashi, K. (2004). Mutagens in surface waters: A review. *Mutation Research/Reviews in Mutation Research*, 567(2-3), 109-149. <https://doi.org/10.1016/j.mrrev.2004.08.003>
- Orsière, T., Sari-Minodier, I., Decome, L., Botta, C., Iarmarcovai, G., & Botta, A. (2005). De la génotoxicologie à la biosurveillance. *Journées Nationales de Santé au Travail dans le BTP, Annales*, 28, 25-28.
- Osman, A. G., Abuel-Fadl, K. Y., & Kloas, W. (2012). *In situ* evaluation of the genotoxic potential of the river Nile: II. Detection of DNA strand-breakage and apoptosis in *Oreochromis niloticus niloticus* (Linnaeus, 1758) and *Clarias gariepinus* (Burchell, 1822). *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 747(1), 14-21. <https://doi.org/10.1016/j.mrgentox.2012.02.013>
- Pathiratne, A., Hemachandra, C. K., & De Silva, N. (2015). Efficacy of *Allium cepa* test system for screening cytotoxicity and genotoxicity of industrial effluents originated from different industrial activities. *Environmental Monitoring and Assessment*, 187(12), 1-12. <https://doi.org/10.1007/s10661-015-4954-z>
- This article is protected by copyright. All rights reserved.

- Qin, R., Jiao, Y., Zhang, S., Jiang, W., & Liu, D. (2010). Effects of aluminium on nucleoli in root tip cells and selected physiological and biochemical characters in *Allium cepa* var. *agrogarum* L. *BMC plant biology*, 10(1), 1-11. <https://doi.org/10.1186/1471-2229-10-225>
- R Development Core Team (2022). *A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing.
- Radić, S., Gregorović, G., Stipaničev, D., Cvjetko, P., Šrut, M., Vujčić, V., Oreščanin, V., & Klobučar, G. I. V. (2013). Assessment of surface water in the vicinity of fertilizer factory using fish and plants. *Ecotoxicology and environmental safety*, 96, 32-40. <https://doi.org/10.1016/j.ecoenv.2013.06.023>.
- Rasoazanany, E.O. (2008). Highlighting of pollutants elements of atmospheric air and industrial wastewater in Antananarivo and Analysis of a forest plants of the east coast of Madagascar. University of Antananarivo.
- Razanakoto, O. R., Raharimalala, S., Sarobidy, E. J. R. F., Rakotondravelo, J.-C., Autfray, P., & Razafimahatratra, H. M. (2021). Why smallholder farms' practices are already agroecological despite conventional agriculture applied on market-gardening. *Outlook on Agriculture*, 50(1), 80–89. <https://doi.org/10.1177/0030727020972120>
- SanJuan-Reyes, N., Gómez-Oliván, L. M., Galar-Martínez, M., García-Medina, S., Islas-Flores, H., González-González, E. D., Cardoso-Vera, J. D., & Jiménez-Vargas, J. M. (2015). NSAID-manufacturing plant effluent induces geno-and cytotoxicity in common carp (*Cyprinus carpio*). *Science of the Total Environment*, 530, 1-10. <https://doi.org/10.1016/j.scitotenv.2015.05.088>
- Scalon, M., Rechenmacher, C., Siebel, A. M., Kayser, M. L., Rodrigues, M. T., Maluf, S. W., Rodrigues M. A. S., & Basso da Silva, L. (2013). Genotoxic potential and physicochemical parameters of Sinos River, southern Brazil. *Scientific World Journal*, 2013, 2097737. <https://doi.org/10.1155/2013/209737>

- Singh, N. P., McCoy, M. T., Tice, R. R., & Schneider, E. L. (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental cell research*, 175(1), 184-191. [https://doi.org/10.1016/0014-4827\(88\)90265-0](https://doi.org/10.1016/0014-4827(88)90265-0)
- Speit, G., Kojima, H., Burlinson, B., Collins, A. R., Kasper, P., Plappert-Helbig, U., Uno, Y., Vasquez, M., Beevers, C., De Boeck, M., Escobar, P. A., Kitamoto, S., Pant, K., Pfuhler, S., Tanaka, J., & Levy, D. D. (2015). Critical issues with the *in vivo* comet assay: a report of the comet assay working group in the 6th International Workshop on Genotoxicity Testing (IWGT). *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 783, 6-12. <https://doi.org/10.1016/j.mrgentox.2014.09.006>
- Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., & Sutton, D. J. (2012). Heavy metal toxicity and the environment. *Molecular, clinical and environmental toxicology*, 101, 133-164. https://doi.org/10.1007/978-3-7643-8340-4_6
- Vijayan, M. M., Pereira, C., Grau, E. G., & Iwama, G. K. (1997). Metabolic responses associated with confinement stress in tilapia: the role of cortisol. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 116(1), 89-95. [https://doi.org/10.1016/S0742-8413\(96\)00124-7](https://doi.org/10.1016/S0742-8413(96)00124-7)
- Vlastos, D., Antonopoulou, M., & Konstantinou, I. K. (2016). Evaluation of toxicity and genotoxicity of 2-Chlorophenol on bacteria, fish and human cells. *Science of the Total Environment*, 551-552, 649-655. <https://doi.org/10.1016/j.scitotenv.2016.02.043>
- Wang, Q.L., Liu, D. H., & Yue, J. Y. (2016). The uptake of cadmium by *Allium cepa* var. *agrogarum* L. and its effects on chromosome and nucleolar behavior in root tip cells. *Phyton-International Journal of Experimental Botany*, 85, 155-161. <https://doi.org/10.32604/phyton.2016.85.155>
- WHO (2017). *Guidelines for*

drinking-water quality, 4th edition, incorporating the 1st addendum. World Health Organization.

Wu, J. Y., Shen, L., Gao, G., Lin, G. F., Ke, X., & Shen, J. H. (2004). A season-dependent variation of genotoxicity of surface water samples from Taihu Lake, Yangzte delta. *Environmental monitoring and assessment*, 98(1), 225-234.
<https://doi.org/10.1023/B:EMAS.0000038188.16088.f3>

Captions to figures




Figure 1. Location of Masay and Viliahazo marshes with respect to the capital city of Antananarivo and land uses in the vicinity. The map showing Masay marsh was modified from Dupuy et al. (2017).




Figure 2. (A) Percentage of damage to the DNA of the erythrocytes of *Oreochromis niloticus* exposed to tap water (negative control), EMS (positive control) and to water taken from the urban (Masay) and rural (Viliahazo) marshes (n = 5 fish per condition, 400 measurements performed per fish) presented as box plots and violin plots (i.e. probability density of the data at different values) and (B) DNA degradation in *O. niloticus* erythrocytes by epifluorescence microscopy. Scale bar: 20 μm. Letters denote significant differences (p < 0.05) between experimental conditions.




Figure 3. (A) Mitotic index of the root meristem cells of *Allium cepa* bulbs (n = 4 per condition, 3 measurements performed per bulb) exposed to tap water (negative

This article is protected by copyright. All rights reserved.

control), ethyl methanesulfonate (positive control) and water from urban (Masay) and rural (Viliahazo) marshes investigated here. (B) 3D image of the meristem region used to calculate the mitotic index from a Z-stack of 109 optical slices and tile scan function (frame size: 1570x1570x110 μm) in the root apex. Insert: magnified view of a portion of meristem cells showing some mitosis. Scale bar: 100 μm . Values are means $\pm$ SD. Letters denote significant differences ($p < 0.05$) between experimental conditions.




Figure 4. (A) Proportion of cells with ≥ 3 nucleoli per nucleus in meristem cells of *Allium cepa* roots (from $n = 4$ bulbs per condition) exposed to tap water (negative control), ethyl methanesulfonate (positive control) and water from urban (Masay) and rural (Viliahazo) marshes investigated here and (B) the nucleus of root meristem cells in each experimental condition. Four bulbs were used per experimental condition. Scale bar: 10 μm . Values are means $\pm$ SD. Letters denote significant differences ($p < 0.05$) between experimental conditions.

Caption to table

Table 1. Summary of water quality parameters and trace elements measured for the tap water (negative control), and water from urban (Masay) and rural (Viliahazo) marshes. Values are means $\pm$ SD. No data were recorded after EMS addition in tap water in the positive control to avoid the contamination of the instruments. Letters denote significant differences ($p < 0.05$) between experimental conditions.

Parameter	n	Treatment		
		Tap water (negative control)	Masay	Viliahazo
Water parameters				
Temperature (°C)	5	22.2 ± 0.6 ^a	22.0 ± 0.5 ^a	22.0 ± 0.4 ^a
DO (mg L ⁻¹)	4	4.7 ± 1.0 ^a	5.1 ± 0.8 ^a	5.4 ± 0.3 ^a
pH	5	6.1 ± 0.2 ^a	6.9 ± 0.2 ^b	5.9 ± 0.2 ^{ac}
Conductivity (µS cm ⁻¹)	5	182 ± 50 ^a	346 ± 38 ^b	89 ± 33 ^c
BDO5 (mg L ⁻¹)	1	1.25	1.80	1.81
Trace elements				
Cr (µg L ⁻¹)	1	0.13	0.24	0.22
Mn (µg L ⁻¹)	1	12.42	57.22	16.43
Fe (µg L ⁻¹)	1	19.14	44.72	14.35
Co (µg L ⁻¹)	1	0.20	0.38	0.11
Ni (µg L ⁻¹)	1	0.46	0.67	0.40
Cu (µg L ⁻¹)	1	1.48	1.19	0.98
Zn (µg L ⁻¹)	1	69.71	10.58	38.46
As (µg L ⁻¹)	1	0.07	0.21	0.05
Pb (µg L ⁻¹)	1	0.76	0.19	0.63