



Development and characterization of microsatellite markers for the French endemic *Angelica heterocarpa* (Apiaceae) and congeneric sympatric species

Emmanuelle Revardel, Olivier Lepais

► To cite this version:

Emmanuelle Revardel, Olivier Lepais. Development and characterization of microsatellite markers for the French endemic *Angelica heterocarpa* (Apiaceae) and congeneric sympatric species. BMC Research Notes, 2022, 15 (1), pp.15. 10.1186/s13104-021-05903-2 . hal-04028545

HAL Id: hal-04028545

<https://hal.inrae.fr/hal-04028545>

Submitted on 14 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH NOTE

Open Access



Development and characterization of microsatellite markers for the French endemic *Angelica heterocarpa* (Apiaceae) and congeneric sympatric species

Emmanuelle Revardel* and Olivier Lepais

Abstract

Objective: *Angelica heterocarpa* (Apiaceae) is a wild endemic French species with special conservation interest in the European Union. It belongs to *Angelica* complex genus which is widespread throughout the north temperate zone, and is sympatric with other congeneric species. The objective of this work is to develop and characterize microsatellite markers as a new tool for understanding the ecology and evolution of *Angelica* species complex.

Results: We identified simple sequence repeat (SSR) regions in a microsatellite-enriched library from *A. heterocarpa* and *A. sylvestris*. All 16 selected SSR regions were found to amplify in these species and were highly polymorphic. Marker transferability was validated in *A. razulii* and *A. archangelica*. These markers will help us to better understand the evolutionary dynamic between rare endemics and widespread sister species, and be useful for conservation of the endemic species. Moreover, they can provide new tools for studying the numerous traditional medicinal herbs of the *Angelica* genus.

Keywords: Plants, Conservation, Population genetics, Genetic diversity, Nuclear SSR, Hybridization, Estuary riverbanks, *Angelica sylvestris*, *Angelica razulii*, *Angelica archangelica*

Introduction

Angelica L. is a large complex genus comprising approximately 110 species confined in the northern hemisphere, with the majority in Eurasia [1]. In this paper, four French native congeneric *Angelica* species are considered. *Angelica heterocarpa* Lloyd is endemic to southwestern French estuary banks. It is protected at national level in France and is listed as priority species in the Habitats Directive of European Union [2]. *Angelica sylvestris* L. is common in open and forest habitats and is widely distributed among Europe and Asia. Despite different ecological niches, *A. heterocarpa* and *A. sylvestris* can live in

neighboring riverbank locations. The observation of morphological intermediates questions the taxonomic relationship and potential for hybridization between these two species. Two other species are present in southwestern France: *Angelica razulii* Gouan, a Pyrenean endemic that shares hydrographic zones with *A. heterocarpa* and *A. sylvestris*, and *Angelica archangelica* L. that occurs naturally in estuaries from northern France and Europe and is cultivated in southwestern France for aromatic and pharmacological interests.

For the effective conservation of *A. heterocarpa*, genetic markers providing resolution at the population level are essential although, until now, not available. Here, we report the characterization of 16 new polymorphic microsatellite markers for *A. heterocarpa* and *A.*

*Correspondence: emmanuelle.revardel@u-bordeaux.fr
University of Bordeaux, INRAE, BIOGECO, 33615 Pessac, France



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Table 1 Characteristics of 16 primer pairs for microsatellites loci developed in *A. heterocarpa* and *A. sylvestris*

Locus	Primer sequences (5' → 3')		Motif	Range (bp)	Label	Multiplex	GenBank
An23	F:	GAAACAAAATCAAATAGTAATCGCA	AC	75–95	PET	1	MZ065562
	R:	AAATGTAATCTGCACGCGGT					
An35	F:	GGTTGCAACTCAGATGCTGG	GT	175–189	FAM	1	MZ065564
	R:	ACCCTGTGCGTATTGCCTAC					
An69	F:	AGCAAGTGAGGCAAGACCAT	AG	144–176	FAM	2	MZ065568
	R:	CAAACCTCTCCTTACCCCAA					
An71	F:	GAGCATCTCGAAGAGATCAA	AG	216–262	FAM	1,2	MZ065569
	R:	GGGACCACTCAGAATTGGAA					
An72	F:	TTGGATTGAGGAGAGGACCA	GAT	66–114	FAM	2	MZ065570
	R:	CTTCTCCAACCAGGATCAG					
An74	F:	GCATTGTAGCTACTGGAGGG	GT	80–134	VIC	2	MZ065571
	R:	GGCAACAGAGTAGACCTTAACCTG					
An89	F:	GCAACCGCAGCTCATTTTAT	GT	88–106	PET	2	MZ065572
	R:	AAACAAAGGACCAGCCGAC					
An91	F:	CTCGCGTTGACAGGGTTTA	GT	129–143	PET	2	MZ065573
	R:	TCGAGTTCTAGTTGGACAGGG					
An93	F:	GGCAGCTAAGTGAAGCAAAA	AC	170–190	VIC	2	MZ065574
	R:	TGCGCATGTTACTAAGGCTG					
S216	F:	TCTCTGAGTATATTTTGGGTGTG	CT	167–179	PET	1	MZ065559
	R:	TCGCAAAATACCCTCATCTC					
An11	F:	GGGACATTCAACACAACATCA	AG	96–128	VIC	1	MZ065561
	R:	CTCTTTCTTGACCCCTTCCA					
An39	F:	TTGGCTGCACTTACATTGTC	CT	212–284	VIC	1	MZ065565
	R:	ATGATAAACCCGGTTGCTTG					
An68	F:	TCCAAAATGCACAGATCCAG	AG	129–171	NED	2	MZ065567
	R:	CTCGTCGAGTTCTACGCTCC					
An32	F:	TGGGTTTCATCAAGATTCAAGG	AG	118–174	NED	1	MZ065563
	R:	GTGTGGTCACTGCAAGCATC					
An41	F:	GGGAAACTGAATTAACCGAGC	AG	263–324	PET	1	MZ065566
	R:	CCACTTGTGGTCTCTAACATGG					
S37	F:	CAAAAGTGGATACTAGTTGTGTG	CT	192–224	NED	1	MZ065560
	R:	GCTCTACCATTAGCAAAACC					

Label: fluorophore dye labeling at 5' end of forward primers

F, forward primer sequence; R, reverse primer sequences; bp, base pair

sylvestris and test their cross-species transferability in *A. razulii* and *A. archangelica*.

Main text

Methods

Plant material was collected across natural populations in France or Germany: two for *A. heterocarpa* (N=78), three for *A. sylvestris* (N=98) and one for *A. razulii* (N=50) and *A. archangelica* (N=3) (Table 4 in Appendix 1). One or two leaflets were collected from each plant and preserved dried in silica gel. DNA was extracted with the Invitex extraction kit (Invitex, Berlin, Germany). Microsatellites markers were developed from sequences obtained from *A. heterocarpa* and *A. sylvestris*

after enrichment by both traditional cloning and high throughput sequencing (GenoScreen, Lille, France) of microsatellite-enriched library [3]. Sequences containing microsatellites were identified using the QDD software [4] and primers were designed using the Primer 3 software [5] using default parameters with 56 °C as annealing temperature. A total of 119 primers pairs for these SSR loci were tested for amplification and genotyping of 4 of each *A. heterocarpa* and *A. sylvestris* individuals using primer extended with M13 sequence for fluorescent labeling [6]. All of them were found to amplify in the both species and among them, 16 showing polymorphisms and consistent peak profile were selected in the final genotyping protocol (Table 1).

Table 2 Genetic diversity characteristics of 16 microsatellites loci for *A. heterocarpa* and *A. sylvestris* populations

Loci	<i>A. heterocarpa</i> —Rézé (N = 40)				<i>A. heterocarpa</i> —Ile-St-Georges (N = 38)				<i>A. sylvestris</i> —Aurice (N = 38)				<i>A. sylvestris</i> —La Brède (N = 30)				<i>A. sylvestris</i> —Le Bonhomme (N = 30)			
	N _A	H _E	H _O	F _{IS}	N _A	H _E	H _O	F _{IS}	N _A	H _E	H _O	F _{IS}	N _A	H _E	H _O	F _{IS}	N _A	H _E	H _O	F _{IS}
An23	6	0.75	0.56	0.25	10	0.82	0.68	0.17	5	0.75	0.76	−0.02	7	0.70	0.50	0.29	4	0.58	0.13	0.79 ^a
An35	5	0.67	0.58	0.14	4	0.53	0.55	−0.05	5	0.54	0.32	0.42	6	0.65	0.70	−0.08	4	0.56	0.57	−0.01
An69	3	0.33	0.30	0.09	9	0.83	0.79	0.05	15	0.88	0.76	0.13	5	0.46	0.33	0.29	6	0.47	0.40	0.15
An71	9	0.76	0.32	0.58 ^a	18	0.91	0.58	0.36 ^a	14	0.88	0.31	0.65 ^a	13	0.89	0.72	0.19	15	0.88	0.59	0.34 ^a
An72	8	0.80	0.69	0.13	8	0.77	0.76	0.01	11	0.79	0.85	−0.07	8	0.85	0.83	0.02	13	0.86	0.87	−0.01
An74	2	0.43	0.10	0.77	4	0.45	0.23	0.49 ^a	4	0.55	0.18	0.67 ^a	4	0.23	0.18	0.24	2	0.04	0.04	0.00
An89	6	0.74	0.71	0.05	5	0.53	0.27	0.49 ^a	8	0.82	0.33	0.60 ^a	6	0.71	0.44	0.39	6	0.65	0.53	0.18
An91	6	0.59	0.53	0.12	6	0.51	0.45	0.12	6	0.61	0.58	0.05	6	0.62	0.73	−0.18	6	0.74	0.69	0.07
An93	5	0.54	0.29	0.47	7	0.69	0.32	0.54 ^a	7	0.77	0.21	0.74 ^a	7	0.79	0.48	0.40 ^a	10	0.89	0.52	0.42 ^a
S216	4	0.58	0.58	0.01	4	0.55	0.50	0.10	4	0.64	0.61	0.05	5	0.70	0.43	0.39	3	0.50	0.53	−0.07
An11	8	0.77	0.75	0.03	9	0.87	0.81	0.07	12	0.84	0.75	0.11	11	0.81	0.67	0.18	8	0.79	0.72	0.09
An39	12	0.82	0.88	−0.07	21	0.92	0.92	0.00	11	0.76	0.69	0.08	12	0.90	0.76	0.16	7	0.78	0.71	0.09
An68	8	0.77	0.72	0.07	11	0.83	0.79	0.05	15	0.83	0.86	−0.03	8	0.77	0.83	−0.09	8	0.74	0.66	0.11
An32	9	0.68	0.74	−0.09	4	0.29	0.19	0.34	8	0.76	0.50	0.34 ^a	6	0.65	0.69	−0.06	8	0.82	0.60	0.27
An41	12	0.89	0.86	0.04	19	0.91	0.66	0.28 ^a	—	—	—	—	10	0.86	0.88	−0.03	—	—	—	—
S37	8	0.84	0.57	0.33	8	0.77	0.24	0.69 ^a	—	—	—	—	6	0.74	0.48	0.36	—	—	—	—
Mean	6.9	0.69	0.57	0.18	9.2	0.70	0.55	0.23	8.9	0.74	0.55	0.27	7.5	0.71	0.60	0.15	7.1	0.66	0.54	0.17

N, number of samples; N_A, number of alleles (N_A); H_O, observed heterozygosity; H_E, expected heterozygosity; F_{IS}, inbreeding coefficient; −, unavailable data

^a Significant deficit in heterozygotes ($P < 0.01$)

Table 3 Number of alleles and allele range of microsatellite loci in *A. razulii* and *A. archangelica*

Loci	<i>A. razulii</i> (N = 50)		<i>A. archangelica</i> (N = 3)	
	N _A	Range (bp)	N _A	Range (bp)
An23	0	0	1	79
An35	3	177–185	3	169–179
An69	7	140–152	2	142–150
An71	5	212–220	2	216–230 (2)
An72	8	63–96	2	69–72
An74	0	0	1	82 (1)
An89	5	92–102 (20)	1	92 (1)
An91	9	125–143	2	133–135 (1)
An93	0	0	2	170–178
S216	0	0	2	169–179
An11	0	0	2	132–138
An39	2	220–224	3	188–232
An68	0	0	—	—
An32	12	124–156	—	—
An41	2	267–271 (12)	—	—
S37	3	151–171 (3)	—	—

Values in () indicates the number of individuals with at least one PCR amplified signal

N, number of samples; N_A, number of alleles; bp, base pairs; —, unavailable data

Multiplexed PCR were achieved using Qiagen Microsatellite Type-It master mix (Qiagen, Hilden, Germany) and cycling conditions were: 15 min at 95 °C; 30 cycles of 30 s at 94 °C, 60 s at 56 °C, 45 s at 72 °C; and 60 °C for 30 min. The fluorescent-labeled PCR products were run on an ABI 3730 DNA Analyzer (Applied Biosystems, Waltham, MA, USA) at the Genome Transcriptome Facility of Bordeaux. Genotype calling was carried out manually using GeneMapper Software (Applied Biosystems, Waltham, MA, USA).

Results and discussion

Overall, the developed microsatellite loci were highly polymorphic with a number of alleles per locus ranged from 2 to 21 with a mean of 8 alleles per locus in the five studied *A. heterocarpa* and *A. sylvestris* populations (Table 2). At the population level, the observed heterozygosity ranged from 0.54 to 0.60, and the expected heterozygosity ranged from 0.66 to 0.74. Four out of the 16 loci showed significant deviations from Hardy–Weinberg equilibrium (HWE) in more than one population after correcting for multiple testing ($P < 0.001$; Table 2).

The cross-species transferability and polymorphism of the developed SSR markers were further tested in the two other congeneric species (Table 3). In *A. razulii*, only

7 out of the 16 loci amplified consistently and were polymorphic. On the 12 markers tested in *A. archangelica*, 7 amplified successfully and were polymorphic. Note that additional markers amplified inconsistently, suggesting presences of null alleles or limited transferability (3 markers for *A. razulii* and 4 for *A. archangelica*, Table 3).

These new polymorphic markers will allow investigating population genetic structure, reproductive system, and potential hybridization within the *Angelica* species complex. Moreover, as numerous *Angelica* species are traditional medicine herbs all over the Eurasia continent, including *A. sylvestris* and *A. archangelica* [7], these microsatellite markers complement the molecular markers previously developed for other congeneric species of particular pharmacological interest [8–10].

Limitations

- Deviation from HWE: Loci showing marked deficits in heterozygotes in most populations are likely affected by null alleles. However, this should be further investigate with larger sample size because the restricted sampled size of the present study hampered additional statistical test. In addition, more biological insight to support the fact that the species is expect to follow HWE is needed because population substructure or hybridization are likely affecting *Angelica* population and would resulted in departure from HWE.
- Sample size limitation for *A. archangelica*: only 3 individuals have been included to test for transferability and polymorphism for this species. Amplification success and polymorphism should be assessed in a larger sample size to confirm monomorphic loci.

Appendix 1

See Table 4

Table 4 Taxonomic, geographic information (municipality, department number, country) and GPS central coordinates of *Angelica* sp. populations sampling (n = number of individuals) represented in this study

Species	Location, country	n	Geographic coordinates	Population name
<i>A. heterocarpa</i>	Isle-Saint-Georges (33). FR	38	N44°43'30"-W0°27'28"	Isle St George
<i>A. heterocarpa</i>	Rézé (44). FR	40	N47°11'28"-W1°36'6"	Rézé
<i>A. sylvestris</i>	La Brède (33). FR	30	N44°40'21"-W0°33'17"	La Brède
<i>A. sylvestris</i>	Aurice (40). FR	38	N43°48'38"-W0°35'36"	Aurice
<i>A. sylvestris</i>	Le Bonhomme (68). FR	30	N48°8'10"-E7°5'19"	Le Bonhomme
<i>A. razulii</i>	Bagnères de Bigorre (64). FR	50	N42°59'48"-E0°7'29"	Bagnères
<i>A. archangelica</i>	Hambourg. DE	3	N53°34'18"-E10°00'09"	Hambourg

FR, France; DE, Germany

Abbreviations

DNA: Deoxyribonucleic acid; HWE: Hardy–Weinberg equilibrium; PCR: Polymerase chain reaction; SSR: Simple sequence repeat.

Acknowledgements

The authors thank Henri Caron, Patrick Léger and Guillaume Lalanne-Tisné for their technical assistance in developing SSR markers and genotyping, and Conservatoire Botanique National Sud Atlantique, particularly Sandrine Lorient, Conservatoire Botanique National de Brest and Jean-Pierre Reduron for sampling natural populations. ABI 3730 capillary electrophoreses were performed at the Genome Transcriptome Facility of Bordeaux (Grants from Investissements d'avenir, Convention attributive d'aide EquipEx Xyloforest ANR-10-EQPX-16-01).

Authors' contributions

ER conceived and designed the project, conducted sampling, experiments and analyses. OL conducted analyses. Both authors contributed to writing. Both authors read and approved the final manuscript.

Funding

Funding were from support of Conservatoire Botanique National Sud Atlantique and University of Bordeaux.

Availability of data and materials

The genotypic dataset supporting the conclusions of this article available in the Data INRAE repository under DOI, <https://doi.org/10.15454/LFMP64>. Sequences are accessible in Genbank under accession number MZ065559-MZ065574.

Declarations

Ethics approval and consent to participate

Plant leaflets of *A. heterocarpa* were collected by the Conservatoire Botanique National Sud-Atlantique and the Conservatoire Botanique National de Brest, which both have national agreement from the French Ministry of the environment with written authorization from the regional legal representatives of the Ministry.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 15 June 2021 Accepted: 24 December 2021

Published online: 10 January 2022

References

1. Liao C, Gao Q, Katz-Downie DS, Downie SR. A systematic study of North American *Angelica* species (Apiaceae) based on nrDNA ITS and cpDNA

- sequences and fruit morphology. *J Syst Evol.* 2021. <https://doi.org/10.1111/jse.12702>.
2. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Eur-Lex.* 1992. <http://data.europa.eu/eli/dir/1992/43/2013-07-01>. Accessed 3 Jan 2022.
 3. Malausa T, Gilles A, Megléc E, Blanquart H, Duthoy S, Costedoat C, et al. High-throughput microsatellite isolation through 454 GS-FLX titanium pyrosequencing of enriched DNA libraries. *Mol Ecol Resour.* 2011;11:638–44. <https://doi.org/10.1111/j.1755-0998.2011.02992.x>.
 4. Megléc E, Costedoat C, Dubut V, Gilles A, Malausa T, Pech N, et al. QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics.* 2010;26:403–4. <https://doi.org/10.1093/bioinformatics/btp670>.
 5. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. In: Misener S, Krawetz SA, editors. *Bioinformatics methods and protocols.* New Jersey: Humana Press; 2000. p. 365–86. <https://doi.org/10.1385/1-59259-192-2:365>.
 6. Schuelke M. An economic method for the fluorescent labeling of PCR fragments. *Nat Biotechnol.* 2000;18:233–4.
 7. Sarker SD, Nahar L. Natural medicine: the genus *Angelica*. *Curr Med Chem.* 2004;11:1479–500. <https://doi.org/10.2174/0929867043365189>.
 8. Hao H-C, Chang J-Y, Chung F-Y. Internal transcribed spacer sequence analysis of *Angelica* from different habitats. *Genom Med Biomark Heal Sci.* 2012;4:43–4. <https://doi.org/10.1016/J.GMBHS.2012.04.017>.
 9. Lu Y, Cheng T, Zhu T, Jiang D, Zhou S, Jin L, et al. Isolation and characterization of 18 polymorphic microsatellite markers for the “Female Ginseng” *Angelica sinensis* (Apiaceae) and cross-species amplification. *Biochem Syst Ecol.* 2015;61:488–92. <https://doi.org/10.1016/J.BSE.2015.07.013>.
 10. Gil J, Um Y, Kim S, Kim O, Koo S, Reddy C, et al. Development of genome-wide SSR markers from *Angelica gigas nakai* using next generation sequencing. *Genes.* 2017;8:238. <https://doi.org/10.3390/genes8100238>.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

