



HAL
open science

An update of evidence for pathogen transmission by ticks of the genus *Hyalomma*

Sarah Bonnet, Stéphane Bertagnoli, Alessandra Falchi, Julie Figoni, Johanna Fite, Thierry Hoch, Elsa Quillery, Sara Moutailler, Alice Raffetin, Magalie René-Martellet, et al.

► To cite this version:

Sarah Bonnet, Stéphane Bertagnoli, Alessandra Falchi, Julie Figoni, Johanna Fite, et al.. An update of evidence for pathogen transmission by ticks of the genus *Hyalomma*. *Pathogens*, 2023, 12 (4), pp.513. 10.3390/pathogens12040513. hal-04040763

HAL Id: hal-04040763

<https://hal.inrae.fr/hal-04040763>

Submitted on 27 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Commentary

An Update of Evidence for Pathogen Transmission by Ticks of the Genus *Hyalomma*

Sarah I. Bonnet ^{1,*}, Stéphane Bertagnoli ², Alessandra Falchi ³, Julie Figoni ⁴, Johanna Fite ⁵, Thierry Hoch ⁶, Elsa Quillery ⁵, Sara Moutailler ⁷, Alice Raffetin ⁸, Magalie René-Martellet ^{9,10}, Gwenaël Vourc'h ^{9,10} and Laurence Vial ¹¹

- ¹ Ecology and Emergence of Arthropod-borne Pathogens unit, Institut Pasteur, Université Paris-cité, Centre National de Recherche Scientifique (CNRS) UMR 2000, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE) USC 1510, 75015 Paris, France
 - ² Interactions Hôtes-Agents Pathogènes unit, Université de Toulouse, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), Ecole Nationale Vétérinaire de Toulouse (ENVT), F-31076 Toulouse, France
 - ³ Bioscope Corse Méditerranée unit UR7310, Faculté de Sciences, Campus Grimaldi, Université de Corse, 20250 Corte, France
 - ⁴ Santé publique France, 94410 Saint-Maurice, France
 - ⁵ Risk Assessment Department, French Agency for Food, Environmental and Occupational Health & Safety (ANSES), 94700 Maisons-Alfort, France
 - ⁶ Biologie Épidémiologie et Analyse de Risque en santé animale unit, Oniris, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), 44300 Nantes, France
 - ⁷ Biologie Moléculaire et immunologie Parasitaire unit, Laboratoire de Santé Animale, Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES), Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), Ecole Nationale Vétérinaire d'Alfort (ENVA), 94700 Maisons-Alfort, France
 - ⁸ Reference Centre for Tick-Borne Diseases, Paris and Northern Region, Department of Infectious Diseases, General Hospital of Villeneuve-Saint-Georges, 94190 Villeneuve-Saint-Georges, France
 - ⁹ Epidémiologie des maladies animales et zoonotiques unit, Université de Lyon, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), VetAgro Sup, 69280 Marcy l'Etoile, France
 - ¹⁰ Epidémiologie des maladies animales et zoonotiques unit, Université Clermont Auvergne, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), VetAgro Sup, 63122 Saint-Genès-Champagnelle, France
 - ¹¹ Animal Santé Territoires Risques Ecosystèmes unit, Centre de coopération internationale en recherche agronomique pour le développement (CIRAD), Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), Université Montpellier II, F-34398 Montpellier, France
- * Correspondence: sarah.bonnet@inrae.fr



Citation: Bonnet, S.I.; Bertagnoli, S.; Falchi, A.; Figoni, J.; Fite, J.; Hoch, T.; Quillery, E.; Moutailler, S.; Raffetin, A.; René-Martellet, M.; et al. An Update of Evidence for Pathogen Transmission by Ticks of the Genus *Hyalomma*.

Pathogens **2023**, *12*, 513. <https://doi.org/10.3390/pathogens12040513>

Academic Editor: Ricardo Pérez-Sánchez

Received: 24 February 2023

Revised: 15 March 2023

Accepted: 22 March 2023

Published: 25 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Current and likely future changes in the geographic distribution of ticks belonging to the genus *Hyalomma* are of concern, as these ticks are believed to be vectors of many pathogens responsible for human and animal diseases. However, we have observed that for many pathogens there are no vector competence experiments, and that the level of evidence provided by the scientific literature is often not sufficient to validate the transmission of a specific pathogen by a specific *Hyalomma* species. We therefore carried out a bibliographical study to collate the validation evidence for the transmission of parasitic, viral, or bacterial pathogens by *Hyalomma* spp. ticks. Our results show that there are very few validated cases of pathogen transmission by *Hyalomma* tick species.

Keywords: ticks; *Hyalomma* sp.; tick-borne pathogens; vectorial competence

In addition to their direct impact as ectoparasites, ticks are of greater importance as vectors of pathogens to animals and are the second most important group of pathogen vectors affecting humans after mosquitoes, mainly due to their transmission of *Borrelia burgdorferi* sensu lato [1]. In addition, these arthropods are capable of transmitting the largest variety of pathogens including bacteria, parasites (protozoa, helminths), and viruses.

However, like all biological vectors, ticks are not simple “syringes”, as is often believed. Each species, or even each tick population, has a vector competence corresponding to its intrinsic ability to acquire the pathogen by feeding on an infected host, allowing the multiplication/development of this agent and its retransmission to a new host during a new blood meal [2]. To this vector competence is added the set of factors that may influence transmission, thus defining the vector capacity—that is, the ability of a vector to transmit a pathogen at a given time and in a defined region, according to extrinsic conditions such as humidity, temperature, but also vector abundance, trophic preferences, etc. [3].

Ticks acquire pathogens during a blood meal taken on infected vertebrate hosts. Several routes of pathogen transmission to vertebrate hosts are then possible on the occasion of a new blood meal via a deposit on the skin of the host and penetration via the bite wound (via feces, by crushing or via coxal fluid excretion), or via the injection of saliva that accompanies the blood meal and represents the predominant route of pathogen transmission for ticks [4]. In addition, within the tick population, a pathogen may persist from one life stage to the next via transstadial transmission (essential for tick-borne transmission by hard ticks that take only one blood meal per life stage), from the female to its offspring via transovarial transmission, from male to female ticks during copulation via sexual transmission, or from an infected tick to a non-infected one via co-feeding when ticks feed adjacent to each other on the same host [2,5].

The unique detection of pathogenic DNA in a tick collected in the environment or on a vertebrate host does not prove vector competence, but only indicates the fact that the tick took a blood meal on an infected animal. Although such DNA detection in unfed hard ticks is more indicative than detection in engorged ones—as, considering that ixodid ticks feed only once per life stage, it suggests transstadial persistence—it should be noted that this does not imply the viability of the concerned pathogen. In fact, the persistence of DNA during the molting process is possible, as detection has been reported of some vertebrate DNA dating back to a blood meal taken by the previous life stage [6]. Furthermore, the DNA of pathogens that are not transmitted by the tick species concerned is often detected simply as a result of feeding on hosts that harbor such pathogens [7]. The detection of mRNA, a priori reflecting the presence of a living organism, presents an additional but still insufficient level of evidence of vectorial transmission. Additionally, the detection of a given pathogen in both ticks and samples from tick-infested vertebrate hosts collected in the same area, the co-occurrence with already known tick-borne pathogens, or a documented infection following a tick bite, can all represent indirect significant evidence of a given pathogen transmission. The demonstration of transstadial and/or transovarial persistence, which validates the existence of a development of the pathogen in ticks, is a strong indication in favor of biological vector transmission. However, conclusive evidence of vector competence for a given pathogen can only be provided by the demonstration of the ability of a tick species to acquire a pathogen on an infected host, to allow its development, and to transmit it to a new host. Unfortunately, very few vector competence experiments have been conducted due to the difficulties encountered in carrying out complete transmission cycles under experimental conditions. Indeed, it requires having pathogen-free tick colonies, vertebrate hosts suitable for both tick engorgement and pathogen replication (or to have effective artificial tick feeding methods combined with optimal cultivation methods of the pathogen), and can require high biosecurity levels depending on the pathogen concerned [8,9].

Ticks from the *Hyalomma* genus are considered to be expanding from some parts of their range, as reported for the invasion of *H. marginatum* into Europe since the late 20th century [10,11]. This is of concern, as these ticks are vectors of many pathogens responsible for human and animal diseases [12] and because there are few measures to control them, in particular during their off-host development [13]. Like other hard ticks, *Hyalomma* species take one blood meal per life stage before molting (larva and nymph) or, after fertilization, laying eggs (female). The majority of *Hyalomma* spp. ticks have a three-host cycle (each of the three life stages must find a new host on which to take a blood meal). However, some are diphasic, such as those of the *marginatum* group (larvae and nymphs taking their blood meals on the

same host), and one species, *Hyalomma scupense*, is monophasic (all three life stages remain on the same host). *Hyalomma* ticks feed on domestic or wild vertebrate hosts, with some species such as *H. marginatum* or *H. rufipes* utilizing a large variety of hosts, which favors pathogen spillover. Humans, by entering the ecosystem of these hosts, can become accidental hosts of ticks, and thus, become exposed to pathogens [14]. The genus *Hyalomma* includes the most xerophilous species among all ticks and may be favored under future climate change [15].

Numerous pathogens—parasitic, viral, or bacterial—have been reported in the scientific literature as transmitted or potentially transmitted by ticks of the genus *Hyalomma*. The synthesis of these studies, with an attributed level of evidence of vectorial transmission, is reported in Table 1. The level of evidence ranges from a simple detection of pathogen DNA or RNA in ticks collected from vertebrate hosts to a formal demonstration of experimental transmission from an infected vertebrate host to a new naïve one, coupled with appropriate epidemiological data. To build this table, we considered the 27 *Hyalomma* species described by Guglielmone et al. in 2010 [16]. Species for which no evidence of a potential vector role has been reported to date have not been included, namely *Hyalomma albiparvum*, *Hyalomma arabica*, *Hyalomma brevipunctatum*, *Hyalomma glabrum*, *Hyalomma hystricis*, *Hyalomma nitidum*, *Hyalomma punctum*, *Hyalomma rhipicephaloides*, *Hyalomma franchinii*, and *Hyalomma kumari*. Our literature review includes the names of *Hyalomma* species that have been used for several past decades but have since been abandoned in favor of the currently used names, namely *Hyalomma plumbeum* (now *H. marginatum*) and *Hyalomma detritum* (now *H. scupense*). Note that the data identified for *Hyalomma savignyi*, now reclassified as *H. lusitanicum*, are not considered here, since *H. savignyi* is now considered to include several subspecies. *Hyalomma savignyi* data could therefore also apply to *H. lusitanicum*, as well as to *H. anatolicum*, *H. impeltatum*, *H. impressum*, *H. marginatum*, or *H. truncatum*. For bibliographic research, a narrative review was performed using the terms “*Hyalomma*” and “[pathogen sought]” (all microorganisms whose transmission by ticks has been reported in the scientific literature with de facto exclusion of symbionts) with the Boolean operator “AND” in the PubMed and Scopus databases without date restriction. The literature search was conducted in English. We retained peer-reviewed research articles and reviews (not including conference proceedings) and book sections. Screening was conducted first on titles, then on abstracts, and finally on the full text when available. After reading the entire articles, the ones that were eliminated corresponded to those that did not have available data or no original data. The number of references found in each of the two databases is shown in Table 1. All references concerning experimental validation and the epidemiological arguments for transmission are mentioned in the table, but the list concerning DNA/RNA detection is not exhaustive.

In conclusion, we observed that there are many missing pathogen vector competence experiments, and that the level of evidence provided by the scientific literature is often not sufficient to validate the existence of vectorial transmission. We conclude that the pathogen/tick associations for which transmission from an infected host to an initially healthy host via tick bite has been experimentally validated are the following:

- Crimean–Congo Hemorrhagic Fever Virus (CCHFv) by *H. dromedarii*, *H. impeltatum*, *H. marginatum*, *H. rufipes*, and *H. truncatum*.
- African Horse Sickness virus by *H. dromedarii*.
- Venezuelan equine encephalitis virus by *H. truncatum*.
- *Theileria annulata* by *H. anatolicum*, *H. dromedarii*, *H. excavatum*, *H. lusitanicum*, and *H. scupense*.
- *Theileria equi* by *H. anatolicum* and *H. excavatum*.
- *Theileria lestoquardi* by *H. anatolicum*.
- *Theileria ovis* by *H. anatolicum*.
- *Babesia occultans* by *H. rufipes*.
- *Coxiella burnetii* by *H. aegyptium*.
- *Anaplasma marginale* by *H. excavatum*.
- *Rickettsia aeschlimannii* by *H. marginatum* and *H. rufipes*.

Table 1. Bibliographic review of evidence in favor of the transmission of pathogens by ticks of the *Hyalomma* genus.

Tick Species	Number of References in PubMed/Scopus	Pathogen Transmitted/ Suspected to Be Transmitted	Detection of DNA/RNA/ Antigen/Pathogen in Ticks	Epidemiological Arguments of Possible Transmission *	Experimental Validation of Transmission **	References
<i>H. aegyptium</i>	96/110	CCHFv	RNA	yes	0	[17]
		<i>Coxiella burnetii</i>	DNA	no	2, 4	[18,19]
		<i>Borrelia turcica</i>	DNA and RNA	no	2	[20–25]
		<i>Borrelia</i> spp.				
		<i>Bartonella bovis</i>	DNA	no	0	[26]
		<i>Ehrlichia canis</i>	DNA	yes	0	[18,25–27]
		<i>Ehrlichia</i> spp.				
		<i>Anaplasma phagocytophilum</i>	DNA	yes	0	[18]
		<i>Rickettsia africae</i>	DNA	no	0	[28]
		<i>Rickettsia aeschlimannii</i>	DNA	yes	0	[26,29,30]
		<i>Rickettsia sibirica mongolitimonae</i>	DNA	no	0	[30,31]
		<i>Rickettsia slovaca</i>	DNA	no	0	[30]
		Meram virus	RNA	no	0	[32]
Tamdy virus	RNA	no	0	[32]		
<i>H. anatolicum</i>	381/546	<i>Babesia caballi</i>	DNA	no	0	[33]
		<i>Theileria equi</i>	DNA, pathogen	yes	2, 4	[33–37]
		<i>Babesia occultans</i>	DNA	no	0	[38–40]
		<i>Babesia bovis</i>	DNA	no	0	[38–41]
		<i>Theileria annulata</i>	DNA, pathogen	yes	2, 4	[35,39,41–51]
		<i>Theileria lestoquardi</i>	DNA	yes	2, 4	[35,44,49,52–56]
		<i>Theileria ovis</i>	DNA	yes	2, 4	[33,35,38,40,57,58]
		<i>Babesia ovis</i>	DNA	no	0	[55]
		CCHFv	RNA, antigen, viral particle	uncertain	1	[59–65]
		Alphavirus	RNA	no	0	[66]
		Zahedan Rhabdovirus	RNA	no	0	[67]
		Tick Borne Encephalitis virus	no	no	5	[68]
		Kadam virus	RNA	no	0	[66]
		Karshi virus	no	no	0, 1	[69]
		Karyana virus	RNA, virus isolation	yes	0	[70]
		Kundal virus	RNA, virus isolation	yes	0	[70]
		Sindbis virus	RNA	no	0	[71]
		<i>Coxiella burnetii</i>	DNA	no	0	[72–74]
		<i>Bartonella</i> spp.	DNA	no	0	[38]
		<i>Borrelia</i> spp.	DNA	no	0	[38]
		<i>Anaplasma marginale</i> , <i>Anaplasma phagocytophilum</i> , <i>Anaplasma ovis</i> , <i>Anaplasma centrale</i> , <i>Ehrlichia</i> spp., <i>Rickettsia massiliae</i> , <i>Rickettsia</i> spp.,	DNA	no	0	[38,41,75]

Table 1. Cont.

Tick Species	Number of References in PubMed/Scopus	Pathogen Transmitted/ Suspected to Be Transmitted	Detection of DNA/RNA/ Antigen/Pathogen in Ticks	Epidemiological Arguments of Possible Transmission *	Experimental Validation of Transmission **	References
<i>H. asiaticum</i>	145/192	<i>Theileria annulata</i>	DNA	no	0	[76–78]
		<i>Babesia occultans</i>	DNA	no	0	[79]
		<i>Babesia caballi</i>	DNA	no	0	[80,81]
		<i>Theileria equi</i>	DNA	no	0	[80]
		CCHFv	RNA, viral particles	yes	1	[65,82–85]
		Chim virus	RNA	no	0	[86]
		Syr-Darya valley fever virus	RNA	no	0	[86]
		Karshi virus	RNA	no	0, 1	[69,87,88]
		Tamdy virus	Virus isolation	yes	0	[89–92]
		<i>Coxiella burnetii</i>	DNA	no	2	[74,93–95]
		<i>Rickettsia sibirica</i>	DNA	no	0	[96]
		<i>Borrelia burgdorferi</i> s.l.	RNA	no	0	[97]
		<i>Rickettsia sibirica mongolitimonae</i>	isolation	yes	0	[98]
<i>H. dromedarii</i>	232/344	<i>Theileria equi</i>	DNA, pathogen in ticks	no	1, 2, 4	[99–102]
		<i>Theileria camelensis</i>	Pathogen in ticks	yes	1, 2, 4	[103–105]
		<i>Theileria annulata</i>	DNA	yes	2, 4	[48,106–112]
		<i>Theileria ovis</i>	DNA	no	0	[40]
		<i>Babesia caballi</i>	DNA	no	0	[101]
		<i>Babesia occultans</i>	DNA	no	0	[101]
		CCHFv	RNA, antigen, viral particles	yes	1, 2, 4	[64,113,114]
		Alphavirus	RNA	no	0	[66]
		Chick Ross virus	RNA	no	0	[66]
		Dera Ghazi Khan virus	RNA	no	0	[115]
		Dhori virus	RNA	no	0	[116]
		Kadam virus	RNA	no	0	[66,117]
		African horse sickness virus	no	no	2, 4	[118]
		Quaranfil virus	RNA	no	0	[119]
		Sindbis virus	RNA	no	0	[66]
		<i>Coxiella burnetii</i>	DNA	no	0	[101,120–124]
		<i>Francisella persica</i>	DNA	no	0	[125]
		<i>Rickettsia aeschlimannii</i>	DNA	no	0	[121,126,127]
		<i>Rickettsia africae</i>	DNA	no	0	[128]
		<i>Anaplasma</i> spp./ <i>Ehrlichia</i> spp.	DNA	no	0	[101]
<i>Bartonella bovis</i> et <i>Bartonella rochalimae</i>	DNA	no	0	[129]		

Table 1. Cont.

Tick Species	Number of References in PubMed/Scopus	Pathogen Transmitted/ Suspected to Be Transmitted	Detection of DNA/RNA/ Antigen/Pathogen in Ticks	Epidemiological Arguments of Possible Transmission *	Experimental Validation of Transmission **	References		
<i>H. excavatum</i>	149/211	<i>Theileria equi</i>	DNA, pathogen	no	2, 4	[34,130,131]		
		<i>Babesia bigemina</i>	DNA	no	0	[41]		
		<i>Babesia bovis</i>	DNA	no	0	[132]		
		<i>Babesia occultans</i>	DNA	no	3	[30,132]		
		<i>Theileria annulata</i>	DNA	uncertain	2, 4	[31,41,51,76,78,132–134]		
		<i>Theileria capreoli</i>	DNA	no	0	[31]		
		<i>Theileria ovis</i>	DNA	no	0	[40,135]		
		<i>Borrelia</i> spp.	DNA	no	0	[136]		
		<i>Coxiella burnetii</i>	DNA	no	0	[121,124,137–139]		
		<i>Rickettsia africae</i>	DNA	no	0	[140]		
		<i>Rickettsia aeschlimannii</i>	DNA	no	0	[140]		
		<i>Anaplasma marginale</i>	DNA	yes	4	[141]		
		<i>Anaplasma centrale</i>	DNA	yes	0	[141]		
		<i>Ehrlichia ruminantium</i>	DNA	no	0	[41]		
		<i>Rickettsia sibirica mongolotimonae</i>	DNA	no	0	[142]		
		CCHFv	RNA, antigen	uncertain	0	[59,61,143]		
<i>H. hussaini</i>	4/7	<i>Coxiella burnetii</i>	DNA	no	0	[144]		
		<i>Rickettsia massiliae</i> , <i>Rickettsia</i> spp.	DNA	no	0	[38]		
<i>H. impeltatum</i>	62/88	<i>Theileria annulata</i>	DNA	no	2, 4	[41,108,112,145]		
		<i>Theileria lestoquardi</i> (<i>Theileria hirci</i>)	no	uncertain	0	[146]		
		<i>Theileria ovis</i>	DNA	no	0	[35]		
		<i>Babesia occultans</i>	DNA	no	0	[101]		
		<i>Babesia bigemina</i>	DNA	no	0	[41]		
		<i>Babesia bovis</i>	DNA	no	0	[41]		
		<i>Babesia pecorum</i>	DNA	no	0	[35]		
				CCHFv	RNA, antigen virus isolation	yes	1, 2, 4 cofeeding	[61,113,147–149]
		<i>Coxiella burnetii</i>	DNA	no	0	[123,124]		
		Alphavirus	RNA	no	0	[66]		
		Dhori virus	RNA	no	0	[66]		
		Sindbis virus	RNA	no	0	[66]		
		<i>Rickettsia africae</i>	DNA	no	0	[140]		
<i>Rickettsia aeschlimannii</i>	DNA	no	0	[121,150,151]				
<i>H. impressum</i>	10/17		CCHFv	antigen	uncertain	0	[64]	
		<i>Theileria annulata</i>	DNA	no	0	[108]		
		<i>Anaplasma/Ehrlichia</i> spp.	DNA	no	0	[101]		
		<i>Rickettsia africae</i>	DNA	no	0	[152]		
<i>H. isaaci</i>	5/5	Kyasanur forest virus	RNA	-	2, 4	[153]		
<i>H. lusitanicum</i>	68/83	<i>Theileria equi</i>	pathogen	no	1, 2, 4	[99,100]		
		<i>Babesia pecorum</i>	No	yes	0	[154]		
		<i>Theileria annulata</i>	No	yes	4	[107,155,156]		
		CCHFv	RNA, antigen	yes	0	[157,158]		
		<i>Anaplasma phagocytophilum</i>	DNA	no	0	[159]		
		<i>Borrelia burgdorferi</i>	DNA	no	0	[160]		
		<i>Borrelia lusitaniae</i>	DNA	no	0	[161]		
<i>Coxiella burnetii</i>	DNA	no	0	[162–165]				

Table 1. Cont.

Tick Species	Number of References in PubMed/Scopus	Pathogen Transmitted/ Suspected to Be Transmitted	Detection of DNA/RNA/ Antigen/Pathogen in Ticks	Epidemiological Arguments of Possible Transmission *	Experimental Validation of Transmission **	References
<i>H. marginatum</i>	451/620	<i>Theileria equi</i>	DNA	yes	0	[130,131,166,167]
		<i>Theileria annulata</i>	DNA	yes	2	[41,51,111]
		<i>Theileria sergenti/orientalis/buffeli</i>	DNA	no	0	[159,166,168]
		<i>Theileria ovis</i>	DNA	no	0	[40,49,169]
		<i>Theileria lestoquardi</i>	DNA	no	0	[170]
		<i>Babesia ovis</i>	DNA, pathogen in ticks	no	1	[55,171]
		<i>Babesia caballi</i>	DNA	yes	0	[130,131,169,172]
		<i>Babesia bigemina</i>	DNA	no	0	[41,167]
		<i>Babesia bovis</i>	DNA	no	0	[41,167]
		<i>Babesia occultans</i>	DNA	yes	2, 3	[30,31,130,134,136, 166,173–176]
		<i>Babesia microti</i>	DNA	no	0	[174]
		<i>Babesia sp. Tavsan1</i>	DNA	no	0	[31]
		CCHFv	RNA, antigen	yes	1, 2, 3, 4	[64,65,143,158,177–188]
		Flavivirus	RNA	no	0	[189]
		Phlebovirus	RNA	no	0	[190]
		Bahig virus	RNA	no	0	[191]
		Batken virus (close to Dhori virus)	RNA	no	0	[192]
		Bhanja virus	RNA	no	0	[193]
		Dhori virus	RNA	no	0	[194,195]
		Tick Borne Encephalitis virus	RNA	no	0	[196]
		Jingmen virus	RNA, virus isolation	yes	0	[197]
		Matruh virus	RNA	no	0	[198]
		Tamdy virus	RNA	no	0	[89]
		Wanowrie virus	RNA	yes	0	[153,199]
		West Nile virus	RNA	no	2, 3	[189,200–202]
		<i>Rickettsia aeschlimannii</i>	DNA	yes	3	[31,136,151,203–208]
		<i>Rickettsia sibirica mongolitimonae</i>	DNA	no	0	[31]
		<i>Anaplasma marginale</i>	DNA	no	0	[31,209]
		<i>Rickettsia africae</i>	DNA	no	0	[210]
		<i>Anaplasma phagocytophilum</i>	DNA	no	0	[204,211]
		<i>Anaplasma platys</i>	DNA	no	0	[211]
		<i>Coxiella burnetii</i>	DNA	no	0	[139,142,211–213]
		<i>Francisella tularensis</i>	DNA	no	0	[214]
		<i>Ehrlichia monacensis (minasensis)</i>	DNA	no	0	[204]
		<i>Ehrlichia ruminantium</i>	DNA	no	0	[41]
		<i>Bartonella</i> spp.	DNA	no	0	[205,213]
<i>Borrelia burgdoferi</i> s.l.	DNA	no	0	[212,213]		
<i>Borrelia</i> spp.	DNA	no	0	[136,152]		

Table 1. Cont.

Tick Species	Number of References in PubMed/Scopus	Pathogen Transmitted/ Suspected to Be Transmitted	Detection of DNA/RNA/ Antigen/Pathogen in Ticks	Epidemiological Arguments of Possible Transmission *	Experimental Validation of Transmission **	References
<i>H. rufipes</i>	189/238	<i>Babesia occultans</i>	DNA, pathogen	no	2, 3, 4	[175,176]
		<i>Theileria ovis</i>	DNA	no	0	[40]
		<i>Theileria annulata</i>	DNA	no	2, 4	[108,109,215]
		CCHFv	RNA, antigen, viral particles	yes	1, 2, 3, 4	[64,216–225]
		Flavivirus	RNA	no	0	[189]
		Dugbe virus	RNA	no	2	[226]
		Alkhurma hemorrhagic fever virus	RNA	no	0	[227]
		St Croix River like virus	RNA	no	0	[228]
		<i>Rickettsia aeschlimannii</i>	DNA	no	0	[35]
		<i>Rickettsia conorii</i>	DNA	no	0	[229]
		<i>Anaplasma marginale, centrale, platys</i>	DNA	no	0	[230]
		<i>Coxiella burnetii</i>	DNA	no	0	[74,212,231,232]
<i>Borrelia burgdorferi</i>	RNA and DNA	no	0	[212,233]		
<i>H. schulzei</i>	17/27	Dhuri virus	RNA	no	0	[66]
<i>H. scupense</i>	34/47 <i>H. detritum</i> : 61/90	<i>Theileria equi</i>	No	no	4	[234,235]
		<i>Theileria annulata</i>	No	yes	2, 4	[107,236–238]
		<i>Babesia ovis</i>	DNA	no	0	[40]
		<i>Theileria ovis</i>	DNA	no	0	[40]
		<i>Rickettsia aeschlimannii</i>	DNA	no	0	[14,205]
		<i>Rickettsia slovaca</i>	DNA	no	0	[205]
		<i>Anaplasma phagocytophilum</i>	DNA	no	0	[205]
		<i>Coxiella burnetii</i>	DNA	no	2, 3	[139,239]
<i>H. somalicum</i>	2/2	<i>R. conorii</i>	DNA	no	0	[83]
<i>H. truncatum</i>	142/193	<i>Theileria equi</i>	DNA	no	0	[101]
		<i>Babesia caballi</i>	No	no	3, 4	[240,241]
		<i>Theileria annulata</i>	DNA	no	0	[108]
		CCHFv	RNA, viral particles	yes	1, 2, 3, 4, 5 Cofeeding	[113,148,149,217,218, 220,222,242–246]
		Bunyamwera virus	RNA	no	0	[247]
		Dugbe virus	RNA	no	0	[248]
		Venezuelan Equine Encephalitis Virus		no	2, 4	[249]
		Kupe virus	RNA	no	0	[248]
		Semliki forest virus	RNA	no	0	[247]
		<i>Coxiella burnetii</i>	DNA	no	0	[152,232,250]
<i>Borrelia</i> spp.	DNA	no	0	[152,232]		
<i>H. turanicum</i>	24/36	CCHFv	RNA	no	0	[187]
		<i>Rickettsia sibirica mongolitimonae</i>	DNA	no	0	[140]

CCHFv: Crimean–Congo Hemorrhagic Fever Virus; * epidemiological arguments of possible transmission: for example co-occurrence of a given pathogen in a tick species and in infested vertebrate hosts of the same area, host co-infection with pathogens known to be transmitted by ticks, or the onset of a disease as a result of tick bites: YES, NO, uncertain. ** Experimental validation. 0: none; 1: pathogen reproduction/replication success in ticks; 2: transstadial transmission; 3: transovarial transmission; 4: transmission to a vertebrate host via a tick bite; 5: sexual transmission between male and female ticks.

Author Contributions: Conceptualization, Methodology, Formal Analysis, Investigation, Review and Editing, S.I.B., S.B., A.F., J.F. (Julie Figoni), J.F. (Johanna Fite), T.H., E.Q., S.M., A.R., M.R.-M., G.V. and L.V.; Original Draft writing and preparation, S.I.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The datasets generated during and/or analyzed during the current study can be found in the main text.

Acknowledgments: This narrative review was conducted by the ad hoc subgroup from the working expert group on the risks related to Hyalomma ticks at the French Agency for Food, Environmental and Occupational Health & Safety (ANSES) commissioned by the French authorities. The authors thank the other experts who were part of this group for stimulating discussions and for their feedback on the expert report: S. Baize and F. Stachurski. We also thank Richard Paul and Jeremy Gray for proofreading the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

- de la Fuente, J.; Estrada-Pena, A.; Venzal, J.M.; Kocan, K.M.; Sonenshine, D.E. Overview: Ticks as Vectors of Pathogens That Cause Disease in Humans and Animals. *Front. Biosci. J. Virtual Libr.* **2008**, *13*, 6938–6946. [[CrossRef](#)] [[PubMed](#)]
- de la Fuente, J.; Antunes, S.; Bonnet, S.; Cabezas-Cruz, A.; Domingos, A.G.; Estrada-Peña, A.; Johnson, N.; Kocan, K.M.; Mansfield, K.L.; Nijhof, A.M.; et al. Tick-Pathogen Interactions and Vector Competence: Identification of Molecular Drivers for Tick-Borne Diseases. *Front. Cell. Infect. Microbiol.* **2017**, *7*, 114. [[CrossRef](#)] [[PubMed](#)]
- De, S.; Kitsou, C.; Sonenshine, D.E.; Pedra, J.H.F.; Fikrig, E.; Kassis, J.A.; Pal, U. Epigenetic Regulation of Tick Biology and Vectorial Capacity. *Trends Genet.* **2021**, *37*, 8–11. [[CrossRef](#)] [[PubMed](#)]
- Šimo, L.; Kazimirova, M.; Richardson, J.; Bonnet, S.I. The Essential Role of Tick Salivary Glands and Saliva in Tick Feeding and Pathogen Transmission. *Front. Cell. Infect. Microbiol.* **2017**, *7*, 281. [[CrossRef](#)] [[PubMed](#)]
- Philip, C.B.; Parker, R.R. Rocky Mountain spotted fever. Investigation of sexual transmission in the wood tick *Dermacentor andersoni*. *Public Health Rep.* **1933**, *48*, 266–272. [[CrossRef](#)]
- Léger, E.; Liu, X.; Masegaglia, S.; Noël, V.; Vourc'h, G.; Bonnet, S.; McCoy, K.D. Reliability of Molecular Host-Identification Methods for Ticks: An Experimental In Vitro Study with *Ixodes ricinus*. *Parasit. Vectors* **2015**, *8*, 433. [[CrossRef](#)] [[PubMed](#)]
- Dwuznik, D.; Mierzejewska, E.J.; Drabik, P.; Kloch, A.; Alsarraf, M.; Behnke, J.M.; Bajer, A. The role of juvenile *Dermacentor reticulatus* ticks as vectors of microorganisms and the problem of 'meal contamination'. *Exp. Appl. Acarol.* **2019**, *78*, 181–202. [[CrossRef](#)] [[PubMed](#)]
- Bonnet, S.I.; Liu, X.Y. Laboratory Artificial Infection of Hard Ticks: A Tool for the Analysis of Tick-Borne Pathogen Transmission. *Acarologia* **2012**, *52*, 453–464. [[CrossRef](#)]
- Bonnet, S.I.; Nadal, C. Experimental Infection of Ticks: An Essential Tool for the Analysis of *Babesia* Species Biology and Transmission. *Pathogens* **2021**, *10*, 1403. [[CrossRef](#)]
- Fernández-Ruiz, N.; Estrada-Peña, A. Towards New Horizons: Climate Trends in Europe Increase the Environmental Suitability for Permanent Populations of *Hyalomma Marginatum* (Ixodidae). *Pathogens* **2021**, *10*, 95. [[CrossRef](#)]
- Bah, M.T.; Grosbois, V.; Stachurski, F.; Muñoz, F.; Duhayon, M.; Rakotoarivony, I.; Appelgren, A.; Calloix, C.; Noguera, L.; Mouillaud, T.; et al. The Crimean-Congo Haemorrhagic Fever Tick Vector *Hyalomma Marginatum* in the South of France: Modelling Its Distribution and Determination of Factors Influencing Its Establishment in a Newly Invaded Area. *Transbound. Emerg. Dis.* **2022**, *69*, e2351–e2365. [[CrossRef](#)] [[PubMed](#)]
- Bakheit, M.A.; Latif, A.A.; Vatansver, Z.; Seitzer, U.; Ahmed, J. The Huge Risks Due to *Hyalomma* Ticks. In *Arthropods as Vectors of Emerging Diseases*; Mehlhorn, H., Ed.; Parasitology Research Monographs; Springer: Berlin/Heidelberg, Germany, 2012; pp. 167–194; ISBN 978-3-642-28842-5.
- Bonnet, S.I.; Vourc'h, G.; Raffetin, A.; Falchi, A.; Figoni, J.; Fite, J.; Hoch, T.; Moutailler, S.; Quillery, E. The Control of *Hyalomma* Ticks, Vectors of the Crimean-Congo Hemorrhagic Fever Virus: Where Are We Now and Where Are We Going? *PLoS Negl. Trop. Dis.* **2022**, *16*, e0010846. [[CrossRef](#)] [[PubMed](#)]
- Kumar, B.; Manjunathachar, H.V.; Ghosh, S. A Review on *Hyalomma* Species Infestations on Human and Animals and Progress on Management Strategies. *Heliyon* **2020**, *6*, e05675. [[CrossRef](#)]
- Estrada-Peña, A.; De La Fuente, J.; Latapia, T.; Ortega, C. The Impact of Climate Trends on a Tick Affecting Public Health: A Retrospective Modeling Approach for *Hyalomma Marginatum* (Ixodidae). *PLoS ONE* **2015**, *10*, e0125760. [[CrossRef](#)]
- Guglielmone, A.A.; Robbins, R.G.; Apanaskevich, D.A.; Petney, T.N.; Estrada-Peña, A.; Horak, I.G.; Shao, R.; Barker, S.C. The Argasidae, Ixodidae and Nuttalliellidae (Acari: Ixodida) of the World: A List of Valid Species Names. *Zootaxa* **2010**, *2528*, 1. [[CrossRef](#)]

17. Kar, S.; Rodriguez, S.E.; Akyildiz, G.; Cajimat, M.N.B.; Bircan, R.; Mears, M.C.; Bente, D.A.; Keles, A.G. Crimean-Congo Hemorrhagic Fever Virus in Tortoises and *Hyalomma Aegyptium* Ticks in East Thrace, Turkey: Potential of a Cryptic Transmission Cycle. *Parasit. Vectors* **2020**, *13*, 201. [[CrossRef](#)] [[PubMed](#)]
18. Paştiu, A.I.; Matei, I.A.; Mihalca, A.D.; D'Amico, G.; Dumitrache, M.O.; Kalmár, Z.; Sándor, A.D.; Lefkaditis, M.; Gherman, C.M.; Cozma, V. Zoonotic Pathogens Associated with *Hyalomma Aegyptium* in Endangered Tortoises: Evidence for Host-Switching Behaviour in Ticks? *Parasit. Vectors* **2012**, *5*, 301. [[CrossRef](#)]
19. Siroký, P.; Kubelová, M.; Modrý, D.; Erhart, J.; Literák, I.; Spitalská, E.; Kocianová, E. Tortoise Tick *Hyalomma Aegyptium* as Long Term Carrier of Q Fever Agent *Coxiella Burnetii*—Evidence from Experimental Infection. *Parasitol. Res.* **2010**, *107*, 1515–1520. [[CrossRef](#)]
20. Brinkmann, A.; Hekimoğlu, O.; Dinçer, E.; Hagedorn, P.; Nitsche, A.; Ergünay, K. A Cross-Sectional Screening by next-Generation Sequencing Reveals *Rickettsia*, *Coxiella*, *Francisella*, *Borrelia*, *Babesia*, *Theileria* and *Hemolivia* Species in Ticks from Anatolia. *Parasit. Vectors* **2019**, *12*, 26. [[CrossRef](#)]
21. Kalmár, Z.; Cozma, V.; Sprong, H.; Jahfari, S.; D'Amico, G.; Mărcuţan, D.I.; Ionică, A.M.; Magdaş, C.; Modrý, D.; Mihalca, A.D. Transstadial Transmission of *Borrelia Turcica* in *Hyalomma Aegyptium* Ticks. *PLoS ONE* **2015**, *10*, e0115520. [[CrossRef](#)]
22. Takano, A.; Goka, K.; Une, Y.; Shimada, Y.; Fujita, H.; Shiino, T.; Watanabe, H.; Kawabata, H. Isolation and Characterization of a Novel *Borrelia* Group of Tick-Borne *Borreliae* from Imported Reptiles and Their Associated Ticks. *Environ. Microbiol.* **2010**, *12*, 134–146. [[CrossRef](#)] [[PubMed](#)]
23. Hepner, S.; Fingerle, V.; Duscher, G.G.; Felsberger, G.; Marosevic, D.; Rollins, R.E.; Okeyo, M.; Sing, A.; Margos, G. Population Structure of *Borrelia Turcica* from Greece and Turkey. *Infect. Genet. Evol. J. Mol. Epidemiol. Evol. Genet. Infect. Dis.* **2020**, *77*, 104050. [[CrossRef](#)] [[PubMed](#)]
24. Keskin, A.; Bursali, A.; Snow, D.E.; Dowd, S.E.; Tekin, S. Assessment of Bacterial Diversity in *Hyalomma Aegyptium*, *H. Marginatum* and *H. Excavatum* Ticks through Tag-Encoded Pyrosequencing. *Exp. Appl. Acarol.* **2017**, *73*, 461–475. [[CrossRef](#)] [[PubMed](#)]
25. Norte, A.C.; Harris, D.J.; Silveira, D.; Nunes, C.S.; Nuncio, M.S.; Martínez, E.G.; Giménez, A.; de Sousa, R.; Lopes de Carvalho, I.; Perera, A. Diversity of Microorganisms in *Hyalomma Aegyptium* Collected from Spur-Thighed Tortoise (*Testudo Graeca*) in North Africa and Anatolia. *Transbound. Emerg. Dis.* **2022**, *69*, 1951–1962. [[CrossRef](#)]
26. Akveran, G.A.; Karasartova, D.; Keskin, A.; Comba, A.; Celebi, B.; Mumcuoglu, K.Y.; Taylan-Ozkan, A. Bacterial and Protozoan Agents Found in *Hyalomma Aegyptium* (L., 1758) (Ixodida: Ixodidae) Collected from *Testudo Graeca* L., 1758 (Reptilia: Testudines) in Corum Province of Turkey. *Ticks Tick-Borne Dis.* **2020**, *11*, 101458. [[CrossRef](#)] [[PubMed](#)]
27. Barradas, P.F.; Lima, C.; Cardoso, L.; Amorim, I.; Gärtner, F.; Mesquita, J.R. Molecular Evidence of *Hemolivia Mauritanica*, *Ehrlichia* spp. and the Endosymbiont *Candidatus Midichloria Mitochondrii* in *Hyalomma Aegyptium* Infesting *Testudo Graeca* Tortoises from Doha, Qatar. *Animals* **2020**, *11*, 30. [[CrossRef](#)]
28. Gargili, A.; Palomar, A.M.; Midilli, K.; Portillo, A.; Kar, S.; Oteo, J.A. *Rickettsia* Species in Ticks Removed from Humans in Istanbul, Turkey. *Vector Borne Zoonotic Dis.* **2012**, *12*, 938–941. [[CrossRef](#)]
29. Bitam, I.; Kernif, T.; Harrat, Z.; Parola, P.; Raoult, D. First Detection of *Rickettsia Aeschlimannii* in *Hyalomma Aegyptium* from Algeria. *Clin. Microbiol. Infect.* **2009**, *15*, 253–254. [[CrossRef](#)]
30. Orkun, Ö. Molecular Investigation of the Natural Transovarial Transmission of Tick-Borne Pathogens in Turkey. *Vet. Parasitol.* **2019**, *273*, 97–104. [[CrossRef](#)]
31. Orkun, Ö.; Emir, H. Identification of Tick-Borne Pathogens in Ticks Collected from Wild Animals in Turkey. *Parasitol. Res.* **2020**, *119*, 3083–3091. [[CrossRef](#)]
32. Ergünay, K.; Dinçer, E.; Kar, S.; Emanet, N.; Yalçınkaya, D.; Polat Dinçer, P.F.; Brinkmann, A.; Hacıoğlu, S.; Nitsche, A.; Özkul, A.; et al. Multiple Orthonairoviruses Including Crimean-Congo Hemorrhagic Fever Virus, Tamdy Virus and the Novel Meram Virus in Anatolia. *Ticks Tick-Borne Dis.* **2020**, *11*, 101448. [[CrossRef](#)] [[PubMed](#)]
33. Li, Y.; Galon, E.M.; Guo, Q.; Rizk, M.A.; Moumouni, P.F.A.; Liu, M.; Li, J.; Ji, S.; Chahan, B.; Xuan, X. Molecular Detection and Identification of *Babesia* spp., *Theileria* spp., and *Anaplasma* spp. in Sheep From Border Regions, Northwestern China. *Front. Vet. Sci.* **2020**, *7*, 630. [[CrossRef](#)] [[PubMed](#)]
34. Moltmann, U.G.; Mehlhorn, H.; Schein, E.; Voigt, W.P.; Friedhoff, K.T. Ultrastructural Study on the Development of *Babesia equi* (Coccidia: Piroplasmia) in the Salivary Glands of Its Vector Ticks. *J. Protozool.* **1983**, *30*, 218–225. [[CrossRef](#)]
35. Springer, A.; Shuaib, Y.A.; Isaa, M.H.; Ezz-Eldin, M.I.-E.; Osman, A.Y.; Yagoub, I.A.; Abdalla, M.A.; Bakiet, A.O.; Mohmed-Noor, S.E.-T.; Schaper, S.; et al. Tick Fauna and Associated *Rickettsia*, *Theileria*, and *Babesia* spp. in Domestic Animals in Sudan (North Kordofan and Kassala States). *Microorganisms* **2020**, *8*, E1969. [[CrossRef](#)]
36. Kumar, S.; Malhotra, D.V.; Sangwan, A.K.; Goel, P.; Kumar, A.; Kumar, S. Infectivity Rate and Transmission Potential of *Hyalomma anatolicum anatolicum* Ticks for *Babesia equi* Infection. *Vet. Parasitol.* **2007**, *144*, 338–343. [[CrossRef](#)]
37. Bhagwan, J.; Kumar, A.; Kumar, R.; Goyal, L.; Goel, P.; Kumar, S. Molecular Evidence of *Theileria equi* Infection in *Hyalomma anatolicum* Ticks Infested on Sero-Positive Indian Horses. *Acta Parasitol.* **2015**, *60*, 322–329. [[CrossRef](#)] [[PubMed](#)]
38. Ghafar, A.; Cabezas-Cruz, A.; Galon, C.; Obregon, D.; Gasser, R.B.; Moutailler, S.; Jabbar, A. Bovine Ticks Harbour a Diverse Array of Microorganisms in Pakistan. *Parasit. Vectors* **2020**, *13*, 1. [[CrossRef](#)]
39. Zeb, J.; Szekeres, S.; Takács, N.; Kontschán, J.; Shams, S.; Ayaz, S.; Hornok, S. Genetic Diversity, Piroplasms and Trypanosomes in *Rhipicephalus microplus* and *Hyalomma anatolicum* Collected from Cattle in Northern Pakistan. *Exp. Appl. Acarol.* **2019**, *79*, 233–243. [[CrossRef](#)]

40. Bekloo, A.J.; Bakhshi, H.; Soufizadeh, A.; Sedaghat, M.M.; Bekloo, R.J.; Ramzgouyan, M.R.; Chegeni, A.H.; Faghihi, F.; Telmadarraiy, Z. Ticks Circulate *Anaplasma*, *Ehrlichia*, *Babesia* and *Theileria* Parasites in North of Iran. *Vet. Parasitol.* **2017**, *248*, 21–24. [[CrossRef](#)]
41. Mossaad, E.; Gaithuma, A.; Mohamed, Y.O.; Sukanuma, K.; Umemiya-Shirafuji, R.; Ohari, Y.; Salim, B.; Liu, M.; Xuan, X. Molecular Characterization of Ticks and Tick-Borne Pathogens in Cattle from Khartoum State and East Darfur State, Sudan. *Pathogens* **2021**, *10*, 580. [[CrossRef](#)]
42. Afshari, A.; Habibi, G.; Abdigoudarzi, M.; Yazdani, F. Establishment and Validation of *Theileria Annulata* Sporozoite Ak-93 Infection in Laboratory-Reared *Hyalomma anatolicum* Tick Using In Vivo and In Vitro Assays. *J. Arthropod-Borne Dis.* **2020**, *14*, 261–269. [[CrossRef](#)] [[PubMed](#)]
43. Aktas, M.; Dumanli, N.; Angin, M. Cattle Infestation by *Hyalomma* Ticks and Prevalence of *Theileria* in *Hyalomma* Species in the East of Turkey. *Vet. Parasitol.* **2004**, *119*, 1–8. [[CrossRef](#)]
44. Al-Fahdi, A.; Alqamashoui, B.; Al-Hamidhi, S.; Kose, O.; Tageldin, M.H.; Bobade, P.; Johnson, E.H.; Hussain, A.-R.; Karagenc, T.; Tait, A.; et al. Molecular Surveillance of *Theileria* Parasites of Livestock in Oman. *Ticks Tick-Borne Dis.* **2017**, *8*, 741–748. [[CrossRef](#)] [[PubMed](#)]
45. Amiri, M.; Yaghfoori, S.; Razmi, G. Molecular Detection of *Theileria annulata* among Dairy Cattle and Vector Ticks in the Herat Area, Afghanistan. *Arch. Razi Inst.* **2021**, *76*, 79–85. [[CrossRef](#)]
46. Dehuri, M.; Panda, M.; Sahoo, N.; Mohanty, B.; Behera, B. Nested PCR Assay for Detection of *Theileria annulata* in *Hyalomma anatolicum* Infesting Cattle from Coastal Odisha, India. *Anim. Biotechnol.* **2022**, *33*, 1229–1234. [[CrossRef](#)]
47. Kartashov, M.Y.; Kononova, Y.V.; Petrova, I.D.; Tupota, N.L.; Mikryukova, T.P.; Ternovoi, V.A.; Tishkova, F.H.; Loktev, V.B. Detection of *Ehrlichia* spp. and *Theileria* spp. in *Hyalomma anatolicum* Ticks Collected in Tajikistan. *Vavilovskii Zhurnal Genet. Sel.* **2020**, *24*, 55–59. [[CrossRef](#)]
48. Omer, S.A.; Alsuwaid, D.F.; Mohammed, O.B. Molecular Characterization of Ticks and Tick-Borne Piroplasms from Cattle and Camel in Hofuf, Eastern Saudi Arabia. *Saudi J. Biol. Sci.* **2021**, *28*, 2023–2028. [[CrossRef](#)]
49. Rahmani-Varmale, M.; Tavassoli, M.; Esmaeilnejad, B. Molecular Detection and Differentiation of *Theileria lestoquardi*, *T. ovis* and *T. annulata* in Blood of Goats and Ticks in Kermanshah Province, Iran. *J. Arthropod-Borne Dis.* **2019**, *13*, 297–309. [[CrossRef](#)]
50. Robinson, P.M. Theileriosis annulata and Its Transmission—A Review. *Trop. Anim. Health Prod.* **1982**, *14*, 3–12. [[CrossRef](#)]
51. Sayin, F.; Dinçer, S.; Karaer, Z.; Cakmak, A.; Inci, A.; Yukari, B.A.; Eren, H.; Vatanserver, Z.; Nalbantoglu, S. Studies on the Epidemiology of Tropical Theileriosis (*Theileria annulata* Infection) in Cattle in Central Anatolia, Turkey. *Trop. Anim. Health Prod.* **2003**, *35*, 521–539. [[CrossRef](#)]
52. Yaghfoori, S.; Mohri, M.; Razmi, G. Experimental *Theileria lestoquardi* Infection in Sheep: Biochemical and Hematological Changes. *Acta Trop.* **2017**, *173*, 55–61. [[CrossRef](#)] [[PubMed](#)]
53. Tajeri, S.; Razmi, G.; Haghparast, A. Establishment of an Artificial Tick Feeding System to Study *Theileria lestoquardi* Infection. *PLoS ONE* **2016**, *11*, e0169053. [[CrossRef](#)] [[PubMed](#)]
54. Taha, K.M.; Elhussein, A.M. Experimental Transmission of *Theileria lestoquardi* by Developmental Stages of *Hyalomma anatolicum* Ticks. *Parasitol. Res.* **2010**, *107*, 1009–1012. [[CrossRef](#)]
55. Abdigoudarzi, M. Detection of Naturally Infected Vector Ticks (Acari: Ixodidae) by Different Species of *Babesia* and *Theileria* Agents from Three Different Enzootic Parts of Iran. *J. Arthropod-Borne Dis.* **2013**, *7*, 164–172.
56. Taha, K.M.; Salih, D.A.; Ahmed, B.M.; Enan, K.A.; Ali, A.M.; ElHussein, A.M. First Confirmed Report of Outbreak of Malignant Ovine Theileriosis among Goats in Sudan. *Parasitol. Res.* **2011**, *109*, 1525–1527. [[CrossRef](#)] [[PubMed](#)]
57. Li, Y.; Guan, G.; Liu, A.; Peng, Y.; Luo, J.; Yin, H. Experimental Transmission of *Theileria ovis* by *Hyalomma anatolicum anatolicum*. *Parasitol. Res.* **2010**, *106*, 991–994. [[CrossRef](#)]
58. Li, Y.; Guan, G.; Ma, M.; Liu, J.; Ren, Q.; Luo, J.; Yin, H. *Theileria ovis* Discovered in China. *Exp. Parasitol.* **2011**, *127*, 304–307. [[CrossRef](#)]
59. Williams, R.J.; Al-Busaidy, S.; Mehta, F.R.; Maupin, G.O.; Wagoner, K.D.; Al-Awaidy, S.; Suleiman, A.J.; Khan, A.S.; Peters, C.J.; Ksiazek, T.G. Crimean-Congo Haemorrhagic Fever: A Seroepidemiological and Tick Survey in the Sultanate of Oman. *Trop. Med. Int. Health* **2000**, *5*, 99–106. [[CrossRef](#)]
60. Bell-Sakyi, L.; Kohl, A.; Bente, D.A.; Fazakerley, J.K. Tick Cell Lines for Study of Crimean-Congo Hemorrhagic Fever Virus and Other Arboviruses. *Vector Borne Zoonotic Dis.* **2012**, *12*, 769–781. [[CrossRef](#)]
61. Khan, A.S.; Maupin, G.O.; Rollin, P.E.; Noor, A.M.; Shurie, H.H.; Shalabi, A.G.; Wasef, S.; Haddad, Y.M.; Sadek, R.; Ijaz, K.; et al. An Outbreak of Crimean-Congo Hemorrhagic Fever in the United Arab Emirates, 1994–1995. *Am. J. Trop. Med. Hyg.* **1997**, *57*, 519–525. [[CrossRef](#)]
62. Petrova, I.D.; Kononova, I.V.; Chausov, E.V.; Shestopalov, A.M.; Tishkova, F.K. Genetic variants of the Crimean-Congo hemorrhagic fever virus circulating in endemic areas of the southern Tajikistan in 2009. *Mol. Genet. Mikrobiol. Virusol.* **2013**, *28*, 29–36. [[CrossRef](#)]
63. Mourya, D.T.; Yadav, P.D.; Shete, A.M.; Gurav, Y.K.; Raut, C.G.; Jadi, R.S.; Pawar, S.D.; Nichol, S.T.; Mishra, A.C. Detection, Isolation and Confirmation of Crimean-Congo Hemorrhagic Fever Virus in Human, Ticks and Animals in Ahmadabad, India, 2010–2011. *PLoS Negl. Trop. Dis.* **2012**, *6*, e1653. [[CrossRef](#)] [[PubMed](#)]
64. Shahid, M.F.; Yaqub, T.; Ali, M.; Ul-Rahman, A.; Bente, D.A. Prevalence and Phylogenetic Analysis of Crimean-Congo Hemorrhagic Fever Virus in Ticks Collected from Punjab Province of Pakistan. *Acta Trop.* **2021**, *218*, 105892. [[CrossRef](#)] [[PubMed](#)]

65. Kayedi, M.H.; Chinikar, S.; Mostafavi, E.; Khakifirooz, S.; Jalali, T.; Hosseini-Chegeni, A.; Naghizadeh, A.; Niedrig, M.; Fooks, A.R.; Shahhosseini, N. Crimean-Congo Hemorrhagic Fever Virus Clade IV (Asia 1) in Ticks of Western Iran. *J. Med. Entomol.* **2015**, *52*, 1144–1149. [[CrossRef](#)]
66. Al-Khalifa, M.S.; Diab, F.M.; Khalil, G.M. Man-Threatening Viruses Isolated from Ticks in Saudi Arabia. *Saudi Med. J.* **2007**, *28*, 1864–1867.
67. Dilcher, M.; Faye, O.; Faye, O.; Weber, F.; Koch, A.; Sadegh, C.; Weidmann, M.; Sall, A.A. Zahedan Rhabdovirus, a Novel Virus Detected in Ticks from Iran. *Viol. J.* **2015**, *12*, 183. [[CrossRef](#)]
68. Chunikhin, S.P.; Stefuktina, L.F.; Korolev, M.B.; Reshetnikov, I.A.; Khozinskaia, G.A. Sexual transmission of the tick-borne encephalitis virus in ixodid ticks (Ixodidae). *Parazitologiya* **1983**, *17*, 214–217.
69. Aristova, V.A.; Gushchina, E.A.; Gromashevskii, V.L.; Gushchin, B.V. Experimental infection of ixodid ticks with Karshi virus. *Parazitologiya* **1986**, *20*, 347–350.
70. Yadav, P.D.; Whitmer, S.L.M.; Sarkale, P.; Fei Fan Ng, T.; Goldsmith, C.S.; Nyayanit, D.A.; Esona, M.D.; Shrivastava-Ranjan, P.; Lakra, R.; Pardeshi, P.; et al. Characterization of Novel Reoviruses Wad Medani Virus (Orbivirus) and Kundal Virus (Coltivirus) Collected from *Hyalomma anatolicum* Ticks in India during Surveillance for Crimean Congo Hemorrhagic Fever. *J. Virol.* **2019**, *93*, e00106-19. [[CrossRef](#)]
71. Kostiuokov, M.A.; Daniyarov, O.; Skvortsova, T.M.; Kondrashina, N.G.; Berezina, L.K. Isolation of the Sindbis virus from *Hyalomma anatolicum* CL Kock 1844 ticks in Tadzhikistan. *Med. Parazitol.* **1981**, *50*, 34–35.
72. Fard, N.S.R.; Khalili, M. PCR-Detection of *Coxiella burnetii* in Ticks Collected from Sheep and Goats in Southeast Iran. *Iran. J. Arthropod-Borne Dis.* **2011**, *5*, 1–6.
73. Fard, N.S.R.; Omid Ghashghaei, O.; Khalili, M.; Sharifi, H. Tick Diversity and Detection of *Coxiella burnetii* in Tick of Small Ruminants Using Nested Trans PCR in Southeast Iran. *Trop. Biomed.* **2016**, *33*, 506–511.
74. Ni, J.; Lin, H.; Xu, X.; Ren, Q.; Aizezi, M.; Luo, J.; Luo, Y.; Ma, Z.; Chen, Z.; Tan, Y.; et al. *Coxiella burnetii* Is Widespread in Ticks (Ixodidae) in the Xinjiang Areas of China. *BMC Vet. Res.* **2020**, *16*, 317. [[CrossRef](#)] [[PubMed](#)]
75. Choubdar, N.; Karimian, F.; Koosha, M.; Nejati, J.; Oshaghi, M.A. *Hyalomma* spp. Ticks and Associated *Anaplasma* spp. and *Ehrlichia* spp. on the Iran-Pakistan Border. *Parasit. Vectors* **2021**, *14*, 469. [[CrossRef](#)] [[PubMed](#)]
76. Yamchi, J.A.; Tavassoli, M. Survey on Infection Rate, Vectors and Molecular Identification of *Theileria annulata* in Cattle from North West, Iran. *J. Parasit. Dis.* **2016**, *40*, 1071–1076. [[CrossRef](#)]
77. Meng, K.; Li, Z.; Wang, Y.; Jing, Z.; Zhao, X.; Liu, J.; Cai, D.; Zhang, L.; Yang, D.; Wang, S. PCR-Based Detection of *Theileria annulata* in *Hyalomma asiaticum* Ticks in Northwestern China. *Ticks Tick-Borne Dis.* **2014**, *5*, 105–106. [[CrossRef](#)]
78. Razmi, G.R.; Hosseini, M.; Aslani, M.R. Identification of Tick Vectors of Ovine Theileriosis in an Endemic Region of Iran. *Vet. Parasitol.* **2003**, *116*, 1–6. [[CrossRef](#)]
79. Sun, M.; Wang, J.; Liu, Z.; Guan, G.; Li, Y.; Liu, J.; Xu, J.; Yin, H.; Luo, J. First Molecular Evidence of *Babesia occultans* and *Theileria separata* Infection in Ticks and Sheep in China. *Exp. Appl. Acarol.* **2019**, *78*, 223–229. [[CrossRef](#)]
80. Narankhajid, M.; Yeruult, C.; Gurbadam, A.; Battsetseg, J.; Aberle, S.W.; Bayartogtokh, B.; Joachim, A.; Duscher, G.G. Some Aspects on Tick Species in Mongolia and Their Potential Role in the Transmission of Equine Piroplasms, *Anaplasma phagocytophilum* and *Borrelia burgdorferi* L. *Parasitol. Res.* **2018**, *117*, 3557–3566. [[CrossRef](#)]
81. Song, R.; Wang, Q.; Guo, F.; Liu, X.; Song, S.; Chen, C.; Tu, C.; Wureli, H.; Wang, Y. Detection of *Babesia* spp., *Theileria* spp. and *Anaplasma ovis* in Border Regions, Northwestern China. *Transbound. Emerg. Dis.* **2018**, *65*, 1537–1544. [[CrossRef](#)]
82. Guo, R.; Shen, S.; Zhang, Y.; Shi, J.; Su, Z.; Liu, D.; Liu, J.; Yang, J.; Wang, Q.; Hu, Z.; et al. A New Strain of Crimean-Congo Hemorrhagic Fever Virus Isolated from Xinjiang, China. *Virol. Sin.* **2017**, *32*, 80–88. [[CrossRef](#)] [[PubMed](#)]
83. Moming, A.; Yue, X.; Shen, S.; Chang, C.; Wang, C.; Luo, T.; Zhang, Y.; Guo, R.; Hu, Z.; Zhang, Y.; et al. Prevalence and Phylogenetic Analysis of Crimean-Congo Hemorrhagic Fever Virus in Ticks from Different Ecosystems in Xinjiang, China. *Virol. Sin.* **2018**, *33*, 67–73. [[CrossRef](#)] [[PubMed](#)]
84. Sun, S.; Dai, X.; Aishan, M.; Wang, X.; Meng, W.; Feng, C.; Zhang, F.; Hang, C.; Hu, Z.; Zhang, Y. Epidemiology and Phylogenetic Analysis of Crimean-Congo Hemorrhagic Fever Viruses in Xinjiang, China. *J. Clin. Microbiol.* **2009**, *47*, 2536–2543. [[CrossRef](#)] [[PubMed](#)]
85. Zhang, Y.; Shen, S.; Fang, Y.; Liu, J.; Su, Z.; Liang, J.; Zhang, Z.; Wu, Q.; Wang, C.; Abudurexiti, A.; et al. Isolation, Characterization, and Phylogenetic Analysis of Two New Crimean-Congo Hemorrhagic Fever Virus Strains from the Northern Region of Xinjiang Province, China. *Virol. Sin.* **2018**, *33*, 74–86. [[CrossRef](#)]
86. L'vov, D.K.; Al'khovskii, S.V.; Shchelkanov, M.I.; Shchetinin, A.M.; Deriabin, P.G.; Gitel'man, A.K.; Aristova, V.A.; Botikov, A.G. Taxonomic status of the Burana virus (BURV) (Bunyaviridae, Nairovirus, Tamdy group) isolated from the ticks *Haemaphysalis punctata* Canestrini et Fanzago, 1877 and *Haem. concinna* Koch, 1844 (Ixodidae, Haemaphysalinae) in Kyrgyzstan. *Vopr. Virusol.* **2014**, *59*, 10–15. [[PubMed](#)]
87. Karimov, S.K.; L'vov, D.K.; Rogovaia, S.G.; Kiriushchenko, T.V.; Ukbueva, T.D. Isolation of the Karshi virus from *Hyalomma asiaticum* ticks in Alma-Ata Province, Kazakh SSR. *Med. Parazitol.* **1978**, *47*, 50–51.
88. Khutoretskaya, N.V.; Aristova, V.A.; Rogovaya, S.G.; Lvov, D.K.; Karimov, S.K.; Skvortsova, T.M.; Kondrashina, N.G. Experimental Study of the Reproduction of Karshi Virus (Togaviridae, Flavivirus) in Some Species of Mosquitoes and Ticks. *Acta Virol.* **1985**, *29*, 231–236.

89. L'vov, D.K.; Sidorova, G.A.; Gromashevsky, V.L.; Kurbanov, M.; Skvoztsova, L.M.; Gofman, Y.P.; Berezina, L.K.; Klimenko, S.M.; Zakharyan, V.A.; Aristova, V.A.; et al. Virus "Tamdy"—A New Arbovirus, Isolated in the Uzbee S.S.R. and Turkmen S.S.R. from Ticks *Hyalomma asiaticum asiaticum* Schullee et Schlottke, 1929, and *Hyalomma plumbeum plumbeum* Panzer, 1796. *Arch. Virol.* **1976**, *51*, 15–21. [[CrossRef](#)]
90. L'vov, D.K.; Sidorova, G.A.; Gromashevskii, V.L.; Skvortsova, T.M.; Aristova, V.A. Isolation of Tamdy virus (Bunyaviridae) pathogenic for man from natural sources in Central Asia, Kazakhstan and Transcaucasia. *Vopr. Virusol.* **1984**, *29*, 487–490.
91. Zhou, H.; Ma, Z.; Hu, T.; Bi, Y.; Mamuti, A.; Yu, R.; Carr, M.J.; Shi, M.; Li, J.; Sharshov, K.; et al. Tamdy Virus in Ixodid Ticks Infesting Bactrian Camels, Xinjiang, China, 2018. *Emerg. Infect. Dis.* **2019**, *25*, 2136–2138. [[CrossRef](#)]
92. Liu, X.; Zhang, X.; Wang, Z.; Dong, Z.; Xie, S.; Jiang, M.; Song, R.; Ma, J.; Chen, S.; Chen, K.; et al. A Tentative Tamdy Orthonavivirus Related to Febrile Illness in Northwestern China. *Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.* **2020**, *70*, 2155–2160. [[CrossRef](#)] [[PubMed](#)]
93. Daiter, A.B. Transovarial and transpermal transmission of *Coxiella burnetii* by the tick *Hyalomma asiaticum* and its role in the ecology of Q rickettsiosis. *Parazitologiya* **1977**, *11*, 403–411. [[PubMed](#)]
94. Daiter, A.B. An experimental infection of *Hyalomma asiaticum* and *Ornithodoros papillipes* ticks with a single and combined infection with *Coxiella burnetii* and *Dermacentroxenus sibiricus*. *Parazitologiya* **1979**, *13*, 8–18. [[PubMed](#)]
95. Batu, N.; Wang, Y.; Liu, Z.; Huang, T.; Bao, W.; He, H.; Geri, L. Molecular Epidemiology of *Rickettsia* sp. and *Coxiella burnetii* Collected from *Hyalomma asiaticum* in Bactrian Camels (*Camelus bactrianus*) in Inner Mongolia of China. *Ticks Tick-Borne Dis.* **2020**, *11*, 101548. [[CrossRef](#)] [[PubMed](#)]
96. Parola, P.; Inokuma, H.; Camicas, J.L.; Brouqui, P.; Raoult, D. Detection and Identification of Spotted Fever Group *Rickettsiae* and *Ehrlichiae* in African Ticks. *Emerg. Infect. Dis.* **2001**, *7*, 1014–1017. [[CrossRef](#)]
97. Wang, W.; Liu, X.; Wang, X.; Dong, H.; Ma, C.; Wang, J.; Liu, B.; Mao, Y.; Wang, Y.; Li, T.; et al. Structural and Functional Diversity of Nairovirus-Encoded Nucleoproteins. *J. Virol.* **2015**, *89*, 11740–11749. [[CrossRef](#)] [[PubMed](#)]
98. Yu, K.F.; Pauls, K.P. Segregation of Random Amplified Polymorphic DNA Markers and Strategies for Molecular Mapping in Tetraploid Alfalfa. *Genome* **1993**, *36*, 844–851. [[CrossRef](#)]
99. Zapf, E.; Schein, E. New Findings in the Development Of *Babesia (Theileria) equi* (Laveran, 1901) in the Salivary Glands of the Vector Ticks, *Hyalomma* Species. *Parasitol. Res.* **1994**, *80*, 543–548. [[CrossRef](#)]
100. Zapf, F.; Schein, E. The Development of *Babesia (Theileria) equi* (Laveran, 1901) in the Gut and the Haemolymph of the Vector Ticks, *Hyalomma* Species. *Parasitol. Res.* **1994**, *80*, 297–302. [[CrossRef](#)]
101. Onyiche, T.E.; Răileanu, C.; Tauchmann, O.; Fischer, S.; Vasić, A.; Schäfer, M.; Biu, A.A.; Ogo, N.I.; Thekisoe, O.; Silaghi, C. Prevalence and Molecular Characterization of Ticks and Tick-Borne Pathogens of One-Humped Camels (*Camelus dromedarius*) in Nigeria. *Parasit. Vectors* **2020**, *13*, 428. [[CrossRef](#)]
102. Scoles, G.A.; Ueti, M.W. Vector Ecology of Equine Piroplasmiasis. *Annu. Rev. Entomol.* **2015**, *60*, 561–580. [[CrossRef](#)] [[PubMed](#)]
103. Hamed, M.I.; Zaitoun, A.M.A.; El-Allawy, T.A.A.; Mourad, M.I. Investigation of *Theileria camelensis* in Camels Infested by *Hyalomma Dromedarii* Ticks in Upper Egypt. *J. Adv. Vet. Res.* **2011**, *1*, 4–7.
104. Kumar, S.; Sudan, V.; Shanker, D.; Devi, A. *Babesia (Theileria) equi* Genotype A among Indian Equine Population. *Vet. Parasitol. Reg. Stud. Rep.* **2020**, *19*, 100367. [[CrossRef](#)] [[PubMed](#)]
105. Hoogstraal, H.; Wassef, H.Y.; Buttiker, W. Ticks (Acarina) of Saudi Arabia Fam *Argasidae* *Ixodidae*. *Fauna Saudi Arab.* **1981**, *3*, 25–110.
106. Mazlum, Z. Transmission of *Theileria annulata* by the Crushed Infected Unfed *Hyalomma dromedarii*. *Parasitology* **1969**, *59*, 597–600. [[CrossRef](#)] [[PubMed](#)]
107. Gharbi, M.; Darghouth, M.A.; Elati, K.; AL-Hosary, A.A.T.; Ayadi, O.; Salih, D.A.; El Hussein, A.M.; Mhadhbi, M.; Khamassi Khbou, M.; Hassan, S.M.; et al. Current Status of Tropical Theileriosis in Northern Africa: A Review of Recent Epidemiological Investigations and Implications for Control. *Transbound. Emerg. Dis.* **2020**, *67*, 8–25. [[CrossRef](#)] [[PubMed](#)]
108. Mamman, A.H.; Lorusso, V.; Adam, B.M.; Dogo, G.A.; Bown, K.J.; Birtles, R.J. Correction to: First Report of *Theileria annulata* in Nigeria: Findings from Cattle Ticks in Zamfara and Sokoto States. *Parasit. Vectors* **2021**, *14*, 353. [[CrossRef](#)]
109. d'Oliveira, C.; van der Weide, M.; Jacquiet, P.; Jongejan, F. Detection of *Theileria annulata* by the PCR in Ticks (Acari: Ixodidae) Collected from Cattle in Mauritania. *Exp. Appl. Acarol.* **1997**, *21*, 279–291. [[CrossRef](#)]
110. Youssef, S.Y.; Yasien, S.; Mousa, W.M.A.; Nasr, S.M.; El-Kelesh, E.A.M.; Mahran, K.M.; Abd-El-Rahman, A.H. Vector Identification and Clinical, Hematological, Biochemical, and Parasitological Characteristics of Camel (*Camelus dromedarius*) Theileriosis in Egypt. *Trop. Anim. Health Prod.* **2015**, *47*, 649–656. [[CrossRef](#)]
111. de Kok, J.B.; d'Oliveira, C.; Jongejan, F. Detection of the Protozoan Parasite *Theileria annulata* in *Hyalomma* Ticks by the Polymerase Chain Reaction. *Exp. Appl. Acarol.* **1993**, *17*, 839–846. [[CrossRef](#)]
112. Mustafa, U.E.H.; Jongejan, F.; Morzaria, S.P. Note on the Transmission of *Theileria annulata* by *Hyalomma* Ticks in the Sudan. *Vet. Q.* **1983**, *5*, 112–113. [[CrossRef](#)] [[PubMed](#)]
113. Logan, T.M.; Linthicum, K.J.; Bailey, C.L.; Watts, D.M.; Dohm, D.J.; Moulton, J.R. Replication of Crimean-Congo Hemorrhagic Fever Virus in Four Species of Ixodid Ticks (Acari) Infected Experimentally. *J. Med. Entomol.* **1990**, *27*, 537–542. [[CrossRef](#)] [[PubMed](#)]
114. Smirnova, S.E.; Mamaev, V.I.; Nepesova, N.M.; Filipenko, P.I.; Kalieva, V.I. Study of the circulation of Crimean hemorrhagic fever virus in Turkmenistan. *J. Microbiol. Epidemiol. Immunobiol.* **1978**, *1*, 92–97.

115. Begum, F.; Wisseman, C.L.; Casals, J. Tick-Borne Viruses of West Pakistan. II. Hazara Virus, a New Agent Isolated from *Ixodes redikorzevi* Ticks from the Kaghan Valley, W. Pakistan. *Am. J. Epidemiol.* **1970**, *92*, 192–194. [[CrossRef](#)]
116. Anderson, C.R.; Casals, J. Dhori Virus, a New Agent Isolated from *Hyalomma dromedarii* in India. *Indian J. Med. Res.* **1973**, *61*, 1416–1420.
117. Wood, O.L.; Moussa, M.I.; Hoogstraal, H.; Büttiker, W. Kadam Virus (Togaviridae, Flavivirus) Infecting Camel-Parasitizing *Hyalomma dromedarii* Ticks (Acari: Ixodidae) in Saudi Arabia. *J. Med. Entomol.* **1982**, *19*, 207–208. [[CrossRef](#)]
118. Awad, F.I.; Amin, M.M.; Salama, S.A.; Khide, S. The Role Played by *Hyalomma dromedarii* in the Transmission of African Horse Sickness Virus in Egypt. *Bull. Anim. Health Prod. Afr. Bull. Sante Prod. Anim. En Afr.* **1981**, *29*, 337–340.
119. Converse, J.D.; Moussa, M.I. Quarantfil Virus from *Hyalomma dromedarii* (Acari: Ixodoidea) Collected in Kuwait, Iraq and Yemen. *J. Med. Entomol.* **1982**, *19*, 209–210. [[CrossRef](#)]
120. Reháček, J.; Brezina, R. Detection of Coxiella Burneti in Saliva of Experimentally Infected Ticks, *Hyalomma dromedarii* Koch. *Bull. World Health Organ.* **1968**, *39*, 974–977.
121. Loftis, A.D.; Reeves, W.K.; Szumlas, D.E.; Abbassy, M.M.; Helmy, I.M.; Moriarity, J.R.; Dasch, G.A. Rickettsial Agents in Egyptian Ticks Collected from Domestic Animals. *Exp. Appl. Acarol.* **2006**, *40*, 67. [[CrossRef](#)]
122. Abdullah, H.H.A.M.; El-Shanawany, E.E.; Abdel-Shafy, S.; Abou-Zeina, H.A.A.; Abdel-Rahman, E.H. Molecular and Immunological Characterization of *Hyalomma dromedarii* and *Hyalomma excavatum* (Acari: Ixodidae) Vectors of Q Fever in Camels. *Vet. World* **2018**, *11*, 1109–1119. [[CrossRef](#)]
123. Selmi, R.; Ben Said, M.; Mamlouk, A.; Ben Yahia, H.; Messadi, L. Molecular Detection and Genetic Characterization of the Potentially Pathogenic *Coxiella burnetii* and the Endosymbiotic *Candidatus midichloria mitochondrii* in Ticks Infesting Camels (*Camelus dromedarius*) from Tunisia. *Microb. Pathog.* **2019**, *136*, 103655. [[CrossRef](#)] [[PubMed](#)]
124. Bellabidi, M.; Benaissa, M.H.; Bissati-Bouafia, S.; Harrat, Z.; Brahmi, K.; Kernif, T. *Coxiella burnetii* in Camels (*Camelus dromedarius*) from Algeria: Seroprevalence, Molecular Characterization, and Ticks (Acari: Ixodidae) Vectors. *Acta Trop.* **2020**, *206*, 105443. [[CrossRef](#)] [[PubMed](#)]
125. Ravi, A.; Ereqat, S.; Al-Jawabreh, A.; Abdeen, Z.; Abu Shamma, O.; Hall, H.; Pallen, M.J.; Nasereddin, A. Metagenomic Profiling of Ticks: Identification of Novel Rickettsial Genomes and Detection of Tick-Borne Canine Parvovirus. *PLoS Negl. Trop. Dis.* **2019**, *13*, e0006805. [[CrossRef](#)] [[PubMed](#)]
126. Morita, C.; El Hussein, A.R.M.; Matsuda, E.; Gabbar, K.M.A.A.; Muramatsu, Y.; Rahman, M.B.A.; Eleragi, A.M.H.; Hassan, S.M.; Chitambo, A.M.; Ueno, H. Spotted Fever Group *Rickettsiae* from Ticks Captured in Sudan. *Jpn. J. Infect. Dis.* **2004**, *57*, 107–109. [[PubMed](#)]
127. Demoncheaux, J.-P.; Socolovschi, C.; Davoust, B.; Haddad, S.; Raoult, D.; Parola, P. First Detection of *Rickettsia aeschlimannii* in *Hyalomma dromedarii* Ticks from Tunisia. *Ticks Tick-Borne Dis.* **2012**, *3*, 398–402. [[CrossRef](#)] [[PubMed](#)]
128. Kernif, T.; Djerbouh, A.; Mediannikov, O.; Ayach, B.; Rolain, J.-M.; Raoult, D.; Parola, P.; Bitam, I. *Rickettsia africae* in *Hyalomma dromedarii* Ticks from Sub-Saharan Algeria. *Ticks Tick-Borne Dis.* **2012**, *3*, 377–379. [[CrossRef](#)]
129. Ereqat, S.; Nasereddin, A.; Vayssier-Taussat, M.; Abdelkader, A.; Al-Jawabreh, A.; Zaid, T.; Azmi, K.; Abdeen, Z. Molecular Evidence of *Bartonella* Species in Ixodid Ticks and Domestic Animals in Palestine. *Front. Microbiol.* **2016**, *7*, 1217. [[CrossRef](#)]
130. Ros-García, A.; M'ghirbi, Y.; Hurtado, A.; Bouattour, A. Prevalence and Genetic Diversity of Piroplasm Species in Horses and Ticks from Tunisia. *Infect. Genet. Evol. J. Mol. Epidemiol. Evol. Genet. Infect. Dis.* **2013**, *17*, 33–37. [[CrossRef](#)]
131. Tirosh-Levy, S.; Gottlieb, Y.; Fry, L.M.; Knowles, D.P.; Steinman, A. Twenty Years of Equine Piroplasmosis Research: Global Distribution, Molecular Diagnosis, and Phylogeny. *Pathogens* **2020**, *9*, 926. [[CrossRef](#)]
132. Al-Hosary, A.; Răileanu, C.; Tauchmann, O.; Fischer, S.; Nijhof, A.M.; Silaghi, C. Tick Species Identification and Molecular Detection of Tick-Borne Pathogens in Blood and Ticks Collected from Cattle in Egypt. *Ticks Tick-Borne Dis.* **2021**, *12*, 101676. [[CrossRef](#)]
133. Hadani, A.; Tsur, I.; Pipano, E.; Zenft, Z. Studies on the Transmission of *Theileria annulata* by Ticks (*Ixodidea*, *Ixodidae*) I. *Hyalomma excavatum*. *J. Protozool.* **1963**, *10*, 35.
134. Aktas, M.; Özübek, S.; Altay, K.; Ipek, N.D.S.; Balkaya, İ.; Utuk, A.E.; Kırbas, A.; Şimsek, S.; Dumanlı, N. Molecular Detection of Tick-Borne Rickettsial and Protozoan Pathogens in Domestic Dogs from Turkey. *Parasit. Vectors* **2015**, *8*, 157. [[CrossRef](#)] [[PubMed](#)]
135. Hashemi-Fesharki, R. Tick-Borne Diseases of Sheep and Goats and Their Related Vectors in Iran. *Parassitologia* **1997**, *39*, 115–117.
136. Orkun, Ö.; Karaer, Z.; Çakmak, A.; Nalbantoğlu, S. Identification of Tick-Borne Pathogens in Ticks Feeding on Humans in Turkey. *PLoS Negl. Trop. Dis.* **2014**, *8*, e3067. [[CrossRef](#)] [[PubMed](#)]
137. Ioannou, I.; Sandalakis, V.; Kassinis, N.; Chochlakis, D.; Papadopoulos, B.; Loukaidis, F.; Tselentis, Y.; Psaroulaki, A. Tick-Borne Bacteria in Mouflons and Their Ectoparasites in Cyprus. *J. Wildl. Dis.* **2011**, *47*, 300–306. [[CrossRef](#)]
138. Abdelkadir, K.; Palomar, A.M.; Portillo, A.; Oteo, J.A.; Ait-Oudhia, K.; Khelef, D. Presence of *Rickettsia aeschlimannii*, “*Candidatus rickettsia barbariae*” and *Coxiella burnetii* in Ticks from Livestock in Northwestern Algeria. *Ticks Tick-Borne Dis.* **2019**, *10*, 924–928. [[CrossRef](#)]
139. Kilicoglu, Y.; Cagirgan, A.A.; Serdar, G.; Kaya, S.; Durmaz, Y.; Gur, Y. Molecular Investigation, Isolation and Phylogenetic Analysis of *Coxiella burnetii* from Aborted Fetus and Ticks. *Comp. Immunol. Microbiol. Infect. Dis.* **2020**, *73*, 101571. [[CrossRef](#)]
140. Kleinerman, G.; Baneth, G.; Mumcuoglu, K.; van Straten, M.; Berlin, D.; Apanaskevich, D.; Abdeen, Z.; Nasereddin, A.; Harrus, S. Molecular Detection of *Rickettsia africae*, *Rickettsia aeschlimannii*, and *Rickettsia sibirica mongolitimonae* in Camels and *Hyalomma* spp. Ticks from Israel. *Vector Borne Zoonotic Dis. Larchmt. N* **2013**, *13*, 851–856. [[CrossRef](#)]

141. Shkap, V.; Kocan, K.; Molad, T.; Mazuz, M.; Leibovich, B.; Krigel, Y.; Michoytchenko, A.; Blouin, E.; de la Fuente, J.; Samish, M.; et al. Experimental Transmission of Field *Anaplasma marginale* and the *A. centrale* Vaccine Strain by *Hyalomma Eecavatum*, *Rhipicephalus sanguineus* and *Rhipicephalus (Boophilus) annulatus* Ticks. *Vet. Microbiol.* **2009**, *134*, 254–260. [[CrossRef](#)]
142. Psaroulaki, A.; Ragiadakou, D.; Kouris, G.; Papadopoulos, B.; Chaniotis, B.; Tselentis, Y. Ticks, Tick-Borne *Rickettsiae*, and *Coxiella burnetii* in the Greek Island of Cephalonia. *Ann. N. Y. Acad. Sci.* **2006**, *1078*, 389–399. [[CrossRef](#)] [[PubMed](#)]
143. Gunes, T.; Poyraz, O.; Vatansever, Z. Crimean-Congo Hemorrhagic Fever Virus in Ticks Collected from Humans, Livestock, and Picnic Sites in the Hyperendemic Region of Turkey. *Vector-Borne Zoonotic Dis.* **2011**, *11*, 1411–1416. [[CrossRef](#)] [[PubMed](#)]
144. Padbidri, V.S.; Rodrigues, J.J.; Shetty, P.S.; Joshi, M.V.; Rao, B.L.; Shukla, R.N. Tick-Borne Rickettsioses in Pune District, Maharashtra, India. *Int. J. Zoonoses* **1984**, *11*, 45–52. [[PubMed](#)]
145. Mamman, A.H.; Lorusso, V.; Adam, B.M.; Dogo, G.A.; Bown, K.J.; Birtles, R.J. First report of *Theileria annulata* in Nigeria: Findings from cattle ticks in Zamfara and Sokoto States. *Parasites Vectors* **2021**, *14*, 242. [[CrossRef](#)] [[PubMed](#)]
146. El-Azazy, O.M.; El-Metenawy, T.M.; Wassef, H.Y. *Hyalomma impeltatum* (Acari: Ixodidae) as a Potential Vector of Malignant Theileriosis in Sheep in Saudi Arabia. *Vet. Parasitol.* **2001**, *99*, 305–309. [[CrossRef](#)]
147. Dohm, D.J.; Logan, T.M.; Linthicum, K.J.; Rossi, C.A.; Turell, M.J. Transmission of Crimean-Congo Hemorrhagic Fever Virus by *Hyalomma impeltatum* (Acari:Ixodidae) after Experimental Infection. *J. Med. Entomol.* **1996**, *33*, 848–851. [[CrossRef](#)]
148. Gordon, S.W.; Linthicum, K.J.; Moulton, J.R. Transmission of Crimean-Congo Hemorrhagic Fever Virus in Two Species of *Hyalomma* Ticks from Infected Adults to Cofeeding Immature Forms. *Am. J. Trop. Med. Hyg.* **1993**, *48*, 576–580. [[CrossRef](#)]
149. Wilson, M.L.; Gonzalez, J.P.; Cornet, J.P.; Camicas, J.L. Transmission of Crimean-Congo Haemorrhagic Fever Virus from Experimentally Infected Sheep to *Hyalomma truncatum* Ticks. *Res. Virol.* **1991**, *142*, 395–404. [[CrossRef](#)]
150. Sadeddine, R.; Diarra, A.Z.; Laroche, M.; Mediannikov, O.; Righi, S.; Benakhla, A.; Dahmana, H.; Raoult, D.; Parola, P. Molecular Identification of Protozoal and Bacterial Organisms in Domestic Animals and Their Infesting Ticks from North-Eastern Algeria. *Ticks Tick-Borne Dis.* **2020**, *11*, 101330. [[CrossRef](#)]
151. Parola, P.; Paddock, C.D.; Socolovschi, C.; Labruna, M.B.; Mediannikov, O.; Kernif, T.; Abdad, M.Y.; Stenos, J.; Bitam, I.; Fournier, P.-E.; et al. Update on Tick-Borne Rickettsioses around the World: A Geographic Approach. *Clin. Microbiol. Rev.* **2013**, *26*, 657–702. [[CrossRef](#)]
152. Ehounoud, C.B.; Yao, K.P.; Dahmani, M.; Achi, Y.L.; Amanzougaghene, N.; N'Douba, A.K.; N'Guessan, J.D.; Raoult, D.; Fenollar, F.; Mediannikov, O. Multiple Pathogens Including Potential New Species in Tick Vectors in Côte d'Ivoire. *PLoS Negl. Trop. Dis.* **2016**, *10*, e0004367. [[CrossRef](#)] [[PubMed](#)]
153. Singh, K.R.; Bhatt, P.N. Transmission of Kyasanur Forest Disease Virus by *Hyalomma marginatum isaaci*. *Indian J. Med. Res.* **1968**, *56*, 610–613. [[PubMed](#)]
154. Jouglin, M.; Fernández-De-Mera, I.G.; de La Cotte, N.; Ruiz-Fons, F.; Gortázar, C.; Moreau, E.; Bastian, S.; de La Fuente, J.; Malandrín, L. Isolation and Characterization of *Babesia pecorum* sp. Nov. from Farmed Red Deer (*Cervus elaphus*). *Vet. Res.* **2014**, *45*, 78. [[CrossRef](#)]
155. Viseras, J.; Hueli, L.E.; Adroher, F.J.; García-Fernández, P. Studies on the Transmission of *Theileria annulata* to Cattle by the Tick *Hyalomma lusitanicum*. *J. Vet. Med. Ser. B* **1999**, *46*, 505–509. [[CrossRef](#)] [[PubMed](#)]
156. Habela, M.; Rol, J.A.; Antón, J.M.; Peña, J.; Corchero, E.; van Ham, I.; Jongejan, E. Epidemiology of Mediterranean Theileriosis in Extremadura Region, Spain. *Parassitologia* **1999**, *41* (Suppl. S1), 47–51.
157. Moraga-Fernández, A.; Ruiz-Fons, F.; Habela, M.A.; Royo-Hernández, L.; Calero-Bernal, R.; Gortazar, C.; de la Fuente, J.; de Mera, I.G.F. Detection of New Crimean-Congo Haemorrhagic Fever Virus Genotypes in Ticks Feeding on Deer and Wild Boar, Spain. *Transbound. Emerg. Dis.* **2021**, *68*, 993–1000. [[CrossRef](#)]
158. Negredo, A.; Habela, M.Á.; Ramírez de Arellano, E.; Diez, F.; Lasala, F.; López, P.; Sarriá, A.; Labiod, N.; Calero-Bernal, R.; Arenas, M.; et al. Survey of Crimean-Congo Hemorrhagic Fever Enzootic Focus, Spain, 2011–2015. *Emerg. Infect. Dis.* **2019**, *25*, 1177–1184. [[CrossRef](#)]
159. Chisu, V.; Loi, F.; Foxi, C.; Chessa, G.; Masu, G.; Rolesu, S.; Masala, G. Coexistence of Tick-Borne Pathogens in Ticks Collected from Their Hosts in Sardinia: An Update. *Acta Parasitol.* **2020**, *65*, 999–1004. [[CrossRef](#)]
160. Toledo, A.; Olmeda, A.S.; Escudero, R.; Jado, I.; Valcárcel, F.; Casado-Nistal, M.A.; Rodríguez-Vargas, M.; Gil, H.; Anda, P. Tick-Borne Zoonotic Bacteria in Ticks Collected from Central Spain. *Am. J. Trop. Med. Hyg.* **2009**, *81*, 67–74. [[CrossRef](#)]
161. Milhano, N.; de Carvalho, I.L.; Alves, A.S.; Arroubé, S.; Soares, J.; Rodriguez, P.; Carolino, M.; Nuncio, M.S.; Piesman, J.; de Sousa, R. Coinfections of *Rickettsia slovaca* and *Rickettsia helvetica* with *Borrelia lusitaniae* in Ticks Collected in a Safari Park, Portugal. *Ticks Tick-Borne Dis.* **2010**, *1*, 172–177. [[CrossRef](#)]
162. Santos-Silva, M.M.; Vatansever, Z. *Hyalomma marginatum* Koch, 1844 (Figs. 139–141). In *Ticks of Europe and North Africa: A Guide to Species Identification*; Estrada-Peña, A., Mihalca, A.D., Petney, T.N., Eds.; Springer International Publishing: Cham, Switzerland, 2017; pp. 349–354; ISBN 978-3-319-63760-0.
163. Bolaños-Rivero, M.; Carranza-Rodríguez, C.; Rodríguez, N.F.; Gutiérrez, C.; Pérez-Arellano, J.-L. Detection of *Coxiella burnetii* DNA in Peridomestic and Wild Animals and Ticks in an Endemic Region (Canary Islands, Spain). *Vector Borne Zoonotic Dis. Larchmt.* **2017**, *17*, 630–634. [[CrossRef](#)] [[PubMed](#)]
164. González, J.; González, M.G.; Valcárcel, F.; Sánchez, M.; Martín-Hernández, R.; Tercero, J.M.; Olmeda, A.S. Transstadial Transmission from Nymph to Adult of *Coxiella burnetii* by Naturally Infected *Hyalomma lusitanicum*. *Pathogens* **2020**, *9*, E884. [[CrossRef](#)] [[PubMed](#)]

165. González, J.; González, M.G.; Valcárcel, F.; Sánchez, M.; Martín-Hernández, R.; Tercero, J.M.; Olmeda, A.S. Prevalence of *Coxiella burnetii* (Legionellales: Coxiellaceae) Infection Among Wildlife Species and the Tick *Hyalomma lusitanicum* (Acari: Ixodidae) in a Meso-Mediterranean Ecosystem. *J. Med. Entomol.* **2020**, *57*, 551–556. [[CrossRef](#)] [[PubMed](#)]
166. Ionita, M.; Mitrea, I.L.; Pfister, K.; Hamel, D.; Silaghi, C. Molecular Evidence for Bacterial and Protozoan Pathogens in Hard Ticks from Romania. *Vet. Parasitol.* **2013**, *196*, 71–76. [[CrossRef](#)] [[PubMed](#)]
167. Iori, A.; Gabrielli, S.; Calderini, P.; Moretti, A.; Pietrobelli, M.; Tampieri, M.P.; Galuppi, R.; Cancrini, G. Tick Reservoirs for Piroplasms in Central and Northern Italy. *Vet. Parasitol.* **2010**, *170*, 291–296. [[CrossRef](#)] [[PubMed](#)]
168. Boularias, G.; Azzag, N.; Galon, C.; Šimo, L.; Boulouis, H.-J.; Moutailler, S. High-Throughput Microfluidic Real-Time PCR for the Detection of Multiple Microorganisms in Ixodid Cattle Ticks in Northeast Algeria. *Pathogens* **2021**, *10*, 362. [[CrossRef](#)] [[PubMed](#)]
169. Köseoğlu, A.E.; Can, H.; Güvendi, M.; Erkunt Alak, S.; Kandemir, Ç.; Taşkın, T.; Demir, S.; Akgül, G.; Değirmenci Döşkaya, A.; Karakavuk, M.; et al. Molecular Investigation of Bacterial and Protozoal Pathogens in Ticks Collected from Different Hosts in Turkey. *Parasit. Vectors* **2021**, *14*, 270. [[CrossRef](#)]
170. Mohammadi, S.M.; Esmailnejad, B.; Jalilzadeh-Amin, G. Molecular Detection, Infection Rate and Vectors of *Theileria lestoquardi* in Goats from West Azerbaijan Province, Iran. *Vet. Res. Forum* **2017**, *8*, 139–144.
171. Razmi, G.R.; Naghibi, A.; Aslani, M.R.; Fathivand, M.; Dastjerdi, K. An epidemiological study on ovine babesiosis in the Mashhad suburb area, province of Khorasan, Iran. *Vet. Parasitol.* **2002**, *108*, 109–115. [[CrossRef](#)]
172. Toma, L.; Di Luca, M.; Mancini, F.; Severini, F.; Mariano, C.; Nicolai, G.; Laghezza Masci, V.; Ciervo, A.; Fausto, A.M.; Cacciò, S.M. Molecular Characterization of *Babesia* and *Theileria* Species in Ticks Collected in the Outskirt of Monte Romano, Lazio Region, Central Italy. *Ann. Ist. Super. Sanita* **2017**, *53*, 30–34. [[CrossRef](#)]
173. Aktas, M. A Survey of Ixodid Tick Species and Molecular Identification of Tick-Borne Pathogens. *Vet. Parasitol.* **2014**, *200*, 276–283. [[CrossRef](#)] [[PubMed](#)]
174. Karasartova, D.; Gureser, A.S.; Gokce, T.; Celebi, B.; Yapar, D.; Keskin, A.; Celik, S.; Ece, Y.; Erenler, A.K.; Usluca, S.; et al. Bacterial and Protozoal Pathogens Found in Ticks Collected from Humans in Corum Province of Turkey. *PLoS Negl. Trop. Dis.* **2018**, *12*, e0006395. [[CrossRef](#)] [[PubMed](#)]
175. Gray, J.S.; De Vos, A.J. Studies on a Bovine *Babesia* Transmitted by *Hyalomma marginatum rufipes* Koch, 1844. *Onderstepoort J. Vet. Res.* **1981**, *48*, 215–223.
176. Dipeolu, O.O.; Amoo, A. The Presence of Kinetes of a *Babesia* Species in the Haemolymph Smears of Engorged *Hyalomma* Ticks in Nigeria. *Vet. Parasitol.* **1984**, *17*, 41–46. [[CrossRef](#)]
177. Akyildiz, G.; Bente, D.; Keles, A.G.; Vatansver, Z.; Kar, S. High Prevalence and Different Genotypes of Crimean-Congo Hemorrhagic Fever Virus Genome in Questing Unfed Adult *Hyalomma marginatum* in Thrace, Turkey. *Ticks Tick-Borne Dis.* **2021**, *12*, 101622. [[CrossRef](#)]
178. Gergova, I.; Kunchev, M.; Kamarinchev, B. Crimean-Congo Hemorrhagic Fever Virus-Tick Survey in Endemic Areas in Bulgaria. *J. Med. Virol.* **2012**, *84*, 608–614. [[CrossRef](#)] [[PubMed](#)]
179. Gevorgyan, H.; Grigoryan, G.G.; Atoyian, H.A.; Rukhkyan, M.; Hakobyan, A.; Zakaryan, H.; Aghayan, S.A. Evidence of Crimean-Congo Haemorrhagic Fever Virus Occurrence in Ixodidae Ticks of Armenia. *J. Arthropod-Borne Dis.* **2019**, *13*, 9–16.
180. Hoogstraal, H. Review Article1: The Epidemiology of Tick-Borne Crimean-Congo Hemorrhagic Fever in Asia, Europe, and Africa23. *J. Med. Entomol.* **1979**, *15*, 307–417. [[CrossRef](#)]
181. Kondratenko, V.F. Importance of ixodid ticks in the transmission and preservation of the causative agent of Crimean hemorrhagic fever in foci of the infection. *Parazitologiya* **1976**, *10*, 297–302.
182. Kondratenko, V.; Blagoveschenskaya, N.; Butenko, A.; Vyshnivetskaya, L.; Zarubina, L.; Milyutin, V.; Kuchin, V.; Novikova, E.; Rabinovich, V.; Shevchenko, S.; et al. Results of virological investigation of ixodid ticks in Crimean hemorrhagic fever focus in Rostov Oblast. *Mater* **1970**, *9*, 29–35.
183. Levi, V.; Vasilenko, S. *Study of the Crimean Hemorrhagic Fever (CHF) Virus Transmission Mechanism in Hyalomma p. plumbeum Ticks*; National Agricultural Library: Beltsville, MD, USA, 1972; pp. 182–185.
184. Meissner, J.D.; Seregin, S.S.; Seregin, S.V.; Yakimenko, N.V.; Vyshemirskii, O.I.; Netesov, S.V.; Petrov, V.S. Complete L Segment Coding-Region Sequences of Crimean Congo Hemorrhagic Fever Virus Strains from the Russian Federation and Tajikistan. *Arch. Virol.* **2006**, *151*, 465–475. [[CrossRef](#)] [[PubMed](#)]
185. Smirnova, S.E.; Karavanov, A.S.; Zimina, I.; Sedova, A.G. The virus carriage of ticks, vectors of the causative agent of Crimean hemorrhagic fever. *Med. Parazitol.* **1991**, *1*, 32–34.
186. Yashina, L.; Petrova, I.; Seregin, S.; Vyshemirskii, O.; Lvov, D.; Aristova, V.; Kuhn, J.; Morzunov, S.; Gutorov, V.; Kuzina, I.; et al. Genetic Variability of Crimean-Congo Haemorrhagic Fever Virus in Russia and Central Asia. *J. Gen. Virol.* **2003**, *84*, 1199–1206. [[CrossRef](#)] [[PubMed](#)]
187. Yesilbag, K.; Aydin, L.; Dincer, E.; Alpay, G.; Girisgin, A.O.; Tuncer, P.; Ozkul, A. Tick Survey and Detection of Crimean-Congo Hemorrhagic Fever Virus in Tick Species from a Non-Endemic Area, South Marmara Region, Turkey. *Exp. Appl. Acarol.* **2013**, *60*, 253–261. [[CrossRef](#)] [[PubMed](#)]
188. Zgurskaya, G.N.; Berezin, V.; Smirnova, S.; Chumakov, M. Investigation of the question of Crimean hemorrhagic fever virus transmission and interepidemic survival in the tick *Hyalomma plumbeum plumbeum* Panzer. *Tr. Inst. Polio. Virus. Entsefalitov. Akad. Med. Nauk.* **1971**, *19*, 217–220.

189. Pascucci, I.; Di Domenico, M.; Capobianco Dondona, G.; Di Gennaro, A.; Polci, A.; Capobianco Dondona, A.; Mancuso, E.; Cammà, C.; Savini, G.; Cecere, J.G.; et al. Assessing the Role of Migratory Birds in the Introduction of Ticks and Tick-Borne Pathogens from African Countries: An Italian Experience. *Ticks Tick-Borne Dis.* **2019**, *10*, 101272. [[CrossRef](#)]
190. Pereira, A.; Figueira, L.; Nunes, M.; Esteves, A.; Cotão, A.J.; Vieira, M.L.; Maia, C.; Campino, L.; Parreira, R. Multiple Phlebovirus (*Bunyaviridae*) Genetic Groups Detected in *Rhipicephalus*, *Hyalomma* and *Dermacentor* Ticks from Southern Portugal. *Ticks Tick-Borne Dis.* **2017**, *8*, 45–52. [[CrossRef](#)]
191. Converse, J.D.; Hoogstraal, H.; Moussa, M.I.; Stek, M.; Kaiser, M.N. Bahig Virus (Tete Group) in Naturally- and Transovarially-Infected *Hyalomma marginatum* Ticks from Egypt and Italy. *Arch. Für Gesamte Virusforsch.* **1974**, *46*, 29–35. [[CrossRef](#)]
192. Al'khovskii, S.V.; L'vov, D.K.; Shchelkanov, M.I.; Shchetinin, A.M.; Deriabin, P.G.; L'vov, D.N.; L'vov, S.S.; Samokhvalov, E.I.; Gitel'man, A.K.; Botikov, A.G. Genetic characterization of the Batken virus (BKNV) (*Orthomyxoviridae*, Thogotovirus) isolated from the Ixodidae ticks *Hyalomma marginatum* Koch, 1844 and the mosquitoes *Aedes caspius* Pallas, 1771, as well as the *Culex hortensis* Ficalbi, 1889 in the Central Asia. *Vopr. Virusol.* **2014**, *59*, 33–37.
193. Hubálek, Z. Biogeography of Tick-Borne Bhanja Virus (*Bunyaviridae*) in Europe. *Interdiscip. Perspect. Infect. Dis.* **2009**, *2009*, 372691. [[CrossRef](#)]
194. Filipe, A.R.; Casals, J. Isolation of Dhori Virus from *Hyalomma marginatum* Ticks in Portugal. *Intervirology* **1979**, *11*, 124–127. [[CrossRef](#)] [[PubMed](#)]
195. Vakalova, E.V.; Butenko, A.M.; Vishnevskaya, T.V.; Dorofeeva, T.E.; Gitelman, A.K.; Kulikova, L.N.; Lvov, D.K.; Alkhovsky, S.V. Results of investigation of ticks in Volga river delta (Astrakhan region, 2017) for Crimean-Congo hemorrhagic fever virus (*Nairoviridae*, Orthonairovirus, CCHFV) and other tick-borne arboviruses. *Vopr. Virusol.* **2019**, *64*, 221–228. [[CrossRef](#)]
196. Yurchenko, O.O.; Dubina, D.O.; Vynograd, N.O.; Gonzalez, J.-P. Partial Characterization of Tick-Borne Encephalitis Virus Isolates from Ticks of Southern Ukraine. *Vector Borne Zoonotic Dis.* **2017**, *17*, 550–557. [[CrossRef](#)] [[PubMed](#)]
197. Dinçer, E.; Hacıoğlu, S.; Kar, S.; Emanet, N.; Brinkmann, A.; Nitsche, A.; Özkul, A.; Linton, Y.-M.; Ergünay, K. Survey and Characterization of Jingmen Tick Virus Variants. *Viruses* **2019**, *11*, 1071. [[CrossRef](#)] [[PubMed](#)]
198. Moussa, M.I.; Imam, I.Z.; Converse, J.D.; El-Karamany, R.M. Isolation of Matruh Virus from *Hyalomma marginatum* Ticks in Egypt. *J. Egypt. Public Health Assoc.* **1974**, *49*, 341–348.
199. Dandawate, C.N.; Shah, K.V.; D'Lima, L.V. Wanowrie Virus: A New Arbovirus Isolated from *Hyalomma marginatum* Isaaci. *Indian J. Med. Res.* **1970**, *58*, 985–989.
200. L'vov, D.N.; Dzharhenov, A.F.; Aristova, V.A.; Kovtunov, A.I.; Gromashevskii, V.L.; Vyshemirskii, O.I.; Galkina, I.V.; Larichev, V.F.; Butenko, A.M.; L'vov, D.K. The isolation of Dhori viruses (*Orthomyxoviridae*, Thogotovirus) and Crimean-Congo hemorrhagic fever virus (*Bunyaviridae*, Nairovirus) from the hare (*Lepus europaeus*) and its ticks *Hyalomma marginatum* in the middle zone of the Volga delta, Astrakhan region, 2001. *Vopr. Virusol.* **2002**, *47*, 32–36.
201. Formosinho, P.; Santos-Silva, M.M. Experimental Infection of *Hyalomma marginatum* Ticks with West Nile Virus. *Acta Virol.* **2006**, *50*, 175–180.
202. Kolodziejek, J.; Marinov, M.; Kiss, B.J.; Alexe, V.; Nowotny, N. The Complete Sequence of a West Nile Virus Lineage 2 Strain Detected in a *Hyalomma marginatum marginatum* Tick Collected from a Song Thrush (*Turdus philomelos*) in Eastern Romania in 2013 Revealed Closest Genetic Relationship to Strain Volgograd 2007. *PLoS ONE* **2014**, *9*, e109905. [[CrossRef](#)]
203. Beati, L.; Meskini, M.; Thiers, B.; Raoult, D. *Rickettsia aeschlimannii* Sp. Nov., a New Spotted Fever Group *Rickettsia* Associated with *Hyalomma marginatum* Ticks. *Int. J. Syst. Bacteriol.* **1997**, *47*, 548–554. [[CrossRef](#)]
204. Cicculi, V.; Capai, L.; Quilichini, Y.; Masse, S.; Fernández-Alvarez, A.; Minodier, L.; Bompard, P.; Charrel, R.; Falchi, A. Molecular Investigation of Tick-Borne Pathogens in Ixodid Ticks Infesting Domestic Animals (Cattle and Sheep) and Small Rodents (Black Rats) of Corsica, France. *Ticks Tick-Borne Dis.* **2019**, *10*, 606–613. [[CrossRef](#)] [[PubMed](#)]
205. Grech-Angelini, S.; Stachurski, F.; Vayssier-Taussat, M.; Devillers, E.; Casabianca, F.; Lancelot, R.; Uilenberg, G.; Moutailler, S. Tick-Borne Pathogens in Ticks (Acari: *Ixodidae*) Collected from Various Domestic and Wild Hosts in Corsica (France), a Mediterranean Island Environment. *Transbound. Emerg. Dis.* **2020**, *67*, 745–757. [[CrossRef](#)]
206. Cicculi, V.; de Lamballerie, X.; Charrel, R.; Falchi, A. First Molecular Detection of *Rickettsia africae* in a Tropical Bont Tick, *Amblyomma variegatum*, Collected in Corsica, France. *Exp. Appl. Acarol.* **2019**, *77*, 207–214. [[CrossRef](#)] [[PubMed](#)]
207. Sentausa, E.; El Karkouri, K.; Michelle, C.; Caputo, A.; Raoult, D.; Fournier, P.-E. Genome Sequence of *Rickettsia tamurae*, a Recently Detected Human Pathogen in Japan. *Genome Announc.* **2014**, *2*, e00838-14. [[CrossRef](#)] [[PubMed](#)]
208. Matsumoto, K.; Parola, P.; Brouqui, P.; Raoult, D. *Rickettsia aeschlimannii* in *Hyalomma* Ticks from Corsica. *Eur. J. Clin. Microbiol. Infect. Dis. Off. Publ. Eur. Soc. Clin. Microbiol.* **2004**, *23*, 732–734. [[CrossRef](#)]
209. Teshale, S.; Kumsa, B.; Menandro, M.L.; Cassini, R.; Martini, M. Anaplasma, Ehrlichia and Rickettsial Pathogens in Ixodid Ticks Infesting Cattle and Sheep in Western Oromia, Ethiopia. *Exp. Appl. Acarol.* **2016**, *70*, 231–237. [[CrossRef](#)]
210. Wallménius, K.; Barboutis, C.; Fransson, T.; Jaenson, T.G.T.; Lindgren, P.-E.; Nyström, F.; Olsen, B.; Salaneck, E.; Nilsson, K. Spotted Fever *Rickettsia* Species in *Hyalomma* and *Ixodes* Ticks Infesting Migratory Birds in the European Mediterranean Area. *Parasit. Vectors* **2014**, *7*, 318. [[CrossRef](#)]
211. Chisu, V.; Foxi, C.; Mannu, R.; Satta, G.; Masala, G. A Five-Year Survey of Tick Species and Identification of Tick-Borne Bacteria in Sardinia, Italy. *Ticks Tick-Borne Dis.* **2018**, *9*, 678–681. [[CrossRef](#)]

212. Toma, L.; Mancini, F.; Di Luca, M.; Cecere, J.G.; Bianchi, R.; Khoury, C.; Quarchioni, E.; Manzia, F.; Rezza, G.; Ciervo, A. Detection of Microbial Agents in Ticks Collected from Migratory Birds in Central Italy. *Vector Borne Zoonotic Dis. Larchmt.* **2014**, *14*, 199–205. [[CrossRef](#)]
213. Mancini, F.; Vescio, F.; Toma, L.; Di Luca, M.; Severini, F.; Cacciò, S.; Mariano, C.; Nicolai, G.; Laghezza Masci, V.; Fausto, A.; et al. Detection of Tick-Borne Pathogens in Ticks Collected in the Suburban Area of Monte Romano, Lazio Region, Central Italy. *Ann. Ist. Super. Sanita* **2019**, *55*, 143–150. [[CrossRef](#)]
214. Pilipenko, V.G.; Derevianchenko, K.I. Detection of *Hyalomma plumbeum* Panz infected with *Pasteurella tularensis* on *Lepus europaeus*. *J. Microbiol. Epidemiol. Immunobiol.* **1955**, *4*, 63–67.
215. Jongejan, F.; Morzaria, S.P.; Mustafa, O.E.H.; Latif, A.A. Infection Rates of *Theileria annulata* in the Salivary Glands of the Tick *Hyalomma marginatum rufipes*. *Vet. Parasitol.* **1983**, *13*, 121–126. [[CrossRef](#)] [[PubMed](#)]
216. Zeller, H.G.; Cornet, J.P.; Camicas, J.L. Crimean-Congo Haemorrhagic Fever Virus Infection in Birds: Field Investigations in Senegal. *Res. Virol.* **1994**, *145*, 105–109. [[CrossRef](#)] [[PubMed](#)]
217. Faye, O.; Cornet, J.P.; Camicas, J.L.; Fontenille, D.; Gonzalez, J.P. Transmission expérimentale du virus de la fièvre hémorragique de Crimée-Congo: Place de trois espèces vectrices dans les cycles de maintenance et de transmission au Sénégal. *Parasite* **1999**, *6*, 27–32. [[CrossRef](#)]
218. Swanepoel, R.; Struthers, J.K.; Shepherd, A.J.; McGillivray, G.M.; Nel, M.J.; Jupp, P.G. Crimean-Congo Hemorrhagic Fever in South Africa. *Am. J. Trop. Med. Hyg.* **1983**, *32*, 1407–1415. [[CrossRef](#)] [[PubMed](#)]
219. Zeller, H.G.; Cornet, J.P.; Diop, A.; Camicas, J.L. Crimean-Congo Hemorrhagic Fever in Ticks (Acari: Ixodidae) and Ruminants: Field Observations of an Epizootic in Bandia, Senegal (1989–1992). *J. Med. Entomol.* **1997**, *34*, 511–516. [[CrossRef](#)]
220. Shepherd, A.J.; Leman, P.A.; Swanepoel, R. Viremia and Antibody Response of Small African and Laboratory Animals to Crimean-Congo Hemorrhagic Fever Virus Infection. *Am. J. Trop. Med. Hyg.* **1989**, *40*, 541–547. [[CrossRef](#)]
221. Zeller, H.G.; Cornet, J.P.; Camicas, J.L. Experimental Transmission of Crimean-Congo Hemorrhagic Fever Virus by West African Wild Ground-Feeding Birds to *Hyalomma marginatum rufipes* Ticks. *Am. J. Trop. Med. Hyg.* **1994**, *50*, 676–681. [[CrossRef](#)]
222. Shepherd, A.J.; Swanepoel, R.; Shepherd, S.P.; Leman, P.A.; Mathee, O. Viraemic Transmission of Crimean-Congo Haemorrhagic Fever Virus to Ticks. *Epidemiol. Infect.* **1991**, *106*, 373–382. [[CrossRef](#)]
223. Mancuso, E.; Toma, L.; Polci, A.; d'Alessio, S.G.; Luca, M.D.; Orsini, M.; Domenico, M.D.; Marcacci, M.; Mancini, G.; Spina, F.; et al. Crimean-Congo Hemorrhagic Fever Virus Genome in Tick from Migratory Bird, Italy. *Emerg. Infect. Dis.* **2019**, *25*, 1418–1420. [[CrossRef](#)]
224. Okorie, T.G. Comparative Studies on the Vector Capacity of the Different Stages of *Amblyomma variegatum fabricius* and *Hyalomma rufipes* Koch for Congo Virus, after Intracoelemic Inoculation. *Vet. Parasitol.* **1991**, *38*, 215–223. [[CrossRef](#)] [[PubMed](#)]
225. Msimang, V.; Weyer, J.; Roux, C.L.; Kemp, A.; Burt, F.J.; Tempia, S.; Grobbelaar, A.; Moolla, N.; Rostal, M.K.; Bagge, W.; et al. Risk Factors Associated with Exposure to Crimean-Congo Haemorrhagic Fever Virus in Animal Workers and Cattle, and Molecular Detection in Ticks, South Africa. *PLoS Negl. Trop. Dis.* **2021**, *15*, e0009384. [[CrossRef](#)]
226. Okorie, T.G.; Fabiyi, A. The Multiplication of Dugbe Virus in the Ixodid Tick, *Hyalomma rufipes* Koch, after Experimental Infection. *Tropenmed. Parasitol.* **1979**, *30*, 439–442. [[PubMed](#)]
227. Hoffman, T.; Lindeborg, M.; Barboutis, C.; Erciyas-Yavuz, K.; Evander, M.; Fransson, T.; Figuerola, J.; Jaenson, T.G.T.; Kiat, Y.; Lindgren, P.-E.; et al. Alkhurma Hemorrhagic Fever Virus RNA in *Hyalomma rufipes* Ticks Infesting Migratory Birds, Europe and Asia Minor. *Emerg. Infect. Dis.* **2018**, *24*, 879–882. [[CrossRef](#)] [[PubMed](#)]
228. Luo, J.; Liu, M.-X.; Ren, Q.-Y.; Chen, Z.; Tian, Z.-C.; Hao, J.-W.; Wu, F.; Liu, X.-C.; Luo, J.-X.; Yin, H.; et al. Micropathogen Community Analysis in *Hyalomma rufipes* via High-Throughput Sequencing of Small RNAs. *Front. Cell. Infect. Microbiol.* **2017**, *7*, 374. [[CrossRef](#)]
229. Mathison, B.A.; Gerth, W.J.; Pritt, B.S.; Baugh, S. Introduction of the Exotic Tick *Hyalomma truncatum* on a Human with Travel to Ethiopia: A Case Report. *Ticks Tick-Borne Dis.* **2015**, *6*, 152–154. [[CrossRef](#)]
230. Dahmani, M.; Davoust, B.; Sambou, M.; Bassene, H.; Scandola, P.; Ameer, T.; Raoult, D.; Fenollar, F.; Mediannikov, O. Molecular Investigation and Phylogeny of Species of the *Anaplasmataceae* Infecting Animals and Ticks in Senegal. *Parasit. Vectors* **2019**, *12*, 495. [[CrossRef](#)]
231. Kumsa, B.; Socolovschi, C.; Almeras, L.; Raoult, D.; Parola, P. Occurrence and Genotyping of *Coxiella burnetii* in Ixodid Ticks in Oromia, Ethiopia. *Am. J. Trop. Med. Hyg.* **2015**, *93*, 1074–1081. [[CrossRef](#)]
232. Diarra, A.Z.; Almeras, L.; Laroche, M.; Berenger, J.-M.; Koné, A.K.; Bocoum, Z.; Dabo, A.; Doumbo, O.; Raoult, D.; Parola, P. Molecular and MALDI-TOF Identification of Ticks and Tick-Associated Bacteria in Mali. *PLoS Negl. Trop. Dis.* **2017**, *11*, e0005762. [[CrossRef](#)]
233. Rollins, R.E.; Schaper, S.; Kahlhofer, C.; Frangoulidis, D.; Strauß, A.F.; Cardinale, M.; Springer, A.; Strube, C.; Bakkes, D.K.; Becker, N.S.; et al. Ticks (Acari: Ixodidae) on birds migrating to the island of Ponza, Italy, and the tick-borne pathogens they carry. *Ticks Tick-borne Dis.* **2020**, *12*, 101590. [[CrossRef](#)]
234. Knuth, P.; Behn, P.; Schulze, P. *Experiments on Equine Piroplasmiasis (Biliary Fever) in 1917*; CABI: Wallingford, UK, 1918; pp. 241–264.
235. Petunin, F. *Hyalomma Scupense* P Sch-a Vector of Nuttalliasis of Horses. *Veterinariia* **1948**, *25*, 14.
236. Samish, M.; Pipano, E. Development of Infectivity in *Hyalomma detritum* (Schulze, 1919) Ticks Infected with *Theileria annulata* (Dchunowsky and Luhs, 1904). *Parasitology* **1978**, *77*, 375–379. [[CrossRef](#)] [[PubMed](#)]
237. Gharbi, M.; Darghouth, M.A. A Review of *Hyalomma scupense* (Acari, Ixodidae) in the Maghreb Region: From Biology to Control. *Parasite* **2014**, *21*, 2. [[CrossRef](#)] [[PubMed](#)]

238. Gharbi, M.; Hayouni, M.E.; Sassi, L.; Dridi, W.; Darghouth, M.A. *Hyalomma scupense* (Acari, Ixodidae) in Northeast Tunisia: Seasonal Population Dynamics of Nymphs and Adults on Field Cattle. *Parasite* **2013**, *20*, 12. [[CrossRef](#)]
239. Pandurov, S.; Zaprianov, M. Studies on the retention of *R. burneti* in *Rh. bursa* and *H. detritum* ticks. *Vet.-Meditsinski Nauki* **1975**, *12*, 43–48.
240. Blouin, E.F.; De Waal, D.T. The Fine Structure of Developmental Stages of *Babesia caballi* in the Salivary Glands of *Hyalomma truncatum*. *Onderstepoort J. Vet. Res.* **1989**, *56*, 189–193.
241. De Waal, D.T. The Transovarial Transmission of *Babesia caballi* by *Hyalomma truncatum*. *Onderstepoort J. Vet. Res.* **1990**, *57*, 99–100.
242. Logan, T.M.; Linthicum, K.J.; Kondig, J.P.; Bailey, C.L. Biology of *Hyalomma impeltatum* (Acari: Ixodidae) Under Laboratory Conditions. *J. Med. Entomol.* **1989**, *26*, 479–483. [[CrossRef](#)]
243. Wilson, M.L.; Gonzalez, J.-P.; LeGuanno, B.; Cornet, J.-P.; Guillaud, M.; Calvo, M.-A.; Digoutte, J.-P.; Camicas, J.-L. Epidemiology of Crimean-Congo Hemorrhagic Fever in Senegal: Temporal and Spatial Patterns. *Arch. Virol. Suppl.* **1990**, *1*, 323–340. [[CrossRef](#)]
244. Gonzalez, J.P.; Camicas, J.L.; Cornet, J.P.; Faye, O.; Wilson, M.L. Sexual and Transovarian Transmission of Crimean-Congo Haemorrhagic Fever Virus in *Hyalomma truncatum* Ticks. *Res. Virol.* **1992**, *143*, 23–28. [[CrossRef](#)]
245. Gonzalez, J.P.; Cornet, J.P.; Wilson, M.L.; Camicas, J.L. Crimean-Congo Haemorrhagic Fever Virus Replication in Adult *Hyalomma truncatum* and *Amblyomma variegatum* Ticks. *Res. Virol.* **1991**, *142*, 483–488. [[CrossRef](#)] [[PubMed](#)]
246. Dickson, D.L.; Turell, M.J. Replication and Tissue Tropisms of Crimean-Congo Hemorrhagic Fever Virus in Experimentally Infected Adult *Hyalomma truncatum* (Acari: Ixodidae). *J. Med. Entomol.* **1992**, *29*, 767–773. [[CrossRef](#)] [[PubMed](#)]
247. Lwande, O.W.; Lutomiah, J.; Obanda, V.; Gakuya, F.; Mutisya, J.; Mulwa, F.; Michuki, G.; Chepkorir, E.; Fischer, A.; Venter, M.; et al. Isolation of Tick and Mosquito-Borne Arboviruses from Ticks Sampled from Livestock and Wild Animal Hosts in Ijara District, Kenya. *Vector-Borne Zoonotic Dis.* **2013**, *13*, 637–642. [[CrossRef](#)] [[PubMed](#)]
248. Lutomiah, J.; Musila, L.; Makio, A.; Ochieng, C.; Koka, H.; Chepkorir, E.; Mutisya, J.; Mulwa, F.; Khamadi, S.; Miller, B.R.; et al. Ticks and Tick-Borne Viruses from Livestock Hosts in Arid and Semiarid Regions of the Eastern and Northeastern Parts of Kenya. *J. Med. Entomol.* **2014**, *51*, 269–277. [[CrossRef](#)] [[PubMed](#)]
249. Linthicum, K.J.; Logan, T.M. Laboratory Transmission of Venezuelan Equine Encephalomyelitis Virus by the Tick *Hyalomma truncatum*. *Trans. R. Soc. Trop. Med. Hyg.* **1994**, *88*, 126. [[CrossRef](#)]
250. Capponi, M.; Chambon, L.; Camicas, J.L.; Dumas, N. 1st isolation of a strain of *Rickettsia (Coxsackia) burneti* from ticks (*Hyalomma truncatum*) in Senegal. *Bull. Soc. Pathol. Exot. Filiales* **1970**, *63*, 530–534.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.