



HAL
open science

Honey bees change the microbiota of pollen

Alberto Prado, Matthieu Barret, Bernard Vaissière, Gloria Torres-Cortes

► **To cite this version:**

Alberto Prado, Matthieu Barret, Bernard Vaissière, Gloria Torres-Cortes. Honey bees change the microbiota of pollen. *Doklady Botanical Sciences*, 2023, 101 (1), pp.127-133. 10.17129/botsci.3125 . hal-04044122

HAL Id: hal-04044122

<https://hal.inrae.fr/hal-04044122>

Submitted on 24 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

HONEY BEES CHANGE THE MICROBIOTA OF POLLEN

LA ABEJA DOMÉSTICA CAMBIA LA MICROBIOTA DEL POLEN

 ALBERTO PRADO^{1*},  MATTHIEU BARRET², BERNARD E. VAISSIÈRE³, AND GLORIA TORRES-CORTES⁴

¹ Escuela Nacional de Estudios Superiores, Unidad Juriquilla, UNAM, Querétaro, México.

² Institut de Recherche en Horticulture et Semences, Agrocampus-Ouest, INRAE, Université d'Angers, Beaucauzé, France.

³ Institut national de recherche pour l'agriculture, l'alimentation et l'environnement, Avignon, France.

⁴ Kimitec, Almería, España.

*Autor para correspondencia: aprado@unam.mx

Abstract

Background: Pollen, as all other plant tissues, harbors different microorganisms. As honey bees (*Apis mellifera*) collect and pack pollen they add regurgitated nectar to moisten and glue the pollen grains, possibly changing the microbial composition.

Hypothesis: We hypothesize that while packing pollen into their corbiculae, honey bees change the pollen microbiota.

Studied species: *Brassica napus* L., *Apis mellifera* L.

Study site and dates: Avignon, France, 2018.

Methods: In this study, we compared the microbiota of clean *Brassica napus* (Brassicaceae) pollen with that of bee-worked corbicular pollen by 16S rRNA gene targeted amplicon sequencing. We also compared these bacterial communities with those present in nectar, the bee gut and the bee surface as potential inoculum sources.

Results: We found that by working pollen, bees increase the bacterial diversity of pollen, by adding honey bee symbionts such as *Bombella*, *Frischella*, *Gilliamella* and *Snodgrassella*, bee pathogens as *Spiroplasma* and nectar dwelling *Lactobacillus* to the new pollen microbiota.

Conclusions: The bee gut is an important source of inoculum of the corbicular pollen microbiota. We discuss the implications of these findings and propose future research avenues.

Keywords: *Apis mellifera*, corbicular pollen, symbionts.

Resumen

Antecedentes: Así como raíces, tallos, hojas y flores, el polen alberga una diversidad de microorganismos. Al momento de recolectar y trasladar el polen a las corbículas, las abejas domésticas (*Apis mellifera*), agregan néctar regurgitado para humidificar y pegar los granos de polen, posiblemente cambiando su composición microbiana.

Hipótesis: Al empaquetar el polen en las corbículas, las abejas cambian la microbiota del polen.

Especies de estudio: *Brassica napus* L., *Apis mellifera* L.

Sitio y años de estudio: Aviñón, Francia, 2018.

Métodos: En este estudio se empleó la secuenciación dirigida del gen 16S rRNA para comparar la microbiota del polen de *Brassica napus* (Brassicaceae) colectado directamente de las anteras, con el polen de las corbículas de abejas trabajadoras. Así mismo, se contrastaron estas dos comunidades bacterianas con las presentes en el néctar, el tracto digestivo y la superficie de la abeja, como posibles fuentes de inóculo.

Resultados: Al coleccionar y trabajar el polen, las abejas incrementan la diversidad bacteriana de este recurso floral. Simbiontes de la abeja como *Bombella*, *Frischella*, *Gilliamella* y *Snodgrassella*, así como patógenos como *Spiroplasma* y bacterias del néctar como *Lactobacillus*, son parte de esta nueva comunidad microbiana del polen.

Conclusiones: El tracto digestivo de la abeja es un inóculo importante de la comunidad del polen presente en la corbícula. En este artículo se discuten las implicaciones de estos resultados y se proponen algunas nuevas líneas de investigación.

Palabras clave: *Apis mellifera*, polen corbicular, simbiontes.

Insect visitors shape floral microbiota by acquiring and depositing microorganisms onto flower surfaces during nectar and pollen collection (Keller *et al.* 2021). The flower microbiota can have important effects on plant (de Vega & Herrera 2012) and pollinator fitness (Adler *et al.* 2018). Therefore, it is crucial to better understand how insects influence floral microbial communities. Honey bees are known to introduce bacteria into the nectar while foraging (Aizenberg-Gershtein *et al.* 2013). In turn, the bacteria vectored to the flower change the chemistry of nectar and affect pollinator behavior and plant reproduction (Vannette *et al.* 2013, Junker *et al.* 2014, Rering *et al.* 2018, Vannette & Fukami 2018). However, less is known about the effects of honey bee foraging on the microbiota of pollen. As a reproductive structure, pollen is of great importance for the biological success of plants because it is directly linked to reproductive output (i.e., fitness). The few studies that have focused on the microbiota of pollen have found complex bacterial communities with a high level of species-specificity (Heydenreich *et al.* 2012, Manirajan *et al.* 2016). The highly sculptured exine surface of pollen grains present cavities filled with a heterogeneous coating of lipid and proteinaceous components that provide a unique habitat for bacteria (Heslop-Harrison & Heslop-Harrison 1985, Aleklett *et al.* 2014). This pollen coat is thicker in insect pollinated plants which could be a particular attractive habitat for microbes (Edlund *et al.* 2004, Aleklett *et al.* 2014). Bacteria on the pollen surface can be found alone or in small colonies forming small biofilms (Zasloff 2017). An interesting aspect of these pollen microbial communities is that they can potentially modulate pollen grain germination on the stigmatic surface (Eisikowitch *et al.* 1990, Christensen *et al.* 2021) and hence influence plant fitness.

The dominant bacterial phyla reported in pollen are Acidobacteria, Actinobacteria, Firmicutes and Proteobacteria (Heydenreich *et al.* 2012, Manirajan *et al.* 2016), at this taxonomic rank no pollen signature can be observed compared to other habitats of the plant. However, at a finer taxonomic rank, differences in microbiota composition can be found between insect- and wind-pollinated tree species, with more similar communities for insect-pollinated species (Manirajan *et al.* 2016). These findings suggest insect vectors affect the microbial composition of pollen by homogenizing the microbial communities across plant species.

Honey bees (*Apis mellifera* L.), one of the most economically important pollinators, harbour a distinct and taxonomically restricted microbiota (Cox-Foster *et al.* 2007, Martinson *et al.* 2011) that includes members of the Orbaceae family, which are all exclusively honey bee and bumble bee symbionts (Engel *et al.* 2013, Kwong & Moran 2013, Li *et al.* 2015). The sociality of *Apis* has been proposed as the key driver in symbiont transmission and the maintenance of a consistent gut microbiota (Martinson *et al.* 2011). Despite the restricted number of species identified in the honey bee gut, symbionts possess distinct functional capabilities linked to host interaction, biofilm formation, and carbohydrate breakdown (Engel *et al.* 2012). The genera previously detected in worker honey bees are *Bartonella*, *Bifidobacterium*, *Bombella*, *Frischella*, *Gilliamella*, *Gluconacetobacter*, *Lactobacillus*, *Serratia*, *Simonsiella* and *Snodgrassella* (Cox-Foster *et al.* 2007, Martinson *et al.* 2011, Engel *et al.* 2012, 2013, Li *et al.* 2015). The distinctive microbiota of honey bees provides an interesting framework to test the effect of pollinators on the pollen microbiota. As an initial approach towards the understanding of how pollinators influence the pollen microbiota, in this study we estimated the composition of pollen-associated bacterial communities of *Brassica napus* before and after it has been collected by the domestic honey bee.

Materials and methods

Detailed methods can be found in Prado *et al.* (2020). Inside an insect proof tunnel *Brassica napus* L. were sown in the soil. Once the plants were in flower and before they had any contact with insects, clean pollen and nectar were collected. Pollen was collected by dissecting closed flower buds and separating the anthers. Anthers were left to dehisce for 4 h at room temperature in glass Petri dishes. To recover the pollen, the dried anthers were placed in a sterile steel tea ball and vibrated using a Vibri Vario pollinator. A total of three samples of anther collected pollen were recovered. Nectar was collected from the flowers with 2 µl capillary tubes between the base of the anthers and transferred to 2 ml Eppendorf tubes (N = 11). Afterward, a small five-frame honey bee (*Apis mellifera* L.) hive was introduced into the insect-proof tunnel, and honey bee foragers were allowed to forage freely on the *B. napus* flow-

ers. Inside the tunnel *B. napus* flowers were the only resource available for bees to forage. Honey bee foragers were captured with a sweeping net and their pollen loads were removed and frozen for further analysis (N = 3 corbicular pollen). Honey bee foragers were sampled to obtain surface and gut microbial assemblages. Entire bees were sonicated in 1 ml of PBS buffer with 0.05 % Tween 20. After removing the liquid (bee surface samples, N = 3), insects were re-suspended in 1 ml of PBS and crushed to recover the microbes inside the bees' gut (gut samples, N = 7). All the suspensions were centrifuged (12,000xg, 20 min, 4 °C) and pellets were stored at -20 °C until DNA extraction.

Total DNA extraction of pollen, corbicular pollen, nectar, bee washes and bee homogenates were performed with the PowerSoil DNA isolation kit (MoBio Laboratories) using the manufacturer's protocol. Amplification, purification, and pooling for amplicon library construction were conducted following the protocol described in Barret *et al.* (2015). PCRs were performed with a high-fidelity polymerase (AccuPrime *Taq* DNA polymerase; Invitrogen) using the manufacturer's protocol and 10 µl of the DNA suspension targeting the V4 of the 16S rRNA gene with the PCR primers 515f/806r. After amplicon purification, a second round of amplification was performed with 5 µl of purified amplicons and primers containing the Illumina adaptors and indexes. All amplicons were purified, quantified, and pooled in equimolar concentrations. Finally, amplicons libraries were mixed with 10 % PhiX control according to Illumina's protocols. Sequencing runs were performed in this study with MiSeq reagent kit V3 (600 cycles).

Data Analysis. Primers sequences of fastq files were first cut off using Cutadapt 1.8. Files were then merged and processed with DADA2 v. 1.8.0 according to the recommendations of the workflow "DADA2 Pipeline Tutorial". The workflow was modified in the truncLen parameter to adjust it to the quality of the sequencing run. The 16S rRNA amplicon sequence variants (ASV) resulting from DADA2 were aligned with a naive Bayesian classifier against the Ribosomal Database Project training set 16 database. Data analyses and visualization were done with RStudio v. 3.3 using the R package *phyloseq* v. 1.24.2.

Results

Microbial assemblages of anther collected pollen consisted of Proteobacteria (42 %), Actinobacteria (42 %), Firmicutes (10 %), Acidobacteria (4 %) and Bacteroidetes (2 %). These microbial communities were mainly characterised by the Families Enterobacteriaceae, Pseudomonadaceae (Proteobacteria) and Microbacteriaceae (Actinobacteria) (Figure 1A). While collecting and packing pollen in their corbiculae, honey bees changed the relative proportion of Bacteria phyla and their overall diversity. Corbicular pollen samples contained Proteobacteria (40 %), Firmicutes (21 %), Actinobacteria (19.7 %), Bacteroidetes (9.8 %), Fusobacteria (2.8 %), Chloroflexi (2.8 %), Armatimonadetes (1.4 %) and Tenericutes (1.4 %). The change in the microbial assemblages of pollen was partly done by incorporating bee gut-associated symbionts in the Families Orbaceae and Neisseriaceae (Proteobacteria), bee pathogens in the Family Spiroplasmataceae (Tenericutes), nectar-associated taxa in the Families Sphingomonadaceae (Proteobacteria) and Lactobacillaceae (Firmicutes) (Figure 1A). The abundance of Families Enterobacteriaceae (Proteobacteria) and Microbacteriaceae (Actinobacteria) was reduced in corbicular pollen as compared to anther collected pollen. Bee-gut symbionts such as *Bombella*, *Frischella*, *Gilliamella* and *Snodgrassella* were completely absent in anther collected pollen but present in corbicular pollen (Figure 1B). ASVs belonging to the genus *Lactobacillus* present in nectar were also incorporated into pollen samples by bees. Genera such as *Escherichia/Shigella* and *Pseudomonas* that were initially abundant in pollen were reduced in corbicular pollen (Figure 1C).

The microbial assemblages in corbicular pollen were richer than those present in the anther collected pollen with an average of 38 and 15.7 ASVs, respectively (Figure 2A). This enrichment represented 38 additional genera and 60 ASVs, and translated into a higher diversity as expressed by the Shannon index and Faith's phylogenetic diversity (Figure 2B-C). Of the 71 ASVs present in the corbicular pollen 16.9 % are only shared with the bee gut while 7 % are only shared with the bee surface, suggesting the potential inoculum sources. The highest richness was observed in nectar and on the honey bee surface, with averages of 54.2 and 53.3 ASVs, respectively. The honey bee gut presented a simpler microbial community with an average of only 19.7 ASVs.

Honey bees change the microbiota of pollen




Figure 1. Changes in the microbial composition of pollen. A) Relative abundance of bacterial families. Only showing Families with more than 3 % relative abundance. B) Average number of reads in pollen samples for bee associated taxa. Honeybees introduce *Bombella*, *Frischella*, *Gilliamella* and *Snodgrassella* into the pollen. C) Average number of reads in pollen samples for genera that strongly decreased due to honey bee processing.




Figure 2. Microbial richness and alpha diversity. (A) Observed richness, (B) Shannon diversity index and (C) Faith's PD phylogenetic diversity.

Discussion

The flower microbiota plays a crucial role on plant (de Vega & Herrera 2012) and pollinator fitness (Adler *et al.* 2018). Therefore, it is essential to understand the ecological processes involved in their assembly. This note focuses on the effect of honey bees on the pollen microbiota. Honey bees aggregate pollen grains by adding regurgitated nectar or diluted honey to transport it on the corbiculae of their hind legs. Unsurprisingly, we found that honey bees add bee and nectar-associated taxa to the pollen microbiota. Nonetheless, these effects could have important implications for plant reproduction and bee health. The pollen microbiota can have both deleterious and beneficial effects on pollen grain germination on the stigmatic surface. The ubiquitous nectar yeast *Metschnikowia reukauffi* inhibits pollen germination of *Asclepias syriaca* (Eisikowitch *et al.* 1990), while the nectar dwelling *Acinetobacter* can induce pollen germination of *Eschscholzia californica* (Christensen *et al.* 2021). By changing the pollen microbiota honey bees could potentially modulate pollen germination and hence plant fitness. However, pollen packed in the corbiculae is no longer available for pollination; only pollen grains that remain on other bee surfaces can be deposited onto the stigma. It is unknown if the pollen that remains on other bee surfaces and later deposited onto the stigma will suffer similar shifts in its microbial assembly as observed in this study. Nonetheless, honey bees have been shown to change the microbiota of seeds issued from bee pollination by increasing the variability of the microbial communities, and by introducing bee-associated taxa (Prado *et al.* 2020) suggesting a similar shift in the microbial assembly in the pollen deposited to the stigma. Here we show that while collecting pollen bees increase the diversity of pollen bacteria by incorporating nectar and gut-associated taxa. Hence, it is possible that the bee contribution to the seed microbiota is a consequence of how honey bees process pollen while foraging. Furthermore, pollinator behaviour (*i.e.*, pollen vs nectar foraging) has been shown to significantly affect the acquisition and deposition of bacteria on to flower surfaces (Russell *et al.* 2019). If we want to understand better the effects of insect pollinators on the plant microbiota, it is crucial to understand the mechanisms involved in the acquisition and deposition of bacteria while insects forage on flowers. Particular attention should be paid to the bacteria vectored to pollen grains and the stigmatic surfaces.

Bee pollen provisions are not consumed immediately after collection; bee larvae usually consume pollen that has been aged. The main effects of pollen ageing are an increase in lactic acid due to the presence of lactic acid bacteria and hence a reduction in pH as well as the opening of the outer shells of the pollen grains (Lipinski 2018). During this process, the microbial diversity in pollen provisions of solitary and social bees changes (Lozo *et al.* 2015, Graystock *et al.* 2017). Microbes present in aged pollen provisions are central to bee health as a major dietary component (Dharampal *et al.* 2019) but also as nutritional mutualists allowing bees to digest some of the recalcitrant components of the pollen wall (Engel *et al.* 2012). Honey bees need to acquire their microbial symbionts twice during their lives, once as larvae and once as adults. Microorganisms acquired by honey bee larvae through food are eliminated during the single defecation prior to pupation. Hence worker bees emerging from the cells are usually devoid of microorganisms and re-acquire their microbiota through food and trophallaxis (Gilliam 1997). Alterations to the adult honey bee gut microbiota can lead to parasite and pathogen prevalence (Schwarz *et al.* 2016). Therefore, the presence of bacterial symbionts in the Orbaceae, such as *Frischella* and *Gilliamella*, and the Neisseriaceae, such as *Snodgrassella*, in the pollen provisions is of great importance for honey bees (Engel *et al.* 2012). Our analysis suggests the bee gut is an important source of inoculum of the corbicular pollen microbiota. In conclusion, the way honey bees forage for pollen affects the pollen microbiota with potential consequences on plant reproduction and honey bee health.

Acknowledgements

We thank Cédric Alaux for his support in planning this experiment, Martial Briand for his help with DADA2 and other bioinformatic analyses. We thank Odile Vilotte, Gilles Aumont and the Agreenskills postdoctoral program for supporting this project. This work was supported by the AgreenSkills+ fellowship program, which received funding from the EU's Seventh Framework Program under grant agreement n° FP7-609398.

Literature cited

- Adler LS, Michaud KM, Ellner SP, McArt SH, Stevenson PC, Irwin RE. 2018. Disease where you dine: Plant species and floral traits associated with pathogen transmission in bumble bees. *Ecology* **99**: 2535-2545. DOI: <https://doi.org/10.1002/ecy.2503>
- Aizenberg-Gershtein Y, Izhaki I, Halpern M. 2013. Do honeybees shape the bacterial community composition in floral nectar? *Plos One* **8**: e67556. DOI: <https://doi.org/10.1371/journal.pone.0067556>
- Aleklett K, Hart M, Shade A. 2014. The microbial ecology of flowers: an emerging frontier in phyllosphere research. *Botany* **92**: 253-266. DOI: <https://doi.org/10.1139/cjb-2013-0166>
- Barret M, Briand M, Bonneau S, Prévieux A, Valière S, Bouchez O, Hunault G, Simoneau P, Jacques MA. 2015. Emergence shapes the structure of the seed microbiota. *Applied and Environmental Microbiology* **81**: 1257-1266. DOI: <https://doi.org/10.1128/AEM.03722-14>
- Christensen SM, Munkres I, Vannette RL. 2021. Nectar bacteria stimulate pollen germination and bursting to enhance microbial fitness. *Current Biology* **31**: 4373-4380. DOI: <https://doi.org/10.1016/j.cub.2021.07.016>
- Cox-Foster DL, Conlan S, Holmes EC, Palacios G, Evans JD, Moran NA, Quan PL, Briese T, Hornig M, Geiser DM. 2007. A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* **318**: 283-287. DOI: <https://doi.org/10.1126/science.1146498>
- de Vega C, Herrera CM. 2012. Relationships among nectar-dwelling yeasts, flowers and ants: patterns and incidence on nectar traits. *Oikos* **121**: 1878-1888. DOI: <https://doi.org/10.1111/j.1600-0706.2012.20295.x>
- Dharampal PS, Carlson C, Currie CR, Steffan SA. 2019. Pollen-borne microbes shape bee fitness. *Proceedings of the Royal Society B*, **286**: 20182894. DOI: <https://doi.org/10.1098/rspb.2018.2894>
- Edlund AF, Swanson R, Preuss D. 2004. Pollen and Stigma Structure and Function: The Role of Diversity in Pollination. *The Plant Cell* **16**: S84-S97. DOI: <https://doi.org/10.1105/tpc.015800>
- Eisikowitch D, Lachance MA, Kevan PG, Willis S, Collins-Thompson DL. 1990. The effect of the natural assemblage of microorganisms and selected strains of the yeast *Metschnikowia reukaufii* in controlling the germination of pollen of the common milkweed *Asclepias syriaca*. *Canadian Journal of Botany* **68**: 1163-1165. DOI: <https://doi.org/10.1139/b90-147>
- Engel P, Martinson VG, Moran NA. 2012. Functional diversity within the simple gut microbiota of the honey bee. *Proceedings of the National Academy of Sciences* **109**: 11002-11007. DOI: <https://doi.org/10.1073/pnas.1202970109>
- Engel P, Kwong WK, Moran NA. 2013. *Frischella perrara* gen. nov., sp. nov., a gammaproteobacterium isolated from the gut of the honeybee, *Apis mellifera*. *International Journal of Systematic and Evolutionary Microbiology* **63**: 3646-3651. DOI: <https://doi.org/10.1099/ijs.0.049569-0>
- Gilliam M. 1997. Identification and roles of non-pathogenic microflora associated with honey bees. *FEMS Microbiology Letters* **155**: 1-10. DOI: <https://doi.org/10.1111/j.1574-6968.1997.tb12678.x>
- Graystock P, Rehan SM, McFrederick QS. 2017. Hunting for healthy microbiomes: determining the core microbiomes of *Ceratina*, *Megalopta*, and *Apis* bees and how they associate with microbes in bee collected pollen. *Conservation Genetics* **18**: 701-711. DOI: <https://doi.org/10.1007/s10592-017-0937-7>
- Heslop-Harrison J, Heslop-Harrison Y. 1985. Surfaces and Secretions in the Pollen–Stigma Interaction: A Brief Review. *Journal of Cell Science* (Supplement 2): 287-300. DOI: https://doi.org/10.1242/jcs.1985.Supplement_2.15
- Heydenreich B, Bellinghausen I, König B, Becker WM, Grabbe S, Petersen A, Saloga J. 2012. Gram-positive bacteria on grass pollen exhibit adjuvant activity inducing inflammatory T cell responses. *Clinical and Experimental Allergy* **42**: 76-84. DOI: <https://doi.org/10.1111/j.1365-2222.2011.03888.x>
- Junker RR, Romeike T, Keller A, Langen D. 2014. Density-dependent negative responses by bumblebees to bacteria isolated from flowers. *Apidologie* **45**: 467-477. DOI: <https://doi.org/10.1007/s13592-013-0262-1>
- Keller A, McFrederick QS, Dharampal P, Steffan S, Danforth BN, Leonhardt SD. 2021. (More than) Hitchhikers through the network: The shared microbiome of bees and flowers. *Current Opinion in Insect Science* **44**: 8-15. DOI: <https://doi.org/10.1016/j.cois.2020.09.007>
- Kwong WK, Moran NA. 2013. Cultivation and characterization of the gut symbionts of honey bees and bumble bees: description of *Snodgrassella alvi* gen. nov., sp. nov., a member of the family Neisseriaceae of the Betaproteobacteria, and

- Gilliamella apicola* gen. nov., sp. nov., a member of Orbaceae fam. nov., Orbales ord. nov., a sister taxon to the order 'Enterobacteriales' of the Gammaproteobacteria. *International Journal of Systematic and Evolutionary Microbiology* **63**: 2008-2018. DOI: <https://doi.org/10.1099/ijs.0.044875-0>
- Lipinski Z. 2018. *Honey bee nutrition and feeding*. United Kingdom: Eva Crane Trust. ISBN: 9788393927906
- Lozo J, Berić T, Terzić-Vidojević A, Stanković S, Fira D, Stanisavljević L. 2015. Microbiota associated with pollen, bee bread, larvae and adults of solitary bee *Osmia cornuta* (Hymenoptera: Megachilidae). *Bulletin of Entomological Research* **105**: 470-476. DOI: <https://doi.org/10.1017/S0007485315000292>
- Li L, Praet J, Borremans W, Nunes OC, Manaia CM, Cleenwerck I, Meeus I, Smagghe G, De Vuyst L, Vandamme P. 2015. *Bombella intestini* gen. nov., sp. nov., an acetic acid bacterium isolated from bumble bee crop. *International Journal of Systematic and Evolutionary Microbiology* **65**: 267-273. DOI: <https://doi.org/10.1099/ijs.0.068049-0>
- Manirajan BA, Ratering S, Rusch V, Schwiertz A, Geissler-Plaum R, Cardinale M, Schnell S. 2016. Bacterial microbiota associated with flower pollen is influenced by pollination type, and shows a high degree of diversity and species-specificity. *Environmental Microbiology* **18**: 5161-5174. DOI: <https://doi.org/10.1111/1462-2920.13524>
- Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran NA. 2011. A simple and distinctive microbiota associated with honey bees and bumble bees. *Molecular Ecology* **20**: 619-628. DOI: <https://doi.org/10.1111/j.1365-294X.2010.04959.x>
- Prado A, Marolleau B, Vaissière BE, Barret M, Torres-Cortes G. 2020. Insect pollination: an ecological process involved in the assembly of the seed microbiota. *Scientific Reports* **10**: 3575. DOI: <https://doi.org/10.1038/s41598-020-60591-5>
- Rering CC, Beck JJ, Hall GW, McCartney MM, Vannette RL. 2018. Nectar-inhabiting microorganisms influence nectar volatile composition and attractiveness to a generalist pollinator. *New Phytologist* **220**: 750-759. DOI: <https://doi.org/10.1111/nph.14809>
- Russell AL, Rebolledo-Gómez M, Shaible TM, Ashman TL. 2019. Movers and shakers: Bumble bee foraging behavior shapes the dispersal of microbes among and within flowers. *Ecosphere* **10**: e02714. DOI: <https://doi.org/10.1002/ecs2.2714>
- Schwarz RS, Moran NA, Evans JD. 2016. Early gut colonizers shape parasite susceptibility and microbiota composition in honey bee workers. *Proceedings of the National Academy of Sciences* **113**: 9345-9350. DOI: <https://doi.org/10.1073/pnas.1606631113>
- Vannette RL, Fukami T. 2018. Contrasting effects of yeasts and bacteria on floral nectar traits. *Annals of Botany* **121**: 1343-1349. DOI: <https://doi.org/10.1093/aob/mcy032>
- Vannette RL, Gauthier MPL, Fukami T. 2013. Nectar bacteria, but not yeast, weaken a plant-pollinator mutualism. *Proceedings of the Royal Society B: Biological Sciences* **280**: 20122601. DOI: <https://doi.org/10.1098/rspb.2012.2601>
- Zasloff M. 2017. Pollen has a microbiome: implications for plant reproduction, insect pollination and human allergies. *Environmental Microbiology* **19**: 1-2. DOI: <https://doi.org/10.1111/1462-2920.13661>

Associate editor: Enrique Jurado

Author contributions: AP, conceptualization, field work, analysis, writing; MB, text revision; BEV, conceptualization, text revision; GTC, conceptualization, lab work, writing.