

# Composting date palm residues promotes circular agriculture in oases

Mustapha El Janati, Paul Robin, Nouraya Akkal-Corfini, Ahmed Bouaziz, Ahmed Sabri, Mohammed Chikhaoui, Zahra Thomas, Abdallah Oukarroum

▶ **To cite this version:**

Mustapha El Janati, Paul Robin, Nouraya Akkal-Corfini, Ahmed Bouaziz, Ahmed Sabri, et al.. Composting date palm residues promotes circular agriculture in oases. Biomass Conversion and Biorefinery, 2022, 10.1007/s13399-022-03387-z . hal-04051151

**HAL Id: hal-04051151**
**https://hal.inrae.fr/hal-04051151**

Submitted on 30 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biomass Conversion and Biorefinery
https://doi.org/10.1007/s13399-022-03387-z

ORIGINAL ARTICLE

# Composting date palm residues promotes circular agriculture in oases

**Mustapha El Janati[1,2,3] · Paul Robin[2] · Nouraya Akkal-Corfini[2] · Ahmed Bouaziz[1] · Ahmed Sabri[4] · Mohammed Chikhaoui[5] · Zahra Thomas[2] · Abdallah Oukarroum[3]**

Received: 14 June 2022 / Revised: 30 September 2022 / Accepted: 9 October 2022
© The Author(s) 2022

**Abstract**
Dry leaves of date palms (DPs) are by-products of date cultivation that are often mismanaged in oasis agroecosystems. Ideally, they should be upcycled in a circular agriculture model before producing and exporting high-value products. Compost inputs enhance soil organic matter and conserve soil. This study investigated composting of four mixtures based on farming system requirements and the availability of organic resources. The DPs were used alone or combined with sheep manure and rock phosphate. We experimented with farm-scale windrow composting to improve understanding and support decision-making. The results showed that temperature increased rapidly to 70 °C at the beginning of composting in the mixtures with manure. Organic matter content decreased by 40% in all windrows during composting, which indicated organic matter mineralisation and loss of volatile solids. Composting DP with or without sheep manure for 136 days with two turnings conserved nitrogen and increased the nitrogen content of compost. Mixing DP with sheep manure and rock phosphate increased the final total phosphorus content of compost to 0.52% (dry matter basis). Our results highlight that composting DP is a feasible alternative to improve circularity in oasis agroecosystems, by increasing organic matter content and fertiliser value, thus paving the way for sustainable production of high-value products from DP.

**Keywords** Composting · Dry palms · Rock phosphate · Nitrogen · Phosphorus losses · Circular agriculture

## 1 Introduction

Date palm (*Phoenix dactylifera* L.) is the main crop in oasis agroecosystems in Saharan and sub-Saharan regions [1]. Unlike the oil palm, it is cultivated in arid regions. In Morocco, the date palm covers 59,613 ha [2] and provides 40–60% of the annual agricultural income of farmers [3]. The Plan Maroc Vert (2009–2020) and Green Generation strategy (2020–2030) paid special attention to the date palm sector given its vital socio-economic impact on the population of oasis areas [4]. The national date palm planting program aims to plant 5 million palm trees in the Green Generation strategy, including 4 million in the Draa-Tafilalet region, with 2.4 million trees for the intensification of traditional oases and 1.6 million trees for extensions (i.e. monocropped date palm) [4].

One of the largest organic waste products (OWPs) in oases is dry date palm (DP) leaves, which remain attached to the tree until they are pruned. Every year, 1.6 million t of DPs are produced worldwide: Agoudjil et al. [5] estimated 105 million date palm trees in the world, each of which produces approximately 15 dry leaves of ca. 1 kg each [6, 7]. In recent years, they have been abandoned in fields, which can cause insect and disease infestation, or burnt, which can cause other environmental issues, especially accidental fires (Fig. 1).

Processing agricultural waste after producing high-value products generates large amounts of OWPs. One way to upcycle them in agriculture is to transform them into a stable organic amendment such as compost to provide stable

✉ Paul Robin
paul.robin@inrae.fr

1 Department of Crop Production, Protection and Biotechnology, Institut Agronomique Et Vétérinaire Hassan II, 10101 Rabat, Morocco

2 INRAE, Institut Agro, SAS, 35000 Rennes, France

3 AgroBioSciences Program, Mohammed VI Polytechnic University, 43150 Benguerir, Morocco

4 National Institute for Agricultural Research, 52000 Errachidia, Morocco

5 Department of Natural Resources & Environment, Institut Agronomique Et Vétérinaire Hassan II, 10101 Rabat, Morocco

Published online: 07 November 2022





**Fig. 1** Current management of dry palm leaves in the Draa-Tafilalet region of Morocco: **a** dry leaves on a palm tree, **b** pruned leaves left beside an irrigation canal, **c** burning leaves, and **d** palm trees after a fire

carbon (C) compounds and fertilising nutrients. Compost produced on farms has replaced commercial compost in recent years [8]. Composting is recognised as an attractive agricultural practice and an economical and sustainable approach to OWP management because it is simple to perform and can be adapted to a wide range of farming systems [9, 10]. However, nutrient losses during composting are a key environmental issue as they reduce the fertiliser value of the final compost [11]. Only a few studies have considered DP in arid areas [12, 13]. Factors such as C availability, the bulk density of dry matter, water content, and aeration influence nutrient losses from compost, particularly nitrogen (N) volatilisation, which differs from immobilisation of N in organic compounds [14–16]. Additives are used during the composting of low-nutrient feedstock to increase the final nutrient content of the compost produced [17]. Composting also helps recycle a variety of OWPs in agriculture, thus increasing nutrient efficiency and reuse. For example, Khiyami et al. [18] mixed 70% DP with 30% shrimp and crab waste to produce a final compost with high nutrient content.

For crop production, phosphorus (P) is generally considered the second limiting nutrient after N in most soils. Composts prepared from OWPs need to be enriched in P in some way [19], while those prepared from animal manure have a higher P:N ratio than most crop requirements [20]. In dry areas, Evans et al. [21] reported that one way to increase P availability to plants is to produce compost from OWPs enriched with either animal manure or rock phosphate (RP). Insufficient P content can limit microbial degradation during composting [22]. Adding RP during composting increases P availability, which increases microbial activity and reproduction and thus increases degradation of OWPs, which results in a stable and mature final compost [23, 24]. Moreover, because adding RP can increase N conservation [25–27], it is an interesting alternative to increase the final nutrient content of compost. Many studies have investigated effects of adding RP on the composting of a variety of feedstock, but none has investigated DP.

Adding organic matter (OM) to soil is the most common practice for maintaining or restoring soil fertility, particularly for soils with degraded structure and low OM content [19, 28]. Other factors interact with compost application, such as climatic conditions, soil properties, and agricultural practices, particularly irrigation and tillage [29, 30]. High irrigation or rainfall amounts can cause nutrients from compost to move deeper into the soil, which increases the potential for them to leach into groundwater. Incorporating compost decreases bulk density and increases water-holding capacity [31], nutrient contents, and biological activity [32–34]. In European experiments, Bruni et al. [35] observed that incorporating 1.93 t C $ha^{-1}$ $year^{-1}$ into the soil increased soil C by more than 4‰ $year^{-1}$. El Janati et al. [6] reported that 1 ha of monocropped date palm produces 2.4 t of DP per year, which could provide ca. half of the amount of C required to increase soil C. They also reported that agricultural soils in oasis regions have poor structure and low OM content (i.e. < 1.5%). Thus, recycling DP through composting in a circular agriculture model is of great interest to promote oasis sustainability.

We hypothesised that composting DP could promote the harvest and reuse of DP, thus avoiding abandoning or burning it in oases and paving the way for further production, biotransformation, and economic development in the regions of date palm production. The originality of our study, compared to previous studies of DP co-composting, was to design and analyse composting processes based on the current farming system of oases in the Draa-Tafilalet region (Morocco). Thus, this study provides a new scientific basis for a "ready-to-use" process for reusing DP, while responding to a range of farmer priorities, such as management of soil OM or fertility.

## 2 Materials and methods

### 2.1 Conceptual framework for choosing composting processes

We used a general approach with three steps (Fig. 2): (1) surveys, to analyse the sources of and needs for organic products to define mixture options; (2) a composting experiment, to test contrasting composting processes adapted to a variety of farmer contexts; and (3) compost-use experiments, to characterise the composts to promote suitable uses for managing either soil OM or fertility. The first and third steps have been discussed by El Janati et al. [6] and El Janati et al. [28] respectively.

The main sources of OWPs available in oases were livestock manure (i.e. sheep, goat, and cattle) and DP, whose characteristics and amounts varied among three types of farming systems (see Table S1 for the amounts of organic C, N, and P produced by the three types). The amount of DP produced was estimated at (i) 1 t $ha^{-1}$ $year^{-1}$ on traditional farms with 53 date palm trees $ha^{-1}$ and (ii) 2.5 t $ha^{-1}$ $year^{-1}$ on modern farms with 160 date palm trees $ha^{-1}$. DP had a high OM content (90.9%) and low N and P contents, thus providing mainly a source of organic C in farming systems. On traditional farms, the amount of livestock manure produced was estimated at 19.4 t $farm^{-1}$ $year^{-1}$ [6]. Manure was a source of N, P, and several other nutrients that increased soil fertility for crop production. It was applied directly on soils. Farmers considered it as a scarce resource for fertilisation. Sheep was one of the main livestock species that contributed to farmers' incomes. Because genetic conservation of the local sheep breed D'man is a current issue of farm sustainability [36], sheep manure (SM) was chosen as the manure feedstock for our experiment.





**Fig. 2** Design of on-farm composting of date palm residues in a circular agriculture model. Step 1: surveys (see details in El Janati et al. [6]); step 2: composting (see details in El Janati et al. [42]); and step 3: compost characterisation (see details in El Janati et al. [28])

Nutrients were exported from farms mainly after harvest (see Table S2 for the organic N and P exported by crops and dates on the three types of farms). Crop residues consumed by livestock represented ca. one-third of total nutrient exports by crops. Modern farms that produced only date palms had no livestock manure; thus, a composting treatment using only DP was one of the mixtures tested.

Manure is a source of nutrients and microflora that increases microbial activity during composting. We decided to use a mixture with SM to provide farmers with an option to increase the area on which compost could be applied. We considered that mixtures needed to contain at least 30% manure to significantly influence the final nutrient content (i.e. impact on crop production) and the biotransformation caused by the initial microbial population (i.e. impact on the composting process), as observed by Abid et al. [12] and Sadik et al. [37]. A 70:30 ratio of DP:SM (v:v) was used for practical reasons.

RP is added to composting mixtures to (i) improve composting by removing the P limitation of microbial activity and (ii) increase the compost's final P content. The P added, which is considered to be a non-volatile element, should be conserved throughout composting. Moreover, the effects of adding RP to such feedstock have not been studied. Thus, we hypothesised that adding RP would increase the temperature of the windrow and increase N conservation by removing the P limitation to microbial growth. Moreover, adding RP was expected to increase the P content of the compost produced, which would increase crop productivity.

We applied the same operations to all composting mixtures. Crushed DP was pre-processed by soaking it in a basin of water before setting up windrows, to stimulate microbial colonisation of DP particles (i.e. "attached microflora") and increase the biodegradability of crushed DP during composting [12]. We turned the windrows to ensure that temperature and humidity changed in order to promote succession of microorganisms on feedstock aggregates and ensure homogenisation of the final products. We limited the number of turnings to limit the need for labour and to create a more acceptable process for farmers. We also hypothesised that limiting windrow turning would decrease water and N and C losses during composting. Thus, we decided to turn the windrows twice, after 5 weeks (day 39) and 11 weeks (day 81). We decided to use a relatively long composting period (> 4 months, 136 d) to degrade the DP.

## 2.2 Experimental design

The experiment was conducted at the experimental station of the National Institute for Agricultural Research of Morocco in Errachidia (31°55′13.0″ N, 4°26′59.0″ W). This station is dedicated to research on oasis agroecosystems. The study site has a dry and continental arid climate, with hot summers (maximum 43 °C), cold winters (minimum − 3 °C), and mean annual rainfall of 116 mm. The soil is a loam [28]. The experiment was performed in autumn and winter, with air temperature ranging from 15 to 35 °C, and total rainfall of 14 mm during the 136 days of composting. The composting feedstock used was DP and SM, which were collected from the experimental station (Table 1). DPs were crushed to a length of ca. 4 cm and then soaked in a basin of water for 7 days. The amounts of each feedstock varied among mixtures (Table 2). The mixture was sampled after mixing the feedstock and before adding water to reach 70% water content (wet mass basis). Because water percolated out of the mixtures before this percentage was reached, however, the initial water content, considered to reflect the water-holding capacity, was determined from the volume of water added to each mixture.

**Table 2** Initial volume and raw material composition of the mixtures (in $m^3$ or kg wet weight $windrow^{-1}$). T1-dry palm leaves; T2-dry palm leaves and rock phosphate; T3-dry palm leaves and sheep manure; T4-dry palm leaves, sheep manure, and rock phosphate. *FAS* free air space

| Treatment | T1 | T2 | T3 | T4 |
|---|---|---|---|---|
| Total volume ($m^3$) | 5.15 | 5.15 | 5.15 | 5.15 |
| Total weight (kg) | 1510 | 1522 | 1766 | 1778 |
| Dry palm leaves (kg) | 1310 | 1310 | 850 | 850 |
| Sheep manure (kg) | 0 | 0 | 516 | 516 |
| Rock phosphate (kg) | 0 | 12 | 0 | 12 |
| Water (kg) | 200 | 200 | 400 | 400 |
| FAS (% total volume) | 74 | 74 | 71 | 71 |

**Table 1** Initial composition of feedstock used in composting mixtures: date palm leaves before (dry) and after (wet) soaking, sheep manure, and rock phosphate (calculated from mixture analysis, not directly observed). Values are expressed as a percentage of wet weight (dry matter) or dry matter (the others)

| Initial feedstock | Dry matter | Organic matter | Total N | Total P | Total K | Total Mg | Total Ca | Total Na |
|---|---|---|---|---|---|---|---|---|
| Dry palm leaves | 93 ± 0 | 88 ± 0 | 0.44 ± 0.01 | 0.03 ± 0.00 | 0.49 ± 0.03 | 0.26 ± 0.01 | 1.01 ± 0.09 | 0.05 ± 0.01 |
| Wet palm leaves | 38 ± 2 | 90 ± 1 | 0.59 ± 0.03 | 0.03 ± 0.01 | 0.12 ± 0.02 | 0.14 ± 0.01 | 0.61 ± 0.03 | 0.05 ± 0.00 |
| Sheep manure | 88 ± 2 | 74 ± 1 | 1.93 ± 0.05 | 0.82 ± 0.04 | 2.34 ± 0.05 | 0.61 ± 0.01 | 2.72 ± 0.56 | 0.64 ± 0.03 |
| Rock phosphate | 100 ± 0 | 0.00 | 0.00 | 6.04 ± 0.03 | 0.00 | 0.00 | 16.12 ± 0.20 | 0.00 ± 0.00 |

The initial water content was 67 ± 2% in treatments T1 and T2 and 56 ± 2% in treatments T3 and T4. During mixing, 12.1 kg of RP was added to one treatment with DP and one treatment with SM. This small amount was determined based on the small P export of intensive date cultivation in monocropped areas (Table S2).

The experimental design was a randomised complete block of mixtures of DP, SM, and RP, with three replicates each (Table 2). The four treatments (12 windrows in total) were 100% crushed DP (T1), 100% crushed DP and RP (T2), 70% crushed DP and 30% SM (T3), and 70% crushed DP and 30% SM and RP (T4). All windrows were set up with the same dimensions (length × width × height) at the beginning and after each turning. The initial dimensions were 3.6 × 2.0 × 1.4 m. On day 39 (first turning), they were 3.5 × 1.9 × 1.3 m. On day 81 (second turning), they were 3.2 × 1.8 × 1.2 m. The final heights (day 136) were 0.98 m for T1 and T2 and 0.93 m for T3 and T4 (see Table S3 for additional information on initial physical and chemical properties of the mixtures).

At each turning, all windrows were weighed and sampled, and their dimensions were recorded. The wet weight was measured using a digital scale. Then, the three windrows of each treatment were mixed together to increase the repeatability within each treatment. The total mass of each treatment was then divided into three identical windrows. Water was added to these replicates to maintain a water content of 55–65%. The volume of water added was the same for the three replicates of each treatment but differed among treatments.

## 2.3 Sampling

Three representative samples (ca. 400 g each) were taken from each windrow on the first and last days and the two turning days. Samples were prepared by mixing six subsamples taken from six sites of each windrow until they were homogenised.

## 2.4 Measurements

The temperatures of each windrow were measured using a metal thermometer (K-type Thermometer probe HI766TR2) each morning (9:00–11:30 am) until day 123. The temperatures were measured at three locations: the top (30 cm from the surface), middle (centre), and bottom (30 cm from the base) layers. The air temperature was also recorded.

After setting up each windrow, but before weighing it, its total volume ($V_{tot}$ in $m^3$) was calculated from the mean height ($h_{avg}$ in m), width ($w_{avg}$ in m), and length ($L_{avg}$ in m) measured at three positions along the windrow:

$$V_{tot} = \frac{h_{avg} \times w_{avg} \times L_{avg}}{2} \quad (1)$$

Compost samples were analysed for water content, pH, electrical conductivity, OM, total N, total P, potassium (K), magnesium (Mg), calcium (Ca), and sodium (Na). Water content, electrical conductivity, and pH were determined in fresh samples. Water content was determined by weight loss after drying at 105 °C [38]. Electrical conductivity and pH were measured using a conductivity electrode or pH metre, respectively, after diluting a sample with distilled water (1:5 sample:water w:v). OM content, also called volatile solids, was determined by loss on ignition at 550 °C for 4 h [39]. OM was converted into organic C by dividing it by 2 [40]. Total N was analysed by the Kjeldahl method [41]. Total P, total K, total Mg, Ca, and Na were analysed by inductively coupled plasma atomic emission spectroscopy (ICP-EOS) after mineralisation with nitric acid and filtering.

## 2.5 Data processing

Measurements of the total dry matter in compost samples varied greatly and showed inconsistencies in initial calculations (e.g. large increases between two turnings), but the P:ash, Mg:ash, and Na:ash ratios remained stable. Therefore, we considered that the dry matter content had likely become biased between windrow sampling and conditioning of the final samples. Thus, the masses of water, dry matter, and the associated elements (N, P, K, Mg, Ca, Na) in windrows were estimated by assuming ash conservation throughout composting. Wet-weight losses were calculated by adding the water input during turnings and rainfall to the measured wet weight of the windrows.

Moisture and free air space (FAS) are known to be critical factors for managing compost [9]. FAS is the fraction of the total volume that is occupied by air and not by solids or water [9]. It was calculated as the total volume ($V_{tot}$ in $m^3$) minus the volumes of water ($V_{water}$ in $m^3$) and dry matter ($V_{solids}$ in $m^3$) and expressed as a percentage of the total volume of the windrow to facilitate comparisons among treatments and dates:

$$FAS = \frac{\left(V_{tot} - V_{solids} - V_{water}\right)}{V_{tot}} \times 100 \quad (2)$$

The water volume was calculated from the water mass ($M_{tot}$: total mass of windrow in kg; $WC$: water content of sample in kg water $kg^{-1}$ wet weight) by assuming a density of water ($\delta_{water}$) equal to 1000 kg $m^{-3}$:

$$V_{water} = \frac{M_{tot} \times WC}{\delta_{water}} \quad (3)$$

The volume of dry matter was calculated from the mass of dry matter (in kg, calculated from $M_{tot}$ and the dry matter content ($DM$ in kg dry matter $kg^{-1}$ wet weight)) and assuming a density of solids ($\delta_{solids}$) of 1600 kg $m^{-3}$ [16]:

$$V_{solids} = \frac{M_{tot} \times DM}{\delta_{solids}} \quad (4)$$

Means of each parameter were tested for statistically significant differences using Welch's *t*-test. All details of raw data and calculations can be found in El Janati et al. [42].

## 3 Results and discussion

### 3.1 Composting processes

#### 3.1.1 Temperature dynamics

Temperature varied during composting (Fig. 3), which reflects the microbial activity of OM degradation and heat loss through passive aeration. All windrows went through mesophilic, thermophilic, and curing phases. For all treatments, windrows had higher temperatures in the centre than near the top or base. In mixtures with DP (T1 and T2), temperatures in the centre of the windrows peaked at 55 °C after 88 days of composting and two turnings. In mixtures with SM (T3 and T4), temperatures in the centre of the windrows peaked at more than 70 °C after 8 days.

Temperature profiles during composting were similar in windrows T1 and T2, as well as in windrows T3 and T4. Adding RP to mixtures T2 and T4 did not change the temperature dynamics, which indicates that P did not limit microbial activity, as shown by Lu et al. [43], who composted pig manure with RP. On the contrary, adding manure clearly accelerated the increase in temperature from the beginning of composting, as observed by Paillat et al. [44]. A rapid increase in temperature was also observed in other green-waste composting experiments [25, 45, 46]. The rapid increase in temperature of mixtures T3 and T4 was due to the high respiration rate of the microbial population, which was enhanced by the increased input of energy, nutrients, and the initial microbial population provided by the SM. As a result, the second temperature peak merged with the





**Fig. 3** Temperature dynamics during composting (for all treatments). T1-dry palm leaves, T2-dry palm leaves and rock phosphate, T3-dry palm leaves and sheep manure, and T4-dry palm leaves, sheep manure, and rock phosphate

first peak to form a longer period of higher temperatures. After the first and second turnings, mixtures T3 and T4 had significantly higher maximum temperatures than mixtures T1 and T2. Therefore, adding manure to DP can be considered an option to increase windrow temperature by 20 °C (from ca. 50 °C to ca. 70 °C), which can be used when crop residues with lower contents of pathogens and weed seeds are desired.

#### 3.1.2 Wet weight dynamics

At the end of composting, the wet mass of each treatment was approximately half of its initial wet mass (Table 3). The decrease in wet mass of mixtures T1, T2, T3, and T4 was 63%, 62%, 60%, and 61%, respectively, much of which was due to evaporation of water. The dry matter loss in mixtures was ca. 42% in T1 and 38% in T2, which was significantly ($p < 0.05$) higher than the 33% in T3 and 31% in T4. Thus, DP had similar biodegradability whether composted alone or with SM. In other studies, Adhikari et al. [47] found that a decrease of 39% and 68% in dry matter corresponded to temperatures of 55 °C and 52 °C, respectively, reached during composting of OWPs (i.e. food waste mixed with chopped wheat straw in a 5:1 and 8.9:1 ratio, respectively). Adding RP to mixtures T2 and T4 did not increase the losses of dry matter, total weight, or water. Thus, P did not limit microbial activity, even in soaked DP.

Wet weight loss did not reflect the temperature dynamics, except during the initial period (days 1–39). During this period, the temperature in windrows T3 and T4 was higher than that in T1 and T2, as was the weight loss (42% and 44% vs. 27% and 30%, respectively). Afterwards, T1 and T2 lost more weight than T3 and T4 did, despite the higher temperatures in T3 and T4, so that the final weight loss was similar among the four treatments. It is likely that T3 and T4 lost less heat due to air convection because the fine particles of SM decreased porosity and lost less water due to evaporation because they contained less water.

**Table 3** Change in wet weight, dry matter, and free air space throughout the composting period

| Composting day | Day 0 | Day 39 | Day 81 | Day 136 |
|---|---|---|---|---|
| Wet weight (kg windrow$^{-1}$) | | | | |
| T1 | 1507 ± 6 | 1206 ± 5 | 960 ± 4 | 667 ± 3 |
| T2 | 1519 ± 6 | 1157 ± 4 | 908 ± 3 | 684 ± 3 |
| T3 | 1763 ± 6 | 1178 ± 4 | 1082 ± 4 | 866 ± 3 |
| T4 | 1775 ± 6 | 1150 ± 4 | 1039 ± 3 | 844 ± 3 |
| Dry matter (kg windrow$^{-1}$) | | | | |
| T1 | 505 ± 20 | 385 ± 49 | 310 ± 63 | 292 ± 75 |
| T2 | 495 ± 36 | 451 ± 33 | 331 ± 37 | 305 ± 44 |
| T3 | 794 ± 48 | 630 ± 29 | 583 ± 36 | 535 ± 31 |
| T4 | 809 ± 43 | 634 ± 55 | 595 ± 51 | 560 ± 49 |
| Free air space (% total volume) | | | | |
| T1 | 74 ± 0 | 76 ± 2 | 75 ± 1 | 81 ± 1 |
| T2 | 74 ± 0 | 76 ± 1 | 79 ± 1 | 81 ± 0 |
| T3 | 72 ± 0 | 79 ± 0 | 74 ± 1 | 77 ± 2 |
| T4 | 71 ± 0 | 79 ± 1 | 77 ± 0 | 78 ± 2 |

#### 3.1.3 Free air space

In all treatments, the FAS increased from the beginning (ca. 70% of total volume) to the end (ca. 80%) of composting (Table 3). The FAS in mixtures T1 and T2 was significantly ($p < 0.05$) higher than that in mixtures T3 and T4, due to their components, especially manure, which has a high content of small particles. Thus, adding SM to mixtures T3 and T4 filled the pores between DP particles, which reduced their FAS. Haug [9] reported that maintaining proper water content and FAS during composting requires balancing many competing forces. Under arid conditions, increased passive aeration due to high FAS is expected to increase evaporation. Therefore, further experiments would be useful to explore ways to decrease FAS and its influence on composting and the final products.

### 3.2 Dynamics of chemical composition

#### 3.2.1 Loss of volatile solids

OM content decreased by 40% in all windrows during composting, which indicated OM mineralisation and loss of volatile solids (Fig. 4). During composting of a mixture of DP (soaked for 7 days before composting) and goat manure, Abid et al. [12] reported a similar loss of OM (37%). El Ouaqoudi et al. [48] observed a smaller loss of OM (29%) after 14 months of DP composted without initial soaking. Sadik et al. [37] observed a loss of OM of ca. 15% in a 1:1 mixture of DP and cow manure. Therefore, DP may contain compounds that resist degradation. Haug [9] and Paillat et al. [44] mentioned that transformation of OM depends on its degradability. OM with recalcitrant components, such as lignin and cellulose, stabilise the remaining organic compounds, which helps increase soil OM further. In our experiment, soaking may have increased the initial microbial activity by decreasing the electrical conductivity, increasing the N content, and filling the xylem of palm fragments with water. The relatively high initial water content (ca. 60%) and the initial period of 39 days without turning allowed hyphae of fungi to develop. They were observed in all windrows during the first turning, which indicated active degradation of cellulose, lignin, and other polysaccharides [49].

Composting of most OM generally has an initial stage of rapid degradation followed by a longer stage of slower degradation [50–52]. In mixtures T3 and T4, most OM was





**Fig. 4** Dynamics of (**a**) the content and (**b**) mass of organic matter during composting: T1-dry palm leaves, T2-dry palm leaves and rock phosphate, T3-dry palm leaves and sheep manure, and T4-dry palm leaves, sheep manure, and rock phosphate. Letters indicate significant differences ($p < 0.05$) between windrows





**Fig. 5** Dynamics of (**a**) the content and (**b**) mass of nitrogen during composting: T1-dry palm leaves, T2-dry palm leaves and rock phosphate, T3-dry palm leaves and sheep manure, and T4-dry palm leaves, sheep manure, and rock phosphate. Letters indicate significant differences ($p < 0.05$) between windrows

lost during the first stage (39 days) of composting. The OM content of mixtures T1 and T2 decreased slowly during composting, which indicates that DP resisted degradation more than SM.

### 3.2.2 Nitrogen conservation

Total N contents were significantly ($p < 0.05$) higher in mixtures T3 and T4 than those in mixtures T1 and T2 (Fig. 5). The total N content in all mixtures increased during composting, with a maximum in T3 and T4. The increase in N content, along with a stable N mass, in all mixtures indicates that mineralisation of DP produced enough available C to supply the immobilisation processes of microorganisms. This result agrees with those of Bernal et al. [53] and Paredes et al. [52], who observed that total N content increased due to the concentration effect if leaching is stopped or minimised. These dynamics also suggest the lack of N losses from the windrows, as reported by many studies [44, 46, 51–53] when composting OWPs with high lignin and cellulose contents.

The effect of RP was not significant, despite a small decrease in N mass in mixture T1 of ca. 10% (from $0.58 \pm 0.03\%$ to $0.53 \pm 0.03\%$ initial dry matter) compared to that in T2, whose N was conserved at $0.59 \pm 0.04\%$ initial dry matter. Thus, composting DP and SM with a lower turning frequency decreases N losses.

### 3.2.3 Dynamics of phosphorus and nutrient contents of the final composts

Compost mixtures with SM (T3 and T4) contained significantly more total P than those without SM (T1 and T2), due to the high P content of the SM added (Fig. 6). Compost mixtures with RP (T2 and T4) had more total P than those without RP (T1 and T3). Adding RP may increase microbial activity and promote P accumulation and preservation, thus increasing the final P content of composts [24, 54]. The final total P content was higher than the initial content in mixture T2, but they were similar in T1, T3, and T4. Adding the same mass of RP to mixtures T2 and T4 led to a lower P





**Fig. 6** Dynamics of (**a**) the content and (**b**) mass of phosphorus during composting: T1-dry palm leaves, T2-dry palm leaves and rock phosphate, T3-dry palm leaves and sheep manure, and T4-dry palm leaves, sheep manure, and rock phosphate. Letters indicate significant differences ($p < 0.05$) between windrows

content in T4 because T4 had a higher initial total mass of dry matter.

The total mass of P decreased significantly by ca. 1.5 kg P $t^{-1}$ initial dry matter, as confirmed by the decrease in mean P content ($2.39 \pm 0.36\%$ to $1.60 \pm 0.14\%$ ash measured in the $2 \times 18$ samples concerned), from days 0–39 in both mixtures with SM, and was not observed in mixtures T1 or T2. This can be explained either by specific biological processes ("real" P loss) or by measurement artefacts ("false" P loss). However, gaseous P compounds could have been associated with manure [55, 56]. We have no evidence to consider one of the two hypotheses as more probable. The risk of P loss should be studied under more controlled conditions to clarify our hypotheses. In the present study, if the final P content of compost is a critical issue, the risk of losing P from manure during composting with DP can be decreased by adding RP to the initial mixture at a rate of 1.5 kg P $t^{-1}$ initial dry matter (25 kg RP). If the loss of P is confirmed by further studies, composting experiments similar to those described here should indicate whether the loss depends or not on the DP:manure ratio or on characteristics of the manure. Initial and final masses of K, Mg, Ca, and Na did not differ significantly ($p < 0.05$) (Table S4).

## 3.3 Adapting composting to the needs of circular agriculture

Different ways to apply the composts produced (Fig. 7) can optimise various properties and help composting meet the needs of circular agriculture.

### 3.3.1 Compost needs in date palm monocropping

Following FAO [57] recommendations, compost is used as an organic fertiliser for young plants, applied per planting hole and supplemented with inorganic fertilisers. Traditionally, 3 kg of compost is mixed with the soil from the hole and then returned to the hole. Our results show that composts T1 and T2 have high OM content, low electrical conductivity, and a pH of 6.5 and should contain more stable compounds. They can provide a long-term increase in soil OM and water-holding capacity. Therefore, larger amounts could be added to increase the expected effects of OM input. El Kinany et al. [58] showed that applying compost to a sandy soil increased the growth of micropropagated date palm plantlets (i.e. fresh and dry biomass, root system, and leaf nutrient content). Applying compost increases aggregate stability and decreases soil density, thus decreasing resistance to root penetration and increasing water-holding capacity [59]. Compost with a high OM content is reported to promote crop growth by influencing the membrane-bound $H^+$-ATPase in the root system, which increases nitrate uptake [60, 61]. Given the properties of oasis soils, the recommended amounts of compost exceed 30 kg per date palm every 2–3 years [6]. Composts T1 and T2 can be applied either as mulch or incorporated into the topsoil, thus minimising water loss from the soil through evaporation and increasing the soil's humus content [62]. Composts T3 and T4 should be incorporated into the topsoil to supply nutrients to the plant in the irrigated zone. Almadini et al. [63] reported that the amount of fertiliser applied strongly influences date palm production in both old and new oases of Al-Ahsa in Saudi Arabia. Moreover, manure is a scarce commodity in traditional oases and is absent in modern oases [6]; thus, using compost produced on farms can offset this lack of organic inputs and promote the production chain of organic dates.

### 3.3.2 Compost needs when date palms are associated with other crops

Because annual crops have a short growing season and need nutrients soon after planting, the compost applied must





**Fig. 7** Adaptation of composting to the needs of circular agriculture of oasis farming systems (practical implications). SOM, soil organic matter; OM, organic matter

provide a sufficient supply of plant-available nutrients. Composts T3 and T4 can be incorporated into the soil during seedbed preparation for these annual crops, particularly cash crops. Each year, farmers apply 2.5 and 2.7 t $ha^{-1}$ of manure to wheat and cumin crops, respectively [6]. Composts T3 and T4 could also be applied to perennial crops at sowing or during growth. Benabderrahim et al. [64] observed that applying DP compost to a loamy sand soil at 30 t $ha^{-1}$ increased both its OM content and water-holding capacity. Using compost to produce forage strengthens the role of livestock in these systems. Maintaining soil water is vital for crop growth and preservation of oasis soils, as soils in arid areas require management to provide sufficient water. Applying compost to the soil as mulch can be an effective way to manage bare soil. All composts produced can be incorporated into the topsoil to increase the formation and stability of soil aggregates.

#### 3.3.3 Compost production to meet OM requirements of cropping systems

El Janati et al. [6] observed that manure is limited and already used in current farming systems, while DPs are insufficiently recycled, and that traditional and mixed farming systems have more C in manure than in DP. Therefore, on the farms that use manure, we suggest combining all available DP leaves, after crushing and soaking, with existing manure in a 70:30 (v:v) ratio to produce composts similar to T3 or T4. Composts T3 and T4 conserved manure nutrients, albeit with a risk of losing some P, which can be compensated by adding RP. In addition, N can be conserved through anaerobic digestion-composting processes [65] and by adding additives to shorten composting times and increase nutrient contents [66].

### 3.4 Potential implementation of the composting technique

We showed that composting is a tool to upcycle DP to meet the need to manage soil OM and fertility in Moroccan oases. Our results confirmed previous studies showing that composts based on DP are suitable for agricultural use [12, 13, 37, 48]. Here, we discuss some of the technical, economic, and sociological issues of this transition pathway.

Identifying the most relevant composting process first depends on the equipment chosen to produce compostable

particles. Equipment categories include chippers, shredders, and grinders. DPs require specific equipment [67]. Composting succeeds when particle sizes vary from 5 mm [12] to 50 cm [48]. Therefore, stakeholders should prioritise socio-economic constraints (e.g. cost, equipment lifetime, maintenance, professional qualifications) and begin sieving feedstock, if necessary. The composting process secondly depends on the feedstock added. The literature shows successful DP composting with feedstock from a variety of sources: urban, livestock, or the food-processing industry. Therefore, stakeholders should maximise short-term agronomic benefits and prioritise the ability to meet needs of production and use operations. The third critical issue is managing salinity. The soaking stage decreased the electrical conductivity of DP. The literature shows that electrical conductivity of DP composts ranges from 1 to 7 mS $cm^{-1}$, depending on the feedstock. Stakeholders should determine how to reuse the soaking water associated with composting.

Underestimating the role of social values and culture in economic functioning can hinder the spread of innovations [68]. Farm managers and agricultural advisors can help meet the growing demand for organic fertilisers and support the spread of organic farming. In addition to windrow composting, small-scale options such as composting in reusable bags can be tested [69, 70]. "Good practices" for safely harvesting, transporting, and upcycling DP, instead of burning it, could be developed and spread as new policies by institutions. Communication also should consider factors that influence long-term dynamics of the OM added [71].

In regions that produce large amounts of DP, a variety of products can be produced while recycling the DP, whether immediately after harvesting, from the liquid produced after the soaking stage or from the DP after soaking. The strategy can be defined by optimising the combination of feed [7], fibre [72], or fuel [73] products desired, depending on the available technologies and economic opportunities, while maximising the social value of these products in the regions concerned. Attention should be paid to the compatibility between new inputs necessary to produce products and agricultural or horticultural uses of the final compost.

## 4 Conclusion

For the first time, our study showed that composting is a feasible alternative for appropriate management of DP in the context of circular agriculture in oasis agroecosystems. We designed, experimented with, and discussed the application of four contrasting composting options that can help resolve the paradox of unused organic residues coexisting with soils' need for organic inputs.

This research provided better understanding of OM decomposition and dynamics of the masses of water, C, N, and P during composting of DP. Our results showed that initial soaking of crushed DP allowed them to be composted without any other additive. They also showed that the quality of final composts could be improved in several ways. Adding manure increased compost temperature, which should increase the sanitation of the initial feedstock. Adding either SM or RP increased the nutrient content of the final compost. Limiting the number of turnings increased the amount of initial N conserved. However, future studies on the effects of water content and lower FAS could help improve the processes.

Discussion of compost uses in a range of cropping systems can help farmers manage available organic resources appropriately by identifying the most suitable composting process and compost uses. The four types of compost met crop requirements for food or feed production. Our discussion highlighted that the composts produced could be used to increase forage production and help improve livestock conservation and sustainable development in such arid agroecosystems, which could increase incomes. Thus, upcycling DP through composting is an effective way to increase the amount of compost produced and increase soil fertility in a circular agriculture model of arid agroecosystems. Moreover, developing harvesting and reuse of DP will facilitate the production of DP-based products, for example for local fuel production.

**Supplementary Information** The online version contains supplementary material available at https://doi.org/10.1007/s13399-022-03387-z.

**Acknowledgements** We are grateful to research grants provided by the Mohammed VI Polytechnic University (UM6P), ERASMUS program, Institut Agro, and INRAE. We are also grateful to the Institut National de la Recherche Agronomique in Errachidia for the technical assistance during the experiment. We thank Safya Menasseri-Aubry for her assistance in initialising and organising the collaboration, Hamza Elfinnou for his valuable help in collecting data, and M. Corson for correcting the English language and his valuable comments.

**Author contribution** Conceptualisation and methodology: P.R., A.B., N.A.-C., and M.E.J.; data curation: P.R. and M.E.J.; software and analysis: P.R., N.A.-C., and M.E.J.; writing: M.E.J., N.A.-C., and P.R.; supervision: A.B., P.R., N.A.-C., A.O., Z.T., A.S., and M.C. All authors have read and agreed to the published version of the manuscript.

**Funding** The experiment was funded by the Agence Nationale pour le Développement des Zones Oasiennes et de l'Arganier (ANDZOA), the OCP Group and Mohammed VI Polytechnic University (UM6P). The affiliated institutions funded the salaries and operating costs of the authors.

**Data availability** All details of raw data and calculations have been published by El Janati et al. [42]. They are freely available at following address: https://doi.org/10.15454/DFIAFE.

## Declarations

**Ethical approval** Not applicable.

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

## References

1. Toutain G, Dollé V (1989) Situation des Systèmes oasiens en régions chaudes. Cahiers de la Recherche-Développement 22:3–14
2. FAOSTAT (2019) Food and Agriculture Organization of the United Nations Statistics Division. http://www.fao.org/faostat/en/#data/QC. Accessed 2 June 2021
3. Sedra M (2012) Guide du Phoeniciculteur: Mise en place et conduite des vergers phoénicicoles (Ar, Fr) Sedra, H. (INRA, Marrakech (Ma) centre Régional de la Recherche Agronomique de Marrakech). INRA, Rabat (Ma), 2012, 311 p., Résumés (Fr, En, Ar), Notes:31 tabl., 136 fig., 67 rèf. https://www.inra.org.ma/fr/content/guide-du-phoeniciculteur-mise-en-place-et-conduite-des-vergerspho%C3%A9nicicoles; https://www.inra.org.ma/sites/default/files/publications/ouvrages/guidephoenicicole1212.pdf
4. MAP (2021) Province d'Errachidia : M. Akhannouch visite des projets de développement agricole. In: Maroc.ma. https://www.maroc.ma/fr/actualites/province-derrachidia-m-akhannouch-visite-des-projets-de-developpement-agricole. Accessed 25 May 2021
5. Agoudjil B, Benchabane A, Boudenne A et al (2011) Renewable materials to reduce building heat loss: characterization of date palm wood. Energy and Buildings 43:491–497. https://doi.org/10.1016/j.enbuild.2010.10.014
6. El Janati M, Akkal-Corfini N, Bouaziz A et al (2021) Benefits of circular agriculture for cropping systems and soil fertility in oases. Sustainability 13:4713. https://doi.org/10.3390/su13094713
7. Barreveld WH (1993) Date palm products. FAO Agricultural Services Bulletin, No 101. http://www.fao.org/docrep/t0681e/t0681e00.htm#con
8. Pane C, Celano G, Piccolo A et al (2015) Effects of on-farm composted tomato residues on soil biological activity and yields in a tomato cropping system. Chem Biol Techn Agric 2:4. https://doi.org/10.1186/s40538-014-0026-9
9. Haug RT (1993) The Practical Handbook of Compost Engineering. Lewis Publishers, Boca Raton, FL
10. Robin P, Paillat J-M, Lorinquer E, Aubert C, Toudic A, Farinet J-L, Cluzeau D, Ponchant P, Levasseur P, Le Bris B, Capdeville J, Oudart D, Akkal-Corfini N, Hassouna Ml (2018) Procédés et traitements par compostage des effluents d'élevages. In: Compostage et composts : Avancées scientifiques et techniques, Guardia Amaury (ed.). Lavoisier Tec et Doc Paris, pp 531–583. https://agritrop.cirad.fr/589434/; https://www.lavoisier.fr/livre/environnement/compostage-et-composts/de-guardia/descriptif-9782743023591
11. Hao X, Chang C, Larney FJ (2004) Carbon, nitrogen balances and greenhouse gas emission during cattle feedlot manure composting. J Environ Qual 33:37–44. https://doi.org/10.2134/jeq2004.3700
12. Abid W, Mahmoud IB, Masmoudi S et al (2020) Physico-chemical and spectroscopic quality assessment of compost issued from date palm waste valorization. J Environ Manage 264:110492. https://doi.org/10.1016/j.jenvman.2020.110492
13. Vico A, Pérez-Murcia MD, Bustamante MA et al (2018) Valorization of date palm (Phoenix dactylifera L.) pruning biomass by co-composting with urban and agri-food sludge. J Environ Manage 226:408–415. https://doi.org/10.1016/j.jenvman.2018.08.035
14. Barrington S (2002) Effect of carbon source on compost nitrogen and carbon losses. Biores Technol 83:189–194. https://doi.org/10.1016/S0960-8524(01)00229-2
15. Bueno P, Tapias R, López F, Díaz MJ (2008) Optimizing composting parameters for nitrogen conservation in composting. Bioresour Technol 99:5069–5077. https://doi.org/10.1016/j.biortech.2007.08.087
16. Oudart D, Robin P, Paillat JM, Paul E (2015) Modelling nitrogen and carbon interactions in composting of animal manure in naturally aerated piles. Waste Manage 46:588–598. https://doi.org/10.1016/j.wasman.2015.07.044
17. Barthod J, Rumpel C, Dignac MF (2018) Composting with additives to improve organic amendments. A review. Agron Sustain Dev 38:17. https://doi.org/10.1007/s13593-018-0491-9
18. Khiyami M, Masmali I, Abu-khuraiba M (2008) Composting a mixture of date palm wastes, date palm pits, shrimp, and crab shell wastes in vessel system. Saudi J Biol Sci 15:7
19. Biswas DR, Narayanasamy G (2006) Rock phosphate enriched compost: an approach to improve low-grade Indian rock phosphate. Bioresour Technol 97:2243–2251. https://doi.org/10.1016/j.biortech.2006.02.004
20. Chabalier P, de Kerchove VV, Macary HS (2006) Guide de la fertilisation organique a La Reunion. CIRAD, Montpellier
21. Evans J, Condon J, Evans J, Condon J (2009) New fertiliser options for managing phosphorus for organic and low-input farming systems. Crop Pasture Sci 60:152–162. https://doi.org/10.1071/CP07153
22. Rashad FM, Saleh WD, Moselhy MA (2010) Bioconversion of rice straw and certain agro-industrial wastes to amendments for organic farming systems: 1. Composting, quality, stability and maturity indices. Bioresour Technol 101:5952–5960. https://doi.org/10.1016/j.biortech.2010.02.103
23. Billah M, Bano A (2015) Role of plant growth promoting rhizobacteria in modulating the efficiency of poultry litter composting with rock phosphate and its effect on growth and yield of wheat. Waste Manage Res 33:63–72. https://doi.org/10.1177/0734242X14559593
24. Zhang L, Sun X (2017) Addition of fish pond sediment and rock phosphate enhances the composting of green waste. Bioresour Technol 233:116–126. https://doi.org/10.1016/j.biortech.2017.02.073
25. Bustamante MA, Ceglie FG, Aly A et al (2016) Phosphorus availability from rock phosphate: combined effect of green waste composting and sulfur addition. J Environ Manage 182:557–563. https://doi.org/10.1016/j.jenvman.2016.08.016
26. Meena D, Biswas RD (2015) Effect of rock phosphate enriched compost and chemical fertilizers on microbial biomass phosphorus and phosphorus fractions. Afr J Microbiol Res 9:1519–1526. https://doi.org/10.5897/AJMR2014.6957
27. Nishanth D, Biswas DR (2008) Kinetics of phosphorus and potassium release from rock phosphate and waste mica enriched compost and their effect on yield and nutrient uptake by wheat (Triticum aestivum). Bioresour Technol 99:3342–3353. https://doi.org/10.1016/j.biortech.2007.08.025

28. El Janati M, Akkal-Corfini N, Robin P, Oukarroum A, Sabri A, Thomas Z, Chikhaoui M, Bouaziz A (2022) Compost from date palm residues increases soil nutrient availability and growth of silage corn (Zea mays L.) in an arid agroecosystem. J Soil Sci Plant Nutr 22:3727–3739. https://doi.org/10.1007/s42729-022-00922-9
29. Amlinger F, Götz B, Dreher P et al (2003) Nitrogen in biowaste and yard waste compost: dynamics of mobilisation and availability—a review. Eur J Soil Biol 39:107–116. https://doi.org/10.1016/S1164-5563(03)00026-8
30. Saleem A, Irshad M, Hassan A et al (2017) Extractability and bioavailability of phosphorus in soils amended with poultry manure co-composted with crop wastes. J Soil Sci Plant Nutr 17:609–623. https://doi.org/10.4067/S0718-95162017000300005
31. Kranz CN, McLaughlin RA, Johnson A et al (2020) The effects of compost incorporation on soil physical properties in urban soils - a concise review. J Environ Manage 261:110209. https://doi.org/10.1016/j.jenvman.2020.110209
32. Alvarenga P, Palma P, Mourinha C et al (2017) Recycling organic wastes to agricultural land as a way to improve its quality: a field study to evaluate benefits and risks. Waste Manage 61:582–592. https://doi.org/10.1016/j.wasman.2017.01.004
33. Biswas DR (2011) Nutrient recycling potential of rock phosphate and waste mica enriched compost on crop productivity and changes in soil fertility under potato–soybean cropping sequence in an Inceptisol of Indo-Gangetic Plains of India. Nutr Cycl Agroecosyst 89:15–30. https://doi.org/10.1007/s10705-010-9372-6
34. Palm CA, Gachengo CN, Delve RJ et al (2001) Organic inputs for soil fertility management in tropical agroecosystems: application of an organic resource database. Agr Ecosyst Environ 83:27–42. https://doi.org/10.1016/S0167-8809(00)00267-X
35. Bruni E, Guenet B, Huang Y et al (2021) Additional carbon inputs to reach a 4 per 1000 objective in Europe: feasibility and projected impacts of climate change based on Century simulations of long-term arable experiments. Biogeosciences 18:3981–4004. https://doi.org/10.5194/bg-18-3981-2021
36. Ouhrouch A, Boitard S, Boyer F, Servin B, Da Silva A, Pompanon F, Haddioui A, Benjelloun B (2021) Genomic uniqueness of local sheep breeds from Morocco. Front Gen 12. https://doi.org/10.3389/fgene.2021.723599
37. Sadik MW, Al Ashhab AO, Zahran MK, Alsaqan FM (2012) Composting mulch of date palm trees through microbial activator in Saudi Arabia. Int J Biochem Biotechnol 1:156–161
38. Fang M, Wong JWC, Ma KK, Wong MH (1999) Co-composting of sewage sludge and coal fly ash: nutrient transformations. Bioresour Technol 67(1):19–24. https://doi.org/10.1016/S0960-8524(99)00095-4
39. Allison LE (1965) Organic Carbon. In: Norman AG (ed) Agronomy Monographs. American Society of Agronomy, Soil Science Society of America, Madison, WI, USA, pp 1367–1378
40. Giroux M, Audesse P (2004) Comparaison de deux méthodes de détermination des teneurs en carbone organique, en azote total et du rapport C/N de divers amendements organiques et engrais de ferme. Agrosol 15(2):107–110. https://irda.blob.core.windows.net/media/2174/giroux-audesse-2004_article_methodes_dosage_c_n_amend_org.pdf
41. Bremner JM (1996) Methods of soil analysis: part 3 chemical methods. Soil Science Society of America, American Society of Agronomy, Madison, WI, USA
42. El Janati M, Robin P, Akkal-Corfini N, Bouaziz A, Sabri A, Chikhaoui M, Zahra T, Oukarroum A (2022) Composting of date palm residues to serve the circular agriculture in oases. Dataset. https://doi.org/10.15454/DFIAFE
43. Lu D, Wang L, Yan B et al (2014) Speciation of Cu and Zn during composting of pig manure amended with rock phosphate. Waste Manage 34:1529–1536. https://doi.org/10.1016/j.wasman.2014.04.008
44. Paillat J-M, Robin P, Hassouna M, Leterme P (2005) Predicting ammonia and carbon dioxide emissions from carbon and nitrogen biodegradability during animal waste composting. Atmos Environ 39:6833–6842. https://doi.org/10.1016/j.atmosenv.2005.07.045
45. Chakroune K, Bouakka M, Hakkou A (2005) Incidence de l'aération sur le traitement par compostage des sous-produits du palmier dattier contaminés par Fusarium oxysporum f.sp. albedinis. Can J Microbiol 51:69–77. https://doi.org/10.1139/w04-109
46. López-González JA, López MJ, Vargas-García MC et al (2013) Tracking organic matter and microbiota dynamics during the stages of lignocellulosic waste composting. Bioresour Technol 146:574–584
47. Adhikari BK, Barrington S, Martinez J, King S (2009) Effectiveness of three bulking agents for food waste composting. Waste Manage 29:197–203. https://doi.org/10.1016/j.wasman.2008.04.001
48. Ouaqoudi FZE, El-Fels L, Lemée L et al (2015) Evaluation of lignocelullose compost stability and maturity using spectroscopic (FTIR) and thermal (TGA/TDA) analysis. Ecol Eng 75:217–222
49. Chang Y, Hudson HJ (1967) The fungi of wheat straw compost: I. Ecological studies. Trans Br Mycol Soc 50:649–666. https://doi.org/10.1016/S0007-1536(67)80097-4
50. Bernal MP, Alburquerque JA, Moral R (2009) Composting of animal manures and chemical criteria for compost maturity assessment. A Rev Bioresour Technol 100:5444–5453. https://doi.org/10.1016/j.biortech.2008.11.027
51. García-Gómez A, Roig A, Bernal MP (2003) Composting of the solid fraction of olive mill wastewater with olive leaves: organic matter degradation and biological activity. Bioresour Technol 86:59–64. https://doi.org/10.1016/S0960-8524(02)00106-2
52. Paredes C, Bernal MP, Cegarra J, Roig A (2002) Bio-degradation of olive mill wastewater sludge by its co-composting with agricultural wastes. Bioresour Technol 85:1–8. https://doi.org/10.1016/s0960-8524(02)00078-0
53. Bernal MP, Navarro AF, Roig A et al (1996) Carbon and nitrogen transformation during composting of sweet sorghum bagasse. Biol Fert Soils 22:141–148. https://doi.org/10.1007/BF00384446
54. Sharif M, Matiullah M et al (2011) Response of fed dung composted with rock phosphate on yield and phosphorus and nitrogen uptake of maize crop. AJB 10:12595–12601. https://doi.org/10.5897/AJB11.1240
55. Gassmann G, Glindemann D (1993) Phosphane (PH3) in the Biosphere. Angew Chem, Int Ed Engl 32:761–763. https://doi.org/10.1002/anie.199307611
56. White AK, Metcalf WW (2007) Microbial metabolism of reduced phosphorus compounds. Annu Rev Microbiol 61:379–400. https://doi.org/10.1146/annurev.micro.61.080706.093357
57. FAO (2002) Date palm cultivation. Chapter XI: date palm technical calendar. In FAO Plant Production and Protection Paper, num. 156, Rev. 1. http://www.fao.org/3/Y4360E/y4360e0f.htm; https://www.fao.org/3/Y4360E/y4360e00.htm#Contents
58. El Kinany S, Achbani E, Faggroud M et al (2019) Effect of organic fertilizer and commercial arbuscular mycorrhizal fungi on the growth of micropropagated date palm cv. Feggouss. J Saudi Soc Agric Sci 18:411–417. https://doi.org/10.1016/j.jssas.2018.01.004
59. Tejada M, García-Martínez AM, Parrado J (2009) Effects of a vermicompost composted with beet vinasse on soil properties, soil losses and soil restoration. CATENA 77:238–247. https://doi.org/10.1016/j.catena.2009.01.004
60. Canellas LP, Olivares FL (2014) Physiological responses to humic substances as plant growth promoter. Chem Biol Technol 1:3. https://doi.org/10.1186/2196-5641-1-3
61. Monda H, Cozzolino V, Vinci G et al (2017) Molecular characteristics of water-extractable organic matter from different

composted biomasses and their effects on seed germination and early growth of maize. Sci Total Environ 590–591:40–49. https://doi.org/10.1016/j.scitotenv.2017.03.026
62. Klein P, Zaid A (2002) Chapter VI: Land preparation, planting operation and fertilisation requirements. http://www.fao.org/3/y4360e/y4360e0a.htm#bm10.1. Accessed 7 Apr 2021
63. Almadini AM, Ismail AIH, Ameen FA (2021) Assessment of farmers practices to date palm soil fertilization and its impact on productivity at Al-Hassa oasis of KSA. Saudi J Biol Sci 28:1451–1458. https://doi.org/10.1016/j.sjbs.2020.11.084
64. Benabderrahim MA, Elfalleh W, Belayadi H, Haddad M (2018) Effect of date palm waste compost on forage alfalfa growth, yield, seed yield and minerals uptake. Int J Recycl Org Waste Agricult 7:1–9. https://doi.org/10.1007/s40093-017-0182-6
65. Yaser AZ, Lamaming J, Suali E et al (2022) Composting and anaerobic digestion of food waste and sewage sludge for campus sustainability: a review. Int J Chem Eng 2022:e6455889. https://doi.org/10.1155/2022/6455889
66. Al Mussa Ugak M, Zahrim Yaser A, Lamaming J et al (2022) Comparative study on passive aerated in-vessel composting of food wastes with the addition of Sabah ragi. Carbon Resources Conversion 5:200–210. https://doi.org/10.1016/j.crcon.2022.05.004
67. Ismail Ghazy MI (2017) Date palm leaves disintegration using drum chipper machine. Misr J Agric Eng 34:53–64. https://doi.org/10.21608/mjae.2017.96879
68. Callois JM (2022) Le renouveau des territoires par la bioéconomie - Les ressources du vivant au cœur d'une nouvelle économie, Book, 1st ed. Editions Quae (224 pages). https://www.quae.com/produit/1727/9782759234929/le-renouveau-des-territoires-par-la-bioeconomie; https://www.quae-open.com/produit/194/9782759234929/le-renouveau-des-territoires-par-la-bioeconomie
69. Mudrunka J, Lyckova B, Kucerova R et al (2019) Composting in Bags Inżynieria Mineralna 21:335–340
70. Altieri R, Esposito A, Nair T (2011) Novel static composting method for bioremediation of olive mill waste. Int Biodeterior Biodegradation 65:786–789. https://doi.org/10.1016/j.ibiod.2011.05.002
71. Diacono M, Montemurro F (2010) Long-term effects of organic amendments on soil fertility. A review Agron Sustain Dev 30:401–422. https://doi.org/10.1051/agro/2009040
72. Faiad A, Alsmari M, Ahmed MMZ et al (2022) Date palm tree waste recycling: treatment and processing for potential engineering applications. Sustainability 14:1134. https://doi.org/10.3390/su14031134
73. Machineni L (2020) Lignocellulosic biofuel production: review of alternatives. Biomass Conv Bioref 10:779–791. https://doi.org/10.1007/s13399-019-00445-x

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.