



HAL
open science

How to pit weeds against parasitic plants. A simulation study with *Phelipanche ramosa*

Olivia Pointurier, Delphine Moreau, Stéphane Cordeau, Stéphanie Gibot-Leclerc, Carole Reibel, Eric Vieren, Nathalie Colbach

► To cite this version:

Olivia Pointurier, Delphine Moreau, Stéphane Cordeau, Stéphanie Gibot-Leclerc, Carole Reibel, et al.. How to pit weeds against parasitic plants. A simulation study with *Phelipanche ramosa*. 19th European Weed Research Society Symposium EWRS, Jun 2022, Athènes, Greece. hal-04060540

HAL Id: hal-04060540

<https://hal.inrae.fr/hal-04060540v1>

Submitted on 30 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

How to pit weeds against parasitic plants.

A simulation study with *Phelipanche ramosa*

O Pointurier, D Moreau, S Cordeau, S Gibot-Leclerc, C Reibel, E Vieren, N Colbach

Agroécologie, INRAE, Institut Agro, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, Dijon, France

Nathalie.Colbach@inrae.fr




Gibot-Leclerc et al. 2012

Branched broomrape (*Phelipanche ramosa* L.) is a parasitic plant that infects crop and weed species in more than 20 families (*Solanaceae*, *Brassicaceae*, *Asteraceae*...). It is a major pest of oilseed rape in France causing up to 90% of yield losses. No curative method is available, the parasite can only be controlled by combining cropping techniques. Simulation models are essential to design such complex management strategies.


Aim Develop a simulation model & use it to design strategies to manage broomrape

Synthesize knowledge on broomrape dynamics & effects in a 3D individual-based model


Experiments to fit functions & estimate parameters




Simulate cropping systems over 30 years x 10 weather repetitions, with and without weeds



Efficient techniques ↓ broomrape up to 79%



Weed ideotype to ↓ broomrape seed bank



There are efficient strategies to control broomrape → need to design strategies adapted to local conditions

Conclusion are only valid if broomrape-attaching weeds reproduce before broomrape does → need to confirm