



HAL
open science

First report of Pepo aphid-borne yellows virus on courgette in Italy

Giuseppe Parrella, E. Troiano, Cecile Desbiez

► **To cite this version:**

Giuseppe Parrella, E. Troiano, Cecile Desbiez. First report of Pepo aphid-borne yellows virus on courgette in Italy. *New Disease Reports*, 2023, 47 (1), pp.e12144. 10.1002/ndr2.12144 . hal-04064123

HAL Id: hal-04064123

<https://hal.inrae.fr/hal-04064123>

Submitted on 11 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

NEW DISEASE REPORT

First report of *Pepo aphid-borne yellows virus* on courgette in Italy

G. Parrella¹  | E. Troiano¹ | C. Desbiez² 

¹Institute for Sustainable Plant Protection of CNR (IPSP-CNR), Portici, (NA), Italy

²INRAE, Pathologie Végétale, Montfavet, France

Correspondence

G. Parrella, Institute for Sustainable Plant Protection of CNR (IPSP-CNR), Portici, Napoli, 80055, Italy.

Email: giuseppe.parrella@ipspp.cnr.it

Funding information

Consiglio Nazionale delle Ricerche, Grant/Award Number: Short Term Mobility 2020

KEYWORDS

PABYV, polerovirus, zucchini

During monitoring for cucurbit viruses in greenhouses in the municipalities of Torre del Greco (Napoli province, Campania) and Aversana (Salerno province, Campania) in September 2019 and October 2022, respectively, samples of commercial hybrids of courgette (*Cucurbita pepo*) showing symptoms of leaf yellowing and mosaics (Figure 1) were collected and stored as dried material. Three samples from 2019 and two samples from 2022, were inoculated mechanically onto courgette. Within two weeks, typical symptoms of *Tomato leaf curl New Delhi virus* (ToLCNDV), consisting of leaf crumpling, yellowing and downward curling, were observed. ToLCNDV infection was confirmed by DAS-ELISA (Bioreba, Switzerland). This is not a new record for Campania since ToLCNDV was detected in courgette in 2016 and causes a serious disease problem for cucurbit crops in this region.

The samples were also checked for polerovirus infection by RT-PCR as follows. Total RNA was extracted from infected tissues using TRI-reagent (Sigma-Aldrich, USA) and RT-PCR was performed with polerovirus primers Pol-G-F and Pol-G-R (Knierim et al., 2014). A 1.1 Kb fragment was obtained from two samples, one sampled in 2019 and the other in 2022. These amplicons were sent to Genoscreen (Lille, France) for direct sequencing. The sequences obtained (GenBank Accession Nos. OP156930 and OP973153, respectively), encompassing part of the polerovirus RNA-dependent RNA polymerase, intergenic region, coat protein and movement protein, shared a close relationship with *Pepo aphid-borne yellows virus* (PABYV) iso-



FIGURE 1 Courgette plant showing yellowing symptoms found to be infected with *Pepo aphid-borne yellows virus* and *Tomato leaf curl New Delhi virus*

lates from different origins. PABYV was first described in 2008 in Mali (Knierim et al., 2014), then in South Africa (Ibaba et al., 2015), Côte d'Ivoire (Kone et al., 2015), in Tanzania and Mayotte (Desbiez et al., 2016), and a polerovirus closely related to PABYV, named “pumpkin polerovirus” was described more recently in Kenya

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *New Disease Reports* published by British Society for Plant Pathology and John Wiley & Sons.




FIGURE 2 Neighbour-joining tree based on the 1.1 Kb fragment of cucurbit-infesting poleroviruses genome (*Cucumber aphid-borne yellows* - Hainan virus, CuABYV-HN; *Cucurbit aphid-borne yellows virus*, CABYV; *Luffa aphid-borne yellows virus*, LABYV; *Melon aphid-borne yellows virus*, MABYV; *Pepo aphid-borne yellows virus*, PABYV; and *Suakwa aphid-borne yellows virus*, SABYV). Bootstrap values ($n = 1000$) above 75% are indicated for each node. The position of the two Italian PABYV isolates within the phylogenetic tree is highlighted with a red box

(Kidanamariam et al., 2019). In the Mediterranean area, PABYV has been detected in Syria in 2009 and in Greece in 2016 (Lotos et al., 2016). The two sequences of the Italian PABYV isolates (IT-236-19 and IT-22, respectively) were 99.2% identical and shared 98.4% identity with the isolate AE142 (LN865082), reported in Greece in 2016 (Lotos et al., 2016), and 98.3% with isolate W-8-1-Sy-09 (KC960436) reported in Syria in 2009. These four isolates formed a distinct group

in a phylogenetic reconstruction (Figure 2), suggesting a common origin for the PABYV isolates spreading in the Mediterranean area.

PABYV belongs to the polerovirus genus which are phloem-restricted RNA plant viruses transmitted by aphids. Nevertheless, *Bemisia tabaci* has been reported recently to transmit a new species of polerovirus, *Pepper whitefly borne vein yellows virus*, in pepper in Israel, as well as a recombinant strain of *Cucurbit aphid-borne yellows*

virus in cucurbits in Brazil. In our case, no aphids were observed during sampling (although previous infestation could not be excluded) but abundant colonies of *B. tabaci* were recorded. The possibility of PABYV transmission by *B. tabaci* requires investigation.

This is the first report of PABYV in Italy. The finding of PABYV in a new geographic area represents further evidence of the active and current spread of the virus, suggesting the need for more extensive sampling of cucurbit crops and the identification of vectors and alternate hosts to better understand the epidemiology of cucurbit-infecting poleroviruses in the Mediterranean and in Africa.

ACKNOWLEDGMENTS

This work was carried out as part of the Short Term Mobility (STM) 2020 programme financed by the National Research Council (CNR) of Italy.

ORCID

G. Parrella  <https://orcid.org/0000-0002-0412-4014>

C. Desbiez  <https://orcid.org/0000-0001-8691-7423>

REFERENCES

Desbiez, C., Millot, P., Wipf-Scheibel, C., Blancard, D., Chesneau, T. & Lecoq, H. (2016) First report of *Pepo aphid-borne yellows virus* in cucurbits in

Tanzania and Mayotte. *New Disease Reports*, 33, 20. [10.5197/j.2044-0588.2016.033.020](https://doi.org/10.5197/j.2044-0588.2016.033.020)

Ibaba, J.D., Laing, M.D. & Gubba, A. (2015) Incidence and phylogeny of viruses infecting cucurbit crops in KwaZulu-Natal, Republic of South Africa. *Crop Protection*, 75, 46–54. [10.1016/j.cropro.2015.04.019](https://doi.org/10.1016/j.cropro.2015.04.019)

Kidanemariam, D.B., Sukal, A.C., Abraham, A.D., Njuguna, J.N., Stomeo, F., Dale, J.L. et al. (2019) Molecular characterisation of a putative new polerovirus infecting pumpkin (*Cucurbita pepo*) in Kenya. *Archives of Virology*, 164, 1717–1721. [10.1007/s00705-019-04219-8](https://doi.org/10.1007/s00705-019-04219-8)

Knierim, D., Tsai, W.S., Maiss, E. and Kenyon, L. (2014) Molecular diversity of poleroviruses infecting cucurbit crops in four countries reveals the presence of members of six distinct species. *Archives of Virology*, 159, 1459–1465. [10.1007/s00705-013-1939-5](https://doi.org/10.1007/s00705-013-1939-5)

Kone, N., Coulibaly, A., Koita, O., Kone, D., Bediako, E.A., Knierim, D., et al. (2015) First report of *Pepo aphid-borne yellows virus* on zucchini in Cote d'Ivoire. *New Disease Reports*, 31, 27. [10.5197/j.2044-0588.2015.031.027](https://doi.org/10.5197/j.2044-0588.2015.031.027)

Lotos, L., Maliogka, V.I. and Katis, N.I. (2016) New poleroviruses associated with yellowing symptoms in different vegetable crops in Greece. *Archives of Virology*, 161, 431–436. [10.1007/s00705-015-2662-1](https://doi.org/10.1007/s00705-015-2662-1)

How to cite this article: Parrella, G., Troiano, E. & Desbiez, C. (2023) First report of *Pepo aphid-borne yellows virus* on courgette in Italy. *New Disease Reports*, 47, e12144. <https://doi.org/10.1002/ndr2.12144>

© 2023. This work is published under <http://creativecommons.org/licenses/by/4.0/>(the “License”). Notwithstanding the ProQuest Terms and Conditions, you may use this content in accordance with the terms of the License.