



HAL
open science

Control theory perspectives on mathematical epidemiology

Pierre-Alexandre Bliman, Alain Rapaport

► **To cite this version:**

Pierre-Alexandre Bliman, Alain Rapaport. Control theory perspectives on mathematical epidemiology. 2023, 22nd World Congress of the International Federation of Automatic Control. hal-04064515

HAL Id: hal-04064515

<https://hal.inrae.fr/hal-04064515v1>

Submitted on 11 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Invited session proposal submitted to IFAC World Congress 2023

Pierre-Alexandre Bliman* Alain Rapaport†

October 14, 2022

Title

Control theory perspectives on mathematical epidemiology

Abstract

Mathematical epidemiology may benefit from techniques and concepts from Control theory. As a matter of fact, understanding and controlling the spread of an infectious epidemic necessitates modeling of the phenomena underlying the propagation, estimation of the characteristic parameters and of the state of the epidemic, scheduling and scaling of the control measures. These are generic questions, for which tools have been developed in Control, admittedly in quite different applicative contexts. The aim of this session is precisely to explore in more details potential profit, as well as limitations, of this cross-fertilization.

Several common features of the problems arising from the application of Control theory to the handling of epidemics are worth mentioning here. First, the transmission of infection is inherently a nonlinear phenomenon. Second, the usual measurements of prevalence leads to a lack of observability and identifiability of the studied state space models at equilibria, a point which complicates the handling of these issues when the state is closed from these singular points (beginning of outbreak or final stabilization). Last, the cost functions of interest are usually non-conventional, typically proportional to the effort made (and not quadratic), and state constraint are not uncommon. All this, plus the important uncertainty present on the models and on the data, makes the corresponding problems rather sensitive. The six contributions that constitute this session offer a partial vision of issues to be addressed. They are more especially related to observability and identifiability (papers 1 and 2), and to optimal control applied to epidemic processes (papers 2 to 6).

Choice of an IFAC technical committee for review

TC 8.2. Biological and Medical Systems

Detailed description of the topic

Paper 1. is concerned with identifiability and observability issues for a compartmental SIR model with quarantine, where the flows of quarantined people are measured. Paper 2. proposes a simultaneous parameter and state estimation scheme for a class of general epidemic models, and solves associated optimal control problems. Paper 3. provides, for a class of positive compartmental systems, general results that characterize the solutions of some pertinent optimal control problems. Paper 4. studies optimal isolation strategies for epidemic models where, differently from the classical exponential distribution, the infectious period obeys an Erlang probability distribution. Paper 5. intends to study in a stochastic framework notions of herd immunity and of viable zones for non-pharmaceutical interventions in presence of constraints on the intensive care occupancy rate. Last, paper 6. presents several qualitative results on the optimal control that minimizes the epidemic final size of SIR model by social distancing.

*pierre-alexandre.bliman@inria.fr

†alain.rapaport@inrae.fr

Titles, authors, and abstracts of all six contributions

1. About the identifiability and observability of the SIR model with quarantine, by Frédéric Hamelin¹, Abderrahman Iggidr², Alain Rapaport³, Gauthier Sallet², Max O. Souza⁴.

We consider a SIRQ model which is an extension of the well-known SIR epidemic model with an additional compartment Q that represents the population that has been removed from the infected population and placed in quarantine. We analyze the identifiability and observability of this model assuming that the flow of individuals placed in quarantine and the size of the quarantine population are known at any time. We also derive some methods to deal with the practical identification of the model parameters and the estimation of the infected population size.

2. Feedback Design for Optimal Control of Epidemic Processes, by Muhammad Umar B. Niaz⁵, Phillip E. Paré⁶, Karl Henrik Johansson⁵.

For reliable epidemic monitoring and control, this paper proposes a feedback mechanism design to effectively cope with data and model uncertainties through. Using the past epidemiological data, we propose a simultaneous parameter and state estimation method of general epidemic models. We provide necessary and sufficient conditions under which the parameter and state estimation problems are well-posed and feasible. Then, we devise optimal control policies by minimizing a predicted cost functional based on the estimated parameters and state of the model, and discuss under what condition the solution turns out to be feasible.

3. Optimal control of compartmental models: the exact solution, by Franco Blanchini⁶, Paolo Bolzern⁷, Patrizio Colaneri⁸, Giuseppe De Nicolao⁹, Giulia Giordano¹⁰.

We formulate a control problem for positive compartmental systems formed by nodes (buffers) and arcs (flows). Our main result is that, on a finite horizon, we can solve the Pontryagin equations in one shot without resorting to trial and error via shooting. As expected, the solution is bang-bang and the switching times can be easily determined. We are also able to find a cost-to-go-function, in an analytic form, by solving a simple nonlinear differential equation. On an infinite horizon, we consider the Hamilton-Jacobi-Bellman theory and we show that the HJB equation can be solved exactly. Moreover, we show that the optimal solution is constant and the cost-to-go function is linear and copositive. This function is the solution of a nonlinear equation. We propose an iterative scheme for solving this equation, which converges in finite time. We also show that an exact solution can be found if there is a positive external disturbance affecting the process and the problem is formulated in a min sup framework. We finally provide illustrative examples related to flood control and epidemiology.

4. Optimal isolation strategies in an SIR model with Erlang-distributed infectious period, by Luca Bolzoni¹¹, Rossella Della Marca¹².

Mathematical models are formal and simplified representations of the knowledge related to a phenomenon. In classical epidemic models, a major simplification consists in assuming that the infectious period is exponentially distributed, then implying that the chance of recovery is independent of the time since infection. Here, we first attempt to investigate the consequences of relaxing this assumption on the performances of time-variant disease control strategies by using optimal control theory. In the

¹Institut Agro, Rennes, France

²Université de Lorraine, CNRS, Inria, IECL, Metz, France

³MISTEA, Univ. Montpellier, INRAE, Institut Agro, Montpellier, France

⁴Instituto de Matemática e Estatística, Universidade Federal Fluminense, Niterói, Brasil

⁵Division of Decision and Control Systems and Digital Futures, KTH Royal Institute of Technology, Sweden

⁶School of Electrical and Computer Engineering, Purdue University, USA

⁷Dipartimento di Scienze Matematiche, Informatiche e Fisiche, Università degli Studi di Udine, Italy

⁸Dipartimento di Elettronica, Informazione e Bioingegneria, Politecnico di Milano, Italy

⁹Dipartimento di Elettronica, Informazione e Bioingegneria, Politecnico di Milano, and IEIIT, CNR, Milano, Italy

¹⁰Department of Computer and System Science, Università degli Studi di Pavia, Italy

¹¹Department of Industrial Engineering, University of Trento, Italy

¹²Risk Analysis and Genomic Epidemiology Unit, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna, Parma, Italy

¹²Mathematics Area, SISSA – International School for Advanced Studies, Trieste, Italy

framework of a basic susceptible–infected–removed (SIR) model, an Erlang distribution of the infectious period is considered and optimal isolation strategies are searched for. The objective functional to be minimized takes into account the cost of the isolation efforts per time unit and the sanitary costs due to the incidence of the epidemic outbreak. Applying the Pontryagin’s minimum principle, we prove that the optimal control problem admits only bang–bang solutions with at most two switches. In particular, the optimal strategy could be postponing the starting intervention time with respect to the beginning of the outbreak. Finally, by means of numerical simulations, we show how the shape of the optimal solutions is affected by the different distributions of the infectious period, by the relative weight of the two cost components, and by the initial conditions.

5. On the Design Techniques for Safety Zones In Brownian-Driven Epidemic Models, by Dan Goreac¹³, Juan Li¹⁴, Yi Wang¹⁵.

This paper is concerned with the mathematical description of *herd immunity* and *feasible zones* for *non-pharmaceutically controlled* Brownian-driven compartmental models when *intensive care units* restrictions are imposed. First, we focus on *consistency* considerations leading to non-negative and sub-unitary components for possibly non-constant population models. Second, based on viscosity approaches, we give a family of conditions allowing to envisage regular-barrier safety zones. Relevant examples of families of noise coefficients are presented.

6. On the problem of minimizing the epidemic final size for SIR model by social distancing, by Alain Rapaport³, Pierre-Alexandre Bliman¹⁷.

We revisit the problem of minimizing the epidemic final size in the SIR model through social distancing. In the existing literature, this problem has been considered imposing a priori interval structure on the time period when interventions are enforced. We show that when considering the more general class of controls with an L^1 constraint on the confinement effort that reduces the infection rate, the support of the optimal control is still a single time interval. This shows that, for this problem, there is no benefit in splitting interventions on several disjoint time periods.

¹³School of Mathematics and Statistics, Shandong University, Weihai, PR China and LAMA, Université Gustave Eiffel, UPEM, Université Paris Est Créteil, CNRS, Marne-la-Vallée, France

¹⁴School of Mathematics and Statistics, Shandong University, Weihai, PR China and Research Center for Mathematics and Interdisciplinary Sciences, Shandong University, Qingdao, PR China

¹⁵School of Mathematics and Statistics, Shandong University, Weihai, PR China

¹⁷Sorbonne Université, Université Paris-Cité, Inria, CNRS, Laboratoire Jacques-Louis Lions, Paris France