



HAL
open science

Mechanical properties of miscanthus rind: relation to tissue structure and composition

Cecile Barron, Paul-Louis Lopez-Marnet, Kasper van der Cruijzen, Cyril Grandjean, Abdelkrim Sadoudi, Herman Höfte, Luisa Trindade

► **To cite this version:**

Cecile Barron, Paul-Louis Lopez-Marnet, Kasper van der Cruijzen, Cyril Grandjean, Abdelkrim Sadoudi, et al.. Mechanical properties of miscanthus rind: relation to tissue structure and composition. 10th International Plant Biomechanics Conference, Aug 2022, Lyon, France. . hal-04066475

HAL Id: hal-04066475

<https://hal.inrae.fr/hal-04066475>

Submitted on 12 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

10th International Plant Biomechanics Conference

<https://plantbiomech.sciencesconf.org/>

Mechanical properties of miscanthus rind : relation to tissue structure and composition.

Barron C.^a, Lopez P.L.^b, van der Cruijssen K.^c, Grandjean C.^b, Sadoudi A.^a, Höfte H.^b and Trindade L. M.^c

^a IATE, Univ Montpellier, INRAE, Institut Agro, Montpellier, France

^b Institut Jean-Pierre Bourgin, INRAE , AgroParisTech, Université Paris-Saclay, Versailles, France

^c Wageningen University & Research, Plant Breeding, Wageningen, The Netherlands

Understanding the mechanical properties of lignocellulosic biomass is of great interest to improve their deconstruction (eg in the pre-grinding operation unit), and their functionality for different applications (eg biomaterials). Ten highly contrasted *Miscanthus sinensis* genotypes were selected based on their overall cell wall composition to cover the variability available within this species, and were harvested in two consecutive years. The rinds from the fifth internode were prepared and the shear force required to cut them was obtained using a dedicated prototype device. Based on the measured mechanical properties a subsample of rinds or associated internode were selected and characterized for histology (by light microscopy after FASGA staining or scanning electron microscopy) and cell wall biochemistry (neutral sugars, Klason lignin, hydroxycinnamid acids). A large variability was revealed for composition and structure (rind thickness and density) allowing to explain differences in the observed mechanical properties.